



# On a Dirichlet problem with $(p, q)$ -Laplacian and parametric concave-convex nonlinearity



Salvatore A. Marano<sup>a</sup>, Greta Marino<sup>a,\*</sup>, Nikolaos S. Papageorgiou<sup>b</sup>

<sup>a</sup> *Dipartimento di Matematica e Informatica, Università degli Studi di Catania, Viale A. Doria 6, 95125 Catania, Italy*

<sup>b</sup> *Department of Mathematics, National Technical University, Zografou Campus, 15780 Athens, Greece*

## ARTICLE INFO

### Article history:

Received 19 December 2018  
Available online 7 March 2019  
Submitted by V. Radulescu

### Keywords:

$(p, q)$ -Laplacian  
Concave-convex nonlinearity  
Positive solution  
Bifurcation-type theorem

## ABSTRACT

A homogeneous Dirichlet problem with  $(p, q)$ -Laplace differential operator and reaction given by a parametric  $p$ -convex term plus a  $q$ -concave one is investigated. A bifurcation-type result, describing changes in the set of positive solutions as the parameter  $\lambda > 0$  varies, is proven. Since for every admissible  $\lambda$  the problem has a smallest positive solution  $\bar{u}_\lambda$ , both monotonicity and continuity of the map  $\lambda \mapsto \bar{u}_\lambda$  are studied.

© 2019 Elsevier Inc. All rights reserved.

## 1. Introduction

Let  $\Omega$  be a bounded domain in  $\mathbb{R}^N$  with a  $C^2$ -boundary  $\partial\Omega$ , let  $1 < \tau < q < p < +\infty$ , and let  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a Carathéodory function. Consider the Dirichlet problem

$$\begin{cases} -\Delta_p u - \Delta_q u = u^{\tau-1} + \lambda f(x, u) & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (P_\lambda)$$

where  $\lambda > 0$  is a parameter while  $\Delta_r$ ,  $r > 1$ , denotes the  $r$ -Laplacian, namely

$$\Delta_r u := \operatorname{div}(|\nabla u|^{r-2} \nabla u) \quad \forall u \in W_0^{1,r}(\Omega).$$

The nonhomogeneous differential operator  $Au := \Delta_p u + \Delta_q u$  that drives  $(P_\lambda)$  is usually called  $(p, q)$ -Laplacian. It stems from a wide range of important applications, including models of elementary particles [8],

\* Corresponding author.

E-mail addresses: [marano@dmf.unict.it](mailto:marano@dmf.unict.it) (S.A. Marano), [greta.marino@dmf.unict.it](mailto:greta.marino@dmf.unict.it) (G. Marino), [npapg@math.ntua.gr](mailto:npapg@math.ntua.gr) (N.S. Papageorgiou).

biophysics [9], plasma physics [26], reaction-diffusion equations [7], elasticity theory [27], etc. That's why the relevant literature looks daily increasing and numerous meaningful works on this subject are by now available; see the survey paper [19] for a larger bibliography.

Since  $\tau < q < p$ , the function  $\xi \mapsto \xi^{\tau-1}$  grows  $(q-1)$ -sublinearly at  $+\infty$ , whereas  $\xi \mapsto f(x, \xi)$  is assumed to be  $(p-1)$ -superlinear near  $+\infty$ , although it need not satisfy the usual (in such cases) Ambrosetti-Rabinowitz condition. So, the reaction in  $(P_\lambda)$  exhibits the competing effects of concave and convex terms, with the latter multiplied by a positive parameter.

The aim of this paper is to investigate how the solution set of  $(P_\lambda)$  changes as  $\lambda$  varies. In particular, we prove that there exists a critical parameter value  $\lambda^* > 0$  for which problem  $(P_\lambda)$  admits

- at least two solutions if  $\lambda \in (0, \lambda^*)$ ,
- at least one solution when  $\lambda = \lambda^*$ , and
- no solution provided  $\lambda > \lambda^*$ .

Moreover, we detect a smallest positive solution  $\bar{u}_\lambda$  for each  $\lambda \in (0, \lambda^*)$  and show that the map  $\lambda \mapsto \bar{u}_\lambda$  turns out left-continuous, besides increasing.

The first bifurcation result for semilinear Dirichlet problems driven by the Laplace operator was established, more than twenty years ago, in the seminal paper [2] and then extended to the  $p$ -Laplacian in [11,16]. These works treat the reaction

$$\xi \mapsto \lambda \xi^{s-1} + \xi^{r-1}, \quad \xi \geq 0,$$

where  $1 < s < p < r < p^*$ ,  $\lambda > 0$ , and  $p^*$  denotes the critical Sobolev exponent. A wider class of nonlinearities has recently been investigated in [22], while [24] deals with Robin boundary conditions. It should be noted that, unlike our case,  $\lambda$  always multiplies the concave term, which changes the analysis of the problem. Finally, [4,14,23] contain analogous bifurcation theorems for problems of a different kind, whereas [20,21] study  $(p, q)$ -Laplace equations having merely concave right-hand side.

Our approach is based on the critical point theory, combined with appropriate truncation and comparison techniques.

## 2. Mathematical background and hypotheses

Let  $(X, \|\cdot\|)$  be a real Banach space. Given a set  $V \subseteq X$ , write  $\overline{V}$  for the closure of  $V$ ,  $\partial V$  for the boundary of  $V$ , and  $\text{int}_X(V)$  or simply  $\text{int}(V)$ , when no confusion can arise, for the interior of  $V$ . If  $x \in X$  and  $\delta > 0$  then

$$B_\delta(x) := \{z \in X : \|z - x\| < \delta\}, \quad B_\delta := B_\delta(0).$$

The symbol  $(X^*, \|\cdot\|_{X^*})$  denotes the dual space of  $X$ ,  $\langle \cdot, \cdot \rangle$  indicates the duality pairing between  $X$  and  $X^*$ , while  $x_n \rightarrow x$  (respectively,  $x_n \rightharpoonup x$ ) in  $X$  means ‘the sequence  $\{x_n\}$  converges strongly (respectively, weakly) in  $X$ ’. We say that  $A : X \rightarrow X^*$  is of type  $(S)_+$  provided

$$x_n \rightharpoonup x \text{ in } X, \quad \limsup_{n \rightarrow +\infty} \langle A(x_n), x_n - x \rangle \leq 0 \quad \implies \quad x_n \rightarrow x.$$

The function  $\Phi : X \rightarrow \mathbb{R}$  is called coercive if  $\lim_{\|x\| \rightarrow +\infty} \Phi(x) = +\infty$  and weakly sequentially lower semicontinuous when

$$x_n \rightharpoonup x \text{ in } X \quad \implies \quad \Phi(x) \leq \liminf_{n \rightarrow \infty} \Phi(x_n).$$

Suppose  $\Phi \in C^1(X)$ . We denote by  $K(\Phi)$  the critical set of  $\Phi$ , i.e.,

$$K(\Phi) := \{x \in X : \Phi'(x) = 0\}.$$

The classical Cerami compactness condition for  $\Phi$  reads as follows:

(C) *Every  $\{x_n\} \subseteq X$  such that  $\{\Phi(x_n)\}$  is bounded and  $(1 + \|x_n\|)\Phi'(x_n) \rightarrow 0$  in  $X^*$  has a convergent subsequence.*

From now on,  $\Omega$  indicates a fixed bounded domain in  $\mathbb{R}^N$  with a  $C^2$ -boundary  $\partial\Omega$ . Let  $u, v : \Omega \rightarrow \mathbb{R}$  be measurable and let  $t \in \mathbb{R}$ . The symbol  $u \leq v$  means  $u(x) \leq v(x)$  for almost every  $x \in \Omega$ ,  $t^\pm := \max\{\pm t, 0\}$ ,  $u^\pm(\cdot) := u(\cdot)^\pm$ . If  $u, v$  belong to a function space, say  $Y$ , then we set

$$[u, v] := \{w \in Y : u \leq w \leq v\}, \quad [u] := \{w \in Y : u \leq w\}.$$

The conjugate exponent  $r'$  of a number  $r \geq 1$  is defined by  $r' := r/(r-1)$ , while  $r^*$  indicates its Sobolev conjugate, namely

$$r^* := \begin{cases} \frac{Nr}{N-r} & \text{when } r < N, \\ +\infty & \text{otherwise.} \end{cases}$$

As usual,

$$\|u\|_r := \left( \int_{\Omega} |u|^r dx \right)^{1/r} \quad \forall u \in L^r(\Omega), \quad \|u\|_{1,r} := \left( \int_{\Omega} |\nabla u|^r dx \right)^{1/r} \quad \forall u \in W_0^{1,r}(\Omega),$$

and  $W^{-1,r'}(\Omega)$  denotes the dual space of  $W_0^{1,r}(\Omega)$ . We will also employ the linear space  $C_0^1(\overline{\Omega}) := \{u \in C^1(\overline{\Omega}) : u|_{\partial\Omega} = 0\}$ , which is complete with respect to the standard  $C^1(\overline{\Omega})$ -norm. Its positive cone

$$C_+ := \{u \in C_0^1(\overline{\Omega}) : u(x) \geq 0 \text{ in } \overline{\Omega}\}$$

has a nonempty interior given by

$$\text{int}(C_+) = \left\{ u \in C_+ : u(x) > 0 \quad \forall x \in \Omega, \quad \frac{\partial u}{\partial n}(x) < 0 \quad \forall x \in \partial\Omega \right\}.$$

Here  $n(x)$  denotes the outward unit normal to  $\partial\Omega$  at  $x$ .

Let  $A_r : W_0^{1,r}(\Omega) \rightarrow W^{-1,r'}(\Omega)$  be the nonlinear operator stemming from the negative  $r$ -Laplacian, i.e.,

$$\langle A_r(u), v \rangle := \int_{\Omega} |\nabla u|^{r-2} \nabla u \cdot \nabla v dx, \quad u, v \in W_0^{1,r}(\Omega).$$

We know [12, Section 6.2] that  $A_r$  is bounded, continuous, strictly monotone, and of type  $(S)_+$ . The Liusternik-Schnirelmann theory gives an increasing sequence  $\{\lambda_{n,r}\}$  of eigenvalues for  $A_r$ . The following assertions can be found in [12, Section 6.2].

(p<sub>1</sub>)  $\lambda_{1,r}$  is positive, isolated, and simple.

(p<sub>2</sub>)  $\|u\|_r^r \leq \frac{1}{\lambda_{1,r}} \|u\|_{1,r}^r$  for all  $u \in W_0^{1,r}(\Omega)$ .

(p<sub>3</sub>)  $\lambda_{1,r}$  admits an eigenfunction  $\phi_{1,r} \in \text{int}(C_+)$  such that  $\|\phi_{1,r}\|_r = 1$ .

Proposition 13 of [6] then ensures that

(p<sub>4</sub>) If  $r \neq \hat{r}$  then  $\phi_{1,r}$  and  $\phi_{1,\hat{r}}$  are linearly independent.

Let  $g : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  be a Carathéodory function satisfying the growth condition

$$|g(x, t)| \leq a(x) (1 + |t|^{s-1}) \quad \text{in } \Omega \times \mathbb{R},$$

where  $a \in L^\infty(\mathbb{R})$ ,  $1 < s \leq p^*$ . Set  $G(x, \xi) := \int_0^\xi g(x, t) dt$  and consider the  $C^1$ -functional  $\varphi : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\varphi(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_\Omega G(x, u(x)) dx, \quad u \in W_0^{1,p}(\Omega).$$

**Proposition 2.1** ([13], Proposition 2.6). *If  $u_0 \in W_0^{1,p}(\Omega)$  is a local  $C_0^1(\overline{\Omega})$ -minimizer of  $\varphi$  then  $u_0 \in C^{1,\alpha}(\overline{\Omega})$  for some  $\alpha \in (0, 1)$  and  $u_0$  turns out to be a local  $W_0^{1,p}(\Omega)$ -minimizer of  $\varphi$ .*

Combining this result with the strong comparison principle below, essentially due to Arcoya-Ruiz [3], shows that certain constrained minimizers actually are ‘global’ critical points. Recall that, given  $h_1, h_2 \in L^\infty(\Omega)$ ,

$$h_1 \prec h_2 \iff \text{ess inf}_K (h_2 - h_1) > 0 \quad \text{for any nonempty compact set } K \subseteq \Omega.$$

**Proposition 2.2.** *Let  $a \in \mathbb{R}_+$ ,  $h_1, h_2 \in L^\infty(\Omega)$ ,  $u_1 \in C_0^1(\overline{\Omega})$ ,  $u_2 \in \text{int}(C_+)$ . Suppose  $h_1 \prec h_2$  as well as*

$$-\Delta_p u_i - \Delta_q u_i + a|u_i|^{p-2} u_i = h_i \quad \text{in } \Omega, \quad i = 1, 2.$$

*Then,  $u_2 - u_1 \in \text{int}(C_+)$ .*

Throughout the paper, ‘for every  $x \in \Omega$ ’ will take the place of ‘for almost every  $x \in \Omega$ ’,  $c_0, c_1, \dots$  indicate suitable positive constants,  $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$  is a Carathéodory function such that  $f(\cdot, t) = 0$  provided  $t \leq 0$ , while  $F(x, \xi) := \int_0^\xi f(x, t) dt$ .

The following hypotheses will be posited.

(h<sub>1</sub>) There exist  $\theta \in [\tau, q]$  and  $r \in (p, p^*)$  such that

$$c_1 t^{p-1} + c_2 t^{q-1} \leq f(x, t) \leq c_0 (t^{\theta-1} + t^{r-1}) \quad \forall (x, t) \in \Omega \times \mathbb{R}_+,$$

where  $c_2 > \lambda_{1,q}$ .

(h<sub>2</sub>)  $\lim_{\xi \rightarrow +\infty} \frac{F(x, \xi)}{\xi^p} = +\infty$  uniformly with respect to  $x \in \Omega$ .

(h<sub>3</sub>)  $\liminf_{\xi \rightarrow +\infty} \frac{f(x, \xi) \xi - p F(x, \xi)}{\xi^\beta} \geq c_3$  uniformly in  $x \in \Omega$ . Here,  $\beta > \tau$  and

$$(r - p) \max \{Np^{-1}, 1\} < \beta < p^*.$$

(h<sub>4</sub>) To every  $\rho > 0$  there corresponds  $\mu_\rho > 0$  such that  $t \mapsto f(x, t) + \mu_\rho t^{p-1}$  is nondecreasing in  $[0, \rho]$  for any  $x \in \Omega$ .

By (h<sub>2</sub>)–(h<sub>3</sub>) the perturbation  $f(x, \cdot)$  is  $(p-1)$ -superlinear at  $+\infty$ . In the literature, one usually treats this case via the well-known Ambrosetti-Rabinowitz condition, namely:

(AR) With appropriate  $M > 0$ ,  $\sigma > p$  one has both  $\operatorname{ess\,inf}_{\Omega} F(\cdot, M) > 0$  and

$$0 < \sigma F(x, \xi) \leq f(x, \xi)\xi, \quad (x, \xi) \in \Omega \times [M, +\infty). \quad (2.1)$$

It easily entails  $c_3 \xi^\sigma \leq F(x, \xi)$  in  $\Omega \times [M, +\infty)$ , which forces (h<sub>2</sub>). However, nonlinearities having a growth rate ‘slower’ than  $t^{\sigma-1}$  at  $+\infty$  are excluded from (2.1). Thus, assumption (h<sub>3</sub>) incorporates in our framework more situations.

**Example 2.3.** Let  $c_2 > \lambda_{1,q}$ . The functions  $f_1, f_2 : \mathbb{R}_+ \rightarrow \mathbb{R}$  defined by

$$f_1(t) := \begin{cases} t^{p-1} + c_2 t^{\tau-1} & \text{if } 0 \leq t \leq 1, \\ t^{r-1} + c_2 t^{q-1} & \text{otherwise,} \end{cases} \quad f_2(t) := t^{p-1} \log(1+t) + c_2 t^{q-1}, \quad t \in \mathbb{R}_+,$$

satisfy (h<sub>1</sub>)–(h<sub>4</sub>). Nevertheless,  $f_1$  alone complies with condition (AR).

### 3. A bifurcation-type theorem

Write  $S_\lambda$  for the set of positive solutions to (P <sub>$\lambda$</sub> ). Lieberman’s nonlinear regularity theory [18, p. 320] and Pucci-Serrin’s maximum principle [25, pp. 111,120] yield

$$S_\lambda \subseteq \operatorname{int}(C_+).$$

Put  $\mathcal{L} := \{\lambda > 0 : S_\lambda \neq \emptyset\}$ . Our first goal is to establish some basic properties of  $\mathcal{L}$ . From now on,  $X := W_0^{1,p}(\Omega)$  and  $\|\cdot\| := \|\cdot\|_{1,p}$ .

**Proposition 3.1.** *Under (h<sub>1</sub>) one has  $\mathcal{L} \neq \emptyset$ .*

**Proof.** Given  $\lambda > 0$ , consider the  $C^1$ -functional  $\Psi_\lambda : W_0^{1,p}(\Omega) \rightarrow \mathbb{R}$  defined by

$$\Psi_\lambda(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} dx \int_0^{u(x)} g_\lambda(t) dt \quad \forall u \in W_0^{1,p}(\Omega),$$

where

$$g_\lambda(t) := (t^+)^{\tau-1} + \lambda c_0 [(t^+)^{\theta-1} + (t^+)^{r-1}], \quad t \in \mathbb{R}.$$

Evidently,  $g_\lambda$  fulfills (2.1) once  $\sigma \in (p, r)$  and  $M > 0$  is big enough. So, condition (C) holds true for  $\Psi_\lambda$ . Moreover,

$$u \in \operatorname{int}(C_+) \implies \lim_{t \rightarrow +\infty} \Psi_\lambda(tu) = -\infty$$

because  $r > p$ . Observe next that if  $s \in [1, p^*]$  then

$$\|u\|_s \leq c \|u\|_{p^*} \leq C \|u\| \quad \forall u \in X,$$

with  $C := C(s, \Omega)$ . This easily leads to

$$\begin{aligned}\Psi_\lambda(u) &\geq \frac{1}{p}\|u\|^p - c_4\|u\|^\tau - \lambda c_5 [\|u\|^\theta + \|u\|^r] \\ &= \left[ \frac{1}{p} - c_4\|u\|^{\tau-p} - \lambda c_5 (\|u\|^{\theta-p} + \|u\|^{r-p}) \right] \|u\|^p, \quad u \in X.\end{aligned}\tag{3.1}$$

Let us set, for any  $t > 0$ ,

$$\gamma_\lambda(t) := c_4 t^{\tau-p} + \lambda c_5 (t^{\theta-p} + t^{r-p}), \quad \hat{\gamma}_\lambda(t) := (c_4 + \lambda c_5) t^{\tau-p} + 2\lambda c_5 t^{r-p}.$$

From  $\tau \leq \theta < p < r$  it follows  $\lambda c_5 t^{\theta-p} \leq \lambda c_5 (t^{\tau-p} + t^{r-p})$ , which implies

$$0 < \gamma_\lambda(t) \leq \hat{\gamma}_\lambda(t) \quad \text{in } (0, +\infty).\tag{3.2}$$

Since  $\lim_{t \rightarrow 0^+} \hat{\gamma}_\lambda(t) = \lim_{t \rightarrow +\infty} \hat{\gamma}_\lambda(t) = +\infty$ , there exists  $t_0 > 0$  satisfying  $\hat{\gamma}'_\lambda(t_0) = 0$ . One has

$$t_0 := t_0(\lambda) := \left[ \frac{(c_4 + \lambda c_5)(p - \tau)}{2\lambda c_5(r - p)} \right]^{\frac{1}{r-\tau}}$$

and, via simple calculations,  $\lim_{\lambda \rightarrow 0^+} \hat{\gamma}_\lambda(t_0) = 0$ . On account of (3.1)–(3.2) we can thus find  $\lambda_0 > 0$  such that

$$\Psi_\lambda(u) \geq m_\lambda > 0 = \Psi_\lambda(0) \quad \text{for all } u \in \partial B(0, t_0), \lambda \in (0, \lambda_0).$$

Pick  $\lambda \in (0, \lambda_0)$ . The mountain pass theorem entails  $\Psi'_\lambda(\bar{u}_\lambda) = 0$  and  $\Psi_\lambda(\bar{u}_\lambda) \geq m_\lambda$  with appropriate  $\bar{u}_\lambda \in X$ . Hence,

$$\langle A_p(\bar{u}_\lambda) + A_q(\bar{u}_\lambda), v \rangle = \int_\Omega [(\bar{u}_\lambda^+)^{\tau-1} + \lambda c_0 ((\bar{u}_\lambda^+)^{\theta-1} + (\bar{u}_\lambda^+)^{r-1})] v \, dx, \quad v \in X,\tag{3.3}$$

and  $\bar{u}_\lambda \neq 0$ . Choosing  $v := -\bar{u}_\lambda^-$  in (3.3) yields  $\|\nabla \bar{u}_\lambda^-\|_p^p + \|\nabla \bar{u}_\lambda^-\|_q^q = 0$ , namely  $\bar{u}_\lambda^- = 0$ . This forces  $\bar{u}_\lambda \geq 0$  while, by (3.3) again,

$$-\Delta_p \bar{u}_\lambda - \Delta_q \bar{u}_\lambda = \bar{u}_\lambda^{\tau-1} + \lambda c_0 (\bar{u}_\lambda^{\theta-1} + \bar{u}_\lambda^{r-1}) \quad \text{in } \Omega.$$

Lieberman's nonlinear regularity theory and Pucci-Serrin's maximum principle finally lead to  $\bar{u}_\lambda \in \text{int}(C_+)$ . Now define, provided  $(x, \xi) \in \Omega \times \mathbb{R}$ ,

$$\bar{f}_\lambda(x, \xi) := \begin{cases} (\xi^+)^{\tau-1} + \lambda f(x, \xi^+) & \text{if } \xi \leq \bar{u}_\lambda(x), \\ \bar{u}_\lambda(x)^{\tau-1} + \lambda f(x, \bar{u}_\lambda(x)) & \text{otherwise,} \end{cases} \quad \bar{F}_\lambda(x, \xi) := \int_0^\xi \bar{f}_\lambda(x, t) \, dt.$$

An easy verification ensures that the associated  $C^1$ -functional

$$\bar{\Phi}_\lambda(u) := \frac{1}{p}\|\nabla u\|_p^p + \frac{1}{q}\|\nabla u\|_q^q - \int_\Omega \bar{F}_\lambda(x, u(x)) \, dx, \quad u \in X,$$

is coercive and weakly sequentially lower semicontinuous. So, it attains its infimum at some point  $u_\lambda \in X$ . Assumption  $(h_1)$  produces

$$\bar{\Phi}_\lambda(u_\lambda) < 0 = \bar{\Phi}_\lambda(0),$$

i.e.,  $u_\lambda \neq 0$ , because  $\tau < q < p$ . As before, from

$$\langle A_p(u_\lambda) + A_q(u_\lambda), v \rangle = \int_{\Omega} \bar{f}_\lambda(x, u_\lambda(x)) v(x) dx \quad \forall v \in X \quad (3.4)$$

we infer  $u_\lambda \geq 0$ . Test (3.4) with  $v := (u_\lambda - \bar{u}_\lambda)^+$ , exploit  $(h_1)$  again, and recall (3.3) to arrive at

$$\begin{aligned} \langle A_p(u_\lambda) + A_q(u_\lambda), (u_\lambda - \bar{u}_\lambda)^+ \rangle &= \int_{\Omega} [\bar{u}_\lambda^{\tau-1} + \lambda f(\cdot, \bar{u}_\lambda)] (u_\lambda - \bar{u}_\lambda)^+ dx \\ &\leq \int_{\Omega} [\bar{u}_\lambda^{\tau-1} + \lambda c_0(\bar{u}_\lambda^{\theta-1} + \bar{u}_\lambda^{\tau-1})] (u_\lambda - \bar{u}_\lambda)^+ dx \\ &= \langle A_p(\bar{u}_\lambda) + A_q(\bar{u}_\lambda), (u_\lambda - \bar{u}_\lambda)^+ \rangle, \end{aligned}$$

which entails  $u_\lambda \leq \bar{u}_\lambda$  by monotonicity. Summing up,  $u_\lambda \in [0, \bar{u}_\lambda] \setminus \{0\}$ . On account of (3.4), one thus has  $u_\lambda \in S_\lambda$  for any  $\lambda \in (0, \lambda_0)$ . This completes the proof.  $\square$

Our next result ensures that  $\mathcal{L}$  is an interval.

**Proposition 3.2.** *Let  $(h_1)$  be satisfied. If  $\hat{\lambda} \in \mathcal{L}$  then  $(0, \hat{\lambda}) \subseteq \mathcal{L}$ .*

**Proof.** Pick  $\hat{u} \in S_{\hat{\lambda}}$ ,  $\lambda \in (0, \hat{\lambda})$ , and define, provided  $(x, \xi) \in \Omega \times \mathbb{R}$ ,

$$\hat{f}_\lambda(x, \xi) := \begin{cases} (\xi^+)^{\tau-1} + \lambda f(x, \xi^+) & \text{if } \xi \leq \hat{u}(x), \\ \hat{u}(x)^{\tau-1} + \lambda f(x, \hat{u}(x)) & \text{otherwise,} \end{cases} \quad \hat{F}_\lambda(x, \xi) := \int_0^\xi \hat{f}_\lambda(x, t) dt.$$

The associated energy functional

$$\hat{\Phi}_\lambda(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} \hat{F}_\lambda(x, u(x)) dx, \quad u \in X,$$

turns out coercive, weakly sequentially lower semicontinuous, besides  $C^1$ . Now, arguing exactly as above yields the conclusion.  $\square$

A careful reading of this proof allows one to state the next ‘monotonicity’ property.

**Corollary 3.3.** *Under hypothesis  $(h_1)$ , for every  $\hat{\lambda} \in \mathcal{L}$ ,  $u_{\hat{\lambda}} \in S_{\hat{\lambda}}$ , and  $\lambda \in (0, \hat{\lambda})$  there exists  $u_\lambda \in S_\lambda$  such that  $u_\lambda \leq u_{\hat{\lambda}}$ .*

Actually, we can prove a more precise assertion.

**Proposition 3.4.** *Suppose  $(h_1)$  and  $(h_4)$  hold. Then to each  $\hat{\lambda} \in \mathcal{L}$ ,  $u_{\hat{\lambda}} \in S_{\hat{\lambda}}$ ,  $\lambda \in (0, \hat{\lambda})$  there corresponds  $u_\lambda \in S_\lambda$  fulfilling  $u_{\hat{\lambda}} - u_\lambda \in \text{int}(C_+)$ .*

**Proof.** Write  $\rho := \|u_{\hat{\lambda}}\|_{\infty}$ . If  $\mu_{\rho}$  is given by  $(h_4)$  while  $u_{\lambda}$  comes from Corollary 3.3 then

$$\begin{aligned} -\Delta_p u_{\hat{\lambda}} - \Delta_q u_{\hat{\lambda}} + \lambda \mu_{\rho} u_{\hat{\lambda}}^{p-1} &= u_{\hat{\lambda}}^{\tau-1} + \hat{\lambda} f(x, u_{\hat{\lambda}}) + \lambda \mu_{\rho} u_{\hat{\lambda}}^{p-1} \\ &= u_{\hat{\lambda}}^{\tau-1} + \lambda f(x, u_{\hat{\lambda}}) + \lambda \mu_{\rho} u_{\hat{\lambda}}^{p-1} + (\hat{\lambda} - \lambda) f(x, u_{\hat{\lambda}}) \\ &\geq u_{\hat{\lambda}}^{\tau-1} + \lambda f(x, u_{\lambda}) + \lambda \mu_{\rho} u_{\lambda}^{p-1} = -\Delta_p u_{\lambda} - \Delta_q u_{\lambda} + \lambda \mu_{\rho} u_{\lambda}^{p-1} \end{aligned} \quad (3.5)$$

because  $u_{\lambda} \leq u_{\hat{\lambda}}$  and  $f(x, t) \geq 0$  once  $t \geq 0$ . The function  $h(x) := (\hat{\lambda} - \lambda) f(x, u_{\hat{\lambda}}(x))$  lies in  $L^{\infty}(\Omega)$ . Indeed, on account of  $(h_1)$ , we have

$$0 \leq h(x) \leq c_0(\hat{\lambda} - \lambda) [\|u\|_{\infty}^{\theta-1} + \|u\|_{\infty}^{r-1}] \quad \forall x \in \Omega.$$

Pick any compact set  $K \subseteq \Omega$ . Recalling that  $u_{\hat{\lambda}} \in \text{int}(C_+)$  and using  $(h_1)$  again gives

$$h(x) \geq (\hat{\lambda} - \lambda) [c_1 u_{\hat{\lambda}}(x)^{p-1} + c_2 u_{\hat{\lambda}}(x)^{q-1}] \geq \left( c_1 \inf_K u_{\hat{\lambda}}^{p-1} + c_2 \inf_K u_{\hat{\lambda}}^{q-1} \right) > 0, \quad x \in \Omega,$$

whence  $0 \prec h$ . Now, (3.5) combined with Proposition 2.2 entails  $u_{\hat{\lambda}} - u_{\lambda} \in \text{int}(C_+)$ .  $\square$

The interval  $\mathcal{L}$  turns out to be bounded.

**Proposition 3.5.** *Let  $(h_1)$  and  $(h_4)$  be satisfied. If  $\lambda^* := \sup \mathcal{L}$  then  $\lambda^* < \infty$ .*

**Proof.** Fix  $\lambda \in \mathcal{L}$ ,  $u_{\lambda} \in S_{\lambda}$ . Note that we can suppose  $\lambda > 1$ , otherwise  $\mathcal{L}$  would be bounded, which of course entails  $\lambda^* < \infty$ . Define

$$g_{\lambda}(x, \xi) := \begin{cases} \lambda [c_1(\xi^+)^{p-1} + c_2(\xi^+)^{q-1}] & \text{if } \xi \leq u_{\lambda}(x), \\ \lambda [c_1 u_{\lambda}(x)^{p-1} + c_2 u_{\lambda}(x)^{q-1}] & \text{otherwise,} \end{cases} \quad G_{\lambda}(x, \xi) := \int_0^{\xi} g_{\lambda}(x, t) dt$$

for every  $(x, \xi) \in \Omega \times \mathbb{R}$ , as well as

$$\Psi_{\lambda}(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} G_{\lambda}(x, u(x)) dx, \quad u \in X.$$

The same arguments employed before yield here a global minimum point, say  $\bar{u}_{\lambda}$ , to  $\Psi_{\lambda}$ . So, in particular,

$$\langle A_p(\bar{u}_{\lambda}) + A_q(\bar{u}_{\lambda}), v \rangle = \int_{\Omega} g_{\lambda}(x, \bar{u}_{\lambda}(x)) v(x) dx \quad \forall v \in X. \quad (3.6)$$

Choosing  $v := -\bar{u}_{\lambda}^-$  first and then  $v := (\bar{u}_{\lambda} - u_{\lambda})^+$  we obtain  $\bar{u}_{\lambda} \in [0, u_{\lambda}]$ ; cf. the proof of Proposition 3.1. Since, by  $(p_3)$  in Section 2,  $u_{\lambda}, \phi_{1,q} \in \text{int}(C_+)$ , through [22, Proposition 1] one has  $t\phi_{1,q} \leq u_{\lambda}$ , with  $t > 0$  small enough. Thus, on account of  $(p_3)$  again,

$$\begin{aligned} \Psi_{\lambda}(t\phi_{1,q}) &= \frac{1}{p} \|\nabla(t\phi_{1,q})\|_p^p + \frac{1}{q} \|\nabla(t\phi_{1,q})\|_q^q - \int_{\Omega} G_{\lambda}(x, t\phi_{1,q}(x)) dx \\ &= \frac{t^p}{p} \|\nabla \phi_{1,q}\|_p^p + \frac{t^q}{q} \|\nabla \phi_{1,q}\|_q^q - \int_{\Omega} \lambda \left( c_1 \frac{t^p}{p} \phi_{1,q}^p + c_2 \frac{t^q}{q} \phi_{1,q}^q \right) dx \\ &= \frac{t^p}{p} \|\nabla \phi_{1,q}\|_p^p + \frac{t^q}{q} \lambda_{1,q} - \lambda c_1 \frac{t^p}{p} \|\phi_{1,q}\|_p^p - \lambda c_2 \frac{t^q}{q} \end{aligned}$$



$$\begin{aligned}
&\leq \frac{t^p}{p} \|\nabla \phi_{1,q}\|_p^p + \frac{t^q}{q} (\lambda_{1,q} - \lambda c_2) \\
&< \frac{t^p}{p} \|\nabla \phi_{1,q}\|_p^p + \frac{t^q}{q} \lambda_{1,q} (1 - \lambda) = c_6 t^p - c_7 t^q.
\end{aligned}$$

Now, recall that  $q < p$  and decrease  $t$  when necessary to achieve

$$\Psi_\lambda(\bar{u}_\lambda) = \min_X \Psi_\lambda \leq \Psi_\lambda(t\phi_{1,q}) < 0 = \Psi_\lambda(0),$$

i.e.,  $\bar{u}_\lambda \neq 0$ . Summing up,  $\bar{u}_\lambda \in [0, u_\lambda] \setminus \{0\}$ , whence, by (3.6), it turns out a positive solution of the equation

$$-\Delta_p u - \Delta_q u = \lambda c_1 |u|^{p-2} u + \lambda c_2 |u|^{q-2} u \quad \text{in } \Omega.$$

Due to [5, Theorem 2.4], this prevents  $\lambda$  from being arbitrary large, as desired.  $\square$

Let us finally prove that  $\mathcal{L} = (0, \lambda^*]$ . From now on,  $\Phi_\lambda : X \rightarrow \mathbb{R}$  will denote the  $C^1$ -energy functional associated with problem  $(P_\lambda)$ . Evidently,

$$\Phi_\lambda(u) = \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \frac{1}{\tau} \|u^+\|_\tau^\tau - \lambda \int_\Omega F(x, u^+(x)) dx \quad \forall u \in X. \quad (3.7)$$

**Proposition 3.6.** *Under  $(h_1)$ ,  $(h_3)$ , and  $(h_4)$  one has  $\lambda^* \in \mathcal{L}$ .*

**Proof.** Pick any  $\{\lambda_n\} \subseteq (0, \lambda^*)$  fulfilling  $\lambda_n \uparrow \lambda^*$ . Via Corollary 3.3, construct a sequence  $\{u_n\} \subseteq X$  such that  $u_n \in S_{\lambda_n}$ ,  $u_n \leq u_{n+1}$ . Then

$$\langle A_p(u_n) + A_q(u_n), v \rangle = \int_\Omega u_n^{\tau-1} v dx + \lambda_n \int_\Omega f(\cdot, u_n) v dx, \quad v \in X. \quad (3.8)$$

We can also assume  $\Phi_\lambda(u_n) < 0$  (see the proof of Proposition 3.1), which means

$$\|\nabla u_n\|_p^p + \frac{p}{q} \|\nabla u_n\|_q^q - \frac{p}{\tau} \|u_n\|_\tau^\tau - \lambda_n \int_\Omega pF(x, u_n(x)) dx < 0. \quad (3.9)$$

Testing (3.8) with  $v := u_n$  gives

$$\|\nabla u_n\|_p^p + \|\nabla u_n\|_q^q = \|u_n\|_\tau^\tau + \lambda_n \int_\Omega f(\cdot, u_n) u_n dx. \quad (3.10)$$

Since  $q < p$  while  $\lambda_1 \leq \lambda_n$ , from (3.9)–(3.10) it follows

$$\int_\Omega [f(\cdot, u_n) u_n - pF(\cdot, u_n)] dx \leq \frac{1}{\lambda_1} \left( \frac{p}{\tau} - 1 \right) \|u_n\|_\tau^\tau \quad \forall n \in \mathbb{N}. \quad (3.11)$$

Observe next that, thanks to  $(h_1)$  and  $(h_3)$ , one has

$$f(x, \xi) \xi - pF(x, \xi) \geq c_8 \xi^\beta - c_9 \quad \text{in } \Omega \times \mathbb{R}_+.$$

Consequently, (3.11) becomes

$$c_8 \|u_n\|_\beta^\beta \leq \frac{1}{\lambda_1} \left( \frac{p}{\tau} - 1 \right) \|u_n\|_\tau^\tau + c_{10} \leq c_{11} \|u_n\|_\beta^\tau + c_{10}, \quad n \in \mathbb{N},$$

because  $\tau < \beta$ . This clearly forces

$$\|u_n\|_\beta \leq c_{12} \quad \forall n \in \mathbb{N}. \quad (3.12)$$

If  $r \leq \beta$  then  $\{u_n\}$  turns out also bounded in  $L^r(\Omega)$ . Using (3.10) besides  $(h_1)$  entails

$$\begin{aligned} \|u_n\|^p &\leq \|\nabla u_n\|_p^p + \|\nabla u_n\|_q^q \leq \|u_n\|_\tau^\tau + \lambda^* \int_\Omega f(\cdot, u_n) u_n \, dx \\ &\leq |\Omega|^{1-\tau/r} \|u_n\|_r^\tau + \lambda^* c_0 \int_\Omega (u_n^\theta + u_n^r) \, dx \\ &\leq |\Omega|^{1-\tau/r} \|u_n\|_r^\tau + \lambda^* c_0 \int_\Omega [(1 + u_n^r) + u_n^r] \, dx, \end{aligned} \quad (3.13)$$

whence  $\{u_n\} \subseteq X$  is bounded. Suppose now  $\beta < r < p^*$ . Two cases may occur.

1)  $p < N$ . Let  $t \in (0, 1)$  satisfy

$$\frac{1}{r} = \frac{1-t}{\beta} + \frac{t}{p^*}. \quad (3.14)$$

The interpolation inequality [12, p. 905] yields  $\|u_n\|_r \leq \|u_n\|_\beta^{1-t} \|u_n\|_{p^*}^t$ . Via (3.12) we thus obtain

$$\|u_n\|_r^r \leq c_{13} \|u_n\|_{p^*}^{tr}, \quad n \in \mathbb{N}. \quad (3.15)$$

Reasoning exactly as before and exploiting (3.15) produce

$$\|u_n\|^p \leq \|\nabla u_n\|_p^p + \|\nabla u_n\|_q^q \leq c_{14} (1 + \|u_n\|_{p^*}^{tr}) \leq c_{15} (1 + \|u_n\|^{tr}). \quad (3.16)$$

Finally, note that  $tr < p$ . Indeed,  $(r-p)\frac{N}{p} < \beta$  due to  $(h_3)$ , while

$$tr < p \iff \frac{r-\beta}{p^*-\beta} < \frac{p}{p^*} \iff (r-p)\frac{N}{p} < \beta;$$

cf. (3.14). Now, the boundedness of  $\{u_n\} \subseteq X$  directly stems from (3.16).

2)  $p \geq N$ , which implies  $p^* = +\infty$ . We will repeat the previous argument with  $p^*$  replaced by any  $\sigma > r$ . Accordingly, if  $t \in (0, 1)$  fulfills  $\frac{1}{r} = \frac{1-t}{\beta} + \frac{t}{\sigma}$  then  $tr = \frac{\sigma(r-\beta)}{\sigma-\beta}$ . Since, thanks to  $(h_3)$  again,

$$\lim_{\sigma \rightarrow +\infty} \frac{\sigma(r-\beta)}{\sigma-\beta} = r - \beta < p,$$

one arrives at  $tr < p$  for  $\sigma$  large enough. This entails  $\{u_n\} \subseteq X$  bounded once more.

Hence, in either case, we may assume

$$u_n \rightharpoonup u^* \text{ in } X \quad \text{and} \quad u_n \rightarrow u^* \text{ in } L^r(\Omega), \quad (3.17)$$

where a subsequence is considered when necessary. Testing (3.8) with  $v := u_n - u^*$  thus yields, as  $n \rightarrow +\infty$ ,

$$\lim_{n \rightarrow +\infty} \langle A_p(u_n) + A_q(u_n), u_n - u^* \rangle = 0,$$

whence, by monotonicity of  $A_q$ ,

$$\limsup_{n \rightarrow +\infty} [\langle A_p(u_n), u_n - u^* \rangle + \langle A_q(u), u_n - u^* \rangle] \leq 0.$$

On account of (3.17) it follows

$$\limsup_{n \rightarrow +\infty} \langle A_p(u_n), u_n - u^* \rangle \leq 0.$$

Recalling that  $A_p$  enjoys the  $(S)_+$ -property, we infer  $u_n \rightarrow u^*$  in  $X$ , besides  $0 \leq u_n \leq u^*$  for all  $n \in \mathbb{N}$ . Finally, let  $n \rightarrow +\infty$  in (3.8) to get

$$\langle A_p(u^*) + A_q(u^*), v \rangle = \int_{\Omega} (u^*)^{\tau-1} v \, dx + \lambda^* \int_{\Omega} f(\cdot, u^*) v \, dx \quad \forall v \in X,$$

i.e.,  $u^* \in S_{\lambda^*}$  and, a fortiori,  $\lambda^* \in \mathcal{L}$ .  $\square$

Some meaningful (bifurcation) properties of the set  $S_{\lambda}$  will now be established.

**Proposition 3.7.** *Suppose  $(h_1)$ – $(h_4)$  hold true. Then, for every  $\lambda \in (0, \lambda^*)$ , problem  $(P_{\lambda})$  admits two solutions  $u_0, \hat{u} \in \text{int}(C_+)$  such that  $u_0 \leq \hat{u}$ . Moreover,  $u_0$  is a local minimizer of the associated energy functional  $\Phi_{\lambda}$ .*

**Proof.** Fix  $\lambda \in (0, \lambda^*)$  and choose  $\eta \in (\lambda, \lambda^*)$ . By Proposition 3.2, there exists  $u_{\eta} \in S_{\eta}$  while Proposition 3.4 provides  $u_0 \in S_{\lambda}$  satisfying

$$u_0 \in \text{int}_{C_0^1(\overline{\Omega})}([0, u_{\eta}]). \quad (3.18)$$

The same reasoning adopted in the proof of Proposition 3.2 ensures here that  $u_0$  is a global minimum point to the functional

$$\Phi_{\lambda, \eta}(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} F_{\lambda, \eta}(x, u(x)) \, dx, \quad u \in X,$$

where  $F_{\lambda, \eta}(x, \xi) := \int_0^{\xi} f_{\lambda, \eta}(x, t) \, dt$ , with

$$f_{\lambda, \eta}(x, \xi) := \begin{cases} (\xi^+)^{\tau-1} + \lambda f(x, \xi^+) & \text{if } \xi \leq u_{\eta}(x), \\ u_{\eta}(x)^{\tau-1} + \lambda f(x, u_{\eta}(x)) & \text{otherwise.} \end{cases}$$

By (3.18),  $u_0$  turns out a local  $C_0^1(\overline{\Omega})$ -minimizer of  $\Phi_{\lambda}$ , because  $\Phi_{\lambda}|_{[0, u_{\eta}]} = \Phi_{\lambda, \eta}|_{[0, u_{\eta}]}$ . Via Proposition 2.1 we then see that this remains valid with  $C_0^1(\overline{\Omega})$  replaced by  $X$ . Set

$$f_0(x, \xi) := \begin{cases} u_0(x)^{\tau-1} + \lambda f(x, u_0(x)) & \text{if } \xi \leq u_0(x), \\ \xi^{\tau-1} + \lambda f(x, \xi) & \text{otherwise,} \end{cases} \quad F_0(x, \xi) := \int_0^{\xi} f_0(x, t) \, dt, \quad (3.19)$$

$(x, \xi) \in \Omega \times \mathbb{R}$ , as well as

$$\Phi_0(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \int_{\Omega} F_0(x, u(x)) dx \quad \forall u \in X. \quad (3.20)$$

From (3.19) and the nonlinear regularity theory it follows  $u_0 \in K(\Phi_0) \subseteq [u_0] \cap \text{int}(C_+)$ . We may thus assume

$$K(\Phi_0) \cap [u_0, u_\eta] = \{u_0\}, \quad (3.21)$$

or else a second solution of  $(P_\lambda)$  bigger than  $u_0$  would exist. Bearing in mind the proof of Proposition 3.6 and making small changes to accommodate the truncation at  $u_0(x)$  shows that  $\Phi_0$  satisfies condition (C). Let us next truncate  $f_0(x, \cdot)$  at  $u_\eta(x)$  to construct a new Carathéodory function  $\tilde{f}$ , with primitive  $\tilde{F}$  and associated functional  $\tilde{\Phi}$ , defined like in (3.20) but replacing  $F_0$  by  $\tilde{F}$ . Evidently,

$$K(\tilde{\Phi}) = K(\Phi_0) \cap [u_0, u_\eta],$$

whence  $K(\tilde{\Phi}) = \{u_0\}$  because of (3.21). Since  $\tilde{\Phi}$  is coercive and weakly sequentially lower semicontinuous, it possesses a global minimum point that must coincide with  $u_0$ . An easy verification gives  $\Phi_0|_{[0, u_\eta]} = \tilde{\Phi}|_{[0, u_\eta]}$ . So, thanks to (3.18),  $u_0$  turns out a local  $C_0^1(\bar{\Omega})$ -minimizer of  $\Phi_0$ . This still holds when  $X$  replaces  $C_0^1(\bar{\Omega})$ ; cf. Proposition 2.1. We may suppose  $K(\Phi_0)$  finite, otherwise infinitely many solutions of  $(P_\lambda)$  bigger than  $u_0$  do exist. Adapting the argument exploited in [1, Proposition 29] provides  $\rho \in (0, 1)$  such that

$$\Phi_0(u_0) < m_0 := \inf\{\Phi_0(u) : \|u - u_0\| = \rho\}. \quad (3.22)$$

Finally, if  $u \in \text{int}(C_+)$  then simple calculations based on  $(h_2)$  entail  $\Phi_0(tu) \rightarrow -\infty$  as  $t \rightarrow +\infty$ . Therefore, the mountain pass theorem can be applied, and there is  $\hat{u} \in X$  fulfilling

$$\hat{u} \in K(\Phi_0), \quad \Phi_0(\hat{u}) \geq m_0. \quad (3.23)$$

Via (3.22)–(3.23) one has  $u_0 \neq \hat{u}$  while the inclusion  $K(\Phi_0) \subseteq [u_0] \cap \text{int}(C_+)$  forces  $u_0 \leq \hat{u}$ , which ends the proof.  $\square$

**Proposition 3.8.** *Under  $(h_1)$ – $(h_4)$ , the solution set  $S_\lambda$  admits a smallest element  $\bar{u}_\lambda$  for every  $\lambda \in \mathcal{L}$ .*

**Proof.** A standard procedure ensures that  $S_\lambda$  turns out downward directed; see, e.g., [10, Section 4]. Lemma 3.10 at p. 178 of [17] yields

$$\text{ess inf } S_\lambda = \inf\{u_n : n \in \mathbb{N}\} \quad (3.24)$$

for some decreasing sequence  $\{u_n\} \subseteq S_\lambda$ . Consequently,  $0 \leq u_n \leq u_1$  and

$$\langle A_p(u_n) + A_q(u_n), v \rangle = \int_{\Omega} [u_n^{\tau-1} + \lambda f(\cdot, u_n)] v dx \quad \forall v \in X. \quad (3.25)$$

Due to  $(h_1)$ , testing (3.25) with  $v := u_n$  we thus obtain

$$\begin{aligned} \|u_n\|^p &\leq \|\nabla u_n\|_p^p + \|\nabla u_n\|_q^q = \int_{\Omega} [u_n^{\tau} + \lambda f(\cdot, u_n) u_n] dx \\ &\leq \int_{\Omega} [u_n^{\tau} + \lambda c_0 (u_n^{\theta} + u_n^r)] dx \leq \int_{\Omega} [u_1^{\tau} + \lambda c_0 (u_1^{\theta} + u_1^r)] dx, \quad n \in \mathbb{N}, \end{aligned}$$

namely  $\{u_n\} \subseteq X$  is bounded. Like before (cf. the proof of Proposition 3.6), this gives  $u_n \rightarrow \bar{u}_\lambda$  in  $X$ , where a subsequence is considered if necessary. So, from (3.25) it easily follows

$$\langle A_p(\bar{u}_\lambda) + A_q(\bar{u}_\lambda), v \rangle = \int_{\Omega} [\bar{u}_\lambda^{\tau-1} + \lambda f(\cdot, \bar{u}_\lambda)] v \, dx \quad \forall v \in X.$$

Showing that  $\bar{u}_\lambda \neq 0$  will entail  $\bar{u}_\lambda \in S_\lambda$ , whence the conclusion by (3.24). To the aim, consider the problem

$$-\Delta_p u - \Delta_q u = u^{\tau-1} \quad \text{in } \Omega, \quad u > 0 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega. \quad (3.26)$$

Its energy functional

$$\Phi_0(u) := \frac{1}{p} \|\nabla u\|_p^p + \frac{1}{q} \|\nabla u\|_q^q - \frac{1}{\tau} \|u^+\|_\tau^\tau, \quad u \in X,$$

turns out coercive and weakly sequentially lower semicontinuous. Hence, there exists  $\tilde{u} \in X$  satisfying  $\Phi_0(\tilde{u}) = \inf_X \Phi_0$ . One has  $\tilde{u}_0 \neq 0$ , because  $\Phi_0(\tilde{u}) < 0 = \Phi_0(0)$  (the argument is like in the proof of Proposition 3.5). Further,  $\Phi'_0(\tilde{u}) = 0$ , i.e.,

$$\langle A_p(\tilde{u}) + A_q(\tilde{u}), v \rangle = \int_{\Omega} (\tilde{u}^+)^{\tau-1} v \, dx \quad \forall v \in X.$$

Choosing  $v := -\tilde{u}^-$  we see that  $u$  is a positive solution to (3.26). Actually,  $\tilde{u} \in \text{int}(C_+)$  and, through a standard procedure [15, Lemma 3.1],  $\tilde{u}$  turns out unique.

**Claim:**  $\tilde{u} \leq u$  for all  $u \in S_\lambda$ .

Indeed, for any fixed  $u \in S_\lambda$ , define

$$\Psi(w) := \frac{1}{p} \|\nabla w\|_p^p + \frac{1}{q} \|\nabla w\|_q^q - \int_{\Omega} dx \int_0^{w(x)} g(x, t) \, dt, \quad w \in X,$$

where

$$g(x, t) := \begin{cases} (t^+)^{\tau-1} & \text{if } t \leq u(x), \\ u(x)^{\tau-1} & \text{otherwise} \end{cases} \quad \forall (x, t) \in \Omega \times \mathbb{R}.$$

The following assertions can be easily verified.

- $\Psi(u^*) = \inf_X \Psi$ , with appropriate  $u^* \in X$ .
- $\Psi(u^*) < 0 = \Psi(0)$ , whence  $u^* \neq 0$ .
- $u^* \in K(\Psi) \subseteq [0, u] \cap C_+$ .

Therefore,  $u^*$  is a positive solution of (3.26). By uniqueness, this implies  $u^* = \tilde{u}$ . Thus, a fortiori,  $\tilde{u} \leq u$ .

The claim brings  $\tilde{u} \leq u_n$ ,  $n \in \mathbb{N}$ , which in turn provides  $0 < \tilde{u} \leq \bar{u}_\lambda$ , as desired.  $\square$

Let us finally come to some meaningful properties of the map

$$k : \lambda \in \mathcal{L} \mapsto \bar{u}_\lambda \in C_0^1(\bar{\Omega}).$$

**Proposition 3.9.** *Suppose  $(h_1)$ – $(h_4)$  hold true. Then the function  $k$  is both*

- (i<sub>1</sub>) strictly increasing, namely  $\bar{u}_{\lambda_2} - \bar{u}_{\lambda_1} \in \text{int}(C_+)$  if  $\lambda_1 < \lambda_2$ , and*
- (i<sub>2</sub>) left-continuous.*

**Proof.** Pick  $\lambda_1, \lambda_2 \in \mathcal{L}$  such that  $\lambda_1 < \lambda_2$ . Since  $\bar{u}_{\lambda_2} \in S_{\lambda_2}$ , Proposition 3.4 yields  $u_{\lambda_1} \in S_{\lambda_1}$  fulfilling  $\bar{u}_{\lambda_2} - u_{\lambda_1} \in \text{int}(C_+)$ , while Proposition 3.8 entails  $\bar{u}_{\lambda_1} \leq u_{\lambda_1}$ . Hence,  $\bar{u}_{\lambda_2} - \bar{u}_{\lambda_1} \in \text{int}(C_+)$ . This shows (i<sub>1</sub>).

If  $\lambda_n \rightarrow \lambda^-$  in  $\mathcal{L}$  then, by (i<sub>1</sub>), the sequence  $\{\bar{u}_{\lambda_n}\}$  turns out increasing. Its boundedness in  $X$  immediately stems from (h<sub>1</sub>); see the previous proof. Now, repeat the argument below (3.17) to arrive at

$$\bar{u}_{\lambda_n} \rightarrow \tilde{u}_\lambda \text{ in } X, \quad (3.27)$$

whence  $\tilde{u}_\lambda \in S_\lambda \subseteq \text{int}(C_+)$ . We finally claim that  $\tilde{u}_\lambda = \bar{u}_\lambda$ . Assume on the contrary

$$\bar{u}_\lambda(x_0) < \tilde{u}_\lambda(x_0) \text{ for some } x_0 \in \Omega. \quad (3.28)$$

Lieberman's nonlinear regularity theory gives  $\{\bar{u}_n\} \subseteq C_0^{1,\alpha}(\bar{\Omega})$  as well as

$$\|\bar{u}_{\lambda_n}\|_{C_0^{1,\alpha}(\bar{\Omega})} \leq c_{16} \quad \forall n \in \mathbb{N}.$$

Since the embedding  $C_0^{1,\alpha}(\bar{\Omega}) \hookrightarrow C_0^1(\bar{\Omega})$  is compact, (3.27) becomes

$$\bar{u}_{\lambda_n} \rightarrow \tilde{u}_\lambda \text{ in } C_0^1(\bar{\Omega}).$$

Because of (3.28), this implies  $\bar{u}_\lambda(x_0) < \bar{u}_{\lambda_n}(x_0)$  for any  $n$  large enough, against (i<sub>1</sub>). Consequently,  $\tilde{u}_\lambda = \bar{u}_\lambda$ , and (i<sub>2</sub>) follows from (3.27).  $\square$

Gathering Propositions 3.1–3.9 together we obtain the following

**Theorem 3.10.** *Let  $(h_1)$ – $(h_4)$  be satisfied. Then, there exists  $\lambda^* > 0$  such that problem  $(P_\lambda)$  admits*

- (j<sub>1</sub>) at least two solutions  $u_0, \hat{u} \in \text{int}(C_+)$ , with  $u_0 \leq \hat{u}$ , for every  $\lambda \in (0, \lambda^*)$ ,*
- (j<sub>2</sub>) at least one solution  $u^* \in \text{int}(C_+)$  when  $\lambda = \lambda^*$ ,*
- (j<sub>3</sub>) no positive solutions for all  $\lambda > \lambda^*$ ,*
- (j<sub>4</sub>) a smallest positive solution  $\bar{u}_\lambda \in \text{int}(C_+)$  provided  $\lambda \in (0, \lambda^*]$ .*

Moreover, the map  $\lambda \in (0, \lambda^*] \mapsto \bar{u}_\lambda \in C_0^1(\bar{\Omega})$  is strictly increasing and left-continuous.

## Acknowledgment

This work is performed within the 2016–2018 Research Plan – Intervention Line 2: ‘Variational Methods and Differential Equations’, and partially supported by INdAM/GNAMPA.

## References

- [1] S. Aizicovici, N.S. Papageorgiou, V. Staicu, Degree theory for operators of monotone type and nonlinear elliptic equations with inequality constraints, *Mem. Amer. Math. Soc.* 196 (2008).
- [2] A. Ambrosetti, H. Brezis, G. Cerami, Combined effects of concave and convex nonlinearities in some elliptic problems, *J. Funct. Anal.* 122 (1994) 519–543.
- [3] D. Arcoya, D. Ruiz, The Ambrosetti-Prodi problem for the  $p$ -Laplace operator, *Comm. Partial Differential Equations* 31 (2006) 849–865.

- [4] G. Barletta, R. Livrea, N.S. Papageorgiou, Bifurcation phenomena for the positive solutions of semilinear elliptic problems with mixed boundary conditions, *J. Nonlinear Convex Anal.* 17 (2016) 1497–1516.
- [5] V. Bobkov, M. Tanaka, On positive solutions for  $(p, q)$ -Laplace equations with two parameters, *Calc. Var. Partial Differential Equations* 54 (2015) 3277–3301.
- [6] V. Bobkov, M. Tanaka, Remarks on minimizers for  $(p, q)$ -Laplace equations with two parameters, *Commun. Pure Appl. Anal.* 17 (2018) 1219–1253.
- [7] L. Cherfils, Y. Ilyasov, On the stationary solutions of generalized reaction diffusion equations with  $(p, q)$ -Laplacian, *Commun. Pure Appl. Anal.* 4 (2005) 9–22.
- [8] G.H. Derrick, Comments on nonlinear wave equations as models for elementary particles, *J. Math. Phys.* 5 (1964) 1252–1254.
- [9] P.C. Fife, *Mathematical Aspects of Reacting and Diffusing Systems*, Lecture Notes in Biomath., vol. 28, Springer, Berlin, 1979.
- [10] M. Filippakis, N.S. Papageorgiou, Multiple constant sign and nodal solutions for nonlinear elliptic equations with the  $p$ -Laplacian, *J. Differential Equations* 245 (2008) 1883–1922.
- [11] J. Garcia Azorero, J. Manfredi, I. Peral Alonso, Sobolev versus Hölder local minimizers and global multiplicity for some quasilinear elliptic equations, *Commun. Contemp. Math.* 2 (2000) 385–404.
- [12] L. Gasiński, N.S. Papageorgiou, *Nonlinear Analysis*, Chapman & Hall, CRC, Boca Raton, FL, 2006.
- [13] L. Gasiński, N.S. Papageorgiou, Multiple solutions for nonlinear coercive problems with a nonhomogeneous differential operator and a nonsmooth potential, *Set-Valued Var. Anal.* 20 (2012) 417–443.
- [14] L. Gasiński, N.S. Papageorgiou, Bifurcation-type results for nonlinear parametric elliptic equations, *Proc. Roy. Soc. Edinburgh Sect. A* 142 (2012) 595–623.
- [15] U. Guarnotta, S.A. Marano, N.S. Papageorgiou, Multiple nodal solutions to a Robin problem with sign-changing potential and locally defined reaction, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* (2019), in press.
- [16] Z. Guo, Z. Zhang,  $W^{1,p}$  versus  $C^1$  local minimizers and multiplicity results for quasilinear elliptic equations, *J. Math. Anal. Appl.* 286 (2003) 32–50.
- [17] S. Hu, N.S. Papageorgiou, *Handbook of Multivalued Analysis*, vol. I: Theory, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1997.
- [18] G. Lieberman, The natural generalization of the natural conditions of Ladyshenskaya and Ural'tseva for elliptic equations, *Comm. Partial Differential Equations* 16 (1991) 311–361.
- [19] S.A. Marano, S. Mosconi, Some recent results on the Dirichlet problem for  $(p, q)$ -Laplace equations, *Discrete Contin. Dyn. Syst. Ser. S* 11 (2018) 279–291.
- [20] S.A. Marano, S. Mosconi, N.S. Papageorgiou, Multiple solutions to  $(p, q)$ -Laplacian problems with resonant concave nonlinearity, *Adv. Nonlinear Stud.* 16 (2016) 51–65.
- [21] S.A. Marano, S. Mosconi, N.S. Papageorgiou, On a  $(p, q)$ -Laplacian problem with concave and asymmetric perturbation, *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.* 29 (2018) 109–125.
- [22] S.A. Marano, N.S. Papageorgiou, Positive solutions to a Dirichlet problem with  $p$ -Laplacian and concave-convex nonlinearity depending on a parameter, *Commun. Pure Appl. Anal.* 12 (2013) 815–829.
- [23] N.S. Papageorgiou, V.D. Radulescu, Bifurcation of positive solutions for nonlinear nonhomogeneous Robin and Neumann problems with competing nonlinearities, *Discrete Contin. Dyn. Syst.* 35 (2016) 5008–5036.
- [24] N.S. Papageorgiou, V.D. Radulescu, D. Repovš, Robin problems with indefinite linear part and competition phenomena, *Commun. Pure Appl. Anal.* 16 (2017) 1293–1314.
- [25] P. Pucci, J. Serrin, *The Maximum Principle*, Birkhäuser, Basel, 2007.
- [26] H. Wilhelmson, Explosive instabilities of reaction-diffusion equations, *Phys. Rev. A* 36 (1987) 965–966.
- [27] V.V. Zhikov, Averaging of functionals of the calculus of variations and elasticity theory, *Izv. Akad. Nauk SSSR Ser. Mat.* 50 (1986) 675–710; English translation in *Math. USSR, Izv.* 29 (1987) 33–66.