

On a q -deformation of modular forms[☆]Victor J.W. Guo^a, Wadim Zudilin^{b,*}^a School of Mathematical Sciences, Huaiyin Normal University, Huai'an 223300, Jiangsu, People's Republic of China^b Department of Mathematics, IMAPP, Radboud University, PO Box 9010, 6500 GL Nijmegen, Netherlands

ARTICLE INFO

Article history:

Received 28 December 2018

Available online 20 March 2019

Submitted by S. Cooper

To Bruce Berndt, with admiration and warm wishes, on his

 $\frac{(q; q)_6}{(1-q)^4(1-q^3)^2}$ birthday, as $q \rightarrow 1$

Keywords:

Ramanujan

 q -Analogue

Cyclotomic polynomial

Basic hypergeometric function

(Super)congruence

ABSTRACT

There are many instances known when the Fourier coefficients of modular forms are congruent to partial sums of hypergeometric series. In our previous work, such partial sums are related to the radial asymptotics of infinite q -hypergeometric sums at roots of unity. Here we combine the two features to construct a hypergeometric q -deformation of two CM modular forms of weight 3 and discuss the corresponding q -congruences.

© 2019 Elsevier Inc. All rights reserved.

1. Introduction

The hypergeometric identity

$$\sum_{k=0}^{\infty} (4k+1) \frac{\left(\frac{1}{2}\right)_k^3}{k!^3} (-1)^k = \frac{2}{\pi}, \quad (1.1)$$

due to G. Bauer [7], is more than 150 years old but still attracts a lot of mathematical interest because of its belonging to a family of the so-called Ramanujan-type identities for $1/\pi$ (see [6,10,23,26,32] and, in particular, [3, Chap. 15] and [9, Chap. 14]). Here $(a)_k = \prod_{j=0}^{k-1} (a+j) = \Gamma(a+k)/\Gamma(a)$ is the Pochhammer notation, so that $(\frac{1}{2})_k/k! = \binom{2k}{k} 2^{-2k}$ for $k = 0, 1, 2, \dots$. One arithmetic manifestation of a special status of (1.1) are supercongruences

[☆] The first author was partially supported by the National Natural Science Foundation of China (grant 11771175).

* Corresponding author.

E-mail addresses: jwguo@hytc.edu.cn (V.J.W. Guo), w.zudilin@math.ru.nl (W. Zudilin).

$$\begin{aligned} \sum_{k=0}^{p-1} (4k+1) \frac{(\frac{1}{2})_k^3}{k!^3} (-1)^k &\equiv \sum_{k=0}^{(p-1)/2} (4k+1) \frac{(\frac{1}{2})_k^3}{k!^3} (-1)^k \\ &\equiv (-1)^{(p-1)/2} p \pmod{p^3} \quad \text{for primes } p > 2, \end{aligned} \quad (1.2)$$

observed by L. Van Hamme [31, Conjecture (B.2)] and proved subsequently by E. Mortenson [24] (see also [33]).

A principal objective of our earlier work [21] was demonstration that the hypergeometric evaluations like (1.1) and congruences for truncated sums like (1.2) may be deduced, in a uniform way, from suitable q -deformations of (1.1). Namely, the asymptotics of such q -deformations as q tends radially to a root of unity governs the behaviour of partial sums related to the degree of that root. Notice that the right-hand sides $a(p) = (-1)^{(p-1)/2} p$ of the congruences (1.2) combine to the Dirichlet generating function

$$L(s) = \prod_{p>2} (1 - a(p)p^{-s})^{-1} = \sum_{m=0}^{\infty} \frac{(-1)^m}{(2m+1)^{s-1}} = L(\chi_{-4}, s-1), \quad (1.3)$$

where $\chi_{-4} = (\frac{-1}{\cdot})$ is the nonprincipal modulo 4 character, and the series evaluates to $\pi/4$ at $s = 2$, so that (1.1) transforms into

$$\sum_{k=0}^{\infty} (4k+1) \frac{(\frac{1}{2})_k^3}{k!^3} (-1)^k = \frac{8L(2)}{\pi^2}.$$

For some other recent progress on q -congruences, we refer the reader to [12–15, 17, 18].

A theme of this note is to give examples of q -deformations of hypergeometric evaluations, which are linked with the coefficients of modular forms rather than Dirichlet characters, and use these q -hypergeometric identities to establish the corresponding (super)congruences.

2. Hypergeometric identities and congruences

A forward player of our exposition is the hypergeometric evaluation

$$\sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^3}{k!^3} = \frac{\pi}{\Gamma(3/4)^4}.$$

Its right-hand side happens to be a (simple multiple of the) period of the CM modular form

$$f_1(\tau) = q \prod_{m=1}^{\infty} (1 - q^{4m})^6 = \sum_{n=1}^{\infty} a_1(n) q^n, \quad \text{where } q = \exp(2\pi i \tau), \quad (2.1)$$

of weight 3:

$$\sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^3}{k!^3} = \frac{\Gamma(1/2)^2}{\Gamma(3/4)^4} = \frac{16L(f_1, 2)}{\pi^2} = \frac{8L(f_1, 1)}{\pi} \quad (2.2)$$

(see [28, Theorem 5]), where $L(f_1, s)$ denotes the Dirichlet L -function of (2.1). This relationship between the hypergeometric series and modular form is somewhat deeper because of the chain of related congruences

$$\sum_{k=0}^{p-1} \frac{(\frac{1}{2})_k^3}{k!^3} \equiv a_1(p) \pmod{p^2} \quad \text{for primes } p > 2 \quad (2.3)$$

(see [25,30]), which link corresponding partial sums with the Fourier coefficients of (2.1). The latter can be given explicitly via

$$a_1(p) = \begin{cases} 2(a^2 - b^2) & \text{if } p = a^2 + b^2, a \text{ odd,} \\ 0 & \text{if } p \equiv 3 \pmod{4}, \end{cases} \quad (2.4)$$

and, in turn, satisfy

$$a_1(p) \equiv \frac{\Gamma_p(1/2)^2}{\Gamma_p(3/4)^4} \pmod{p^2} \quad \text{for } p > 2$$

(see [31, Sect. 1 and Conjecture (A.2)]). This visually makes the series in (2.1) a plausible generating function of the truncations in (2.3). Furthermore, we remark that [30]

$$a_1(p) \equiv \frac{(\frac{1}{2})_m^2}{m!^2} \pmod{p^2} \quad \text{if } m = (p-1)/4 \in \mathbb{Z}.$$

We should stress that not every formal q -analogue of a hypergeometric summation, $|q| < 1$, may suit for application of the general machinery from our earlier work [21]. An example of q -deforming (2.2) is given by

$$\sum_{k=0}^{\infty} (1 + q^{4k+1}) \frac{(q^2; q^4)_k^3}{(q^4; q^4)_k^3} q^k = \frac{(q^2; q^4)_{\infty}^2 (q^3; q^4)_{\infty}^2}{(q; q^4)_{\infty}^2 (q^4; q^4)_{\infty}^2} \quad (2.5)$$

(apply [21, eq. (52)] with $a = b = c = 1$), where the q -notation $(a; q)_k$ stands for $\prod_{j=0}^{k-1} (1 - aq^j)$ for $k = 0, 1, 2, \dots, \infty$. Identity (2.5) originates however from

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{1 - q^{4k+1}}{1 - q} \frac{(q^2; q^4)_k^3}{(q^4; q^4)_k^3} (-q)^k &= \frac{(q^2; q^4)_{\infty}^2 (-q^3; q^4)_{\infty}^2}{(1 - q) (-q; q^4)_{\infty}^2 (q^4; q^4)_{\infty}^2} \\ &= \frac{(q^5; q^4)_{\infty}^4 (q^6; q^8)_{\infty}^4}{(q; q^2)_{\infty}^2 (q^4; q^4)_{\infty}^2} \end{aligned} \quad (2.6)$$

(replace q with $-q$), which makes it a q -analogue of Bauer's formula (1.1). This 'true' origin (1.1) makes the asymptotics of (2.5) at roots of unity q related to the truncated sums of (1.1) rather than (2.2).

In this note, we give a q -extension of (2.2) that accommodates the related congruences (2.3) and, by these means, implicitly provides a q -generalisation of the generating function (2.1). We also provide a similar q -extension of the formula

$${}_3F_2\left(\begin{matrix} \frac{1}{2}, \frac{1}{2}, \frac{1}{2} \\ 1, 1 \end{matrix} \middle| -1\right) = \frac{\Gamma(1/2)^2}{\sqrt{2}\Gamma(5/8)^2\Gamma(7/8)^2} = \frac{12\sqrt{2}L(f_2, 2)}{\pi^2} = \frac{12L(f_2, 1)}{\pi} \quad (2.7)$$

(see [28, Theorem 5]), which underlies (1.1) and is attached to the weight 3 CM modular form

$$f_2(\tau) = \sum_{n=1}^{\infty} a_2(n)q^n = q \prod_{m=1}^{\infty} (1 - q^m)^2 (1 - q^{2m}) (1 - q^{4m}) (1 - q^{8m})^2.$$

In this case, we have [25]

$$a_2(p) = \begin{cases} 2(a^2 - 2b^2) & \text{if } p = a^2 + 2b^2, \\ 0 & \text{if } p \equiv 5, 7 \pmod{8}, \end{cases}$$

as well as

$$(-1)^{(p-1)/2} \sum_{k=0}^{p-1} \frac{(\frac{1}{2})_k^3}{k!^3} (-1)^k \equiv \frac{\Gamma_p(1/2)^2}{\Gamma_p(5/8)^2 \Gamma_p(7/8)^2} \equiv a_2(p) \pmod{p^2} \quad (2.8)$$

for primes $p > 2$ and

$$a_2(p) \equiv \frac{(\frac{1}{4})_m^2}{m!^2} \times \begin{cases} 1 & \text{if } m = (p-1)/8 \in \mathbb{Z} \\ -\frac{1}{2} & \text{if } m = (3p-1)/8 \in \mathbb{Z} \end{cases} \pmod{p^2}.$$

3. A q -Clausen identity

F.H. Jackson's generalisation of Clausen's identity [22] (see also [29, eq. (3.2)]) implies

$${}_2\phi_1 \left[\begin{matrix} qa, q/a \\ q^4 \end{matrix}; q^4, z \right] {}_2\phi_1 \left[\begin{matrix} qa, q/a \\ q^4 \end{matrix}; q^4, q^2 z \right] = \sum_{k=0}^{\infty} \frac{(aq; q^2)_k (q/a; q^2)_k (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} z^k, \quad (3.1)$$

where the *basic hypergeometric series* is defined as

$${}_{r+1}\phi_r \left[\begin{matrix} a_1, a_2, \dots, a_{r+1} \\ b_1, b_2, \dots, b_r \end{matrix}; q, z \right] = \sum_{k=0}^{\infty} \frac{(a_1; q)_k (a_2; q)_k \cdots (a_{r+1}; q)_k z^k}{(q; q)_k (b_1; q)_k (b_2; q)_k \cdots (b_r; q)_k}.$$

Clearly, if one takes $a = 1$ and $z = q^2$ in (3.1), then the first series on the left is q -Gauss-summable,

$${}_2\phi_1 \left[\begin{matrix} q, q \\ q^4 \end{matrix}; q^4, q^2 \right] = \frac{(q^3, q^3; q^4)_{\infty}}{(q^2, q^4; q^4)_{\infty}},$$

so that this leads to a ‘natural’ reduction of the series on the right and makes no surprise from divisibility of the result by some cyclotomic polynomials

$$\Phi_n(q) = \prod_{\substack{j=1 \\ \gcd(j,n)=1}}^n (q - e^{2\pi i j/n}).$$

Theorem 1. *For any positive integer n , we have*

$$\sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} q^{2k} \equiv \begin{cases} \frac{(q^3; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} \pmod{\Phi_n(q)} & \text{if } n \equiv 1 \pmod{4}, \\ 0 \pmod{\Phi_n(q)} & \text{if } n \equiv 3 \pmod{4}. \end{cases} \quad (3.2)$$

Proof. From the proof of [18, Theorem 4.1] we know that

$$(q; q^4)_k^2 \equiv (q^{1-3n}; q^4)_k (q^{1+3n}; q^4)_k \pmod{\Phi_n(q)^2}.$$

Therefore, for $n \equiv 3 \pmod{4}$ we have, modulo $\Phi_n(q)^2$,

$$\sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} q^{2k} \equiv \sum_{k=0}^{(3n-1)/4} \frac{(q^{1-3n}; q^4)_k (q^{1+3n}; q^4)_k}{(q^4; q^4)_k^2} q^{2k} = \frac{(q^{3-3n}; q^4)_{(3n-1)/4}}{(q^4; q^4)_{(3n-1)/4}}$$

by the q -Chu–Vandermonde summation formula [11, Appendix (II.7)]:

$${}_2\phi_1 \left[\begin{matrix} a, q^{-n} \\ c \end{matrix}; q, \frac{cq^n}{a} \right] = \frac{(c/a; q)_n}{(c; q)_n}. \quad (3.3)$$

The proof then follows from the fact that $(q^{3-3n}; q^4)_{(3n-1)/4}$ contains the factor $1 - q^{-2n}$.

Similarly, for $n \equiv 1 \pmod{4}$, the result follows from

$$(q; q^4)_k^2 \equiv (q^{1-n}; q^4)_k (q^{1+n}; q^4)_k \pmod{\Phi_n(q)^2}. \quad \square$$

Making the substitution $q \rightarrow q^{-1}$ in (3.2), we obtain

$$\sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} q^{4k} \equiv \begin{cases} q^{(n^2-1)/4} \frac{(q^3; q^4)_{(n-1)/4}}{(q^4; q^4)_{(n-1)/4}} \pmod{\Phi_n(q)} & \text{if } n \equiv 1 \pmod{4}, \\ 0 \pmod{\Phi_n(q)} & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Remarkably, the truncations

$$\sum_{k=0}^{n-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k}$$

of the series on the right-hand side in (3.1) when $a = 1$ and $z = q^2$ are divisible by more cyclotomic polynomials, always squared, than the corresponding sums in (3.2) for all $n \geq 2$; the congruence

$$\sum_{k=0}^{n-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} \equiv 0 \pmod{\Phi_n(q)^2}$$

for $n \equiv 3 \pmod{4}$, not necessarily prime, discussed in [19, Corollary 1.2] is a particular instance of this high divisibility. We numerically observe that

$$\sum_{k=0}^{n-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} \equiv \begin{cases} q^{(n-1)/2} \frac{(q^2; q^2)_{(n-1)/2}^2}{(q^4; q^4)_{(n-1)/4}^2} & \text{if } n \equiv 1 \pmod{4}, \\ 0 & \text{if } n \equiv 3 \pmod{4}, \end{cases}$$

is true not just modulo $\Phi_n(q)^2$ but also modulo $\prod_{\ell \equiv 3 \pmod{4}, \ell | n} \Phi_\ell(q)^2$ for $n \equiv 1 \pmod{4}$ and even modulo $\prod_{\ell \equiv 3 \pmod{4}, \ell < n} \Phi_\ell(q)^2$ if $n \equiv 3 \pmod{4}$.

Theorem 2. *Modulo $\Phi_n(q)^2$,*

$$\sum_{k=0}^{n-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} \equiv \begin{cases} q^{(n-1)/2} \frac{(q^2; q^4)_{(n-1)/4}^2}{(q^4; q^4)_{(n-1)/4}^2} & \text{if } n \equiv 1 \pmod{4}, \\ 0 & \text{if } n \equiv 3 \pmod{4}. \end{cases} \quad (3.4)$$

Proof. Although the method of proving [19, Corollary 1.2] can also be used to establish (3.4), here we give a somewhat different argument. We shall demonstrate a parametric generalisation of (3.4), namely, that

$$\sum_{k=0}^{n-1} \frac{(aq; q^2)_k (q/a; q^2)_k (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} \equiv \begin{cases} q^{(n-1)/2} \frac{(q^2; q^4)_{(n-1)/4}^2}{(q^4; q^4)_{(n-1)/4}^2} & \text{if } n \equiv 1 \pmod{4}, \\ 0 & \text{if } n \equiv 3 \pmod{4}, \end{cases} \quad (3.5)$$

holds true modulo $(1 - aq^n)(a - q^n)$. For $a = q^{-n}$ or $a = q^n$, the left-hand side of (3.5) is equal to

$$\sum_{k=0}^{(n-1)/2} \frac{(q^{1-n}; q^2)_k (q^{1+n}; q^2)_k (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} = {}_4\phi_3 \left[\begin{matrix} q^{1-n}, q^{n+1}, q, -q \\ q^2, -q^2, q^2 \end{matrix}; q^2, q^2 \right]. \quad (3.6)$$

By Andrews' terminating q -analogue of Watson's formula (see [2] or [11, Appendix (II.17)]),

$${}_4\phi_3 \left[\begin{matrix} q^{-m}, a^2 q^{m+1}, b, -b \\ aq, -aq, b^2 \end{matrix}; q, q \right] = \begin{cases} \frac{b^m (q, a^2 q^2 / b^2; q^2)_{m/2}}{(a^2 q^2, b^2 q; q^2)_{m/2}} & \text{if } m \text{ is even,} \\ 0 & \text{if } m \text{ is odd,} \end{cases}$$

we conclude that the right-hand side of (3.6) is just that of (3.5). Finally, letting $a \rightarrow 1$ in (3.5), we are led to (3.4). \square

Remark. The $n \equiv 3 \pmod{4}$ case can also be deduced from [15, Theorem 1.1]. For $n \equiv 1 \pmod{4}$, we can also prove (3.4) without using Andrews' formula as follows. By (3.1) and (3.3), we have, modulo $\Phi_n(q)^2$,

$$\begin{aligned} \sum_{k=0}^{n-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} &\equiv \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^2)_k (q^{1+n}; q^2)_k (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} \\ &= \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^4)_k (q^{1+n}; q^4)_k}{(q^4; q^4)_k^2} q^{2k} \\ &\quad \times \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^4)_k (q^{1+n}; q^4)_k}{(q^4; q^4)_k^2} q^{4k} \\ &= q^{(n^2-1)/4} \frac{(q^{3-n}; q^4)_{(n-1)/4}^2}{(q^4; q^4)_{(n-1)/4}^2}, \end{aligned}$$

which is in fact the same as (3.4).

The consideration above corresponds to a q -deformation of (2.3); our q -analogue of (2.8) is somewhat similar but weaker.

We first prove the following result.

Theorem 3. For any positive odd integer n , we have, modulo $\Phi_n(q)$,

$$\sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^{3k} \equiv \begin{cases} \frac{(q^5, q^7; q^8)_{(n-1)/8}}{(q^4; q^4)_{(n-1)/4}} & \text{if } n \equiv 1 \pmod{8}, \\ \frac{(q^5, q^7; q^8)_{(3n-1)/8}}{(q^4; q^4)_{(3n-1)/4}} & \text{if } n \equiv 3 \pmod{8}, \\ 0, & \text{if } n \equiv 5, 7 \pmod{8}. \end{cases} \quad (3.7)$$

Proof. We use the q -Kummer (Bailey–Daum) summation formula [11, Appendix (II.9)]:

$${}_2\phi_1 \left[\begin{matrix} a, b \\ aq/b \end{matrix}; q, \frac{-q}{b} \right] = \frac{(-q; q)_\infty (aq; q^2)_\infty (aq^2/b^2; q^2)_\infty}{(-q/b; q)_\infty (aq/b; q)_\infty}. \quad (3.8)$$

For $n \equiv 1 \pmod{8}$, by (3.8) we obtain

$$\begin{aligned}
{}_2\phi_1 \left[\begin{matrix} q^{1-n}, q \\ q^{4-n} \end{matrix}; q^4, -q^3 \right] &= \frac{(-q^4; q^4)_\infty (q^{5-n}; q^8)_\infty (q^{7-n}; q^8)_\infty}{(-q^3; q^4)_\infty (q^{4-n}; q^4)_\infty} \\
&= \frac{(q^{5-n}; q^8)_{(n-1)/8} (q^{7-n}; q^8)_{(n-1)/8}}{(q^{4-n}; q^4)_{(n-1)/4}}.
\end{aligned} \tag{3.9}$$

Similarly, for $n \equiv 3 \pmod{8}$, we have

$${}_2\phi_1 \left[\begin{matrix} q^{1-3n}, q \\ q^{4-3n} \end{matrix}; q^4, -q^3 \right] = \frac{(q^{5-3n}; q^8)_{(3n-1)/8} (q^{7-3n}; q^8)_{(3n-1)/8}}{(q^{4-3n}; q^4)_{(3n-1)/4}}. \tag{3.10}$$

Furthermore,

$$\sum_{k=0}^{n-1} \frac{(q^{1-n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-n}; q^4)_k} (-q)^{3k} = 0 \quad \text{for } n \equiv 5 \pmod{8}, \tag{3.11}$$

$$\sum_{k=0}^{n-1} \frac{(q^{1-3n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-3n}; q^4)_k} (-q)^{3k} = 0 \quad \text{for } n \equiv 7 \pmod{8}. \tag{3.12}$$

Since $q^{3n} \equiv q^n \equiv 1 \pmod{\Phi_n(q)}$, we deduce the desired q -congruences (3.7) from (3.9)–(3.12). \square

We complement Theorem 3 with the following related result.

Theorem 4. *For any positive odd integer n , we have, modulo $\Phi_n(q)$,*

$$\begin{aligned}
&\sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^k \\
&\equiv \begin{cases} \frac{(q, q^2, -q^3, -q^4; q^4)_{(n-1)/8}}{(q^4; q^4)_{(n-1)/4}} q^{(1-n^2)/8} & \text{if } n \equiv 1 \pmod{8}, \\ \frac{(q, q^2, -q^3, -q^4; q^4)_{(3n-1)/8}}{(q^4; q^4)_{(3n-1)/4}} q^{(1-n^2)/8} & \text{if } n \equiv 3 \pmod{8}, \\ -\frac{(1-q^2)(-q^3, -q^4, q^5, q^6; q^4)_{(n-5)/8}}{(q^4; q^4)_{(n-5)/4}} q^{(9-n^2)/8} & \text{if } n \equiv 5 \pmod{8}, \\ -\frac{(1-q^2)(-q^3, -q^4, q^5, q^6; q^4)_{(3n-5)/8}}{(q^4; q^4)_{(3n-5)/4}} q^{(9-n^2)/8} & \text{if } n \equiv 7 \pmod{8}. \end{cases}
\end{aligned}$$

Proof. Letting $q \rightarrow q^4$, $b = q$ and $c = -q^{2-4N}$ in Jackson's terminating q -analogue of Dixon's sum (see [11, Appendix (II.15)]),

$${}_3\phi_2 \left[\begin{matrix} q^{-2N}, b, c \\ q^{1-2N}/b, q^{1-2N}/c \end{matrix}; q, \frac{q^{2-N}}{bc} \right] = \frac{(b, c; q)_N (q, bc; q)_{2N}}{(q, bc; q)_N (b, c; q)_{2N}}, \tag{3.13}$$

we obtain

$${}_2\phi_1 \left[\begin{matrix} q^{-8N}, q \\ q^{3-8N} \end{matrix}; q^4, -q^5 \right] = \frac{(q, -q^{2-4N}; q^4)_N (q^4, -q^{3-4N}; q^4)_{2N}}{(q^4, -q^{3-4N}; q^4)_N (q, -q^{2-4N}; q)_{2N}}.$$

Taking $q \rightarrow q^{-1}$ in the above identity, we are led to

$${}_2\phi_1 \left[\begin{matrix} q^{-8N}, q \\ q^{3-8N} \end{matrix}; q^4, -q \right] = \frac{(q, -q^{2-4N}; q^4)_N (q^4, -q^{3-4N}; q^4)_{2N} q^{-4N}}{(q^4, -q^{3-4N}; q^4)_N (q, -q^{2-4N}; q)_{2N}}.$$

It follows that, for $n \equiv 1 \pmod{8}$,

$$\begin{aligned} \sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^k &\equiv \sum_{k=0}^{(n-1)/4} \frac{(q^{1-n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-n}; q^4)_k} (-q)^k \\ &= \frac{(q, -q^{(5-n)/2}; q^4)_{(n-1)/8} (q^4, -q^{(7-n)/2}; q^4)_{(n-1)/4} q^{(1-n)/2}}{(q^4, -q^{(7-n)/2}; q^4)_{(n-1)/4} q^{(1-n)/8} (q, -q^{(5-n)/2}; q^4)_{(n-1)/4}} \\ &= \frac{(q, q^2, -q^3, -q^4; q^4)_{(n-1)/8}}{(q^{4-n}; q^4)_{(n-1)/4}} q^{(1-n^2)/8}. \end{aligned}$$

Since $q^n \equiv 1 \pmod{\Phi_n(q)}$, we complete the proof of the theorem for the first case. The other three cases follow in a similar way: modulo $\Phi_n(q)$,

$$\begin{aligned} \sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^k &\equiv \sum_{k=0}^{(3n-1)/4} \frac{(q^{1-3n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-3n}; q^4)_k} (-q)^k \quad \text{for } n \equiv 3 \pmod{8}, \\ \sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^k &\equiv \sum_{k=0}^{(n-1)/4} \frac{(q^{1-n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-n}; q^4)_k} (-q)^k \quad \text{for } n \equiv 5 \pmod{8}, \\ \sum_{k=0}^{n-1} \frac{(q; q^4)_k^2}{(q^4; q^4)_k^2} (-q)^k &\equiv \sum_{k=0}^{(3n-1)/4} \frac{(q^{1-3n}; q^4)_k (q; q^4)_k}{(q^4; q^4)_k (q^{4-3n}; q^4)_k} (-q)^k \quad \text{for } n \equiv 7 \pmod{8}, \end{aligned}$$

and for the last two instances, we use the ‘odd version’ of Jackson’s q -analogue of Dixon’s sum (which follows from [5, eq. (2.3)]),

$${}_3\phi_2 \left[\begin{matrix} q^{1-2N}, b, c \\ q^{2-2N}/b, q^{1-2N}/c \end{matrix} ; q, \frac{q^{2-N}}{bc} \right] = \frac{(b, c; q)_N (q, bc; q)_{2N-1}}{(q; q)_{N-1} (bc; q)_N (b; q)_{2N-1} (c; q)_{2N}},$$

instead of (3.13). \square

Theorem 5. For any positive integer $n \equiv 1 \pmod{4}$, modulo $\Phi_n(q)$, we have

$$\begin{aligned} &\sum_{k=0}^{(n-1)/2} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} (-q)^k \\ &\equiv \begin{cases} \frac{(q, q^2; q^4)_{(n-1)/8} (-q; q)_{(n-1)/2}}{(q^4; q^4)_{(n-1)/4}} q^{-(1-n)^2/4} & \text{if } n \equiv 1 \pmod{8}, \\ 0 & \text{if } n \equiv 5 \pmod{8}. \end{cases} \end{aligned}$$

Proof. For $n \equiv 1 \pmod{8}$, in Jackson’s q -Clausen identity (3.1) we take $a = q^{-n}$ and $z = -q$ to obtain

$$\begin{aligned} &\sum_{k=0}^{(n-1)/2} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} (-q)^k \\ &\equiv \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^2)_k (q^{1+n}; q^2)_k (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} (-q)^k \\ &= \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^4)_k (q^{1+n}; q^4)_k}{(q^4; q^4)_k^2} (-q)^k \end{aligned}$$

$$\times \sum_{k=0}^{n-1} \frac{(q^{1-n}; q^4)_k (q^{1+n}; q^4)_k}{(q^4; q^4)_k^2} (-q)^{3k} \pmod{\Phi_n(q)^2}. \quad (3.14)$$

By (3.7), we see that the second sum is congruent to

$$\begin{aligned} \frac{(q^5, q^7; q^8)_{(n-1)/8}}{(q^4; q^4)_{(n-1)/4}} &\equiv \frac{(q^{5-n}, q^{7-n}; q^8)_{(n-1)/8}}{(q^4; q^4)_{(n-1)/4}} \\ &= \frac{(q^2, q^4; q^8)_{(n-1)/8} q^{-(n-1)(n-3)/8}}{(q^4; q^4)_{(n-1)/4}} \pmod{\Phi_n(q)}, \end{aligned}$$

while by Theorem 4 the first sum is congruent to

$${}_2\phi_1 \left[\begin{matrix} q^{1-n}, q \\ q^{4-n}, q^4, -q \end{matrix} \right] \equiv \frac{(q, q^2, -q^3, -q^4; q^4)_{(n-1)/8}}{(q^4; q^4)_{(n-1)/4}} q^{(1-n^2)/8} \pmod{\Phi_n(q)}.$$

This establishes the $n \equiv 1 \pmod{8}$ case of the theorem after some simplifications.

For $n \equiv 5 \pmod{8}$, we again have (3.14). Since $q^n \equiv 1 \pmod{\Phi_n(q)}$, we know that the second sum on the right-hand side of (3.14) is congruent to 0 modulo $\Phi_n(q)$ by Theorem 3, and so is the right-hand side of (3.14). This proves the theorem for $n \equiv 5 \pmod{8}$. \square

We leave the related cases when $n \equiv 3 \pmod{4}$ of Theorem 5 as an open problem to the reader.

Problem 1. For any positive integer $n \equiv 7 \pmod{8}$, show that

$$\sum_{k=0}^{(n-1)/2} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} (-q)^k \equiv 0 \pmod{\Phi_n(q)}.$$

Give a related q -congruence for $n \equiv 3 \pmod{8}$.

4. Conclusion and open questions

We have the following generalisation of Theorem 2 for $n \equiv 3 \pmod{4}$, which (partly) forms the grounds of the arithmetic observations preceding the statement of the theorem.

Conjecture 1. For $n \equiv 3 \pmod{4}$ and any positive integer r , we have

$$\begin{aligned} \sum_{k=0}^{rn-1} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} &\equiv 0 \pmod{\Phi_n(q)^2}, \\ \sum_{k=0}^{rn+(n-1)/2} \frac{(q; q^2)_k^2 (q^2; q^4)_k}{(q^2; q^2)_k^2 (q^4; q^4)_k} q^{2k} &\equiv 0 \pmod{\Phi_n(q)^2}. \end{aligned}$$

We also give a related generalisation of [21, Conjecture 4.13].

Conjecture 2. For $n \equiv 3 \pmod{4}$ and any positive integer r , we have

$$\sum_{k=0}^{rn-1} \frac{(1 + q^{4k+1})(q^2; q^4)_k^3}{(1 + q)(q^4; q^4)_k^3} q^k \equiv 0 \pmod{\Phi_n(q)^2},$$

$$\sum_{k=0}^{rn+(n-1)/2} \frac{(1+q^{4k+1})(q^2; q^4)_k^3}{(1+q)(q^4; q^4)_k^3} q^k \equiv 0 \pmod{\Phi_n(q)^2}.$$

Note that, although similar congruences with a parameter a modulo $(1-aq^n)(a-q^n)$ can be deduced, we cannot take the limit as $a \rightarrow 1$ to accomplish the proof of Conjectures 1 and 2 this time. Using the q -Lucas theorem, we can show that all the congruences in Conjectures 1 and 2 are true modulo $\Phi_n(q)$. Moreover, the following similar congruence in [21, Theorem 4.14],

$$\sum_{k=0}^{(n-1)/2} \frac{(q; q^2)_k^2}{(q^2; q^2)_k (q^4; q^4)_k} q^{2k} \equiv 0 \pmod{\Phi_n(q)^2} \quad \text{for } n \equiv 3 \pmod{4},$$

does not have such a generalisation. For this reason, we believe that Conjectures 1 and 2 are not easy to prove.

The q -extension (2.6) of Bauer's formula (1.1) is not unique. In [16,20] we mention a 'more classical' version

$$\sum_{k=0}^{\infty} \frac{1-q^{4k+1}}{1-q} \frac{(q; q^2)_k^3}{(q^2; q^2)_k^3} (-q)^{k^2} = \frac{(q; q^2)_{\infty} (q^3; q^2)_{\infty}}{(q; q^2)_{\infty}^2}. \quad (4.1)$$

In spite of (4.1) (in fact, its parametric modification) being suitable for proving the congruences (1.2) on the basis of our method from [21], dropping off the factor $(1-q^{4k+1})/(1-q)$ here does not lead to a 'suitable' q -analogue of (2.7). Namely, a numerical check suggests no congruences for the truncated sums of the resulting series.

Finally, we notice that the sequence $a_1(p)$ in (2.4) is ultimately linked with a remarkable classics, the two-square theorem due to Fermat and Gauss (see [1, Chap. 4], [8] and also the unpublished portion of Ramanujan's paper [27], reproduced in [4, Chap. 10] and relating the two-square generating function to the Dirichlet L -function in (1.3)). However, no reasonable q -analogues of this result have been recorded yet.

References

- [1] M. Aigner, G.M. Ziegler, *Proofs from the Book*, 6th ed., Springer, Berlin, 2018.
- [2] G.E. Andrews, On q -analogues of the Watson and Whipple summations, *SIAM J. Math. Anal.* 7 (1976) 332–336.
- [3] G.E. Andrews, B.C. Berndt, *Ramanujan's Lost Notebook, Part II*, Springer, New York, 2009.
- [4] G.E. Andrews, B.C. Berndt, *Ramanujan's Lost Notebook, Part III*, Springer, New York, 2012.
- [5] W.N. Bailey, On the analogue of Dixon's theorem for bilateral basic hypergeometric series, *Quart. J. Math. (Oxford) Ser. (2)* 1 (1950) 318–320.
- [6] N.D. Baruah, B.C. Berndt, H.H. Chan, Ramanujan's series for $1/\pi$: a survey, *Amer. Math. Monthly* 116 (7) (2009) 567–587.
- [7] G. Bauer, Von den Coefficienten der Reihen von Kugelfunctionen einer Variablen, *J. Reine Angew. Math.* 56 (1859) 101–121.
- [8] F.W. Clarke, W.N. Everitt, L.L. Littlejohn, S.J.R. Vorster, H.J.S. Smith and the Fermat two squares theorem, *Amer. Math. Monthly* 106 (7) (1999) 652–665.
- [9] S. Cooper, *Ramanujan's Theta Functions*, Springer, Cham, 2017.
- [10] S.B. Ekhad, D. Zeilberger, A WZ proof of Ramanujan's formula for π , in: J.M. Rassias (Ed.), *Geometry, Analysis, and Mechanics*, World Scientific, Singapore, 1994, pp. 107–108.
- [11] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, 2nd ed., *Encyclopedia Math. Appl.*, vol. 96, Cambridge Univ. Press, Cambridge, 2004.
- [12] V.J.W. Guo, A q -analogue of the (I.2) supercongruence of Van Hamme, *Int. J. Number Theory* 15 (1) (2019) 29–36.
- [13] V.J.W. Guo, Proof of a q -congruence conjectured by Tauraso, *Int. J. Number Theory* 15 (1) (2019) 37–41.
- [14] V.J.W. Guo, A q -analogue of a curious supercongruence of Guillera and Zudilin, *J. Difference Equ. Appl.* 25 (3) (2019) 342–350.
- [15] V.J.W. Guo, Some q -congruences with parameters, *Acta Arith.* (2019), in press, preprint: arXiv:1804.10963 [math.NT].
- [16] V.J.W. Guo, q -Analogues of three Ramanujan-type formulas for $1/\pi$, *Ramanujan J.* (2019), <https://doi.org/10.1007/s11139-018-0096-6>, in press.
- [17] V.J.W. Guo, M.J. Schlosser, Some new q -congruences for truncated basic hypergeometric series, *Symmetry* 11 (2) (2019), Art. 268.

- [18] V.J.W. Guo, M.J. Schlosser, Some q -supercongruences from transformation formulas for basic hypergeometric series, preprint, arXiv:1812.06324 [math.NT], December 2018, 39 pp.
- [19] V.J.W. Guo, J. Zeng, Some q -supercongruences for truncated basic hypergeometric series, *Acta Arith.* 171 (4) (2015) 309–326.
- [20] V.J.W. Guo, W. Zudilin, Ramanujan-type formulae for $1/\pi$: q -analogues, *Integral Transforms Spec. Funct.* 29 (7) (2018) 505–513.
- [21] V.J.W. Guo, W. Zudilin, A q -microscope for supercongruences, *Adv. Math.* 346 (2019) 329–358.
- [22] F.H. Jackson, Certain q -identities, *Quart. J. Math. (Oxford)* 12 (1941) 167–172.
- [23] P. Levrie, Using Fourier–Legendre expansions to derive series for $1/\pi$ and $1/\pi^2$, *Ramanujan J.* 22 (2) (2010) 221–230.
- [24] E. Mortenson, A p -adic supercongruence conjecture of Van Hamme, *Proc. Amer. Math. Soc.* 136 (12) (2008) 4321–4328.
- [25] K. Ono, Values of Gaussian hypergeometric series, *Trans. Amer. Math. Soc.* 350 (3) (1998) 1205–1223.
- [26] S. Ramanujan, Modular equations and approximations to π , *Quart. J. Math. (Oxford) Ser. (2)* 45 (1914) 350–372.
- [27] S. Ramanujan, Highly composite numbers, *Proc. Lond. Math. Soc. (2)* 14 (1915) 347–409.
- [28] M. Rogers, J.G. Wan, I.J. Zucker, Moments of elliptic integrals and critical L -values, *Ramanujan J.* 37 (1) (2015) 113–130.
- [29] M.J. Schlosser, q -Analogues of two product formulas of hypergeometric functions by Bailey, in: Z. Nashed, X. Li (Eds.), *Frontiers in Orthogonal Polynomials and q -Series*, World Scientific, Singapore, 2018, pp. 445–449.
- [30] Z.-H. Sun, Congruences concerning Legendre polynomials II, *J. Number Theory* 133 (2013) 1950–1976.
- [31] L. Van Hamme, Some conjectures concerning partial sums of generalized hypergeometric series, in: *p -Adic Functional Analysis*, Nijmegen, 1996, in: *Lect. Notes Pure Appl. Math.*, vol. 192, Dekker, New York, 1997, pp. 223–236.
- [32] W. Zudilin, Ramanujan-type formulae for $1/\pi$: a second wind?, in: N. Yui, et al. (Eds.), *Modular Forms and String Duality*, Banff, June 3–8, 2006, in: *Fields Inst. Commun. Ser.*, vol. 54, Amer. Math. Soc., Providence, RI, 2008, pp. 179–188.
- [33] W. Zudilin, Ramanujan-type supercongruences, *J. Number Theory* 129 (2009) 1848–1857.