



A q -Clausen–Orr type formula and its applications



Hong-Fang Guo^{a,b}, Victor J.W. Guo^{b,*}, Jiang Zeng^c

^a Department of Mathematics, East China Normal University, Shanghai 200241, People's Republic of China

^b School of Mathematical Sciences, Huaiyin Normal University, Huai'an, Jiangsu 223300, People's Republic of China

^c Univ Lyon, Université Claude Bernard Lyon 1, CNRS UMR 5208, Institut Camille Jordan, 43, blvd. du 11 novembre 1918, F-69622 Villeurbanne Cedex, France

ARTICLE INFO

Article history:

Received 13 August 2016

Available online 14 April 2017

Submitted by M.J. Schlosser

Keywords:

q -Delannoy numbers

q -analogue of Clausen's formula

q -binomial theorem

q -Chu–Vandermonde

q -Pfaff–Saalschütz

ABSTRACT

We show that certain terminating ${}_6\phi_5$ series can be factorized into a product of two ${}_3\phi_2$ series. As applications we prove a summation formula for a product of two q -Delannoy numbers along with some congruences for sums involving q -Delannoy numbers. This confirms three recent conjectures of the second author.

© 2017 Elsevier Inc. All rights reserved.

1. Introduction

Clausen's formula

$$\left\{ {}_2F_1 \left[\begin{matrix} a, b \\ a + b + 1/2 \end{matrix} ; z \right] \right\}^2 = {}_3F_2 \left[\begin{matrix} 2a, 2b, a + b \\ a + b + 1/2, 2a + 2b \end{matrix} ; z \right]$$

plays a central role in Ramanujan's derivation for various series for $1/\pi$. See [1,3] for some recent developments of this formula. More general formulas connecting products of two hypergeometric series as a single series were obtained by Orr in 1899 (see [15, p. 75]). This paper was motivated by a recent paper of the second author [7], where he proved some congruences of sums involving even powers of Delannoy numbers and raised some problems of finding the q -analogues. Recall that the Delannoy numbers $D(m, n)$ count lattice paths from $(0, 0)$ to (m, n) consisting of horizontal $(1, 0)$, vertical $(0, 1)$, and diagonal $(1, 1)$ steps, and have the following explicit formulas in terms of binomial coefficients:

* Corresponding author.

E-mail addresses: ghfkeji@126.com (H.-F. Guo), jwguo@hytc.edu.cn (V.J.W. Guo), zeng@math.univ-lyon1.fr (J. Zeng).

$$D(m, n) = \sum_{k=0}^n \binom{n}{k} \binom{n+m-k}{n} = \sum_{k=0}^n \binom{n}{k} \binom{m}{k} 2^k. \quad (1.1)$$

The reader is referred to Dziemiańczuk [4] and the references therein for how to generalize Delannoy numbers via counting weighted lattice paths. Recall that the *basic hypergeometric series* ${}_r\phi_s$ is defined as

$${}_r\phi_s \left[\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, \dots, b_s \end{matrix}; q, z \right] = \sum_{k=0}^{\infty} \frac{(a_1; q)_k (a_2; q)_k \cdots (a_r; q)_k}{(q; q)_k (b_1; q)_k \cdots (b_s; q)_k} \left((-1)^k q^{\binom{k}{2}} \right)^{1+s-r} z^k,$$

where $(a; q)_n = (1-a)(1-aq) \cdots (1-aq^{n-1})$ for $n = 1, 2, \dots$, and $(a; q)_0 = 1$. Aiming to answer the q -problems in [7], we are led to prove the following q -Clausen–Orr type formula.

Theorem 1.1. *Let n be a non-negative integer. Then*

$${}_3\phi_2 \left[\begin{matrix} q^{-n}, a, x \\ c, 0 \end{matrix}; q, q \right] {}_3\phi_2 \left[\begin{matrix} q^{-n}, a, c/x \\ c, 0 \end{matrix}; q, q \right] = a^n {}_6\phi_5 \left[\begin{matrix} q^{-n}, cq^n, a, c/a, x, c/x \\ c, \sqrt{c}, -\sqrt{c}, \sqrt{cq}, -\sqrt{cq} \end{matrix}; q, q \right]. \quad (1.2)$$

It is interesting to compare (1.2) with Jackson's q -analogue of Clausen's formula [9,10]:

$${}_2\phi_1 \left[\begin{matrix} a, b \\ abq^{\frac{1}{2}} \end{matrix}; q, z \right] {}_2\phi_1 \left[\begin{matrix} a, b \\ abq^{\frac{1}{2}} \end{matrix}; q, zq^{\frac{1}{2}} \right] = {}_4\phi_3 \left[\begin{matrix} a, b, a^{\frac{1}{2}}b^{\frac{1}{2}}, -a^{\frac{1}{2}}b^{\frac{1}{2}} \\ ab, a^{\frac{1}{2}}b^{\frac{1}{2}}q^{\frac{1}{4}}, -a^{\frac{1}{2}}b^{\frac{1}{2}}q^{\frac{1}{4}} \end{matrix}; q^{\frac{1}{2}}, z \right], \quad |z| < 1,$$

and the following q -analogue of Clausen's formula [5, Appendix (III.22)]:

$$\left\{ {}_4\phi_3 \left[\begin{matrix} a, b, abz, ab/z \\ abq^{\frac{1}{2}}, -abq^{\frac{1}{2}}, -ab \end{matrix}; q, q \right] \right\}^2 = {}_5\phi_4 \left[\begin{matrix} a^2, b^2, ab, abz, ab/z \\ abq^{\frac{1}{2}}, -abq^{\frac{1}{2}}, -ab, a^2b^2 \end{matrix}; q, q \right], \quad (1.3)$$

where both series are supposed to be terminated. Indeed, letting $c = x^2$, the identity (1.2) reduces to the following formula, which seems to be new.

Corollary 1.2. *Let n be a non-negative integer. Then*

$$\left\{ {}_3\phi_2 \left[\begin{matrix} q^{-n}, a, x \\ x^2, 0 \end{matrix}; q, q \right] \right\}^2 = a^n {}_5\phi_4 \left[\begin{matrix} q^{-n}, x^2q^n, a, x^2/a, x \\ x^2, -x, xq^{\frac{1}{2}}, -xq^{\frac{1}{2}} \end{matrix}; q, q \right]. \quad (1.4)$$

In particular, the right-hand side of (1.4) is non-negative for real a, x , and q . Furthermore, if n is even, then by (1.3), the right-hand side of (1.4) may be written as

$$a^n \left\{ {}_4\phi_3 \left[\begin{matrix} q^{-\frac{n}{2}}, xq^{\frac{n}{2}}, a, x^2/a \\ xq^{\frac{1}{2}}, -xq^{\frac{1}{2}}, -x \end{matrix}; q, q \right] \right\}^2.$$

Writing $n = 2m$ in (1.4) and taking the square root we obtain an identity between two polynomials in a of degree $2m$, where the sign is determined by comparing the coefficients of a^{2m} . We record the resulting formula as the second corollary.

Corollary 1.3. *Let m be a non-negative integer. Then*

$${}_3\phi_2 \left[\begin{matrix} q^{-2m}, a, x \\ x^2, 0 \end{matrix}; q, q \right] = a^m {}_4\phi_3 \left[\begin{matrix} q^{-m}, xq^m, a, x^2/a \\ xq^{\frac{1}{2}}, -xq^{\frac{1}{2}}, -x \end{matrix}; q, q \right].$$

For some other q -Clausen type formulas, the reader is referred to Gasper and Rahman [5, Exercise 8.17] and Schlosser [13]. On the other hand, in their study of some q -supercongruences for certain truncated basic hypergeometric series related to [16,17], Guo and Zeng [8] stumbled on the following q -Clausen–Orr type formula:

$$\begin{aligned} & \left(\sum_{k=s}^n \frac{(q^{-2n}; q^2)_k (x; q)_k q^k}{(q; q)_{k-s} (q; q)_{k+s}} \right) \left(\sum_{k=s}^n \frac{(q^{-2n}; q^2)_k (q/x; q)_k q^k}{(q; q)_{k-s} (q; q)_{k+s}} \right) \\ &= \frac{(-1)^n (q^2; q^2)_n^2 q^{-n^2}}{(q^2; q^2)_{n-s} (q^2; q^2)_{n+s}} \sum_{k=s}^n \frac{(-1)^k (q^2; q^2)_{n+k} (x; q)_k (q/x; q)_k q^{k^2-2nk}}{(q^2; q^2)_{n-k} (q; q)_{k-s} (q; q)_{k+s} (q; q)_{2k}}. \end{aligned} \quad (1.5)$$

Noticing that $(c; q)_{2k} = (\sqrt{c}; q)_k (-\sqrt{c}; q)_k (\sqrt{cq}; q)_k (-\sqrt{cq}; q)_k$, we can rewrite (1.2) as

$$\begin{aligned} & \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (a; q)_k (x; q)_k q^k}{(q; q)_k (c; q)_k} \right) \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (a; q)_k (c/x; q)_k q^k}{(q; q)_k (c; q)_k} \right) \\ &= a^n \sum_{k=0}^n \frac{(q^{-n}; q)_k (cq^n; q)_k (a; q)_k (c/a; q)_k (x; q)_k (c/x; q)_k q^k}{(q; q)_k (c; q)_k (c; q)_{2k}}. \end{aligned} \quad (1.6)$$

Clearly (1.6) is an extension of (1.5). In the same vein we shall establish the following result.

Theorem 1.4. *Let n be a non-negative integer. Then*

$$\begin{aligned} & \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (x; q^2)_k q^k}{(q; q)_k (c; q)_k} \right) \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (x; q^2)_k c^k q^{n k - \binom{k}{2}}}{(q; q)_k (c; q)_k x^k} \right) \\ &= \sum_{k=0}^n \frac{(q^{-n}; q)_k (cq^n; q)_k (x; q^2)_k (c^2/x; q^2)_k q^k}{(q; q)_k (c; q)_k (c; q)_{2k}}. \end{aligned} \quad (1.7)$$

In this paper we shall consider two q -analogues of $D(m, n)$. We first recall some standard q -notation (see [5]). The q -binomial coefficients are given by

$$\begin{bmatrix} n \\ k \end{bmatrix} := \begin{cases} \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}} & \text{if } n \geq k \geq 0, \\ 0 & \text{otherwise.} \end{cases}$$

The following two natural q -analogues of Delannoy numbers were introduced in [4, p. 30] and [12]:

$$D_q(m, n) := \sum_{k=0}^n q^{\binom{k}{2}} \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} n+m-k \\ n \end{bmatrix}, \quad (1.8)$$

$$D_q^*(m, n) := \sum_{k=0}^n q^{\binom{k+1}{2}} \begin{bmatrix} n \\ k \end{bmatrix} \begin{bmatrix} n+m-k \\ n \end{bmatrix}. \quad (1.9)$$

Note that $D_q^*(m, n) := q^{mn} D_{q^{-1}}(m, n)$. We first show that both of the polynomials $D_q(m, n)$ and $D_q^*(m, n)$ have a q -analogue of the second expression in (1.1) and provide a q -analogue of [7, (3.1)], which was asked in [7, Problem 5.2].

Theorem 1.5. *Let m and n be non-negative integers. Then*

$$D_q(m, n) = \sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} (-1; q)_k, \quad (1.10)$$

$$D_q^*(m, n) = \sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} (-q; q)_k. \quad (1.11)$$

Moreover,

$$D_q(m, n) D_q^*(m, n) = \sum_{k=0}^n q^{(m-k)(n-k)} \begin{bmatrix} n+k \\ 2k \end{bmatrix} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} m+k \\ k \end{bmatrix} (-1; q)_k (-q; q)_k. \quad (1.12)$$

Applying the formula (1.12), we shall prove the following two results originally conjectured by the second author [7, Conjectures 5.3 and 5.4].

Theorem 1.6. *Let p be an odd prime and m a positive integer. Then*

$$\begin{aligned} & \sum_{k=0}^{p-1} \frac{1 - q^{2k+1}}{1 - q} D_q(m, k) D_{q^{-1}}(m, k) q^{-k} \\ & \equiv \begin{cases} \frac{1 - q^{-2m}}{1 - q^2} q \pmod{[p]^2} & \text{if } m \equiv 0 \pmod{p}, \\ \frac{1 - q^{2m+2}}{1 - q^2} q \pmod{[p]^2} & \text{if } m \equiv -1 \pmod{p}, \\ 0 \pmod{[p]^2} & \text{otherwise,} \end{cases} \end{aligned} \quad (1.13)$$

where $[p] = 1 + q + \cdots + q^{p-1}$, and the congruences are understood in the polynomial ring $\mathbb{Z}[q]$.

Theorem 1.7. *Let m , n , and r be positive integers. Then all of*

$$\sum_{k=0}^{n-1} \frac{(1 - q^m)(1 - q^{m+1})(1 - q^{2k+1})}{(1 - q^2)(1 - q^n)^2} D_q(m, k) D_{q^{-1}}(m, k) q^{-k}, \quad (1.14)$$

$$\sum_{k=0}^{n-1} \frac{1 - q^{2k+1}}{1 - q^n} D_q(m, k)^r D_{q^{-1}}(m, k)^r q^{-k}, \quad (1.15)$$

$$\sum_{k=0}^{n-1} (-1)^{n-k-1} \frac{1 - q^{2k+1}}{1 - q^n} D_q(m, k)^r D_{q^{-1}}(m, k)^r q^{\binom{k}{2}} \quad (1.16)$$

are Laurent polynomials in q with non-negative integer coefficients.

Note that Theorem 1.7 is a q -analogue of [7, Theorem 1.1] for the first three polynomials. The rest of the paper is organized as follows. We shall give three lemmas in Section 2 and prove Theorem 1.1 in Section 3. In Sections 4 and 5 we prove Theorems 1.4 and 1.5. In Sections 6 and 7, by using Theorem 1.5 we give proofs of Theorems 1.6 and 1.7, respectively.

2. Three lemmas

The following three lemmas are crucial ingredients of our proof of [Theorem 1.1](#).

Lemma 2.1. *Let n and h be positive integers and let m be a non-negative integer with $h \leq n - m$. Then*

$$\begin{aligned} & \sum_{j=0}^m \sum_{k=0}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_k (x; q)_j (x; q)_k (q^{j-m-h+1}; q)_{h-1} (q^{k-m-h+1}; q)_{h-1} (1 - q^{k-j}) q^{2j+k}}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k} \\ &= \frac{(q; q)_n (q; q)_{h-1} (x; q)_{m+h} (c/x; q)_{n-h} x^{n-h} q^{\frac{m^2+3m}{2} - mn - mh - h^2 + h}}{(-1)^{m-1} (q; q)_m (c; q)_m (c; q)_n (q; q)_{n-m-h}}. \end{aligned} \quad (2.1)$$

Proof. Note that both sides of (2.1) are polynomial in x of degree $m + n$ with the same leading coefficient. Therefore, to prove (2.1), it suffices to prove that both sides have the same roots as polynomials in x . Denote the left-hand side of (2.1) by $L_{m,n}(x)$. We first assert that

$$\begin{aligned} L_{m,n}(x) &= \sum_{j=0}^m \sum_{k=m+1}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_k (x; q)_j (x; q)_k}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k} \\ &\quad \times (q^{j-m-h+1}; q)_{h-1} (q^{k-m-h+1}; q)_{h-1} (1 - q^{k-j}) q^{2j+k}. \end{aligned} \quad (2.2)$$

In fact, since $(1 - q^{j-k}) q^{2k+j} = -(1 - q^{k-j}) q^{k+2j}$, we have $\sum_{j=0}^m \sum_{k=0}^m = 0$ for the summands in $L_{m,n}(x)$. We now consider the following two cases.

- For $x = q^{-r}$ with $0 \leq r \leq m + h - 1$, we have

$$\begin{aligned} L_{m,n}(q^{-r}) &= \sum_{j=0}^m \sum_{k=0}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_k (q^{-r}; q)_j (q^{-r}; q)_k}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k} \\ &\quad \times (q^{j-m-h+1}; q)_{h-1} (q^{k-m-h+1}; q)_{h-1} (1 - q^{k-j}) q^{2j+k}. \end{aligned}$$

If $r \leq m$, then $L_{m,n}(q^{-r}) = 0$ by the antisymmetry of j and k in $L_{m,n}(q^{-r})$. If $r \geq m + 1$, then $h \geq r - m + 1$, i.e., $r - m - h + 1 \leq 0$, and so $(q^{k-m-h+1}; q)_{h-1} = 0$ for $m + 1 \leq k \leq r$. Hence, by (2.2), we again get $L_{m,n}(q^{-r}) = 0$.

- For $x = cq^r$ with $0 \leq r \leq n - h - 1$, we shall prove that

$$\sum_{k=0}^n \frac{(q^{-n}; q)_k (cq^r; q)_k (q^{k-m-h+1}; q)_{h-1} (1 - q^{k-j}) q^{2j+k}}{(q; q)_k (c; q)_k} = 0. \quad (2.3)$$

In fact, we can rewrite the left-hand side of (2.3) as

$$\sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{-nk + \binom{k}{2}} R_k, \quad (2.4)$$

where

$$R_k = \frac{(cq^r; q)_k (q^{k-m-h+1}; q)_{h-1} (1 - q^{k-j}) q^{2j+k}}{(c; q)_k}.$$

Since

$$\frac{(cq^r; q)_k}{(c; q)_k} = \frac{(cq^k; q)_r}{(c; q)_r},$$

we see that R_k is a polynomial in q^k of degree $r + h - 1 + 2 \leq n$ with no constant term. By the q -binomial theorem (see, for example, [2, Theorem 3.3])

$$\sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{k}{2}} x^k = (x; q)_n, \quad (2.5)$$

we have

$$\sum_{k=0}^n (-1)^k \begin{bmatrix} n \\ k \end{bmatrix} q^{\binom{k}{2}} q^{-ik} = 0 \quad \text{for } 0 \leq i \leq n-1.$$

It follows that the expression (2.4) is equal to 0. Namely, the identity (2.3) holds.

Hence, we see that all the $m + n$ roots of $L_{m,n}(x)$ are the same as those of the right-hand side of (2.1). \square

Lemma 2.2. *Let n and h be positive integers and let m be a non-negative integer with $h \leq n - m$. Then*

$$\begin{aligned} & \sum_{j=0}^m \sum_{k=m+h}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_k (a; q)_j (a; q)_k (1 - q^{k-j}) q^{j+k+jh} \begin{bmatrix} k-m-1 \\ h-1 \end{bmatrix} \begin{bmatrix} m+h-j-1 \\ h-1 \end{bmatrix}}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k} \\ &= \frac{(q; q)_n (a; q)_{m+h} (c/a; q)_{n-h} a^{n-h} q^{\frac{m^2+m-h^2+h}{2} - mn}}{(-1)^{m-h} (q; q)_m (c; q)_m (c; q)_n (q; q)_{h-1} (q; q)_{n-m-h}}. \end{aligned} \quad (2.6)$$

Proof. It is easy to see that $\begin{bmatrix} k-m-1 \\ h-1 \end{bmatrix} = 0$ for $m+1 \leq k < m+h$. Therefore, the left-hand side of (2.6) remains unchanged when we replace $\sum_{k=m+h}^n$ by $\sum_{k=m+1}^n$. Moreover,

$$\begin{bmatrix} k-m-1 \\ h-1 \end{bmatrix} \begin{bmatrix} m+h-j-1 \\ h-1 \end{bmatrix} = \frac{(q^{j-m-h+1}; q)_{h-1} (q^{k-m-h+1}; q)_{h-1} q^{(m-j)(h-1) + \binom{h}{2}}}{(-1)^{h-1} (q; q)_{h-1}^2}.$$

The proof then follows from (2.1) and (2.2) with $x = a$. \square

The following result has been proved in [8, (3.5)].

Lemma 2.3 ([8]). *Let n be a positive integer. Then*

$$(x; q)_n + (a/x; q)_n = (x; q)_n (a/x; q)_n + (a; q)_n + \sum_{k=1}^{n-1} (x; q)_k (a/x; q)_k B_{n,k}(a), \quad (2.7)$$

where

$$B_{n,k}(a) := (1 - q^n) \sum_{h=1}^{n-k} (-1)^h \begin{bmatrix} n-k-1 \\ h-1 \end{bmatrix} \begin{bmatrix} k+h-1 \\ h-1 \end{bmatrix} \frac{q^{\binom{h}{2} + kh} a^h}{1 - q^h}.$$

3. Proof of Theorem 1.1

Recall that (1.2) can be rewritten as (1.6). As

$$\left(\sum_{k=0}^n a_k\right) \left(\sum_{j=0}^n b_j\right) = \sum_{k=0}^n a_k b_k + \sum_{0 \leq j < k \leq n} (a_k b_j + a_j b_k),$$

the left-hand side of (1.6) is equal to

$$\begin{aligned} & \sum_{k=0}^n \frac{(q^{-n}; q)_k^2 (a; q)_k^2 q^{2k}}{(q; q)_k^2 (c; q)_k^2} (x; q)_k (c/x; q)_k \\ & + \sum_{0 \leq j < k \leq n} \frac{(q^{-n}; q)_j (q^{-n}; q)_k (a; q)_j (a; q)_k q^{j+k} ((x; q)_j (c/x; q)_k + (x; q)_k (c/x; q)_j)}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k}. \end{aligned} \quad (3.1)$$

For $0 \leq j < k$, from (2.7) we deduce that

$$\begin{aligned} & (x; q)_j (c/x; q)_k + (x; q)_k (c/x; q)_j \\ & = (x; q)_j (c/x; q)_j ((xq^j; q)_{k-j} + (cq^j/x; q)_{k-j}) \\ & = (x; q)_k (c/x; q)_k + (x; q)_j (c/x; q)_j (cq^{2j}; q)_{k-j} + \sum_{i=1}^{k-j-1} (x; q)_{j+i} (c/x; q)_{j+i} B_{k-j,i}(cq^{2j}) \\ & = (x; q)_k (c/x; q)_k + (x; q)_j (c/x; q)_j + \sum_{i=0}^{k-j-1} (x; q)_{j+i} (c/x; q)_{j+i} B_{k-j,i}(cq^{2j}), \end{aligned}$$

where we have used the q -binomial theorem (2.5) in the last step:

$$(cq^{2j}; q)_{k-j} = 1 + \sum_{h=1}^{k-j} (-1)^h \begin{bmatrix} k-j \\ h \end{bmatrix} q^{\binom{h}{2} + 2jh} c^h.$$

It follows that (3.1) can be written as $\sum_{m=0}^n \alpha_m (x; q)_m (c/x; q)_m$, where

$$\begin{aligned} \alpha_m &= \sum_{j=0}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_m (a; q)_j (a; q)_m q^{j+m}}{(q; q)_j (q; q)_m (c; q)_j (c; q)_m} \\ & + \sum_{j=0}^m \sum_{k=m+1}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_k (a; q)_j (a; q)_k q^{j+k}}{(q; q)_j (q; q)_k (c; q)_j (c; q)_k} B_{k-j,m-j}(cq^{2j}). \end{aligned} \quad (3.2)$$

By the q -Chu–Vandermonde summation formula [5, Appendix (II.6)]:

$${}_2\phi_1 \left[\begin{matrix} a, q^{-n} \\ c \end{matrix}; q, q \right] = \frac{(c/a; q)_n a^n}{(c; q)_n}, \quad (3.3)$$

we have

$$\sum_{j=0}^n \frac{(q^{-n}; q)_j (q^{-n}; q)_m (a; q)_j (a; q)_m q^{j+m}}{(q; q)_j (q; q)_m (c; q)_j (c; q)_m} = (-1)^m \frac{(q; q)_n (a; q)_m (c/a; q)_n q^{\frac{m^2+m}{2} - mn}}{(q; q)_m (c; q)_m (c; q)_n (q; q)_{n-m}} a^n. \quad (3.4)$$

Substituting (3.4) and (2.6) into (3.2), we obtain

$$\alpha_m = \frac{(-1)^m (q; q)_n q^{\frac{m^2+m}{2} - mn}}{(q; q)_m (c; q)_m (c; q)_n} \sum_{h=0}^{n-m} \frac{(a; q)_{m+h} (c/a; q)_{n-h} a^{n-h} q^{mh} c^h}{(q; q)_h (q; q)_{n-m-h}}. \quad (3.5)$$

The last sum can be simplified again by the q -Chu–Vandermonde formula (3.3) and is equal to

$$\frac{(a; q)_m (c/a; q)_m (cq^{2m}; q)_{n-m} a^n}{(q; q)_{n-m}}. \quad (3.6)$$

It follows from (3.5) and (3.6) that α_m is just the coefficient of $(x; q)_m (c/x; q)_m$ on the right-hand side of (1.6).

Remark. Letting $c = q^{2s+1}$, $a = aq^s$, $x = xq^s$, and replacing n by $n - s$ in (1.6) ($0 \leq s \leq n$), we get the following result:

$$\begin{aligned} & \left(\sum_{k=s}^n \frac{(q^{-n}; q)_k (a; q)_k (x; q)_k q^k}{(q; q)_{k-s} (q; q)_{k+s}} \right) \left(\sum_{k=s}^n \frac{(q^{-n}; q)_k (a; q)_k (q/x; q)_k q^k}{(q; q)_{k-s} (q; q)_{k+s}} \right) \\ &= \frac{(q^{-n}; q)_s (a; q)_s a^{n-s} q^{(n+1)s-s^2}}{(q^{n+1}; q)_s (q/a; q)_s} \sum_{k=s}^n \frac{(q^{-n}; q)_k (q^{n+1}; q)_k (a; q)_k (q/a; q)_k (x; q)_k (q/x; q)_k q^k}{(q; q)_{k-s} (q; q)_{k+s} (q; q)_{2k}}. \end{aligned} \quad (3.7)$$

It is clear that the $a = -q^{-n}$ case of (3.7) reduces to (1.5).

4. Proof of Theorem 1.4

We need a special case of Theorem 1.1. Letting $a = -x$ in (1.6), we are led to

$$\begin{aligned} & \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (x^2; q^2)_k q^k}{(q; q)_k (c; q)_k} \right) \left(\sum_{k=0}^n \frac{(q^{-n}; q)_k (-x; q)_k (c/x; q)_k q^k}{(q; q)_k (c; q)_k} \right) \\ &= (-x)^n \sum_{k=0}^n \frac{(q^{-n}; q)_k (cq^n; q)_k (x^2; q^2)_k (c^2/x^2; q^2)_k q^k}{(q; q)_k (c; q)_k (c; q)_{2k}}. \end{aligned} \quad (4.1)$$

We also need the following result.

Lemma 4.1. *Let n be a non-negative integer. Then*

$$\sum_{k=0}^n \frac{(q^{-n}; q)_k (x; q)_k (y; q)_k q^k}{(q; q)_k (c; q)_k} = \sum_{k=0}^n (-1)^k \frac{(q^{-n}; q)_k (x; q)_k (c/y; q)_k x^{n-k} y^k q^{n-k-\binom{k}{2}}}{(q; q)_k (c; q)_k}. \quad (4.2)$$

Proof. This follows from combining Jackson's two transformations of terminating ${}_2\phi_1$ series [5, Appendix (III.7) and (III.8)]:

$${}_3\phi_2 \left[\begin{matrix} q^{-n}, b, bq^{-n}/c \\ bq^{1-n}/c, 0 \end{matrix}; q, q \right] = b^n {}_3\phi_1 \left[\begin{matrix} q^{-n}, b, q/z \\ bq^{1-n}/c \end{matrix}; q, \frac{z}{c} \right]$$

with $b \rightarrow x$, $c \rightarrow xq^{1-n}/c$ and $z \rightarrow qy/c$. \square

Replacing x and y by $-x$ and c/x respectively in (4.2), we are led to

$$\sum_{k=0}^n \frac{(q^{-n}; q)_k (-x; q)_k (c/x; q)_k q^k}{(q; q)_k (c; q)_k} = (-x)^n \sum_{k=0}^n \frac{(q^{-n}; q)_k (x^2; q^2)_k c^k q^{n-k-\binom{k}{2}}}{(q; q)_k (c; q)_k x^{2k}}. \quad (4.3)$$

Combining (4.1) and (4.3) (also $x \rightarrow \sqrt{x}$), we obtain Theorem 1.4.

Remark. Letting $c = q^{2s+1}$, $x \rightarrow xq^{2s}$, and replacing n by $n - s$ ($0 \leq s \leq n$) in (1.7), we get the following identity

$$\begin{aligned} & \left(\sum_{k=s}^n \frac{(q^{-n}; q)_k (x; q^2)_k q^k}{(q; q)_{k-s} (q; q)_{k+s}} \right) \left(\sum_{k=s}^n \frac{(q^{-n}; q)_k (x; q^2)_k q^{(n+1)k - \binom{k}{2}}}{(q; q)_{k-s} (q; q)_{k+s} x^k} \right) \\ &= \frac{(-1)^s (q; q)_n^2 (x; q^2)_s q^s}{(q; q)_{n-s} (q; q)_{n+s} (q^2/x; q^2)_s x^s} \sum_{k=s}^n \frac{(q^{-n}; q)_k (q^{n+1}; q)_k (x; q^2)_k (q^2/x; q^2)_k q^k}{(q; q)_{k-s} (q; q)_{k+s} (q; q)_{2k}}, \end{aligned}$$

which was originally conjectured in a preliminary version (arXiv:1408.0512v1) of [8].

5. Proof of Theorem 1.5

If we replace n by $n - i$ in the q -Chu–Vandermonde summation formula (3.3) with $a = q^{-m+i}$ and $c = q^{i+1}$ then

$$\sum_{k=i}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n-i \\ k-i \end{bmatrix} = \begin{bmatrix} n+m-i \\ n \end{bmatrix}.$$

Hence, by the q -binomial theorem (2.5) we have

$$\begin{aligned} \sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} (x; q)_k &= \sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} \sum_{i=0}^k (-1)^i \begin{bmatrix} k \\ i \end{bmatrix} q^{\binom{i}{2}} x^i \\ &= \sum_{i=0}^m (-1)^i \begin{bmatrix} n \\ i \end{bmatrix} q^{\binom{i}{2}} x^i \sum_{k=i}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n-i \\ k-i \end{bmatrix} \\ &= \sum_{i=0}^m q^{\binom{i}{2}} \begin{bmatrix} n \\ i \end{bmatrix} \begin{bmatrix} n+m-i \\ n \end{bmatrix} (-x)^i. \end{aligned}$$

When $x = -1$ and $x = -q$, we obtain (1.10) and (1.11), respectively. Now, letting $c = q$ and $a = q^{-m}$ in (1.6), we get

$$\begin{aligned} & \left(\sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} (x; q)_k \right) \left(\sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} n \\ k \end{bmatrix} (q/x; q)_k \right) \\ &= \sum_{k=0}^m q^{(m-k)(n-k)} \begin{bmatrix} n+k \\ 2k \end{bmatrix} \begin{bmatrix} m \\ k \end{bmatrix} \begin{bmatrix} m+k \\ k \end{bmatrix} (x; q)_k (q/x; q)_k. \end{aligned}$$

This entails the identity (1.12) by taking $x = -1$ or $-q$.

6. Proof of Theorem 1.6

The following identity can be easily proved by induction.

$$\sum_{k=j}^{n-1} (1 - q^{2k+1}) \begin{bmatrix} k+j \\ 2j \end{bmatrix} q^{-(j+1)k} = \frac{(1 - q^n)(1 - q^{n-j})}{1 - q^{j+1}} \begin{bmatrix} n+j \\ 2j \end{bmatrix} q^{-(j+1)(n-1)}. \quad (6.1)$$

By (1.12) and (6.1), the left-hand side of (1.13) is equal to

$$\begin{aligned}
& \sum_{k=0}^{p-1} \frac{1-q^{2k+1}}{1-q} \sum_{j=0}^k q^{j^2-mj-(j+1)k} \begin{bmatrix} k+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} (-1; q)_j (-q; q)_j \\
&= \sum_{j=0}^{p-1} q^{j^2-mj-(j+1)(p-1)} \frac{(1-q^p)(1-q^{p-j})}{(1-q)(1-q^{j+1})} \begin{bmatrix} p+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} (-1; q)_j (-q; q)_j. \tag{6.2}
\end{aligned}$$

By [7, Theorem 2.1], we know that

$$\frac{(1-q^m)(1-q^{m+1})(1-q^{p-j})}{(1-q)(1-q^p)(1-q^{j+1})} \begin{bmatrix} p+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} = \frac{(1-q^{p-j})(1-q^{j+1})}{(1-q)(1-q^p)} \begin{bmatrix} p+j \\ 2j \end{bmatrix} \begin{bmatrix} m+1 \\ j+1 \end{bmatrix} \begin{bmatrix} m+j \\ j+1 \end{bmatrix}$$

is a polynomial in q with non-negative integer coefficients. Since $[p] = (1-q^p)/(1-q)$ is an irreducible polynomial in q for any prime p and $\gcd(q^m-1, q^n-1) = (q^{\gcd(m,n)}-1)$, we conclude that

$$\frac{1-q^{p-j}}{1-q^{j+1}} \begin{bmatrix} p+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} \equiv 0 \pmod{[p]} \quad \text{if } m \not\equiv 0, -1 \pmod{p},$$

and so the right-hand side of (6.2) is congruent to 0 modulo $[p]^2$ in this case.

On the other hand, if $m \equiv 0, -1 \pmod{p}$, then

$$\frac{1-q^{p-j}}{1-q^{j+1}} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} \equiv 0 \pmod{[p]} \quad \text{if } j = 0, 1, \dots, p-2.$$

Therefore, if $m \equiv 0 \pmod{p}$, then the right-hand side of (6.2) is congruent to

$$\begin{aligned}
q \begin{bmatrix} 2p-1 \\ 2p-2 \end{bmatrix} \begin{bmatrix} m \\ p-1 \end{bmatrix} \begin{bmatrix} m+p-1 \\ p-1 \end{bmatrix} (-1; q)_{p-1} (-q; q)_{p-1} &\equiv q \frac{1-q^{2p-1}}{1-q} \cdot \frac{1-q^m}{1-q^{p-1}} (-q^{-1}) \cdot \frac{2q}{1+q} \\
&\equiv -\frac{2q(1-q^m)}{1-q^2} \\
&\equiv \frac{1-q^{-2m}}{1-q^2} q \pmod{[p]^2},
\end{aligned}$$

where we have used the congruence $(-q; q)_{p-1} \equiv 1 \pmod{[p]}$ (see [6, (1.6)] or [11]); while if $m \equiv -1 \pmod{p}$, then the right-hand side of (6.2) is congruent to

$$\frac{1-q^{2p-1}}{1-q} \cdot \frac{1-q^{m+1}}{1-q^{p-1}} \cdot \frac{2q}{1+q} \equiv \frac{2q(1-q^{m+1})}{1-q^2} \equiv \frac{1-q^{2m+2}}{1-q^2} q \pmod{[p]^2}.$$

7. Proof of Theorem 1.7

Similarly to (6.2), the left-hand side of (1.14) is equal to

$$\sum_{j=0}^{n-1} q^{j^2-mj-(j+1)(n-1)} \frac{(1-q^m)(1-q^{m+1})(1-q^{n-j})}{(1-q^2)(1-q^n)(1-q^{j+1})} \begin{bmatrix} n+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} (-1; q)_j (-q; q)_j.$$

It is easy to see that

$$\frac{(1-q^m)(1-q^{m+1})(1-q^{n-j})}{(1-q^2)(1-q^n)(1-q^{j+1})} \begin{bmatrix} n+j \\ 2j \end{bmatrix} \begin{bmatrix} m \\ j \end{bmatrix} \begin{bmatrix} m+j \\ j \end{bmatrix} (-q; q)_j$$

$$= \begin{cases} \frac{(1-q^m)(1-q^{m+1})}{(1-q^2)(1-q)} & \text{if } j = 0, \\ \frac{(1-q^{n-j})(1-q^{j+1})}{(1-q)(1-q^n)} \begin{bmatrix} n+j \\ 2j \end{bmatrix} \begin{bmatrix} m+1 \\ j+1 \end{bmatrix} \begin{bmatrix} m+j \\ j+1 \end{bmatrix} (-q^2; q)_{j-1} & \text{if } j \geq 1 \end{cases}$$

is a polynomial in q with non-negative integer coefficients by [7, Theorem 2.1]. We conclude that (1.14) is the desired Laurent polynomials in q .

Let

$$S_n(x_0, \dots, x_n; q) = \sum_{k=0}^n \begin{bmatrix} n+k \\ 2k \end{bmatrix} \begin{bmatrix} 2k \\ k \end{bmatrix} q^{-nk} x_k.$$

To prove that (1.15) and (1.16) also have the same properties, we first establish the following result.

Lemma 7.1. *Let n and r be positive integers. Then both*

$$\sum_{k=0}^{n-1} \frac{1-q^{2k+1}}{1-q^n} S_k(x_0, \dots, x_k)^r q^{-k} \text{ and } \sum_{k=0}^{n-1} (-1)^{n-k-1} \frac{1-q^{2k+1}}{1-q^n} S_k(x_0, \dots, x_k)^r q^{\binom{k}{2}}$$

are polynomials in x_0, \dots, x_{n-1}, q and q^{-1} with non-negative integer coefficients.

Proof. Recall the identity

$$\begin{bmatrix} k+i \\ 2i \end{bmatrix} \begin{bmatrix} 2i \\ i \end{bmatrix} \begin{bmatrix} k+j \\ 2j \end{bmatrix} \begin{bmatrix} 2j \\ j \end{bmatrix} = \sum_{s=i}^{i+j} \begin{bmatrix} i+j \\ i \end{bmatrix} \begin{bmatrix} j \\ s-i \end{bmatrix} \begin{bmatrix} s \\ j \end{bmatrix} \begin{bmatrix} k+s \\ 2s \end{bmatrix} \begin{bmatrix} 2s \\ s \end{bmatrix} q^{(i+j-s)(k-s)},$$

which can be proved by using the q -Pfaff-Saalschütz identity (see [14, Lemma 2.1]). It follows that

$$\begin{aligned} S_k(x_0, \dots, x_k)^r &= \sum_{0 \leq i_1, \dots, i_r \leq k} \prod_{j=1}^r \begin{bmatrix} k+i_j \\ 2i_j \end{bmatrix} \begin{bmatrix} 2i_j \\ i_j \end{bmatrix} q^{-ki_j} x_{i_j} \\ &= \sum_{0 \leq i_1, \dots, i_r \leq k} x_{i_1} \cdots x_{i_r} \sum_{s=i_1}^{i_1+\dots+i_r} P(i_1, \dots, i_r, s) \begin{bmatrix} k+s \\ 2s \end{bmatrix} \begin{bmatrix} 2s \\ s \end{bmatrix} q^{-ks}, \end{aligned} \quad (7.1)$$

where $P(i_1, \dots, i_r, s)$ is a Laurent polynomial in q independent of k with non-negative integer coefficients. Therefore, by (6.1), we see that

$$\begin{aligned} &\sum_{k=0}^{n-1} \frac{1-q^{2k+1}}{1-q^n} S_k(x_0, \dots, x_k)^r q^{-k} \\ &= \sum_{0 \leq i_1, \dots, i_r \leq n-1} x_{i_1} \cdots x_{i_r} \sum_{s=i_1}^{i_1+\dots+i_r} P(i_1, \dots, i_r, s) \frac{1-q^{n-s}}{1-q^{s+1}} \begin{bmatrix} n+s \\ 2s \end{bmatrix} \begin{bmatrix} 2s \\ s \end{bmatrix} q^{-(s+1)(n-1)} \end{aligned}$$

is a polynomial in x_0, \dots, x_{n-1}, q and q^{-1} with non-negative integer coefficients since

$$\frac{1-q^{n-s}}{1-q^{s+1}} \begin{bmatrix} n+s \\ 2s \end{bmatrix} \begin{bmatrix} 2s \\ s \end{bmatrix} = \begin{bmatrix} n+s \\ s \end{bmatrix} \begin{bmatrix} n \\ s+1 \end{bmatrix}.$$

Similarly, since

$$\sum_{k=s}^{n-1} (-1)^{n-k-1} \frac{1-q^{2k+1}}{1-q^n} \begin{bmatrix} k+s \\ 2s \end{bmatrix} \begin{bmatrix} 2s \\ s \end{bmatrix} q^{\binom{k}{2}-sk} = \begin{bmatrix} n-1 \\ s \end{bmatrix} \begin{bmatrix} n+s \\ s \end{bmatrix} q^{\binom{n}{2}-sn},$$

we deduce from (7.1) that

$$\begin{aligned} & \sum_{k=0}^{n-1} (-1)^{n-k-1} \frac{1-q^{2k+1}}{1-q^n} S_k(x_0, \dots, x_k)^r q^{\binom{k}{2}} \\ &= \sum_{0 \leq i_1, \dots, i_r \leq n-1} x_{i_1} \cdots x_{i_r} \sum_{s=i_1}^{i_1+\dots+i_r} P(i_1, \dots, i_r, s) \begin{bmatrix} n-1 \\ s \end{bmatrix} \begin{bmatrix} n+s \\ s \end{bmatrix} q^{\binom{n}{2}-sn} \end{aligned}$$

is a polynomial in x_0, \dots, x_{n-1}, q and q^{-1} with non-negative integer coefficients. \square

For $k = 0, \dots, n-1$, let

$$x_k = \begin{bmatrix} m+k \\ 2k \end{bmatrix} (-1; q)_k (-q; q)_k q^{k^2-mk}.$$

Then the identity (1.12) may be rewritten as

$$D_q(m, n) D_{q^{-1}}(m, n) = \sum_{k=0}^n \begin{bmatrix} n+k \\ 2k \end{bmatrix} \begin{bmatrix} 2k \\ k \end{bmatrix} q^{-nk} x_k.$$

It is clear that x_0, \dots, x_{n-1} are Laurent polynomials in q with non-negative integer coefficients. By Lemma 7.1, so are the expressions (1.15) and (1.16).

Acknowledgments

The authors would like to thank the referees and the editor for helpful comments on a previous version of this paper. The second author was partially sponsored by the National Natural Science Foundation of China (grant 11371144), the Natural Science Foundation of Jiangsu Province (grant BK20161304), and the Qing Lan Project of Education Committee of Jiangsu Province.

References

- [1] G. Almkvist, D. van Straten, W. Zudilin, Generalizations of Clausen's formula and algebraic transformations of Calabi–Yau differential equations, *Proc. Edinb. Math. Soc.* (2) 54 (2011) 273–295.
- [2] G.E. Andrews, *The Theory of Partitions*, Cambridge University Press, Cambridge, 1998.
- [3] H.H. Chan, Y. Tanigawa, Y. Yang, W. Zudilin, New analogues of Clausen's identities arising from the theory of modular forms, *Adv. Math.* 228 (2011) 1294–1314.
- [4] M. Dziemiańczuk, Generalizing Delannoy numbers via counting weighted lattice paths, *Integers* 13 (2013), #A54.
- [5] G. Gasper, M. Rahman, *Basic Hypergeometric Series*, second edition, *Encyclopedia Math. Appl.*, vol. 96, Cambridge University Press, Cambridge, 2004.
- [6] V.J.W. Guo, Some congruences related to the q -Fermat quotients, *Int. J. Number Theory* 11 (2015) 1049–1060.
- [7] V.J.W. Guo, Proof of Sun's conjectures on integer-valued polynomials, *J. Math. Anal. Appl.* 444 (2016) 182–191.
- [8] V.J.W. Guo, J. Zeng, Some q -supercongruences for truncated basic hypergeometric series, *Acta Arith.* 171 (2015) 309–326.
- [9] F.H. Jackson, The q^θ equations whose solutions are products of solutions of q^θ equations of lower order, *Quart. J. Math.*, Oxford Ser. 11 (1940) 1–17.
- [10] F.H. Jackson, Certain q -identities, *Quart. J. Math.*, Oxford Ser. 12 (1941) 167–172.
- [11] H. Pan, A q -analogue of Lehmer's congruence, *Acta Arith.* 128 (2007) 303–318.
- [12] H. Pan, A Lucas-type congruence for q -Delannoy numbers, preprint, arXiv:1508.02046, 2015.
- [13] M.J. Schlosser, q -Analogues of two product formulas of hypergeometric functions by Bailey, preprint, arXiv:1612.07284.
- [14] A.L. Schmidt, Generalized q -Legendre polynomials, *J. Comput. Appl. Math.* 49 (1993) 243–249.
- [15] L.J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [16] Z.-H. Sun, Generalized Legendre polynomials and related supercongruences, *J. Number Theory* 143 (2014) 293–319.
- [17] L. van Hamme, Some conjectures concerning partial sums of generalized hypergeometric series, in: *p-Adic Functional Analysis, Nijmegen, 1996, in: *Lect. Notes Pure Appl. Math.*, vol. 192, Dekker, 1997, pp. 223–236.*