



On the classes of higher-order Jensen-convex functions and Wright-convex functions

Kazimierz Nikodem, Teresa Rajba, Szymon Wąsowicz*

Department of Mathematics and Computer Science, University of Bielsko-Biała, Willowa 2, 43–309 Bielsko-Biała, Poland

ARTICLE INFO

Article history:

Received 19 January 2012

Available online 23 June 2012

Submitted by M. Laczko

Keywords:

Convex functions of higher order

Jensen-convex functions of higher order

Wright-convex functions of higher order

Forward difference

Backward difference

Hamel basis

Dirac measure

ABSTRACT

The classes of n -Wright-convex functions and n -Jensen-convex functions are compared with each other. It is shown that for any odd natural number n the first one is the proper subclass of the second one. To reach this aim new tools connected with measure theory are developed.

© 2012 Elsevier Inc. All rights reserved.

1. Introduction

Let $\mathcal{I} \subset \mathbb{R}$ be the interval and $f : \mathcal{I} \rightarrow \mathbb{R}$. The usual *forward difference* operator is denoted by

$$\Delta_h f(x) = f(x+h) - f(x),$$

where $x \in \mathcal{I}$ and $h \in \mathbb{R}$ with $x+h \in \mathcal{I}$. Its iterates we define by the usual way, i.e.

$$\Delta_{h_1 \dots h_n h_{n+1}} f(x) = \Delta_{h_1 \dots h_n} (\Delta_{h_{n+1}} f(x))$$

for $n \in \mathbb{N}$, $x \in \mathcal{I}$ and $h_1, \dots, h_n, h_{n+1} \in \mathbb{R}$ with all needed arguments belonging to \mathcal{I} (sometimes we will not write the evident assumptions of this kind). If all increments are equal, $h_1 = \dots = h_n = h$, then we use the standard notation

$$\Delta_h^n f(x) = \Delta_{h \dots h} f(x),$$

where the increment h is taken n times. It is not difficult to check that

$$\begin{aligned} \Delta_{h_1 \dots h_{n+1}} f(x) &= f(x+h_1+\dots+h_{n+1}) - \sum_{1 \leq j_1 < \dots < j_n \leq n+1} f(x+h_{j_1}+\dots+h_{j_n}) \\ &\quad + \sum_{1 \leq j_1 < \dots < j_{n-1} \leq n+1} f(x+h_{j_1}+\dots+h_{j_{n-1}}) \\ &\quad \vdots \\ &\quad + (-1)^n \sum_{1 \leq j_1 \leq n+1} f(x+h_{j_1}) + (-1)^{n+1} f(x). \end{aligned} \quad (1.1)$$

* Corresponding author.

E-mail addresses: knikodem@ath.bielsko.pl (K. Nikodem), trajba@ath.bielsko.pl (T. Rajba), swasowicz@ath.bielsko.pl (S. Wąsowicz).

In this paper also the *backward difference* will be used. It is defined by

$$\nabla_h f(x) = f(x) - f(x - h), \quad (1.2)$$

where $x \in \mathcal{I}$ and $h \in \mathbb{R}$ with $x - h \in \mathcal{I}$. Its iterates are defined similarly to those of the forward differences. Obviously $\nabla_h f(x + h) = \Delta_h f(x)$ and using (1.1), by the induction argument we arrive at

$$\nabla_{h_1 \dots h_{n+1}} f(x + h_1 + \dots + h_{n+1}) = \Delta_{h_1 \dots h_{n+1}} f(x). \quad (1.3)$$

Recall that f is called *Jensen-convex of order n* (n -Jensen-convex for short), if

$$\Delta_h^{n+1} f(x) \geq 0 \quad (1.4)$$

for all $x \in \mathcal{I}$ and $h > 0$ with $x + (n + 1)h \in \mathcal{I}$ (cf. e.g. [1]). Obviously for $n = 1$ we arrive at the condition

$$\Delta_h^2 f(x) = f(x + 2h) - 2f(x + h) + f(x) \geq 0$$

for all $x \in \mathcal{I}$ and $h > 0$ with $x + h \in \mathcal{I}$, which is equivalent to

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2}, \quad x, y \in \mathcal{I},$$

i.e. to the Jensen-convexity of f .

A function f is called *Wright-convex* (cf. [2]), if

$$f(tx + (1 - t)y) + f((1 - t)x + ty) \leq f(x) + f(y)$$

for all $x, y \in \mathcal{I}$, $t \in [0, 1]$. This condition is equivalent to

$$\Delta_{h_1 h_2} f(x) \geq 0$$

for all $x \in \mathcal{I}$, $h_1, h_2 > 0$ with $x + h_1 + h_2 \in \mathcal{I}$ (see [3]). Following this observation, in [4,3], higher order Wright-convexity was defined: a function f is *Wright-convex of order n* (n -Wright-convex for short), if

$$\Delta_{h_1 \dots h_{n+1}} f(x) \geq 0 \quad (1.5)$$

for all $x \in \mathcal{I}$ and $h_1, \dots, h_{n+1} > 0$ with $x + h_1 + \dots + h_{n+1} \in \mathcal{I}$. Of course, setting above $h_1 = \dots = h_{n+1} = h$, we obtain $\Delta_h^{n+1} f(x) \geq 0$, which means that every n -Wright-convex function is n -Jensen-convex.

Then the natural question arises, whether the converse is also true, i.e. whether n -Jensen-convex functions are n -Wright-convex. For $n = 1$ the negative answer is not too difficult to give. Namely, the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = |a(x)|$, where $a : \mathbb{R} \rightarrow \mathbb{R}$ is a discontinuous additive function, is Jensen-convex and it is not Wright-convex (cf. [5]). Indeed, by the well-known Ng's representation (cf. [6]), if f was Wright-convex, it would be the sum of an additive function and a convex one. Then either f would be continuous, or its graph would be dense on the whole plane (cf. e.g. [1]). But neither f is continuous, nor the graph of f is dense on the whole plane.

In the series of papers [4,7,3] rather extensive study of higher-order Wright-convexity was given. However, the above mentioned problem was not considered. In this paper we fill this gap by delivering the negative answer for any odd positive integer n . Let us emphasize that for (odd) $n > 1$ the appropriate counterexample is not easy to construct, as it was for $n = 1$, i.e. in the case of the ordinary Jensen-convexity and Wright-convexity. To reach our goal we develop new tools of measure-theoretical nature, which, we hope, could be also useful for some future research. Let us also mention that for even natural numbers n the considered problem still remains open.

The paper is organized in the following way. In Section 2 we formulate our main result and we prove a part of it. In the next section, to throw some light to the nature of our main problem, we consider the case of n -Jensen-convexity and n -Wright-convexity for $n = 3$. We also perform some considerations for $n = 2$ to show that for even values of n our problem seems to be rather difficult. The nontrivial part of the proof of the main result is postponed to the last section.

2. Main result

Recall that, for $x \in \mathbb{R}$, we have $x_+ = \max\{x, 0\} = \frac{x+|x|}{2}$ and $x_+^n = (x_+)^n$. We start with the following, well-known, lemma.

Lemma 2.1. *Let $n \in \mathbb{N}$, $c \geq 0$. The function $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ given by $\varphi(x) = cx_+^n$ is n -Jensen-convex.*

Proof. It is easy to see that $\varphi^{(n-1)}(x) = cn!x_+$ is a convex function, whence φ is so-called n -convex function, which is obviously n -Jensen-convex (cf. [1,8]). \square

Corollary 2.2. *If $a : \mathbb{R} \rightarrow \mathbb{R}$ is an additive function and n is an odd natural number, then $f(x) = a(x)_+^n$ is n -Jensen-convex.*

Proof. Let $\varphi(x) = x_+^n$. Then $f(x) = \varphi(a(x))$. Using the well-known formula (cf. [1, Corollary 15.1.2]), see also (1.1) we obtain

$$\begin{aligned}\Delta_h^{n+1}f(x) &= \sum_{i=0}^{n+1} \binom{n+1}{i} (-1)^i f(x + (n+1-i)h) \\ &= \sum_{i=0}^{n+1} \binom{n+1}{i} (-1)^i \varphi(a(x) + (n+1-i)a(h)) = \Delta_{a(h)}^{n+1}\varphi(a(x)) \geq 0,\end{aligned}$$

because φ is an n -convex function (for instance, by Lemma 2.1) and n is an odd number (if φ is n -convex and n is odd, then $\Delta_k^{n+1}\varphi(y) \geq 0$ for any $y \in \mathbb{R}$ and any increment $k \in \mathbb{R}$, cf. [1, p. 429], a comment before Lemma 15.3.1). By virtue of (1.4) the proof is finished. \square

In the rest of this paper we use the following idea. An additive map $a : \mathbb{R} \rightarrow \mathbb{R}$ is a linear functional over the vector space of real numbers over the field of rational numbers. Then the function a is uniquely determined by its values on a Hamel basis (cf. e.g. [1]).

Now we are in a position to state our main result.

Theorem 2.3. *Let $n \in \mathbb{N}$ be an odd number. Take a Hamel basis H containing positive numbers h_1, \dots, h_{n+1} . Let $a : \mathbb{R} \rightarrow \mathbb{R}$ be any additive function satisfying*

$$a(h_1) = -1, \quad a(h_2) = \dots = a(h_{n+1}) = 1.$$

The function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) = (a(x))_+^n$$

is n -Jensen-convex whereas it is not n -Wright-convex.

Proof. By Corollary 2.2 the function f is n -Jensen convex. To prove that f is not n -Wright-convex, it is enough to show that $\Delta_{h_1 \dots h_{n+1}} f(0) = -1$ (see (1.5)). However, this is not a trivial job. It requires to develop new tools, and, on the other hand, it is rather long. For these reasons we postpone the rest of the proof to the last section. \square

Since every n -Wright convex function is n -Jensen convex, by the above theorem we obtain immediately the following.

Corollary 2.4. *For any odd $n \in \mathbb{N}$ the class of n -Wright-convex functions is properly contained in the class of n -Jensen-convex functions.*

If $n \in \mathbb{N}$ is even, the question whether the above inclusion is proper, remains an open problem.

3. Two particular cases

3.1. The case $n = 3$

As we mentioned in the Introduction, in the general case the proof of Theorem 2.3 is difficult. In this subsection we deliver some simpler proof for the case $n = 3$.

Take the Hamel basis H such that $h_1, h_2, h_3, h_4 \in H$ are distinct and positive. Let $a : \mathbb{R} \rightarrow \mathbb{R}$ be the additive function such that $a(h_1) = -1, a(h_2) = a(h_3) = a(h_4) = 1$. Let $f(x) = (a(x))_+^3$. Due to Corollary 2.2 the function $f : \mathbb{R} \rightarrow \mathbb{R}$ is 3-Jensen-convex. We will show that f is not 3-Wright-convex. To this end we will check that $\Delta_{h_1 h_2 h_3 h_4} f(0) = -1 < 0$, so the inequality (1.5) does not hold for $n = 3$. We have

$$f(0 + h_1 + h_2 + h_3 + h_4) = (a(0) + a(h_1) + a(h_2) + a(h_3) + a(h_4))_+^3 = 8.$$

Similarly

$$f(0 + h_1 + h_2 + h_3) = f(0 + h_1 + h_2 + h_4) = f(0 + h_1 + h_3 + h_4) = 1,$$

$$f(0 + h_2 + h_3 + h_4) = 27,$$

$$f(0 + h_1 + h_2) = f(0 + h_1 + h_3) = f(0 + h_1 + h_4) = 0,$$

$$f(0 + h_2 + h_3) = f(0 + h_2 + h_4) = f(0 + h_3 + h_4) = 8,$$

$$f(0 + h_1) = 0,$$

$$f(0 + h_2) = f(0 + h_3) = f(0 + h_4) = 1,$$

$$f(0) = 0.$$

Then, having in mind the formula (1.1), we arrive at

$$\begin{aligned}\Delta_{h_1 h_2 h_3 h_4}^4 f(x) &= f(0 + h_1 + h_2 + h_3 + h_4) - [f(0 + h_1 + h_2 + h_3) + f(0 + h_1 + h_2 + h_4) \\ &\quad + f(0 + h_1 + h_3 + h_4) + f(0 + h_2 + h_3 + h_4)] \\ &\quad + [f(0 + h_1 + h_2) + f(0 + h_1 + h_3) + f(0 + h_1 + h_4) \\ &\quad + f(0 + h_2 + h_3) + f(0 + h_2 + h_4) + f(0 + h_3 + h_4)] \\ &\quad - [f(0 + h_1) + f(0 + h_2) + f(0 + h_3) + f(0 + h_4)] \\ &\quad + f(0) = 8 - 30 + 24 - 3 + 0 = -1.\end{aligned}$$

In a similar way this proof was also repeated for $n \in \{5, 7, 9, 11\}$, however, the computations were done by the computer.

3.2. The case $n = 2$

Now we discuss the case $n = 2$ to convince the reader that for even values of n our problem is not easy to solve. Precisely, we will try to compare the classes of 2-Jensen-convex functions with the class of 2-Wright-convex ones.

Looking at the example given in the Introduction we could suppose that the function $f : \mathbb{R} \rightarrow \mathbb{R}$ given by $f(x) = |Q(x)|$, where $Q : \mathbb{R} \rightarrow \mathbb{R}$ fulfils the quadratic functional equation

$$Q(x+y) + Q(x-y) = 2Q(x) + 2Q(y), \quad (3.1)$$

could be a good example of a 2-Jensen-convex function which is not 2-Wright-convex. Unfortunately, f need not to be 2-Jensen-convex. To see this take the Hamel basis H containing the vectors $1, \sqrt{2}$ and $\sqrt[4]{2}$. Next take the additive function $a : \mathbb{R} \rightarrow \mathbb{R}$ defined on H by $a(1) = -9$, $a(\sqrt{2}) = 4$ and $a(h) = 0$ for $h \in H \setminus \{1, \sqrt{2}\}$. Then the function $Q(x) = a(x^2)$ fulfils (3.1). Finally, for $x = 1$, $h = \sqrt[4]{2} - 1 > 0$ we have $Q(x) = -9$, $Q(x+h) = 4$, $Q(x+2h) = 7$, $Q(x+3h) = 0$, whence

$$\Delta_h^3 f(x) = \Delta_h^3 |Q(x)| = |Q(x+3h)| - 3|Q(x+2h)| + 3|Q(x+h)| - |Q(x)| = -18 < 0,$$

which, according to (1.4), proves our claim.

Having in mind Theorem 2.3, it is reasonable to expect that the function $f(x) = (a(x))_+^2$ (for some properly chosen additive function $a : \mathbb{R} \rightarrow \mathbb{R}$) could be the nice example of a 2-Jensen-convex function which is not 2-Wright-convex. However, such a function is not 2-Jensen-convex for any discontinuous additive function a and for any additive function of the form $a(x) = cx$ with $c < 0$. If $a(x) = cx$ with some $c \geq 0$, then f is continuous and, as we will show, f is 2-Jensen-convex. Hence, by continuity, f is also 2-Wright-convex (cf. [1, Theorem 15.7.1]), so it is not a good candidate for our counterexample.

Proposition 3.1. *If $a : \mathbb{R} \rightarrow \mathbb{R}$ is a discontinuous additive function, then $f(x) = (a(x))_+^2$ is not 2-Jensen-convex.*

Proof. Since a is a discontinuous additive function, its graph is dense on the whole plane (cf. e.g. [1]). Then close to the point $(0, 1)$ there exists a point $(x, a(x))$. We can claim, for example, that

$$0.9 < a(x) < 1.1. \quad (3.2)$$

Similarly, close to the point $(1, -2)$ there exists the point $(h, a(h))$. We can claim that $h > 0$ and

$$-2.1 < a(h) < -1.9.$$

Therefore

$$\begin{aligned}-5.4 < a(x+3h) &= a(x) + 3a(h) < -4.6 \implies (a(x+3h))_+ = 0, \\ -3.3 < a(x+2h) &= a(x) + 2a(h) < -2.7 \implies (a(x+2h))_+ = 0, \\ -1.2 < a(x+h) &= a(x) + a(h) < -0.8 \implies (a(x+h))_+ = 0.\end{aligned}$$

By (3.2) we get $(a(x))_+ = a(x) > 0$. Hence, by $f(x) = (a(x))_+^2$,

$$\Delta_h^3 f(x) = f(x+3h) - 3f(x+2h) + 3f(x+h) - f(x) = -(a(x))^2 < 0,$$

so the inequality (1.4) does not hold for $n = 2$ and for any $x \in \mathbb{R}$, $h > 0$. \square

Proposition 3.2. *If $a(x) = cx$ for some $c \geq 0$, then $f(x) = (a(x))_+^2$ is 2-Jensen-convex.*

Proof. Since

$$f(x) = (a(x))_+^2 = \left(\frac{cx + |cx|}{2} \right)^2 = c^2 \left(\frac{x + |x|}{2} \right)^2 = c^2 x_+^2,$$

then f is 2-Jensen-convex by Lemma 2.1. \square

Proposition 3.3. If $a(x) = cx$ for some $c < 0$, then $f(x) = (a(x))_+^2$ is not 2-Jensen-convex.

Proof. If $c < 0$, then we have

$$f(x) = (a(x))_+^2 = \left(\frac{cx + |cx|}{2} \right)^2 = \left(\frac{cx - c|x|}{2} \right)^2 = c^2 \left(\frac{x - |x|}{2} \right)^2.$$

Therefore

$$f(-x) = c^2 \left(\frac{-x - |x|}{2} \right)^2 = c^2 \left(\frac{x + |x|}{2} \right)^2 = c^2 x_+^2$$

and $f(x) = c^2 (-x)_+^2$. Setting $x = -1$, $h = 1$ we obtain $\Delta_h^3 f(x) = -c^2 < 0$, so f is not 2-Jensen-convex. \square

4. Proof of Theorem 2.3

In this section we develop new tools connected with the measure theory and we use them to prove that the function f defined in Theorem 2.3 is not n -Wright-convex. According to our best knowledge this approach was not used so far.

4.1. Notations and basic facts

By $\mathcal{B}(\mathbb{R})$ we denote the σ -field of Borel subsets of \mathbb{R} . By Borel measure we mean any measure defined on $\mathcal{B}(\mathbb{R})$. It is known that the distribution function $F_\mu(x) = \mu((-\infty, x))$ determines μ i.e. to know the value the Borel measure, it is enough to know its values on the intervals $(-\infty, x)$ for any $x \in \mathbb{R}$ (cf. [9, Sections 12, 14]).

Throughout this section we deal only with the functions $f : \mathbb{R} \rightarrow \mathbb{R}$. In addition to the backward difference operator ∇_h given by (1.2) we also consider the *backward translation operator*

$$\tau_h f(x) = f(x - h), \quad x, h \in \mathbb{R}.$$

Let $\mathcal{M}(\mathbb{R})$ be the set of all Borel measures ν on $\mathcal{B}(\mathbb{R})$ such that $\nu((-\infty, x)) < \infty$, $x \in \mathbb{R}$.

Remark 4.1. If $\nu \in \mathcal{M}(\mathbb{R})$, then $\lim_{x \rightarrow -\infty} \nu((-\infty, x)) = 0$.

Proof. It is an easy consequence of the general property of the measure: if $(A_k : k \in \mathbb{N})$ is a descending sequence of measurable sets with $\nu(A_1) < \infty$, then $\nu(\bigcap_{k \in \mathbb{N}} A_k) = \lim_{k \rightarrow \infty} \nu(A_k)$. \square

We will consider the operators τ_h and ∇_h defined not only for the functions, but also for the measures $\nu \in \mathcal{M}(\mathbb{R})$ (such an approach is frequently used in the Measure Theory):

$$\tau_h \nu(B) = \nu(B - h), \quad \nabla_h \nu(B) = \nu(B) - \tau_h \nu(B) = \nu(B) - \nu(B - h)$$

for $B \in \mathcal{B}(\mathbb{R})$ with $\nu(B) < \infty$.

Let $\nu \in \mathcal{M}(\mathbb{R})$. We define

$$\mathcal{J}_h \nu(B) = \sum_{n=0}^{\infty} \tau_h^n \nu(B), \quad h > 0, B \in \mathcal{B}(\mathbb{R}),$$

where $\tau_h^0 \nu(B) = \nu(B)$, $\tau_h^{n+1}(B) = \tau_h(\tau_h^n \nu)(B)$. It is not difficult to check that

$$\mathcal{J}_h \nu \in \mathcal{M}(\mathbb{R}) \iff \sum_{n=0}^{\infty} F_\nu(x - nh) < \infty, \quad x \in \mathbb{R} \quad \text{and} \quad \lim_{x \rightarrow -\infty} \sum_{n=0}^{\infty} F_\nu(x - nh) = 0.$$

For these notations, see also [10, 11].

Proposition 4.2. Let $\mu \in \mathcal{M}(\mathbb{R})$ and $h > 0$.

1. If

$$\nabla_h \mu \geq 0, \tag{4.1}$$

then there exists $\nu \in \mathcal{M}(\mathbb{R})$ such that μ has the form

$$\mu = \mathcal{J}_h \nu. \tag{4.2}$$

Moreover,

$$\nu = \nabla_h \mu. \tag{4.3}$$

2. If μ has the form (4.2) with $\nu \in \mathcal{M}(\mathbb{R})$, then the conditions (4.1) and (4.3) hold.

Proof. 1. Let μ fulfils (4.1). Using the definition of ∇_h we have $\nabla_h \mu = \mu - \tau_h \mu$, whence $\mu = \nabla_h \mu + \tau_h \mu$. Then

$$\begin{aligned}\tau_h \mu &= \tau_h \nabla_h \mu + \tau_h^2 \mu, \\ \tau_h^2 \mu &= \tau_h^2 \nabla_h \mu + \tau_h^3 \mu, \\ &\vdots \\ \tau_h^n \mu &= \tau_h^n \nabla_h \mu + \tau_h^{n+1} \mu.\end{aligned}$$

Hence

$$\mu = \nu + \tau_h \nu + \cdots + \tau_h^n \nu + \tau_h^{n+1} \mu,$$

where $\nu = \nabla_h \mu$, $n = 1, 2, \dots$. Taking into account Remark 4.1 we infer that

$$\tau_h^{n+1} \mu((-\infty, x)) = \nu((-\infty, x - (n+1)h)) \xrightarrow{n \rightarrow \infty} 0$$

whence the distribution function of the measure μ , i.e. $F_\mu(x) = \mu((-\infty, x))$ ($x \in \mathbb{R}$), is equal to the distribution function of a measure $\mathcal{J}_h \nu$, where ν is given by (4.3). Then these measures are equal (cf. e.g. [9, Sections 12, 14]), which finishes the proof of 1.

2. Let μ has the form (4.2) with $\nu \in \mathcal{M}(\mathbb{R})$. Then

$$\mu = \mathcal{J}_h \nu = \sum_{n=0}^{\infty} \tau_h^n \nu = \nu + \sum_{n=1}^{\infty} \tau_h^n \nu. \quad (4.4)$$

Using the definition of τ_h and τ_h^n we get

$$\begin{aligned}\tau_h(\mathcal{J}_h \nu)(B) &= \mathcal{J}_h \nu(B-h) = \sum_{n=0}^{\infty} \tau_h^n \nu(B-h) \\ &= \sum_{n=0}^{\infty} \tau_h(\tau_h^n \nu)(B) = \sum_{n=0}^{\infty} \tau_h^{n+1} \nu(B) = \sum_{n=1}^{\infty} \tau_h^n \nu(B).\end{aligned}$$

Therefore $\tau_h \mu = \sum_{n=1}^{\infty} \tau_h^n \nu$, which, together with (4.4), yields $\mu = \nu + \tau_h \mu$, which implies $\nu = \mu - \tau_h \mu$ and the proof of 2. is finished. \square

For $\nu \in \mathcal{M}(\mathbb{R})$ and $h_1, \dots, h_n > 0$ denote $\mathcal{J}_{h_1 h_2 \dots h_n} \nu = \mathcal{J}_{h_1} \mathcal{J}_{h_2} \dots \mathcal{J}_{h_n} \nu$. As the immediate consequence of Proposition 4.2 we obtain the following.

Proposition 4.3. Let $h_1, \dots, h_n > 0$.

- (a) If $\nu \in \mathcal{M}(\mathbb{R})$ fulfils the condition $\mathcal{J}_{h_1 \dots h_n} \nu \in \mathcal{M}(\mathbb{R})$, then $\nabla_{h_1 \dots h_n}(\mathcal{J}_{h_1 \dots h_n} \nu) = \nu$.
- (b) If $\mu \in \mathcal{M}(\mathbb{R})$ fulfils the condition $\nabla_{h_1 \dots h_n} \mu \geq 0$, then $\mathcal{J}_{h_1 \dots h_n}(\nabla_{h_1 \dots h_n} \mu) = \mu$.

4.2. Preparation to the proof of Theorem 2.3

Fix $n \in \mathbb{N}$ and consider the Hamel basis $H \subset \mathbb{R}$ such that $h_1, \dots, h_{n+1} \in H$ are distinct and positive. We keep this convention throughout the whole section. Recall that if $x \in \mathbb{R}$, then δ_x denotes the Dirac measure, i.e. $\delta_x(B) = 1$ if $x \in B$ and $\delta_x(B) = 0$, $x \notin B$, where $B \subset \mathbb{R}$. Define the measures $\mu_1, \dots, \mu_{n+1} \in \mathcal{M}(\mathbb{R})$ by

$$\mu_i = \mathcal{J}_{h_1 \dots h_{n+1}} \delta_{h_i}, \quad i = 1, \dots, n+1. \quad (4.5)$$

Then define the signed measure μ by

$$\mu = \mu_2 + \cdots + \mu_{n+1} - \mu_1. \quad (4.6)$$

Being the elements of the Hamel basis, h_1, \dots, h_{n+1} are incommensurable, and it is not difficult to check the formula

$$\mu_i = \sum_{j_1, \dots, j_{n+1}=0}^{\infty} \delta_{h_i + j_1 h_1 + \dots + j_{n+1} h_{n+1}}, \quad i = 1, \dots, n+1. \quad (4.7)$$

Next take the sets $A, A_1, \dots, A_{n+1} \subset \mathbb{R}$ defined by

$$\begin{aligned}A_i &= \left\{ h_i + \sum_{\substack{j=1 \\ j \neq i}}^{n+1} \varepsilon_j h_j : \varepsilon_j \in \{0, 1\}, j = 1, \dots, n+1 \right\}, \quad i = 1, \dots, n+1, \\ A &= A_1 \cup \cdots \cup A_{n+1}.\end{aligned}$$

We will use the frequent notation $\mu(x) = \mu(\{x\})$.

Lemma 4.4. Let $i \in \{1, \dots, n+1\}$. Then

- (a) $\mu_i(x) = 1$ for $x \in A_i$,
- (b) $\mu_i(x) = 0$ for $x \in A \setminus A_i$,
- (c) $\mu|_A(x) < 0 \iff x = h_1$,
- (d) $\mu(h_1) = \mu_1(h_1) = -1$,
- (e) $\mu_+|_A = \mu|_A + \delta_{h_1}$.

Proof. It is enough to use (4.5), (4.6), (4.7). We omit a standard and easy proof. \square

Recall that in Theorem 2.3 we defined the function $f(x) = (a(x))_+^n$ ($x \in \mathbb{R}$), where $a : \mathbb{R} \rightarrow \mathbb{R}$ is the additive function such that $a(h_1) = -1$, $a(h_2) = \dots = a(h_{n+1}) = 1$. Now we prove the crucial property of this function f . Let us notice that the function f could be, of course, defined for any $n \in \mathbb{N}$ and the result below is not dependent on the evenness of n .

Theorem 4.5. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined as above and μ be a signed measure given by (4.6). Then

$$f(x) = (\mu + \delta_{h_1})^n(x) \quad \text{for every } x \in A. \quad (4.8)$$

In particular,

$$\nabla_{h_1 \dots h_{n+1}} f(h_1 + \dots + h_{n+1}) = \nabla_{h_1 \dots h_{n+1}} (\mu + \delta_{h_1})^n(h_1 + \dots + h_{n+1}). \quad (4.9)$$

Proof. To prove (4.8) it is enough to show that

$$a(x) = \mu(x) \quad \text{for every } x \in A. \quad (4.10)$$

Indeed, then for any $x \in A$ we have $a_+(x) = \mu_+(x)$ and trivially $f(x) = (a_+(x))^n = (\mu_+(x))^n$. Taking into account Lemma 4.4(e) we get (4.8).

To prove (4.10) fix $x \in A$. Then $x = \varepsilon_1 h_1 + \dots + \varepsilon_{n+1} h_{n+1}$, where $\varepsilon_i \in \{0, 1\}$, $i = 1, \dots, n+1$ and $\varepsilon_1 + \dots + \varepsilon_{n+1} > 0$. Two cases are possible.

CASE 1. $\varepsilon_1 = 1$

If $\varepsilon_2 = \dots = \varepsilon_{n+1} = 0$, then $a(x) = a(h_1) = -1$ and by Lemma 4.4(d) $\mu(x) = \mu(h_1) = -1$, so (4.10) holds. If $\varepsilon_j \neq 0$ for some $j \in \{2, \dots, n+1\}$, then without loss of generality we may assume that $x = h_1 + \dots + h_k$ for some $k \in \{2, \dots, n+1\}$. Since $x \in A_1 \cap \dots \cap A_k$ and $x \notin A_{k+1}, \dots, x \notin A_{n+1}$, we have by Lemma 4.4(a), (b)

$$\mu_1(x) = \dots = \mu_k(x) = 1, \quad \mu_{k+1}(x) = \dots = \mu_{n+1}(x) = 0.$$

Hence, by virtue of (4.6), $\mu(x) = k - 2$. By additivity

$$a(x) = a(h_1 + \dots + h_k) = a(h_1) + \dots + a(h_k) = k - 2,$$

which proves that $a(x) = \mu(x)$.

CASE 2. $\varepsilon_1 = 0$

Without loss of generality we may assume that $x = h_2 + \dots + h_k$ for some $k \in \{2, \dots, n+1\}$. Arguing exactly in the same way as in the previous case, we arrive at $\mu(x) = k - 1 = a(x)$. \square

4.3. Proof of Theorem 2.3

Theorem 4.5 allows us to work with measures instead of the original function f . We present below three useful formulae. We will prove them after the proof of Theorem 2.3.

Lemma 4.6. Let $\mu = \mu_2 + \dots + \mu_{n+1} - \mu_1$ be the signed measure given by (4.6). Then

$$(\mu + \delta_{h_1})^n(x) = \mu^n(x) - (-1)^n \delta_{h_1}(x) \quad \text{for any } x \in A, \quad (4.11)$$

$$\nabla_{h_1 \dots h_{n+1}} \mu^n(h_1 + \dots + h_{n+1}) = 0, \quad (4.12)$$

$$\nabla_{h_1 \dots h_{n+1}} \delta_{h_1}(h_1 + \dots + h_{n+1}) = (-1)^n. \quad (4.13)$$

The Final Step of the Proof of Theorem 2.3. Recall that $n \in \mathbb{N}$ was odd and we have chosen the Hamel basis $H \subset \mathbb{R}$ such that $h_1, \dots, h_{n+1} \in H$ were positive. We took the additive function $a : \mathbb{R} \rightarrow \mathbb{R}$ such that $a(h_1) = -1$ and $a(h_2) = \dots = a(h_{n+1}) = 1$. Then we defined the function $f : \mathbb{R} \rightarrow \mathbb{R}$ by $f(x) = (a(x))_+^n$ and we have shown that f is n -Jensen-convex. It was left to prove that f is not n -Wright-convex. It is enough to check that $\Delta_{h_1 \dots h_{n+1}} f(0) = -1$. By (1.3)

this is equivalent to $\nabla_{h_1 \dots h_{n+1}} f(h_1 + \dots + h_{n+1}) = -1$. Using (4.9) and (4.11) we obtain

$$\begin{aligned} \nabla_{h_1 \dots h_{n+1}} f(h_1 + \dots + h_{n+1}) &= \nabla_{h_1 \dots h_{n+1}} (\mu + \delta_{h_1})^n (h_1 + \dots + h_{n+1}) \\ &= \nabla_{h_1 \dots h_{n+1}} (\mu^n (h_1 + \dots + h_{n+1}) - (-1)^n \delta_{h_1} (h_1 + \dots + h_{n+1})) \\ &= \nabla_{h_1 \dots h_{n+1}} \mu^n (h_1 + \dots + h_{n+1}) + \nabla_{h_1 \dots h_{n+1}} \delta_{h_1} (h_1 + \dots + h_{n+1}) = -1 \end{aligned}$$

due to (4.12) and (4.13). This finishes the proof. \square

4.4. Proof of Lemma 4.6

Proof of Formula (4.11). Let $x \in A$. Of course $\delta_{h_1}^j = \delta_{h_1}$ ($j \in \mathbb{N}$). Therefore

$$\begin{aligned} (\mu + \delta_{h_1})^n(x) &= \mu^n(x) + \delta_{h_1}^n(x) + \sum_{k=1}^{n-1} \binom{n}{k} \mu^k(x) \delta_{h_1}^{n-k}(x) \\ &= \mu^n(x) + \delta_{h_1}(x) + \sum_{k=1}^{n-1} \binom{n}{k} \mu^k(x) \delta_{h_1}(x). \end{aligned} \quad (4.14)$$

Put $\lambda = \mu_2 + \dots + \mu_{n+1}$. Then $\mu = \lambda - \mu_1$ and for $k = 1, \dots, n-1$ we get

$$\mu^k(x) = \sum_{j=1}^{k-1} \binom{k}{j} \lambda^j(x) (-\mu_1^{k-j}(x)) + \lambda^k(x) + (-1)^k \mu_1^k(x). \quad (4.15)$$

It is easy to see that

- (a) $\lambda^j(x) \delta_{h_1}(x) = 0, j = 1, \dots, k$,
- (b) $\mu_1^k(x) \delta_{h_1}(x) = \delta_{h_1}(x)$.

For, notice that λ is concentrated on the set $A_2 \cap \dots \cap A_{n+1}$ (see 4.4(a), (b)), which gives (a), while (b) is trivial. Then (4.15) yields

$$\mu^k(x) \delta_{h_1}(x) = (-1)^k \delta_{h_1}(x), \quad k = 1, \dots, n-1,$$

and, consequently,

$$\begin{aligned} \sum_{k=1}^{n-1} \binom{n}{k} \mu^k(x) \delta_{h_1}(x) &= \sum_{k=1}^{n-1} \binom{n}{k} (-1)^k \delta_{h_1}(x) = \delta_{h_1}(x) \left[\sum_{k=1}^{n-1} \binom{n}{k} (-1)^k \right] \\ &= \delta_{h_1}(x) \left[\sum_{k=0}^n \binom{n}{k} (-1)^k - \binom{n}{0} (-1)^0 - \binom{n}{n} (-1)^n \right] \\ &= \delta_{h_1}(x) [0 - 1 - (-1)^n]. \end{aligned}$$

We conclude the proof putting this last equation into (4.14). \square

Proof of Formula (4.12). Let $x \in A$. Applying the Multinomial Theorem to $\mu = \mu_2 + \dots + \mu_{n+1} - \mu_1$ we arrive at

$$\mu^n(x) = \sum_{j_1 + \dots + j_{n+1} = n} \binom{n}{j_1, j_2, \dots, j_{n+1}} \mu_2^{j_2}(x) \cdot \dots \cdot \mu_{n+1}^{j_{n+1}}(x) \cdot (-\mu_1^{j_1}(x)), \quad (4.16)$$

where

$$\binom{n}{j_1, j_2, \dots, j_{n+1}} = \frac{n!}{j_1! \cdot j_2! \cdot \dots \cdot j_{n+1}!}$$

are the *multinomial coefficients*. Due to Lemma 4.4(a), (b) we have

$$\mu_j^k(x) = \begin{cases} \mu_j(x) & \text{for } k = 1, 2, \dots, \\ 1 & \text{for } k = 0 \end{cases} \quad (4.17)$$

with the convention $0^0 = 1$. Next we will prove that

$$\mu_{j_1}(x) \cdot \mu_{j_2}(x) \cdot \dots \cdot \mu_{j_k}(x) = \mathcal{J}_{h_1 \dots h_{n+1}} \delta_{h_{j_1} + \dots + h_{j_k}}(x), \quad (4.18)$$

where $1 \leq j_1 < j_2 < \dots < j_k \leq n+1$. For simplicity we will only check that

$$\mu_1(x) \cdot \mu_2(x) = \mathcal{J}_{h_1 \dots h_{n+1}} \delta_{h_1 + h_2}(x), \quad (4.19)$$

the proof in the general case is analogous. By (4.7)

$$\mu_1(x) = \sum_{j_1, \dots, j_{n+1}=0}^{\infty} \delta_{h_1+j_1 h_1+\dots+j_{n+1} h_{n+1}}(x),$$

$$\mu_2(x) = \sum_{j_1, \dots, j_{n+1}=0}^{\infty} \delta_{h_2+j_1 h_1+\dots+j_{n+1} h_{n+1}}(x).$$

If $x \in A_1 \cap A_2$, then $x = h_1 + h_2 + j_1 h_1 + \dots + j_{n+1} h_{n+1}$. Because h_1, \dots, h_{n+1} belong to the Hamel basis, this representation is unique and both the above sums are equal to 1. By the same argument, also

$$\mathcal{J}_{h_1 \dots h_{n+1}} \delta_{h_1+h_2}(x) = \sum_{j_1, \dots, j_{n+1}=0}^{\infty} \delta_{h_1+h_2+j_1 h_1+\dots+j_{n+1} h_{n+1}}(x) = 1.$$

By Lemma 4.4(a) we infer that $\mu_1(x)\mu_2(x) = 1$ and (4.19) holds. The remaining case $x \in A \setminus (A_1 \cup A_2)$ we handle in the similar way, also using Lemma 4.4(b).

Taking into account (4.17) and (4.18) we obtain that

$$\mu_2^{j_2}(x) \cdot \dots \cdot \mu_{n+1}^{j_{n+1}}(x) \cdot (-\mu_1(x)^{j_1}) = (-1)^{j_1} \mathcal{J}_{h_1 \dots h_{n+1}} \delta_{\varepsilon_2 h_2 + \dots + \varepsilon_{n+1} h_{n+1} + \varepsilon_1 h_1}(x), \quad (4.20)$$

where

$$\varepsilon_k = \begin{cases} 0 & \text{for } j_k = 0, \\ 1 & \text{for } j_k > 0 \end{cases} \quad (4.21)$$

for $k = 1, \dots, n+1$. By Proposition 4.3(a) we have

$$\nabla_{h_1 \dots h_{n+1}} (\mathcal{J}_{h_1 \dots h_{n+1}} \delta_{\varepsilon_2 h_2 + \dots + \varepsilon_{n+1} h_{n+1} + \varepsilon_1 h_1})(x) = \delta_{\varepsilon_2 h_2 + \dots + \varepsilon_{n+1} h_{n+1} + \varepsilon_1 h_1}(x). \quad (4.22)$$

Consequently, by (4.16), (4.20) and (4.22) we get

$$\nabla_{h_1 \dots h_{n+1}} \mu^n(x) = \sum_{j_1 + \dots + j_{n+1} = n} \binom{n}{j_1, j_2, \dots, j_{n+1}} (-1)^{j_1} \delta_{\varepsilon_1 h_1 + \dots + \varepsilon_{n+1} h_{n+1}}(x). \quad (4.23)$$

Observe that, for $x = h_1 + \dots + h_{n+1}$, there is $\varepsilon_1 = \dots = \varepsilon_{n+1} = 1$, so, by (4.21), $j_1 + \dots + j_{n+1} \geq n+1$ and, in the sum (4.23), there is no component $\delta_{h_1 + \dots + h_{n+1}}(h_1 + \dots + h_{n+1})$. Since h_1, \dots, h_{n+1} belong to the Hamel basis, every component of this sum equals 0, so $\nabla_{h_1 \dots h_{n+1}} \mu^n(h_1 + \dots + h_{n+1}) = 0$ and the formula (4.12) is true. \square

Proof of Formula (4.13). By (1.3) we have

$$\nabla_{h_1 \dots h_{n+1}} \delta_{h_1}(h_1 + \dots + h_{n+1}) = \Delta_{h_1 \dots h_{n+1}} \delta_{h_1}(0).$$

We compute this term using (1.1). Notice that (by the choice of h_1, \dots, h_{n+1} as distinct elements of the Hamel basis) the only non-zero component of the sum occurring there is $(-1)^n \delta_{h_1}(h_1) = (-1)^n$, which appears in the penultimate line. \square

References

- [1] M. Kuczma, An Introduction to the Theory of Functional Equations and Inequalities, Birkhäuser, Basel, 2009.
- [2] A.W. Roberts, D.E. Varberg, Convex Functions, Academic Press, New York, London, 1973.
- [3] G. Maksa, Zs. Páles, Decomposition of higher-order Wright-convex functions, J. Math. Anal. Appl. 359 (2) (2009) 439–443.
- [4] A. Gilányi, Zs. Páles, On convex functions of higher order, Math. Inequal. Appl. 11 (2) (2008) 271–282.
- [5] K. Nikodem, On some class of midconvex functions, Ann. Polon. Math. 50 (2) (1989) 145–151.
- [6] C.T. Ng, Functions generating Schur-convex sums, in: General Inequalities, 5 (Oberwolfach, 1986), in: Internat. Schriftenreihe Numer. Math., vol. 80, Birkhäuser, Basel, 1987, pp. 433–438.
- [7] A. Gilányi, Zs. Páles, On Dinghas-type derivatives and convex functions of higher order, Real Anal. Exchange 27 (2) (2001/02) 485–493.
- [8] T. Popoviciu, Sur quelques propriétés des fonctions d'une ou de deux variables réelles, Mathematica, Cluj 8 (1934) 1–85.
- [9] P. Billingsley, Probability and Measure, third ed., John Wiley and Sons, New York, 1995.
- [10] T. Rajba, An application of the Choquet theorem to the study of randomly-superinvariant measures, Opuscula Math. 32 (2) (2012) 317–326.
- [11] T. Rajba, A generalization of multiple Wright-convex functions via randomization, J. Math. Anal. Appl. 388 (1) (2012) 548–565.