



R-bounded approximating sequences and applications to semigroups [☆]

M. Hoffmann,¹ N. Kalton,^{*} and T. Kucherenko

Department of Mathematics, University of Missouri-Columbia, Columbia, MO 65211, USA

Received 15 December 2002

Available online 24 April 2004

Submitted by R. Triggiani

Abstract

It is shown that on certain Banach spaces, including $C[0, 1]$ and $L_1[0, 1]$, there is no strongly continuous semigroup $(T_t)_{0 < t < 1}$ consisting of weakly compact operators such that $(T_t)_{0 < t < 1}$ is an R-bounded family. More general results concerning approximating sequences are included and some variants of R-boundedness are also discussed.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Strongly continuous semigroups; R-boundedness; Approximation property

1. Introduction

Recent work on semigroup theory [13,24] has highlighted the importance of the concept of R-boundedness. Let us recall the definition of R-bounded families of operators (cf. [2, 7,9]).

Definition 1.1. A family \mathcal{T} of operators in $\mathcal{L}(X, Y)$ is called *R-bounded* with R-boundedness constant $C > 0$ if letting $(\epsilon_k)_{k=1}^\infty$ be a sequence of independent Rademachers on some probability space then for every $x_1, \dots, x_n \in X$ and $T_1, \dots, T_n \in \mathcal{T}$ we have

[☆] The authors acknowledge support from NSF grant DMS-9870027.

^{*} Corresponding author.

E-mail addresses: hoffmann@math.sc.edu (M. Hoffmann), nigel@math.missouri.edu (N. Kalton), tamara@math.missouri.edu (T. Kucherenko).

¹ Current address: Department of Mathematics, University of South Carolina, Columbia, SC 29208, USA.

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k T_k x_k \right\|^2 \right)^{1/2} \leq C \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2}. \quad (1.1)$$

By the Kahane–Khintchine inequality we can replace 2 above by any other exponent $1 \leq p < \infty$ to obtain an equivalent definition. We will also need the following definition introduced in [13].

Definition 1.2. A family \mathcal{T} of operators in $\mathcal{L}(X, Y)$ is called *WR-bounded* with WR-boundedness constant $C > 0$ if for every $x_1, \dots, x_n \in X$, $y_1^*, \dots, y_n^* \in Y^*$ and $T_1, \dots, T_n \in \mathcal{T}$ we have

$$\sum_{k=1}^n |(T_k x_k, y_k^*)| \leq C \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k y_k^* \right\|^2 \right)^{1/2}. \quad (1.2)$$

It is clear by the Cauchy–Schwarz inequality that R-boundedness implies WR-boundedness. The converse is not true in general, but it holds for spaces with non-trivial type [13, 20].

In [13] it was shown that no reasonable differential operator on L_1 can have an H^∞ -calculus. In this note we consider the related question whether a differential-type operator on L_1 can generate an R-bounded semigroup. Note that if A is an R-sectorial operator (cf. [13]) with R-sectoriality angle less than $\pi/2$ then the semigroup $(e^{-tA})_{0 < t < 1}$ is necessarily R-bounded. In general, one expects a semigroup generated by a differential operator on a bounded domain to consist of weakly compact operators. We are thus led to consider the question whether one can have a strongly continuous semigroup $(T_t)_{0 < t < 1}$ on L_1 such that each T_t is weakly compact (or equivalently compact, since L_1 has the Dunford–Pettis property) and such that the family $(T_t)_{0 < t < 1}$ is R-bounded. In fact this leads to considering versions of the approximation property; the only property of the semigroup needed is commutativity. We consider the general question whether on a given separable Banach space one can find an R-bounded sequence $(T_n)_{n \in \mathbb{N}}$ of commuting weakly compact operators such that $\lim_{n \rightarrow \infty} T_n x = x$ for all $x \in X$. Our main results show that for the spaces $L_1[0, 1]$, $C(K)$ (except c_0) and the disk algebra $A(\mathbb{D})$ this is impossible. These results may be regarded as extensions of classical results that the spaces $L_1, C(K)$ do not have unconditional bases [15].

In the case of L_1 we are led to consider a natural weakening of R-boundedness, where we use the definition (1.1) but only for single vectors.

Definition 1.3. A family \mathcal{T} of operators in $\mathcal{L}(X, Y)$ is called *semi-R-bounded* if there is a constant $C > 0$ such that for every $x \in X$, $a_1, \dots, a_n \in \mathbb{C}$ and $T_1, \dots, T_n \in \mathcal{T}$ we have

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k a_k T_k x \right\|^2 \right)^{1/2} \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|x\|. \quad (1.3)$$

We note that semi-R-boundedness is equivalent to R-boundedness for operators on L_1 . In Theorem 2.2 we actually characterize all spaces where semi-R-boundedness is equivalent to R-boundedness as spaces which are either Hilbert spaces or GT-spaces of cotype 2 in the terminology of Pisier [19].

2. R-boundedness and WR-boundedness

In this section, we make some remarks about R-boundedness and related notions.

Note that in a space of type 2, any uniformly bounded collection $\mathcal{T} \subset L(X, X)$ is semi-R-bounded. The converse is also true:

Proposition 2.1. *A Banach space X has type 2 if and only if uniform boundedness is equivalent to semi-R-boundedness.*

Proof. Suppose that every uniformly bounded family of operators is already semi-R-bounded. Pick any $x \in X$ and $x^* \in X^*$ such that $\|x\| = \|x^*\| = 1$ and $x^*(x) = 1$. Notice that the family $\mathcal{T} = \{x^* \otimes u : \|u\| = 1\}$ is uniformly bounded with constant one and hence semi-R-bounded by assumption. Let C be the semi-R-boundedness constant of \mathcal{T} . Select any $x_1, \dots, x_n \in X$ and write $x_k = \|x_k\|u_k$, where $\|u_k\| = 1$. Then $\{x^* \otimes u_k : k = 1, \dots, n\} \subset \mathcal{T}$ and

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2} &= \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| (x^* \otimes u_k)x \right\|^2 \right)^{1/2} \\ &\leq C \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| x \right\|^2 \right)^{1/2} = C \|x\| \left(\sum_{k=1}^n \|x_k\|^2 \right)^{1/2} \\ &= C \left(\sum_{k=1}^n \|x_k\|^2 \right)^{1/2}. \end{aligned}$$

Thus, X has type 2. \square

For some spaces, semi-R-boundedness is equivalent to R-boundedness and we are able to completely characterize these spaces in the next theorem. Let us recall that a Banach space X is called a *GT-space* if every bounded operator $T : X \rightarrow \ell_2$ is absolutely summing. Examples of GT-spaces of cotype 2 are L_1 , the quotient of L_1 by a reflexive subspace [14,19], and L_1/H_1 [8]. It is unknown whether every GT-space has cotype 2.

Theorem 2.2. *Suppose X is separable. Then the following are equivalent:*

- (i) *Every semi-R-bounded family of operators on X is R-bounded.*
- (ii) *X is isomorphic to ℓ_2 or X is a GT-space of cotype 2.*

Proof. First we prove that (i) implies (ii). Suppose that every semi-R-bounded family of operators on X is R-bounded. Let us note that this implies the existence of a constant K so that if \mathcal{T} has semi-R-boundedness constant C then it has R-boundedness constant KC ; for otherwise we could find a sequence \mathcal{T}_n of families with semi-R-boundedness constant one and R-boundedness constant at least 4^n ; then the family $\bigcup_{n \geq 1} 2^{-n} \mathcal{T}_n$ contradicts our

assumption. Fix $M > 1$ and take $x \in X$. Choose $n \in \mathbb{N}$. By Dvoretzky's theorem [16] we can find $e_1, \dots, e_n \in X$ such that for any $a_1, \dots, a_n \in \mathbb{C}$ we have

$$M^{-1} \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \leq \left\| \sum_{k=1}^n a_k e_k \right\| \leq M \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}.$$

Consider the family of operators $\mathcal{T}_n = \{u^* \otimes e_k: \|u^*\| = 1, k = 1, \dots, n\}$. Then each \mathcal{T}_n is semi-R-bounded with constant M as follows. A finite subfamily of \mathcal{T}_n is of the form $\{u_{kj}^* \oplus e_k: 1 \leq k \leq n, 1 \leq j \leq m_k\}$ for some $m_1, \dots, m_n \in \mathbb{N}$. Then for every $a_{11}, \dots, a_{nm_n} \in \mathbb{C}$ we have (letting ϵ_{kj} denote independent Rademachers)

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{k=1}^n \sum_{j=1}^{m_k} \epsilon_{kj} a_{kj} u_{kj}^*(x) e_k \right\|^2 \right)^{1/2} &\leq M \left(\mathbb{E} \sum_{k=1}^n \left| \sum_{j=1}^{m_k} \epsilon_{kj} u_{kj}^*(x) a_{kj} \right|^2 \right)^{1/2} \\ &\leq M \left(\sum_{k=1}^n \mathbb{E} \left| \sum_{j=1}^{m_k} \epsilon_{kj} u_{kj}^*(x) a_{kj} \right|^2 \right)^{1/2} \leq M \left(\sum_{k=1}^n \sum_{j=1}^{m_k} |a_{kj}|^2 \right)^{1/2} \|x\|. \end{aligned}$$

Our assumption implies that each \mathcal{T}_n is R-bounded with constant KM . Let $x_1, \dots, x_n \in X$ and write $x_k = \|x_k\| u_k$, where $\|u_k\| = 1$. Choose $u_k^* \in X^*$ such that $u_k^*(u_k) = 1$ and $\|u_k^*\| = 1$. Now we have

$$\begin{aligned} \left(\sum_{k=1}^n \|x_k\|^2 \right)^{1/2} &\leq M \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| e_k \right\|^2 \right)^{1/2} \\ &= M \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| u_k^*(u_k) e_k \right\|^2 \right)^{1/2} \\ &= M \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| (u_k^* \otimes e_k)(u_k) \right\|^2 \right)^{1/2} \\ &\leq KM^2 \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| u_k \right\|^2 \right)^{1/2} = KM^2 \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2}. \end{aligned}$$

This shows that X has cotype 2.

Let us assume that X has non-trivial type. Then by results of Pisier [19] and also by Figiel and Tomczak-Jaegermann [12], ℓ_2^n is uniformly complemented in X . Thus, for some constant C , for every $n \in \mathbb{N}$ we can choose a biorthogonal system $\{(e_k, e_k^*): k = 1, \dots, n\}$ in $X \times X^*$ such that

$$\left\| \sum_{k=1}^n a_k e_k \right\| \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}$$

and

$$\left\| \sum_{k=1}^n a_k e_k^* \right\| \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2}$$

for all $a_1, \dots, a_n \in \mathbb{C}$. Note that for any $x \in X$ and $a_1, \dots, a_n \in \mathbb{C}$,

$$\left| \sum_{k=1}^n a_k e_k^*(x) \right| \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|x\|$$

and so

$$\left(\sum_{k=1}^n |e_k^*(x)|^2 \right)^{1/2} \leq C \|x\|.$$

Consider the family of operators $\mathcal{T}_n = \{e_k^* \otimes u : \|u\| = 1, k = 1, \dots, n\}$. Let $x \in X$. Then for any $a_1, \dots, a_n \in \mathbb{C}$ and every $u_1, \dots, u_n \in X$ of norm one we have

$$\begin{aligned} \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k a_k (e_k^* \otimes u_k)(x) \right\| &= \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k a_k e_k^*(x) u_k \right\| \leq \sum_{k=1}^n \|a_k e_k^*(x) u_k\| \\ &= \sum_{k=1}^n |a_k| |e_k^*(x)| \leq \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \left(\sum_{k=1}^n |e_k^*(x)|^2 \right)^{1/2} \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|x\|. \end{aligned}$$

We conclude that \mathcal{T}_n is semi-R-bounded with constant C and hence \mathcal{T}_n is R-bounded for constant KC independent of n . This implies that X has type 2 as follows. Choose any $x_1, \dots, x_n \in X$ and write $x_k = \|x_k\| u_k$, where $\|u_k\| = 1$. Then

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2} &= \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| u_k \right\|^2 \right)^{1/2} \\ &= \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| e_k^*(e_k) u_k \right\|^2 \right)^{1/2} \\ &= \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| (e_k^* \otimes u_k)(e_k) \right\|^2 \right)^{1/2} \\ &\leq KC \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \|x_k\| e_k \right\|^2 \right)^{1/2} \leq KC^2 \left(\sum_{k=1}^n \|x_k\|^2 \right)^{1/2}. \quad (2.1) \end{aligned}$$

Now, X has type 2 and cotype 2 and is therefore isomorphic to ℓ_2 by Kwapien's theorem [25].

Now suppose on the contrary that X has trivial type. We will show that X is a GT-space, i.e., any $T : X \rightarrow \ell_2$ is 1-summing. Fix $T : X \rightarrow \ell_2$ of norm one. Since X has cotype 2 we can equivalently show that any such T is 2-summing [11]. It suffices to check that for any $n \in \mathbb{N}$ and operator $S : \ell_2^n \rightarrow X$ such that $\|S\| \leq 1$ we have $\pi_2(TS) \leq C$, where C does not depend on n [25]. One can assume that $TS : \ell_2^n \rightarrow \ell_2^n$ and that TS is diagonal with respect to the canonical orthonormal basis (e_k) in ℓ_2^n , i.e., $TSe_k = \lambda_k e_k$ for some $\lambda_1, \dots, \lambda_n$. Then it suffices to show uniform boundedness of the Hilbert–Schmidt norms $\|TS\|_{\text{HS}} = (\sum_{k=1}^n \|TSe_k\|^2)^{1/2}$. Write $f_k^* = T^* e_k^* \in X^*$ and $f_k = Se_k \in X$. Consider $\{f_k^* \otimes u : k = 1, \dots, n, \|u\| = 1\}$. We will show that this family is semi-R-bounded

with constant one. Take $u_1, \dots, u_n \in X$ of norm one and $a_1, \dots, a_n \in \mathbb{C}$. Then for $x \in X$ we have

$$\begin{aligned} \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k a_k f_k^*(x) u_k \right\| &\leq \sum_{k=1}^n |a_k| |f_k^*(x)| \leq \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \left(\sum_{k=1}^n |e_k^*(Tx)|^2 \right)^{1/2} \\ &= \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|Tx\| \leq \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|x\|. \end{aligned}$$

Therefore, $\{f_k^* \otimes u: k = 1, \dots, n, \|u\| = 1\}$ is R-bounded with constant K .

Since X has trivial type, it contains ℓ_1^n uniformly [19]. Hence, for fixed $M > 1$ and every $n \in \mathbb{N}$ there are $y_1, \dots, y_n \in X$ with $\|y_k\| = 1$ for $1 \leq k \leq n$ such that

$$\sum_{k=1}^n |a_k| \leq M \left\| \sum_{k=1}^n a_k y_k \right\|. \quad (2.2)$$

Choose any scalars b_1, \dots, b_n . Now using R-boundedness and Kahane's inequality for $p = 1$ with constant A we have

$$\begin{aligned} \sum_{k=1}^n |b_k| |\lambda_k| &= \sum_{k=1}^n |b_k| |f_k^*(f_k)| \leq M \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k b_k f_k^*(f_k) y_k \right\|^2 \right)^{1/2} \\ &= M \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k b_k (f_k^* \otimes y_k)(f_k) \right\|^2 \right)^{1/2} \leq KM \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k b_k f_k \right\|^2 \right)^{1/2} \\ &\leq KM \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k b_k e_k \right\|^2 \right)^{1/2} \leq KM \left(\sum_{k=1}^n |b_k|^2 \right)^{1/2}. \end{aligned}$$

Thus,

$$\left(\sum_{k=1}^n |\lambda_k|^2 \right)^{1/2} \leq KM,$$

and so $\|TS\|_{\text{HS}} \leq KM$. Therefore, any operator $T: X \rightarrow \ell_2$ is 2-summing. This completes the proof of (i) implies (ii).

Now we will show that (ii) implies (i). Suppose that X is a GT-space of cotype 2, and that \mathcal{T} is a family of semi-R-bounded operators. We will show that \mathcal{T} is R-bounded. Since X is separable, there is a quotient map $Q: \ell_1 \rightarrow X$. First, we show that any semi-R-bounded family of operators from ℓ_1 into X is already R-bounded. Let \mathcal{S} be such a family with semi-R-boundedness constant one. Suppose $S_1, \dots, S_n \in \mathcal{S}$ and $x_1, \dots, x_n \in \ell_1$. Then $x_k = \sum_{j=1}^\infty \xi_{jk} e_j$, where (e_j) is the canonical basis of ℓ_1 .

Let us denote by C the constant in the Kahane–Khintchine inequality for any Banach space:

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|^2 \right)^{1/2} \leq C \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|.$$

Thus

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k S_k x_k \right\|^2 \right)^{1/2} \leq C \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k S_k x_k \right\|.$$

Then

$$\begin{aligned} \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k S_k x_k \right\| &= \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k S_k \sum_{j=1}^{\infty} \xi_{jk} e_j \right\| \leq \sum_{j=1}^{\infty} \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \xi_{jk} S_k e_j \right\| \\ &\leq \sum_{j=1}^{\infty} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \xi_{jk} S_k e_j \right\|^2 \right)^{1/2} \leq \sum_{j=1}^{\infty} \left(\sum_{k=1}^n |\xi_{jk}|^2 \right)^{1/2}. \end{aligned}$$

Combining and using the Khintchine inequality again we obtain

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k S_k x_k \right\|^2 \right)^{1/2} &\leq C^2 \sum_{j=1}^{\infty} \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \xi_{jk} \right\| = C^2 \mathbb{E} \left(\sum_{j=1}^{\infty} \left\| \sum_{k=1}^n \epsilon_k \xi_{jk} \right\| \right) \\ &= C^2 \mathbb{E} \left\| \sum_{j=1}^{\infty} \sum_{k=1}^n \epsilon_k \xi_{jk} e_j \right\|_{\ell_1} = C^2 \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k \sum_{j=1}^{\infty} \xi_{jk} e_j \right\| \\ &= C^2 \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|. \end{aligned}$$

Combining the previous two computations gives that \mathcal{S} is R-bounded.

Now let \mathcal{T} be a family of operators on X with semi-boundedness constant one. Let $Q: \ell_1 \rightarrow X$ be a quotient map and note that the family $\mathcal{S} = \{TQ: T \in \mathcal{T}\}$ is R-bounded with some constant B by the above calculation.

We will apply a characterization of GT-spaces of cotype 2 due to Pisier [19].

Proposition 2.3 (Pisier). *X is a GT-space of cotype 2 if and only if there is a constant $C > 0$ such that for any $n \in \mathbb{N}$, $x_1, \dots, x_n \in X$, there are $y_1, \dots, y_n \in \ell_1$ such that $Qy_k = x_k$, $k = 1, \dots, n$, and*

$$\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k y_k \right\| \leq C \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|. \quad (2.3)$$

Now take $n \in \mathbb{N}$, $T_1, \dots, T_n \in \mathcal{T}$ and $x_1, \dots, x_n \in X$. Choose $y_1, \dots, y_n \in \ell_1$ according to Proposition 2.3. Then

$$\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k T_k x_k \right\| = \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k T_k Q y_k \right\| \leq B \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k y_k \right\| \leq CB \mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k \right\|.$$

Thus, \mathcal{T} is R-bounded. The proof is complete. \square

For a set \mathcal{T} of bounded linear operators we will use the notation $\mathcal{T}^* = \{T^*: T \in \mathcal{T}\}$.

Lemma 2.4.

- (i) If \mathcal{T} is R -bounded then \mathcal{T}^{**} is R -bounded (with the same constant).
- (ii) If \mathcal{T} is WR -bounded then \mathcal{T}^* and \mathcal{T}^{**} are WR -bounded (with the same constant).
- (iii) If \mathcal{T} is semi- R -bounded then \mathcal{T}^{**} is semi- R -bounded (with the same constant).

Proof. The proofs of (i) and (iii) are similar. For (i) suppose $T_1, \dots, T_n \in \mathcal{T}$ and that \mathcal{T} has R -boundedness constant one. Let $\Omega = \{-1, 1\}^n$ with \mathbb{P} normalized counting measure on Ω . Let ϵ_k be the sequence of coordinate maps on Ω . Let $\text{Rad}(\Omega; X)$ be the subspace of $L_2(\Omega, \mathbb{P}; X)$ generated by the functions $\epsilon_k \otimes x$ for $1 \leq k \leq n$ and $x \in X$ (this space is isomorphic to X^n). Then $\text{Rad}(\Omega; X^{**})$ can be identified naturally with a subspace of $\text{Rad}(\Omega; X)^{**}$. Consider the map $\mathbf{T}: \text{Rad}(\Omega; X) \rightarrow \text{Rad}(\Omega; X)$ defined by

$$\mathbf{T}\left(\sum_{k=1}^n \epsilon_k \otimes x_k\right) = \sum_{k=1}^n \epsilon_k \otimes T_k x_k.$$

Then $\|\mathbf{T}\| \leq 1$ and so $\|\mathbf{T}^{**}\| \leq 1$ and (i) follows.

Let us now prove (ii). Suppose \mathcal{T} is WR -bounded with constant one and $T_1, \dots, T_n \in \mathcal{T}$. Suppose $x_1^*, \dots, x_n^* \in X^*$ are such that

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x_k^* \right\|^2\right)^{1/2} \leq 1.$$

Then, using the identification of $\text{Rad}(\Omega, X^{**})$ as the bidual of $\text{Rad}(\Omega, X)$ we observe that the set of functions of the form $\sum_{k=1}^n \epsilon_k x_k^{**}$ in $\text{Rad}(\Omega, X^{**})$ such that

$$\sum_{k=1}^n |\langle T_k^* x_k^*, x_k^{**} \rangle| \leq 1$$

is weak*-closed and contains the unit ball of $\text{Rad}(\Omega, X)$. By Goldstine's theorem it contains the unit ball of $\text{Rad}(\Omega, X^{**})$ and this implies that \mathcal{T}^* is WR -bounded with constant one. \square

Now it is time to give an example of a family of operators that is uniformly bounded but not WR -bounded. The previous lemma will imply that the corresponding dual family is semi- R -bounded but not WR -bounded.

Example. Let $X = \ell_p$, $1 \leq p < 2$. Pick any non-zero element $x \in X$ and choose $u^* \in X^*$ of norm one such that $u^*(x) \neq 0$. Define $T_k = u^* \otimes e_k$, where (e_k) is the canonical basis of X . The family $\{T_k\}$ is uniformly bounded, $\|T_k\| = 1$, but we will show that it is not WR -bounded. Consider the dual basis (e_k^*) in $(\ell_p)^*$. Then

$$\sum_{k=1}^n |\langle T_k x, e_k^* \rangle| = \sum_{k=1}^n |\langle u^*(x) e_k, e_k^* \rangle| = n |u^*(x)|. \quad (2.4)$$

On the other hand, we have

$$\left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x \right\|^2 \right)^{1/2} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k e_k^* \right\|^2 \right)^{1/2} = \|x\| n^{1/2} n^{1/q}. \quad (2.5)$$

Here q satisfies $1/p + 1/q = 1$. If $p < 2$ then $q > 2$ and $1/2 + 1/q < 1$, so for $1 \leq p < 2$ the family $\{T_k\}$ cannot be WR-bounded.

We have $T_k^* = e_k^{**} \otimes u^*$ on $X^* = \ell_q$, where $2 < q \leq \infty$. Consider $q \neq \infty$. Since by reflexivity $T_k^{**} = T_k$ and using Lemma 2.4 we see that $\{T_k^*\}$ is not WR-bounded. However, X^* has type 2 and hence $\{T_k^*\}$ is semi-R-bounded by Proposition 2.1.

3. The main results

Suppose X is any Banach space. We shall say that a sequence $\mathcal{T} = (T_k)_{k=1}^\infty$ is an *approximating sequence* if $\lim_{k \rightarrow \infty} \|x - T_k x\| = 0$ for every $x \in X$. We will say that \mathcal{T} is compact (relatively, weakly compact) if each T_k is compact (relatively, weakly compact). We will say that \mathcal{T} is commuting if we have $T_k T_l = T_l T_k$ for $l, k \in \mathbb{N}$.

If \mathcal{T} is a commuting approximating sequence, let us define the subspace $E_{\mathcal{T}}$ of X^* to be the closed linear span of $\bigcup_k T_k^*(X^*)$. The following lemma is trivial.

Lemma 3.1. *If \mathcal{T} is a commuting approximating sequence then $E_{\mathcal{T}}$ is a norming subspace of X^* , i.e., for some C we have*

$$\|x\| \leq C \sup_{x^* \in E_{\mathcal{T}}} |x^*(x)|, \quad x \in X,$$

and, if \mathcal{T} is weakly compact, $\lim_{n \rightarrow \infty} T_n^* x^* = x^*$ weakly for $x^* \in E_{\mathcal{T}}$.

Let us recall that a Banach space X has property (V) of Pełczyński if every unconditionally converging operator $T: X \rightarrow Y$ is weakly compact. The spaces $C(K)$ have property (V) [17] and more generally any C^* -algebra has property (V) [18]. The disk algebra $A(\mathbb{D})$ also has property (V) [10,14]; see also [23]. We also recall that a Banach space X is said to have property (V^*) if whenever (x_n) is a bounded sequence in X then either

- (i) (x_n) has a subsequence which is weakly Cauchy or
- (ii) (x_n) has a subsequence (y_n) such that for some sequence (y_n^*) in X^* and $\delta > 0$ we have $|y_n^*(y_n)| \geq \delta$ and

$$\left\| \sum_{k=1}^n a_k y_k^* \right\| \leq \max_{1 \leq k \leq n} |a_k|, \quad a_1, \dots, a_n \in \mathbb{C}, \quad n \in \mathbb{N}.$$

Property (V^*) was introduced by Pełczyński [17]. We note that Bombal [4] shows that every Banach lattice not containing c_0 has property (V^*) . Any subspace of a space with property (V^*) also has property (V^*) .

Lemma 3.2. Let X, Y be Banach spaces and let $\mathcal{T} = (T_k)_{k=1}^\infty$ be any sequence of operators in $\mathcal{L}(X, Y)$. Suppose either

- (i) \mathcal{T} is semi-R-bounded or
- (ii) \mathcal{T} is WR-bounded and Y has property (V^*) .

Then for every $x \in X$ the sequence $(T_k x)_{k=1}^\infty$ has a weakly Cauchy subsequence.

Proof. If not, by passing to a subsequence we can suppose $(T_k x)_{k=1}^\infty$ is equivalent to the canonical ℓ_1 -basis [21,22]. If \mathcal{T} is semi-R-bounded we observe that for some C we have

$$\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k a_k T_k x \right\| \leq C \left(\sum_{k=1}^n |a_k|^2 \right)^{1/2} \|x\|, \quad a_1, \dots, a_n \in \mathbb{C}, \quad n \in \mathbb{N}.$$

This gives a contradiction.

In case (ii), we can pass to a subsequence and assume the existence of $y_n^* \in Y^*$ such that

$$\left\| \sum_{k=1}^n a_k y_k^* \right\| \leq \max_{1 \leq k \leq n} |a_k|, \quad a_1, \dots, a_n \in \mathbb{C}, \quad n \in \mathbb{N},$$

and $|y_n^*(T_n x)| \geq \delta > 0$ for all n . Then

$$n\delta \leq \sum_{k=1}^n |y_k^*(T_k x)| \leq C \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k x \right\|^2 \right)^{1/2} \left(\mathbb{E} \left\| \sum_{k=1}^n \epsilon_k y_k^* \right\|^2 \right)^{1/2} \leq C\sqrt{n}.$$

This also yields a contradiction. \square

Theorem 3.3. Let X be a Banach space with a commuting weakly compact approximating sequence \mathcal{T} . Suppose either that

- (i) \mathcal{T} is semi-R-bounded and X is weakly sequentially complete or
- (ii) \mathcal{T} is WR-bounded and X has property (V^*) .

Then X is isomorphic to a dual space.

Proof. In either case we consider the family $\mathcal{T}^{**} \subset \mathcal{L}(X^{**}, X)$. By Lemma 3.2 for each $x^{**} \in X^{**}$ we can find a subsequence $T_{k_n}^{**} x^{**}$ so that $T_{k_n}^{**}(x^{**})$ is weakly convergent to some $y \in X$. Then for $x^* \in X^*$,

$$x^*(T_k y) = \lim_{n \rightarrow \infty} x^*(T_k T_{k_n}^{**} x^{**}) = \lim_{n \rightarrow \infty} x^*(T_{k_n} T_k^{**} x^{**})$$

so that $T_k y = T_k^{**} x^{**}$. Hence $\lim_{k \rightarrow \infty} \|y - T_k^{**} x^{**}\| = 0$.

We now show that $E_{\mathcal{T}}^*$ can be identified with X . Clearly X canonically embeds in $E_{\mathcal{T}}^*$ since $E_{\mathcal{T}}$ is norming. If $f^* \in E_{\mathcal{T}}^*$ then by the Hahn–Banach theorem there exists $x^{**} \in X^{**}$ with $\|x^{**}\| = \|f^*\|$ and $x^{**}(x^*) = f^*(x^*)$ for $x^* \in E_{\mathcal{T}}$. Let $y = \lim_{k \rightarrow \infty} T_k^{**} x^{**}$. Then for $x^* \in E_{\mathcal{T}}$,

$$x^*(y) = \lim_{k \rightarrow \infty} x^*(T_k^{**} x^{**}) = \lim_{k \rightarrow \infty} x^{**}(T_k^* x^*) = f^*(x^*).$$

Hence $E_{\mathcal{T}}^* = X$. \square

Theorem 3.4. *The space $L_1(0, 1)$ does not have a commuting weakly compact approximating sequence which is either semi-R-bounded or WR-bounded.*

Proof. L_1 is not a dual space [25]. \square

Of course a semi-R-bounded sequence in L_1 is actually R-bounded.

Theorem 3.5. *Let X be a separable Banach space with property (V). If X has a commuting weakly compact approximating sequence $(T_n)_{n=1}^\infty$ which is WR-bounded, then X^* is separable, and has a WR-bounded commuting weakly compact approximating sequence.*

Proof. Since X has (V), it follows that X^* has property (V^*) . We show that $\lim_{n \rightarrow \infty} T_n^* x^* = x^*$ weakly for $x^* \in X^*$. Indeed $T_n^* x^*$ converges weak* to x^* and it must have a weakly convergent subsequence by Lemma 3.2. Hence $x^* \in E_{\mathcal{T}}$ so $X^* = E_{\mathcal{T}}$. Now $T_n^*(B_{X^*})$ is weakly compact by Gantmacher's theorem also and weak*-metrizable, hence norm separable. Thus X^* is separable, and so by Mazur's theorem, and a diagonal argument, we can find a sequence of convex combinations $(S_n^*)_{n=1}^\infty$ of $(T_n^*)_{n=1}^\infty$ which is an approximating sequence. \square

Corollary 3.6. *If K is an uncountable compact metric space then $C(K)$ has no WR-bounded commuting weakly compact approximating sequence. The disk algebra has no WR-bounded weakly compact approximating sequence.*

We now consider $C(K)$ when K is countable. In this case $C(K)$ is homeomorphic to a space $C(\alpha) = C([1, \alpha])$, where α is a countable ordinal. There is a characterization of such $C(K)$ due to Bessaga and Pełczyński [3].

Theorem 3.7 (Bessaga–Pełczyński). *If $\alpha < \beta$, $C(\omega^\alpha \cdot k)$ is isomorphic to $C(\omega^\beta \cdot n)$ if and only if $\beta < \alpha \cdot \omega$. Consequently, $C(\omega^\gamma)$, $0 \leq \gamma < \omega_1$, is a complete list of representatives of the isomorphism classes of $C(K)$ for K a countable compact metric space.*

The following lemma can be obtained as an applications of ℓ_1 -indices [1,5,6]. However, for convenience of the reader we will give a direct proof by construction.

Lemma 3.8. *Let α be a countable ordinal with $\alpha \geq \omega$. Then there exists $f \in C(\alpha)^{**}$ so that whenever $f_n \in C(\alpha)$ converges to $f \in C(\alpha)^{**}$ weak* then for any $m \in \mathbb{N}$ there exist $n_1, \dots, n_m \in \mathbb{N}$ such that*

$$\left\| \sum_{k=1}^m \epsilon_k f_{n_k} \right\| \geq \frac{1}{2}m, \quad \epsilon_k = \pm 1, \quad k = 1, 2, \dots, m.$$

Proof. In this case $C(\alpha)^{**}$ can be identified with $\ell_\infty(\alpha)$. It is easy to see that it suffices to consider the case $\alpha = \omega^\omega$.

Consider $f \in X^{**}$ defined by $f(\sum_{k=0}^N \omega^k l_k) = (-1)^{\sum_{k=0}^N l_k}$ and $f(\omega^\omega) = 1$. Writing K for the space $[1, \omega^\omega]$ let $K^{(p)}$ denote the p th derived set of K . Then $K^{(p)}$ consists of all ordinals of the form $\sum_{k=p}^n \omega^k l_k$ together with ω^ω . For each $p \in \mathbb{N}$, $K^{(p)}$ is non-empty. Furthermore for each $\alpha \in K^{(p)}$ and every open neighborhood V of α we have that f takes both values ± 1 on $V \cap K^{(p-1)}$.

Let $f_n \in C(K)$ be any sequence such that (f_n) converges to f weak*.

Fix $0 < \delta < 1/2$ and $m \in \mathbb{N}$. We construct $(f_{n_1}, \dots, f_{n_m})$ inductively. We start from $K^{(m)}$. By definition of f we can pick $\alpha_1^1, \alpha_2^1 \in K^{(m)}$ such that $f(\alpha_j^1) = (-1)^j$ for $j = 1, 2$. Then find $n_1 \in \mathbb{N}$ such that $|f_{n_1}(\alpha_j^1) - (-1)^j| < \delta$. Since f_{n_1} is continuous we can choose open neighborhoods U_j^1 of α_j^1 such that $|f_{n_1}(\alpha) - (-1)^j| < \delta$ for all $\alpha \in U_j^1$.

For the inductive step, suppose that $(n_j)_{j=1}^k, (\alpha_j^k)_{j=1}^{2^k}$ and open sets $(U_j^k)_{j=1}^{2^k}$ have been chosen so that $\alpha_j^k \in U_j^k$. Then for $i = 1, \dots, 2^k$ find points $\alpha_{2i-1}^{k+1}, \alpha_{2i}^{k+1} \in U_i^k \cap K^{(m-k+1)}$ with $f(\alpha_j^{k+1}) = (-1)^j$. By pointwise convergence, we can select $n_{k+1} > n_k$ such that $|f_{n_{k+1}}(\alpha_j^{k+1}) - (-1)^j| < \delta$. Since $f_{n_{k+1}}$ is continuous, there are neighborhoods $U_{2i-1}^{k+1}, U_{2i}^{k+1} \subset U_i^k, i = 1, \dots, 2^k$, such that for all $\alpha \in U_j^{k+1}$ we have $|f_{n_{k+1}}(\alpha) - (-1)^j| < \delta$.

In the m th iteration this will give 2^m neighborhoods and m functions f_{n_1}, \dots, f_{n_m} so that for any $\epsilon_1, \dots, \epsilon_m \in \{-1, +1\}$ there is α contained in one of these neighborhoods such that $|f_k(\alpha) - \epsilon_k| < \delta$ for all $k = 1, \dots, m$. Hence

$$\left\| \sum_{k=1}^m \epsilon_k f_{n_k} \right\| \geq (1 - \delta)m. \quad \square$$

Theorem 3.9. Let K be a compact metric space. Suppose there is an R -bounded commuting weakly compact approximating sequence in $C(K)$. Then $C(K)$ is isomorphic to c_0 .

Proof. By Corollary 3.6 we need only consider the case when K is countable. By Theorem 3.7 it suffices to consider the case when $K = [1, \alpha]$, where $\alpha \geq \omega^\omega$. Pick $f \in C(K)^{**}$ satisfying the hypotheses of Lemma 3.8.

Suppose (T_n) is an R -bounded weakly compact approximating sequence for $C(K)$. Then (T_n^*) is an approximating sequence for $C(K)^*$ by Theorem 3.5 and hence $T_n^{**}f$ converges to f weak*. It follows that for any m we can choose n_1, \dots, n_m so that

$$\left\| \sum_{k=1}^m \epsilon_k T_{n_k}^{**} f \right\| \geq \frac{1}{2}m, \quad \epsilon_k = \pm 1.$$

Hence

$$\left(\mathbb{E} \left\| \sum_{k=1}^m \epsilon_k T_{n_k}^{**} f \right\|^2 \right)^{1/2} \geq \frac{1}{2}m.$$

This contradicts the fact that T_n is R -bounded (or even semi- R -bounded). \square

Remark. We can replace the assumption of R -boundedness by the assumption that (T_n) and (T_n^*) are both semi- R -bounded. By Theorem 2.2 this hypothesis would imply that (T_n^*) is actually R -bounded and hence that (T_n) is WR -bounded. We only used the fact that (T_n) is both semi- R -bounded and WR -bounded.

Let us conclude by stating our main result with respect to semigroups. (Actually our results are somewhat stronger than stated below.)

Theorem 3.10. *Let X be a separable Banach space with an R -bounded strongly continuous semigroup $(T_t)_{t>0}$ consisting of weakly compact operators. Then if*

- (1) $X = L_1(\mu)$ for some measure μ then X is isomorphic to ℓ_1 (i.e., μ is purely atomic).
- (2) $X = C(K)$ then X is isomorphic to c_0 .

Acknowledgment

The authors thank Dale Alspach for drawing their attention to the notion of the ℓ_1 index of sequences.

References

- [1] D. Alspach, R. Judd, E. Odell, The Szlenk index and local ℓ_1 -indices, Positivity, in press.
- [2] E. Berkson, T.A. Gillespie, Spectral decompositions and harmonic analysis on UMD spaces, *Studia Math.* 112 (1994) 13–49.
- [3] C. Bessaga, A. Pełczyński, Spaces of continuous functions (IV), *Studia Math.* 19 (1960) 53–62.
- [4] F. Bombal, On (V^*) sets and Pełczyński's property (V^*) , *Glasgow Math. J.* 32 (1990) 109–120.
- [5] J. Bourgain, On convergent sequences of continuous functions, *Bull. Soc. Math. Belg.* 32 (1980) 235–249.
- [6] J. Bourgain, Remarks on the double dual of a Banach space, *Bull. Soc. Math. Belg.* 32 (1980) 171–178.
- [7] J. Bourgain, Some remarks on Banach spaces in which martingale difference sequences are unconditional, *Ark. Mat.* 21 (1983) 163–168.
- [8] J. Bourgain, W.J. Davis, Martingale transforms and complex uniform convexity, *Trans. Amer. Math. Soc.* 294 (1986) 501–515.
- [9] P. Clément, B. de Pagter, F.A. Sukochev, H. Witvliet, Schauder decomposition and multiplier theorems, *Studia Math.* 138 (2000) 135–163.
- [10] F. Delbaen, Weakly compact operators on the disc algebra, *J. Algebra* 45 (1977) 284–294.
- [11] J. Diestel, H. Jarchow, A. Tonge, Absolutely Summing Operators, in: *Cambridge Studies in Advanced Mathematics*, vol. 43, Cambridge Univ. Press, Cambridge, 1995.
- [12] T. Figiel, N. Tomczak-Jaegermann, Projections onto Hilbertian subspaces of Banach spaces, *Israel J. Math.* 33 (1979) 155–171.
- [13] N. Kalton, L. Weis, The H^∞ -calculus and sums of closed operators, *Math. Ann.* 321 (2001) 319–345.
- [14] S.V. Kisliakov, The Dunford–Pettis, Pełczyński and Grothendieck conditions, *Dokl. Akad. Nauk SSSR* 225 (1975) 1252–1255 (in Russian).
- [15] J. Lindenstrauss, L. Tzafriri, *Classical Banach Spaces I*, Springer, Berlin, 1977.
- [16] V.D. Milman, G. Schechtman, Asymptotic Theory of Finite-Dimensional Normed Spaces, in: *Springer Lecture Notes*, vol. 1200, 1986.
- [17] A. Pełczyński, Banach spaces on which every unconditionally converging operator is weakly compact, *Bull. Acad. Polon. Sci. Sr. Sci. Math. Astronom. Phys.* 10 (1962) 641–648.
- [18] H. Pfitzner, Weak compactness in the dual of a C^* -algebra is determined commutatively, *Math. Ann.* 298 (1994) 349–371.

- [19] G. Pisier, Factorization of Linear Operators and Geometry of Banach Spaces, in: CBMS Regional Conference Series, vol. 60, American Mathematical Society, Providence, RI, 1986.
- [20] G. Pisier, Holomorphic semigroups and the geometry of Banach spaces, *Ann. Math.* 115 (1982) 375–392.
- [21] H.P. Rosenthal, A characterization of Banach spaces containing l^1 , *Proc. Nat. Acad. Sci. USA* 71 (1974) 2411–2413.
- [22] E. Odell, H.P. Rosenthal, A double-dual characterization of separable Banach spaces containing l^1 , *Israel J. Math.* 20 (1975) 375–384.
- [23] S.F. Saccone, The Pełczyński property for tight subspaces, *J. Funct. Anal.* 148 (1997) 86–116.
- [24] L. Weis, Operator-valued Fourier multiplier theorems and maximal L_p -regularity, *Math. Ann.* 319 (2001) 735–758.
- [25] P. Wojtaszczyk, Banach Spaces for Analysts, in: Cambridge Studies in Advanced Mathematics, vol. 25, Cambridge Univ. Press, Cambridge, 1991.