



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa



Non-smooth atomic decomposition of 2-microlocal spaces and application to pointwise multipliers ☆

Helena F. Gonçalves, Henning Kempka *

ARTICLE INFO

Article history:

Received 17 April 2015

Available online xxxx

Submitted by P.G. Lemarie-Rieusset

Keywords:

2-microlocal spaces

Besov and Triebel–Lizorkin spaces

Variable integrability

Atomic decomposition

Pointwise multipliers

ABSTRACT

We provide non-smooth atomic characterizations for 2-microlocal Besov and Triebel–Lizorkin spaces with variable integrability $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$. These spaces cover the classical Besov and Triebel–Lizorkin spaces as well as spaces of variable smoothness and integrability. As an application, we state a pointwise multipliers result for these spaces.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

In this article we generalize the atomic decomposition theorem for 2-microlocal Besov and Triebel–Lizorkin spaces $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and present an application to pointwise multipliers.

The 2-microlocal function spaces initially appeared in the book of Peetre [20] and have also been studied by Bony [2] in connection with pseudodifferential operators. Later on, they were investigated by Jaffard [9] as well as Jaffard and Meyer [10]. In [16] and [17], Lévy Véhel and Seuret showed that they are a useful tool to measure local regularity and to describe the oscillatory behavior of functions near singularities.

Spaces of variable integrability, also known as variable exponent function spaces $L_{p(\cdot)}(\mathbb{R}^n)$, can be traced back to Orlicz [19] 1931, but the modern development started with the papers [15] of Kováčik and Rákosník as well as [3] of Diening. The spaces $L_{p(\cdot)}(\mathbb{R}^n)$ have interesting applications in fluid dynamics, image processing, PDE and variational calculus, see the introduction of [6]. For an overview we refer to [5].

The concept of function spaces with variable smoothness and the concept of variable integrability were firstly mixed up by Diening, Hästö and Roudenko in [6]. They defined Triebel–Lizorkin spaces $F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$

☆ The authors were supported by the German science foundation (DFG) within the project KE 1847/1-1.

* Corresponding author.

E-mail addresses: helena.goncalves@mathematik.tu-chemnitz.de (H.F. Gonçalves), henning.kempka@mathematik.tu-chemnitz.de (H. Kempka).

and from the trace theorem on \mathbb{R}^{n-1} it became clear why it is natural to have all parameters variable. Due to

$$\mathrm{Tr} F_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n) = F_{p(\cdot),p(\cdot)}^{s(\cdot)-\frac{1}{p(\cdot)}}(\mathbb{R}^{n-1}), \text{ with } s(\cdot) - \frac{1}{p(\cdot)} > (n-1) \max\left(\frac{1}{p(\cdot)} - 1, 0\right),$$

(Theorem 3.13 in [6]) we see the necessity of taking s and q variable if p is not constant. A similar interplay between smoothness and integrability has also been observed by Vybíral in [27] by proving Sobolev embeddings for variable exponent spaces. Moreover, Almeida and Hästö also introduced Besov spaces $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ with all three indices variable in [1].

The scale we consider here – mixing 2-microlocal weights with variable integrability – was introduced in [12,13] and provides a unified approach that covers many spaces related with variable smoothness and generalized smoothness. Many results have been studied regarding these spaces, in particular the possibility of decomposing functions $f \in B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ or $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ as linear combinations of smooth atoms, which are the building blocks for atomic decompositions.

The study of smooth atomic decompositions for Besov spaces with p and q variable was firstly done by Drihem in [7], where he proved the result for Besov spaces with variable smoothness and integrability $B_{p(\cdot),q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$. Following the same ideas, we present in Section 2 the smooth atomic decomposition result for 2-microlocal Besov spaces $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$.

In Section 3 we develop non-smooth atomic decompositions for 2-microlocal Besov and Triebel–Lizorkin spaces $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$. We show that we can replace the usual atoms used in smooth atomic decompositions by more general ones, whose assumptions on the smoothness are relaxed and, nevertheless, we keep all the crucial information compared to smooth atomic decompositions. This modification appears in [26], where Triebel and Winkeloß suggested the use of these weaker conditions to define function spaces intrinsically on domains.

In this direction, recently Schneider and Vybíral in [23] and Scharf in [22] proved some results about non-smooth atoms for the classical Besov space $B_{p,q}^s(\mathbb{R}^n)$, but using different approaches. For the case of function spaces with variable exponents this is the first work concerning non-smooth atomic decompositions. Our approach follows Scharf in [22] and generalizes and extends the smooth atomic decomposition results previously referred.

In Section 4, we use the non-smooth atomic representation theorem to deal with pointwise multipliers in the function spaces $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$. Our result covers the corresponding results obtained in [23] and [22] for classical Besov spaces $B_{p,q}^s(\mathbb{R}^n)$.

2. Notation and definitions

We shall adopt the following general notation: \mathbb{N} denotes the set of all natural numbers, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, \mathbb{Z} denotes the set of integers, \mathbb{R}^n for $n \in \mathbb{N}$ denotes the n -dimensional real Euclidean space with $|x|$, for $x \in \mathbb{R}^n$, denoting the Euclidean norm of x . For a real number a , let $a_+ := \max(a, 0)$.

If $s \in \mathbb{R}$, then there are uniquely determined $[s]^- \in \mathbb{Z}$ and $\{s\}^+ \in (0, 1]$ with $s = [s]^- + \{s\}^+$.

Definition 2.1. Let $s > 0$. Then the Hölder space with index s is defined as

$$\mathcal{C}^s = \left\{ f \in C^{[s]^-} : \|f\|_{\mathcal{C}^s} := \sum_{|\alpha| \leq [s]^-} \sup_{x \in \mathbb{R}^n} |D^\alpha f(x)| + \sum_{|\alpha| = [s]^-} \sup_{x, y \in \mathbb{R}^n, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^{\{s\}^+}} < \infty \right\}.$$

If $s = 0$, then we set $\mathcal{C}^0 = L_\infty$.

For $q \in (0, \infty]$, ℓ_q stands for the linear space of all complex sequences $f = (f_j)_{j \in \mathbb{N}_0}$ endowed with the quasi-norm

$$\|f\|_{\ell_q} = \left(\sum_{j=0}^{\infty} |f_j|^q \right)^{1/q},$$

with the usual modification if $q = \infty$. By c, c_1, c_2 , etc. we denote positive constants independent of appropriate quantities. For two non-negative expressions (i.e., functions or functionals) \mathcal{A}, \mathcal{B} , the symbol $\mathcal{A} \lesssim \mathcal{B}$ (or $\mathcal{A} \gtrsim \mathcal{B}$) means that $\mathcal{A} \leq c\mathcal{B}$ (or $c\mathcal{A} \geq \mathcal{B}$), for some $c > 0$. If $\mathcal{A} \lesssim \mathcal{B}$ and $\mathcal{A} \gtrsim \mathcal{B}$, we write $\mathcal{A} \sim \mathcal{B}$ and say that \mathcal{A} and \mathcal{B} are equivalent.

In order to define 2-microlocal Besov and Triebel–Lizorkin spaces with variable integrability, we start by recalling the definition of admissible weight sequences. We follow [13].

Definition 2.2. Let $\alpha \geq 0$ and $\alpha_1, \alpha_2 \in \mathbb{R}$ with $\alpha_1 \leq \alpha_2$. A sequence of non-negative measurable functions in \mathbb{R}^n $w = (w_j)_{j \in \mathbb{N}_0}$ belongs to the class $\mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ if the following conditions are satisfied:

(i) There exists a constant $c > 0$ such that

$$0 < w_j(x) \leq c w_j(y) (1 + 2^j |x - y|)^\alpha \quad \text{for all } j \in \mathbb{N}_0 \text{ and all } x, y \in \mathbb{R}^n.$$

(ii) For all $j \in \mathbb{N}_0$ it holds

$$2^{\alpha_1} w_j(x) \leq w_{j+1}(x) \leq 2^{\alpha_2} w_j(x) \quad \text{for all } x \in \mathbb{R}^n.$$

Such a system $(w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ is called admissible weight sequence.

Properties of admissible weights may be found in [11, Remark 2.4].

Before introducing the function spaces under consideration we still need to recall some notation. By $\mathcal{S}(\mathbb{R}^n)$ we denote the Schwartz space of all complex-valued rapidly decreasing infinitely differentiable functions on \mathbb{R}^n and by $\mathcal{S}'(\mathbb{R}^n)$ the dual space of all tempered distributions on \mathbb{R}^n . For $f \in \mathcal{S}'(\mathbb{R}^n)$ we denote by \widehat{f} the Fourier transform of f and by f^\vee the inverse Fourier transform of f .

Let $\varphi_0 \in \mathcal{S}(\mathbb{R}^n)$ be such that

$$\varphi_0(x) = 1 \quad \text{if } |x| \leq 1 \quad \text{and} \quad \text{supp } \varphi_0 \subset \{x \in \mathbb{R}^n : |x| \leq 2\}. \quad (2.1)$$

Now define $\varphi(x) := \varphi_0(x) - \varphi_0(2x)$ and set $\varphi_j(x) := \varphi(2^{-j}x)$ for all $j \in \mathbb{N}$. Then the sequence $(\varphi_j)_{j \in \mathbb{N}_0}$ forms a smooth dyadic partition of unity.

By $\mathcal{P}(\mathbb{R}^n)$ we denote the class of exponents, which are measurable functions $p : \mathbb{R}^n \rightarrow (c, \infty]$ for some $c > 0$. Let $p \in \mathcal{P}(\mathbb{R}^n)$. Then, $p^+ := \text{ess-sup}_{x \in \mathbb{R}^n} p(x)$, $p^- := \text{ess-inf}_{x \in \mathbb{R}^n} p(x)$ and $L_{p(\cdot)}(\mathbb{R}^n)$ is the variable exponent Lebesgue space, which consists of all measurable functions f such that for some $\lambda > 0$ the modular $\varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f/\lambda)$ is finite, where

$$\varrho_{L_{p(\cdot)}(\mathbb{R}^n)}(f) := \int_{\mathbb{R}_0^n} |f(x)|^{p(x)} dx + \text{ess-sup}_{x \in \mathbb{R}_\infty^n} |f(x)|.$$

Here \mathbb{R}_∞^n denotes the subset of \mathbb{R}^n where $p(x) = \infty$ and $\mathbb{R}_0^n = \mathbb{R}^n \setminus \mathbb{R}_\infty^n$. The Luxemburg norm of a function $f \in L_{p(\cdot)}(\mathbb{R}^n)$ is given by

$$\|f\|_{L_{p(\cdot)}(\mathbb{R}^n)} := \inf \left\{ \lambda > 0 : \varrho_{L_{p(\cdot)}(\mathbb{R}^n)} \left(\frac{f}{\lambda} \right) \leq 1 \right\}.$$

In order to define the mixed spaces $\ell_{q(\cdot)}(L_{p(\cdot)})$, we need to define another modular. For $p, q \in \mathcal{P}(\mathbb{R}^n)$ and a sequence $(f_\nu)_{\nu \in \mathbb{N}_0}$ of complex-valued Lebesgue measurable functions on \mathbb{R}^n , we define

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}(f_\nu) = \sum_{\nu=0}^{\infty} \inf \left\{ \lambda_\nu > 0 : \varrho_{p(\cdot)} \left(\frac{f_\nu}{\lambda_\nu^{1/q(\cdot)}} \right) \leq 1 \right\}. \quad (2.2)$$

If $q^+ < \infty$, then we can replace (2.2) by the simpler expression

$$\varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})}(f_\nu) = \sum_{\nu=0}^{\infty} \left\| |f_\nu|^{q(\cdot)} \mid L_{\frac{p(\cdot)}{q(\cdot)}} \right\|. \quad (2.3)$$

The (quasi-)norm in the $\ell_{q(\cdot)}(L_{p(\cdot)})$ spaces is defined as usual by

$$\|f_\nu \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\| = \inf \left\{ \mu > 0 : \varrho_{\ell_{q(\cdot)}(L_{p(\cdot)})} \left(\frac{f_\nu}{\mu} \right) \leq 1 \right\}. \quad (2.4)$$

For the sake of completeness, we state also the definition of the space $L_{p(\cdot)}(\ell_{q(\cdot)})$. At first, one just takes the norm $\ell_{q(\cdot)}$ of $(f_\nu(x))_{\nu \in \mathbb{N}_0}$ for every $x \in \mathbb{R}^n$ and then the $L_{p(\cdot)}$ -norm with respect to $x \in \mathbb{R}^n$, i.e.

$$\|f_\nu \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\| = \left\| \left(\sum_{\nu=0}^{\infty} |f_\nu(x)|^{q(x)} \right)^{1/q(x)} \mid L_{p(\cdot)}(\mathbb{R}^n) \right\|.$$

We introduce now the Hardy–Littlewood maximal operator \mathcal{M}_t , which is defined for a locally integrable function $f \in L_1^{loc}$ and for $0 < t \leq 1$ by

$$\mathcal{M}_t(f)(x) = \left(\sup_{x \in Q} \int_Q |f(y)|^t dy \right)^{1/t},$$

and $\mathcal{M}(f)(x) = \mathcal{M}_1(f)(x)$. In order to guarantee the boundedness of \mathcal{M}_t in $L_{p(\cdot)}(\mathbb{R}^n)$ for non-constant exponent $p(\cdot)$ we require certain regularity conditions on $p \in \mathcal{P}(\mathbb{R}^n)$. Let us define the important classes.

Definition 2.3. Let $g \in C(\mathbb{R}^n)$. We say that g is locally log-Hölder continuous, abbreviated $g \in C_{loc}^{\log}(\mathbb{R}^n)$, if there exists $c_{\log}(g) > 0$ such that

$$|g(x) - g(y)| \leq \frac{c_{\log}(g)}{\log(e + 1/|x - y|)} \quad \text{for all } x, y \in \mathbb{R}^n. \quad (2.5)$$

We say that g is globally log-Hölder continuous, abbreviated $g \in C^{\log}(\mathbb{R}^n)$, if g is locally log-Hölder continuous and there exists $g_\infty \in \mathbb{R}$ such that

$$|g(x) - g_\infty| \leq \frac{c_{\log}}{\log(e + |x|)} \quad \text{for all } x \in \mathbb{R}^n. \quad (2.6)$$

We use the notation $p \in \mathcal{P}^{\log}(\mathbb{R}^n)$ if $p \in \mathcal{P}(\mathbb{R}^n)$ and $1/p \in C^{\log}(\mathbb{R}^n)$. It was proved in [4] that the maximal operator \mathcal{M} is bounded in $L_{p(\cdot)}(\mathbb{R}^n)$ provided that $p \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and $1 < p^- \leq p^+ \leq \infty$.

The definitions of the spaces below were given in [14]. The restriction $q^+ < \infty$ in the F -case comes from the use of Lemma 3.8, which is essential for studying Triebel–Lizorkin spaces with variable exponents.

Definition 2.4. Let $(\varphi_j)_{j \in \mathbb{N}_0}$ be a partition of unity as above, $\mathbf{w} = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$.

(i) The space $B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ is defined as the collection of all $f \in \mathcal{S}'(\mathbb{R}^n)$ such that

$$\|f \mid B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)\|_\varphi := \|(w_j (\varphi_j \widehat{f})^\vee)_{j \in \mathbb{N}_0} \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\|$$

is finite.

(ii) If $p^+, q^+ < \infty$, then the space $F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ is defined as the collection of all $f \in \mathcal{S}'(\mathbb{R}^n)$ such that

$$\|f \mid F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)\|_\varphi := \|(w_j (\varphi_j \widehat{f})^\vee)_{j \in \mathbb{N}_0} \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\|$$

is finite.

Remark 2.5. These spaces include very well-known spaces. For $p = \text{const}$ and $w_j(x) = 2^{js}$ we get back to the classical Besov and Triebel–Lizorkin spaces $B_{p,q}^s(\mathbb{R}^n)$ and $F_{p,q}^s(\mathbb{R}^n)$.

Also the spaces of generalized smoothness are contained in this approach (see [8,18]) by taking

$$w_j(x) = 2^{js}\Psi(2^{-j}), \quad \text{or more general } w_j(x) = \sigma_j.$$

Here, $\{\sigma_j\}_{j \in \mathbb{N}_0}$ is an admissible sequence, which means that there exist $d_0, d_1 > 0$ with $d_0\sigma_j \leq \sigma_{j+1} \leq d_1\sigma_j$ and Ψ is a slowly varying function.

Moreover, these 2-microlocal spaces also cover the spaces of variable smoothness and integrability $B_{p(\cdot), q(\cdot)}^{s(\cdot)}$ and $F_{p(\cdot), q(\cdot)}^{s(\cdot)}$, introduced in [6] and [1]. If $s \in C_{\text{loc}}^{\log}(\mathbb{R}^n)$ (which is the standard condition on $s(\cdot)$), then $\mathbf{w} = (w_j(x))_{j \in \mathbb{N}_0} = (2^{js(x)})_{j \in \mathbb{N}_0}$ belongs to $\mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ with $\alpha_1 = s^-$, $\alpha_2 = s^+$ and $\alpha = c_{\log}(s)$, where $c_{\log}(s)$ is the constant for $s(\cdot)$ from (2.5).

For $p, q \in \mathcal{P}(\mathbb{R}^n)$, we put

$$\sigma_p := n \left(\frac{1}{p^-} - 1 \right)_+ \quad \text{and} \quad \sigma_{p,q} := n \left(\frac{1}{\min(1, p^-, q^-)} - 1 \right).$$

2.1. Smooth atomic decompositions of $B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ and $F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$

In [13] it was presented the atomic decomposition theorems for $F_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ and, with $q = \text{const}$, for $B_{p(\cdot), q}^{\mathbf{w}}(\mathbb{R}^n)$. The case of having both parameters p and q variable in the Besov scale was studied by Drihem in [7], where the atomic decomposition of Besov spaces with variable smoothness and integrability $B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$ was stated. Here we present the atomic decomposition for 2-microlocal Besov spaces $B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$, whose proof follows mainly from [7] using Remark 2 in [14].

At first, we shall introduce some notation. Let \mathbb{Z}^n stand for the lattice of all points in \mathbb{R}^n with integer-valued components, $Q_{\nu, m}$ denotes a cube in \mathbb{R}^n with sides parallel to the axes of coordinates, centered at $2^{-\nu}m = (2^{-\nu}m_1, \dots, 2^{-\nu}m_n)$ and with side length $2^{-\nu}$, where $m = (m_1, \dots, m_n) \in \mathbb{Z}^n$ and $\nu \in \mathbb{N}_0$. If Q is a cube in \mathbb{R}^n and $r > 0$ then rQ is the cube in \mathbb{R}^n concentric with Q and with side length r times the side length of Q . By $\chi_{\nu, m}$ we denote the characteristic function of the cube $Q_{\nu, m}$.

We define now smooth atoms, which are the building blocks for atomic decompositions.

Definition 2.6. Let $K, L \in \mathbb{N}_0$ and let $d > 1$. A K -times continuously differentiable complex-valued function $a \in C^K(\mathbb{R}^n)$ is called $[K, L]$ -atom centered at $Q_{\nu, m}$, for all $\nu \in \mathbb{N}_0$ and $m \in \mathbb{Z}^n$, if

- (i) $\text{supp } a \subset dQ_{\nu, m}$,
- (ii) $|D^\beta a(x)| \leq 2^{|\beta|\nu}$ for $|\beta| \leq K$,
- (iii) $\int_{\mathbb{R}^n} x^\beta a(x) dx = 0$ for $0 \leq |\beta| < L$ and $\nu \geq 1$.

Remark 2.7. If an atom a is centered at $Q_{\nu, m}$, i.e., if it fulfills (i), then we denote it by $a_{\nu, m}$. If $L = 0$ or $\nu = 0$, then no moment conditions (iii) are required.

We shall now introduce the sequence spaces $b_{p(\cdot), q(\cdot)}^w$ and $f_{p(\cdot), q(\cdot)}^w$, whose use will become clear in the following.

Definition 2.8. Let $w = (w_\nu)_{\nu \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}(\mathbb{R}^n)$.

- (i) The sequence space $b_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ consists of those complex-valued sequences $\lambda = (\lambda_{\nu, m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ such that

$$\|\lambda \mid b_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)\| := \left\| \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu, m}| w_\nu(2^{-\nu} m) \chi_{\nu, m} \right)_{\nu \in \mathbb{N}_0} \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\|$$

is finite.

- (ii) If $p^+, q^+ < \infty$, then the sequence space $f_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ consists of those complex-valued sequences $\lambda = (\lambda_{\nu, m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ such that

$$\|\lambda \mid f_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)\| := \left\| \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu, m}| w_\nu(2^{-\nu} m) \chi_{\nu, m} \right)_{\nu \in \mathbb{N}_0} \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n)) \right\|$$

is finite.

We present now the atomic decomposition results for the scale of 2-microlocal spaces with variable p and q . While for the Triebel–Lizorkin space $F_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ the result was stated in [13], the equivalent result for Besov spaces $B_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ is new. However we do not present the proof here, since the idea goes back to [7] where the atomic decomposition was stated for $B_{p(\cdot), q(\cdot)}^{s(\cdot)}(\mathbb{R}^n)$, see also Remark 2 in [14].

Theorem 2.9. (See Corollary 5.6 in [13].) Let $w = (w_\nu)_{\nu \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ with $0 < p^- \leq p^+ < \infty$ and $0 < q^- \leq q^+ < \infty$. Furthermore, let $d > 1$, $K, L \in \mathbb{N}_0$ with $K > \alpha_2$ and $L > \sigma_{p, q} - \alpha_1$ be fixed. Then every $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $F_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ if and only if it can be represented as

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu, m} a_{\nu, m}, \quad \text{convergence being in } \mathcal{S}'(\mathbb{R}^n), \quad (2.7)$$

for $(a_{\nu, m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ $[K, L]$ -atoms according to Definition 2.6 and $\lambda \in f_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$. Moreover,

$$\|f \mid F_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)\| \sim \inf \|\lambda \mid f_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)\|,$$

where the infimum is taken over all possible representations of f .

Theorem 2.10. Let $w = (w_\nu)_{\nu \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$. Furthermore, let $d > 1$, $K, L \in \mathbb{N}_0$ with $K > \alpha_2$ and $L > \sigma_p - \alpha_1 + c_{\log}(1/q)$ be fixed. Then every $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $B_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ if and only if it can be represented as

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu, m} a_{\nu, m}, \quad \text{convergence being in } \mathcal{S}'(\mathbb{R}^n), \quad (2.8)$$

for $(a_{\nu, m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ $[K, L]$ -atoms according to Definition 2.6 and $\lambda \in b_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$. Moreover,

$$\|f\|_{B_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)} \sim \inf \|\lambda\|_{b_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)},$$

where the infimum is taken over all possible representations of f .

3. Non-smooth atomic decompositions

3.1. Non-smooth atoms

At first, we present the concept of *non-smooth atoms*, slightly adapted from [22] to our scale. Note that the usual parameters K and L are now non-negative real numbers instead of natural numbers.

Definition 3.1. Let $K, L \geq 0$, $d > 1$ and $c > 0$. A function $a : \mathbb{R}^n \rightarrow \mathbb{R}$ is called a non-smooth $[K, L]$ -atom centered at $Q_{\nu, m}$, for all $\nu \in \mathbb{N}_0$ and $m \in \mathbb{Z}^n$, if

- (i) $\text{supp } a \subset dQ_{\nu, m}$,
- (ii) $\|a(2^{-\nu} \cdot) | \mathcal{C}^K(\mathbb{R}^n)\| \leq c$,
- (iii) and for every $\psi \in \mathcal{C}^L(\mathbb{R}^n)$ it holds

$$\left| \int_{dQ_{\nu, m}} \psi(x) a(x) dx \right| \leq c 2^{-\nu(L+n)} \|\psi\|_{\mathcal{C}^L(\mathbb{R}^n)}.$$

Remark 3.2.

- (a) As in the smooth case, if $L = 0$, then condition (iii) can be ignored since it follows from conditions (i) and (ii) with $K = 0$. If $K = 0$, then by Definition 2.1 we only require a to be suitable bounded.
- (b) The modification of condition (ii) here was suggested in [26] (with some minor adjustments) and one can see that the usual formulation as in Definition 2.6(ii) follows from this one if K is a natural number, since $C^K(\mathbb{R}^n) \hookrightarrow \mathcal{C}^K(\mathbb{R}^n)$.
- (c) Regarding condition (iii), the modification here was suggested by Skrzypczak in [24] for natural numbers $L + 1$ (replacing $\mathcal{C}^L(\mathbb{R}^n)$ by $C^L(\mathbb{R}^n)$). Here, as in [22], we extended this definition to general positive numbers L .

3.2. Local means

Let us recall the characterization of $B_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot), q(\cdot)}^w(\mathbb{R}^n)$ by local means, proved in Theorem 6 of [14] and Theorem 2.2 of [12], respectively. For each system $(\Psi_k)_{k \in \mathbb{N}_0} \subset \mathcal{S}(\mathbb{R}^n)$, for each distribution $f \in \mathcal{S}'(\mathbb{R}^n)$ and to each number $a > 0$, we define the Peetre maximal operator by

$$(\Psi_k^* f)_a(x) = \sup_{y \in \mathbb{R}^n} \frac{|(\Psi_k * f)(y)|}{1 + |2^k(y - x)|^a}, \quad \text{where } k \in \mathbb{N}_0 \text{ and } x \in \mathbb{R}^n.$$

We start with two given functions $\psi_0, \psi \in \mathcal{S}(\mathbb{R}^n)$. We define

$$\psi_j(x) = \psi(2^{-j}x), \quad \text{for } x \in \mathbb{R}^n \text{ and } j \in \mathbb{N}. \quad (3.1)$$

Furthermore, for all $k \in \mathbb{N}_0$ we write $\Psi_k = \hat{\psi}_k$.

Remark 3.3. Before presenting the results, we would like to highlight that a careful look into the proof of the Theorem 6 in [14] shows that the condition $a > \frac{n+c_{\log}(1/q)}{p^-} + \alpha$ should be replaced by

$$a > \frac{n}{p^-} + c_{\log}(1/q) + \alpha.$$

Proposition 3.4. Let $\mathbf{w} = (w_k)_{k \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$, $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$ and let $a > 0$, $R \in \mathbb{N}_0$ with $R > \alpha_2$. Further, let ψ_0, ψ belong to $\mathcal{S}(\mathbb{R}^n)$ with

$$D^\beta \psi(0) = 0, \quad \text{for } 0 \leq |\beta| < R, \quad (3.2)$$

and

$$|\psi_0(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : |x| < \varepsilon\} \quad (3.3)$$

$$|\psi(x)| > 0 \quad \text{on } \{x \in \mathbb{R}^n : \varepsilon/2 < |x| < 2\varepsilon\} \quad (3.4)$$

for some $\varepsilon > 0$.

(i) For $a > \frac{n}{p^-} + c_{\log}(1/q) + \alpha$ and for all $f \in \mathcal{S}'(\mathbb{R}^n)$ we have

$$\|f \mid B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)\| \sim \|(\Psi_k * f)w_k \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\| \sim \|(\Psi_k^* f)_a w_k \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\|$$

(ii) If $p^+, q^+ < \infty$, then for $a > \frac{n}{\min(p^-, q^-)} + \alpha$ and for all $f \in \mathcal{S}'(\mathbb{R}^n)$ we have

$$\|f \mid F_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)\| \sim \|(\Psi_k * f)w_k \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\| \sim \|(\Psi_k^* f)_a w_k \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\|$$

Remark 3.5.

- (a) This result shows that the definition of the 2-microlocal spaces of variable integrability given in Definition 2.4 is independent of the resolution of unity used.
- (b) The conditions (3.2) are usually called *moment conditions*, while (3.3) and (3.4) are the so-called *Tauberian conditions*.
- (c) If $R = 0$, then no moment conditions (3.2) on ψ are required.
- (d) The notation $c_{\log}(1/q)$ stands for the constant from (2.5) with $1/q(\cdot)$.

3.2.1. Helpful lemmas

Before presenting the main theorem of this section, we provide some technical lemmas which will be useful later on. The first result is an adaptation of Lemma 3.8 in [22] and it shows that local means are related to non-smooth $[K, L]$ -atoms.

Lemma 3.6. Let $j \in \mathbb{N}_0$. If $\psi_j = \psi(2^{-j}\cdot)$ is a local mean as in Proposition 3.4, then $2^{-jn}\psi_j$ is a non-smooth $[K, L]$ -atom centered at $Q_{j,0}$ for arbitrary large $K > 0$ and for $L \leq R + 1$ with R from (3.2).

The next result can be obtained following the steps of the proof of Theorem 3.12 in [22] with our normalization. The idea is to use the fact that not only $a_{\nu,m}$ but also Ψ_j can be understood as atoms in the sense of Definition 3.1. In this way, we can use estimates of the type (ii) and (iii) of Definition 3.1 in both functions and get this result.

Lemma 3.7. *Let $j \in \mathbb{N}_0$, $\psi_j = \psi(2^{-j}\cdot)$ be a local mean as in Proposition 3.4 and let $(a_{\nu,m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ be non-smooth $[K, L]$ -atoms. Then*

$$\left| \int_{\mathbb{R}^n} \Psi_j(y) a_{\nu,m}(x-y) dy \right| \leq c 2^{-(j-\nu)K} \chi(c Q_{\nu,m})(x), \quad \text{for } j \geq \nu$$

and

$$\left| \int_{\mathbb{R}^n} \Psi_j(x-y) a_{\nu,m}(y) dy \right| \leq c 2^{-(\nu-j)(L+n)} \chi(c 2^{\nu-j} Q_{\nu,m})(x), \quad \text{for } j < \nu.$$

Since the maximal operator is of no use in the case when q is a variable function (see [1] and [6]), we have to work with another tool. We use convolution inequalities from [6] and [14] for $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$, respectively. Therefore, we introduce the functions

$$\eta_{\nu,R}(x) = \frac{2^{n\nu}}{(1 + 2^\nu |x|)^R},$$

for $\nu \in \mathbb{N}_0$ and $R > 0$.

Lemma 3.8. (See Theorem 3.2 in [6].) *Let $p, q \in C^{\log}(\mathbb{R}^n)$ with $1 < p^- \leq p^+ < \infty$ and $1 < q^- \leq q^+ < \infty$. Then the inequality*

$$\|\eta_{\nu,R} * f_\nu \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\| \leq c \|f_\nu \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\|$$

holds for every sequence $(f_\nu)_{\nu \in \mathbb{N}_0}$ of $L_1^{\text{loc}}(\mathbb{R}^n)$ functions and constant $R > n$.

Lemma 3.9. (See Lemma 10 in [14].) *Let $p, q \in \mathcal{P}(\mathbb{R}^n)$ with $p(\cdot) \geq 1$. For all $R > n + c_{\log}(1/q)$, there exists a constant $c > 0$ such that for all sequences $(f_\nu)_{\nu \in \mathbb{N}_0} \in \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))$ it holds*

$$\|\eta_{\nu,R} * f_\nu \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\| \leq c \|f_\nu \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\|.$$

Lemma 3.10. (See Lemma 9 in [14].) *Let $p, q \in \mathcal{P}(\mathbb{R}^n)$ and $\delta > 0$. Let $(g_k)_{k \in \mathbb{Z}}$ be a sequence of non-negative measurable functions on \mathbb{R}^n and denote*

$$G_\nu(x) = \sum_{k \in \mathbb{Z}} 2^{-|\nu-k|\delta} g_k(x), \quad \text{for } x \in \mathbb{R}^n \text{ and } \nu \in \mathbb{Z}.$$

Then there exist constants $C_1, C_2 > 0$, depending on $p(\cdot)$, $q(\cdot)$ and δ , such that

$$\|G_\nu \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\| \leq C_1 \|g_k \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))\|$$

and

$$\|G_\nu \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\| \leq C_2 \|g_k \mid L_{p(\cdot)}(\ell_{q(\cdot)}(\mathbb{R}^n))\|.$$

Remark 3.11. Naturally, [Lemma 3.10](#) holds true also if the indices k and ν run only over natural numbers.

Lemma 3.12. (See Lemma 5.5 in [\[13\]](#).) Let $0 < t \leq 1, j, \nu \in \mathbb{N}_0$ and $(h_{\nu,m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ be non-negative real numbers. Then we have for $R > 0$ and $x \in \mathbb{R}^n$ that

$$\sum_{m \in \mathbb{Z}^n} h_{\nu,m} (1 + 2^j |x - 2^{-\nu} m|)^{-R} \leq c \max(1, 2^{(\nu-j)R}) \left(\left[\eta_{\nu,Rt} * \left(\sum_{m \in \mathbb{Z}^n} h_{\nu,m}^t \chi_{\nu,m}(\cdot) \right) \right] (x) \right)^{1/t}.$$

3.3. A more general atomic representation theorem

In this section we shall prove the more general atomic decomposition theorem. We will follow the approach of the proof of Theorem 3.12 in [\[22\]](#). But first we will prove the convergence of the atomic series in $\mathcal{S}'(\mathbb{R}^n)$.

Theorem 3.13. Let $w = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{log}(\mathbb{R}^n)$. Let $K, L \geq 0$ with $K > \alpha_2$ and $L > \sigma_p - \alpha_1$. Then

$$\sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m} \quad (3.5)$$

converges unconditionally in $\mathcal{S}'(\mathbb{R}^n)$, where $a_{\nu,m}$ are non-smooth $[K, L]$ -atoms located at $Q_{\nu,m}$ and $\lambda \in b_{p(\cdot), q(\cdot)}^w$ or $\lambda \in f_{p(\cdot), q(\cdot)}^w$.

Proof. According to [\[1\]](#) and [\[13\]](#) (see also the explanation before Proposition 3.9 in [\[13\]](#)), for $t < \min(1, p^-)$ we have the following sequences of embeddings

$$b_{p(\cdot), q(\cdot)}^w \hookrightarrow b_{\frac{p(\cdot)}{t}, q(\cdot)}^{\varrho} \hookrightarrow b_{\frac{p(\cdot)}{t}, \infty}^{\varrho}$$

and

$$f_{p(\cdot), q(\cdot)}^w \hookrightarrow f_{\frac{p(\cdot)}{t}, q(\cdot)}^{\varrho} \hookrightarrow b_{\frac{p(\cdot)}{t}, \infty}^{\varrho},$$

where $\varrho_j(x) = w_j(x) 2^{-j \frac{t}{p(x)} \sigma_t}$. Since $2^{-j \frac{t}{p(x)} \sigma_t} =: 2^{-js(x) \sigma_t}$ is again a 2-microlocal weight sequence with $s(\cdot) \in C^{log}(\mathbb{R}^n)$ and $0 \leq s(x) < 1$, then ϱ is also an admissible weight sequence and it can be shown that $\varrho \in \mathcal{W}_{\alpha_1 - \sigma_t, \alpha_2}^\beta(\mathbb{R}^n)$ with $\beta \geq 0$ large enough. In fact, it can be shown that $\beta = \alpha + t \sigma_t c_{log}(1/p)$, where the constant $c_{log}(1/p)$ comes from [\(2.5\)](#).

Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$. We can assume that $\lambda \in b_{\frac{p(\cdot)}{t}, \infty}^{\varrho}$, having in mind the above embeddings. From the conditions (i) and (iii) of the [Definition 3.1](#) we obtain

$$\left| \int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m}(x) \varphi(x) \right| \leq c 2^{-\nu(L+n)} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \|\varphi(\cdot) \psi(2^\nu \cdot - m)\| \mathcal{C}^L(\mathbb{R}^n), \quad (3.6)$$

where $\psi \in C^\infty(\mathbb{R}^n)$, $\psi(x) = 1$ for $x \in dQ_{0,0}$ and $\text{supp } \psi \subset (d+1)Q_{0,0}$. Since $\varphi \in \mathcal{S}(\mathbb{R}^n)$, we can estimate the norm in [\(3.6\)](#) from above in the following way:

$$\|\varphi(\cdot) \psi(2^\nu \cdot - m)\| \mathcal{C}^L(\mathbb{R}^n) \leq C_M (1 + |2^{-\nu} m|)^{-M} \sim C_M (1 + |y|)^{-M}, \quad y \in Q_{\nu,m},$$

where $M \in \mathbb{N}_0$ is at our disposal and C_M does not depend on ν and m . According to the properties of the admissible weight sequence [\(3.6\)](#) becomes

$$\begin{aligned}
\left| \int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m}(x) \varphi(x) dx \right| &\leq c 2^{-\nu(L+n)} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(y) (1+|y|)^{-M} \chi_{\nu,m}(y) \varrho_{\nu}(y)^{-1} \\
&\leq c 2^{-\nu(L+n)} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) (1+2^{\nu}|y-2^{-\nu}m|)^{-\beta} 2^{\nu(\alpha_1-\sigma_t)} (1+|y|)^{-M+\beta} \chi_{\nu,m}(y) \\
&\sim c 2^{-\nu(L+\alpha_1-\sigma_t)} \int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) (1+2^{\nu}|x-2^{-\nu}m|)^{-\beta} (1+|x|)^{-M+\beta} \chi_{\nu,m}(x) dx.
\end{aligned}$$

We estimate now the integral and from Hölder's inequality for $L_{p(\cdot)}(\mathbb{R}^n)$ with $p(\cdot)/t > 1$ and choosing M large enough we get

$$\begin{aligned}
\int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) (1+2^{\nu}|x-2^{-\nu}m|)^{-\beta} (1+|x|)^{-M+\beta} \chi_{\nu,m}(x) dx \\
\leq c \left\| \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) (1+2^{\nu}|x-2^{-\nu}m|)^{-\beta} \chi_{\nu,m}(\cdot) \right\|_{L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n)}.
\end{aligned}$$

Finally, we use Lemma 3.7 of [13] with $\beta > \frac{n}{t}$ and the boundedness of the maximal operator and obtain

$$\begin{aligned}
\left| \int_{\mathbb{R}^n} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m}(x) \varphi(x) \right| &\leq c 2^{-\nu(L+\alpha_1-\sigma_t)} \left\| \mathcal{M} \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) \chi_{\nu,m}(\cdot) \right) \right\|_{L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n)} \\
&\leq c 2^{-\nu(L+\alpha_1-\sigma_t)} \left\| \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| \varrho_{\nu}(2^{-\nu}m) \chi_{\nu,m}(\cdot) \right\|_{L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n)}. \quad (3.7)
\end{aligned}$$

Summing up over $\nu \in \mathbb{N}_0$ and since $L > \sigma_t - \alpha_1 > \sigma_p - \alpha_1$, we arrive at

$$\sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \left| \int_{\mathbb{R}^n} \lambda_{\nu,m} a_{\nu,m}(x) \varphi(x) dx \right| \leq C \|\lambda\|_{b_{\frac{p(\cdot)}{t},\infty}^{\mathbf{e}}},$$

from which the absolute convergence of (3.5) and the unconditional convergence in $\mathcal{S}'(\mathbb{R}^n)$ follow. \square

Now we state the atomic decomposition result.

Theorem 3.14. Let $\mathbf{w} = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^{\alpha}(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{\log}(\mathbb{R}^n)$.

- (i) Let $K, L \geq 0$ with $K > \alpha_2$ and $L > \sigma_p - \alpha_1 + c_{\log}(1/q)$. Then $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$ if and only if it can be represented as

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m}, \quad \text{convergence being in } \mathcal{S}'(\mathbb{R}^n), \quad (3.8)$$

for $(a_{\nu,m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ non-smooth $[K, L]$ -atoms according to Definition 3.1 and $\lambda \in b_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)$. Moreover,

$$\|f\|_{B_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)} \sim \inf \|\lambda\|_{b_{p(\cdot), q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n)},$$

where the infimum is taken over all possible representations of f .

- (ii) Let $K, L \geq 0$ with $K > \alpha_2$ and $L > \sigma_{p,q} - \alpha_1$. If $p^+, q^+ < \infty$, then $f \in \mathcal{S}'(\mathbb{R}^n)$ belongs to $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ if and only if it can be represented as

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m}, \quad \text{convergence being in } \mathcal{S}'(\mathbb{R}^n),$$

for $(a_{\nu,m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ non-smooth $[K, L]$ -atoms according to [Definition 3.1](#) and $\lambda \in f_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$. Moreover,

$$\|f\|_{F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)} \sim \inf \|\lambda\|_{f_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)},$$

where the infimum is taken over all possible representations of f .

Proof. We rely on the proof of Theorem 3.12 in [\[22\]](#).

There are two directions we shall prove. Observing that every smooth $[K, L]$ -atom is a non-smooth $[K, L]$ -atom one direction follows directly from [Theorems 2.10 and 2.9](#) (see the proof of Theorem 3.12 in [\[22\]](#)). We will focus on the other direction, which is the main part of the proof. We need to show that, even with weaker conditions on the atoms, we still get an element of $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ or $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ when considering a linear combination of these non-smooth atoms. For this, we will use the equivalent characterization by local means given in [Proposition 3.4](#). As usual, we divide the following summation in dependence on $j \in \mathbb{N}_0$ into two parts

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m} = \sum_{\nu=0}^j \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m} + \sum_{\nu=j+1}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu,m} a_{\nu,m} = f_j + f^j.$$

We have

$$\begin{aligned} \|f\|_{B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)} &\leq c \left(\|f_j\|_{B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)} + \|f^j\|_{B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)} \right) \\ &= c \|(\Psi_j * f_j)w_j\|_{\ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))} + c \|(\Psi_j * f^j)w_j\|_{\ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n))} \\ &= \text{I} + \text{II}. \end{aligned} \tag{3.9}$$

In both parts of the summation, the crucial part is the estimate of

$$\left| \int_{\mathbb{R}^n} \Psi_j(y) a_{\nu,m}(x-y) dy \right|,$$

which we already have from [Lemma 3.7](#). As stated before, we use that not only $a_{\nu,m}$ but also Ψ_j can be interpreted as non-smooth atoms and admit estimates as (ii) and (iii) in [Definition 3.1](#).

Let us first consider the case $j \geq \nu$. We have, from [Lemma 3.7](#), the estimate

$$\left| \int_{\mathbb{R}^n} \Psi_j(y) a_{\nu,m}(x-y) dy \right| \leq c 2^{-(j-\nu)K} \chi(c Q_{\nu,m})(x),$$

and from the properties of the weight sequence in [Definition 2.2](#),

$$w_j(x) \leq C 2^{(j-\nu)\alpha_2} w_\nu(2^{-\nu}m)(1 + 2^\nu|x - 2^{-\nu}m|)^\alpha.$$

Putting both together, we have

$$|(\Psi_j * f_j)(x) w_j(x)| \leq c \sum_{\nu=0}^j \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| 2^{-(j-\nu)(K-\alpha_2)} w_\nu(2^{-\nu}m) \chi(cQ_{\nu,m})(x),$$

using that $|x - 2^{-\nu}m| \lesssim 2^{-\nu}$ for $x \in cQ_{\nu,m}$. Further, using [Lemma 3.10](#) with $\delta = K - \alpha_2 > 0$ we get

$$\begin{aligned} \text{I} &\leq c \left\| \sum_{\nu=0}^j 2^{-(j-\nu)(K-\alpha_2)} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| w_\nu(2^{-\nu}m) \chi(cQ_{\nu,m}) \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \\ &\leq c' \left\| \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| w_\nu(2^{-\nu}m) \chi(cQ_{\nu,m}) \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \\ &= c' \left\| \lambda \mid b_{p(\cdot),q(\cdot)}^{\mathbf{w}}(\mathbb{R}^n) \right\|. \end{aligned}$$

Now, let $j < \nu$. [Lemma 3.7](#) gives us the estimate

$$\left| \int_{\mathbb{R}^n} \Psi_j(x-y) a_{\nu,m}(y) dy \right| \leq c 2^{-(\nu-j)(L+n)} \chi(c2^{\nu-j}Q_{\nu,m})(x)$$

and using the properties of the weight sequence again, we get

$$w_j(x) \leq C 2^{-(\nu-j)\alpha_1} w_\nu(2^{-\nu}m) (1 + 2^j|x - 2^{-\nu}m|)^\alpha.$$

Note that the following assertion is true: for every $x \in \mathbb{R}^n$ and every $M > 0$, there exists a constant $C > 0$ such that

$$\chi(c2^{\nu-j}Q_{\nu,m})(x) \leq C (1 + 2^j|x - 2^{-\nu}m|)^{-M}.$$

Combining the last three estimates, we have

$$|(\Psi_j * f^j)(x) w_j(x)| \leq c \sum_{\nu=j+1}^{\infty} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| 2^{-(\nu-j)(L+n+\alpha_1)} w_\nu(2^{-\nu}m) (1 + 2^j|x - 2^{-\nu}m|)^{\alpha-M}.$$

We can now estimate II. Let $0 < t \leq \min(1, p^-)$. By [Lemma 3.12](#) with $M - \alpha > 0$ we get

$$\begin{aligned} \text{II} &\leq c \left\| \sum_{\nu=j+1}^{\infty} \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| 2^{-(\nu-j)(L+n+\alpha_1)} w_\nu(2^{-\nu}m) (1 + 2^j|x - 2^{-\nu}m|)^{\alpha-M} \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \\ &\leq c \left\| \sum_{\nu=j+1}^{\infty} 2^{-(\nu-j)\delta} \left[\eta_{\nu,(M-\alpha)t} * \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}|^t w_\nu^t(2^{-\nu}m) \chi_{\nu,m}(\cdot) \right) \right]^{1/t} \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \end{aligned}$$

with $\delta := L + n + \alpha_1 - M + \alpha$ and by [Lemma 3.10](#) with $\delta \geq 0$ we obtain

$$\begin{aligned} &\leq c' \left\| \left[\eta_{\nu,(M-\alpha)t} * \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}|^t w_\nu^t(2^{-\nu}m) \chi_{\nu,m}(\cdot) \right) \right]^{1/t} \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \\ &= c' \left\| \eta_{\nu,(M-\alpha)t} * \left(\sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}|^t w_\nu^t(2^{-\nu}m) \chi_{\nu,m}(\cdot) \right) \mid \ell_{\frac{q(\cdot)}{t}}(L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n)) \right\|^{1/t}. \end{aligned}$$

This can be further estimated using [Lemma 3.9](#) with $M - \alpha > \frac{n}{t} + c_{\log}(1/q)$ (note that we are applying the result for $\ell_{\frac{q(\cdot)}{t}}(L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n))$)

$$\begin{aligned} &\leq c'' \left\| \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}|^t w_{\nu}^t(2^{-\nu}m) \chi_{\nu,m}(\cdot) \mid \ell_{\frac{q(\cdot)}{t}}(L_{\frac{p(\cdot)}{t}}(\mathbb{R}^n)) \right\|^{1/t} \\ &= c'' \left\| \sum_{m \in \mathbb{Z}^n} |\lambda_{\nu,m}| w_{\nu}(2^{-\nu}m) \chi_{\nu,m}(\cdot) \mid \ell_{q(\cdot)}(L_{p(\cdot)}(\mathbb{R}^n)) \right\| \\ &= c'' \left\| \lambda \mid b_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n) \right\|. \end{aligned}$$

In the F -case we do the same estimates as above, use the second part of [Lemma 3.10](#) and apply, instead of [Lemma 3.9](#), [Lemma 3.8](#) for $L_{\frac{p(\cdot)}{t}}(\ell_{\frac{q(\cdot)}{t}})$ with $t < \min(1, p^-, q^-)$ and $R = (M - \alpha)t > n$. \square

4. Pointwise multipliers

Let φ be a bounded function on \mathbb{R}^n . The question is under which conditions the mapping $f \mapsto \varphi \cdot f$ makes sense and generates a bounded operator in a given space $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ or $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$.

For the classical spaces $B_{p,q}^s(\mathbb{R}^n)$ and $F_{p,q}^s(\mathbb{R}^n)$, Triebel studied this problem in Section 4.2 of [\[25\]](#), where two different approaches were followed: via atoms and via local means. The first idea of taking an atomic decomposition of f required the non-existence of moment conditions as in [Definition 2.6\(iii\)](#) since the moment conditions are in general destroyed by multiplication with φ . A more general result was then obtained by Triebel using local means. A good overview on pointwise multipliers in constant exponent spaces $B_{p,q}^s(\mathbb{R}^n)$ and $F_{p,q}^s(\mathbb{R}^n)$ can be found in Chapter 4 in [\[21\]](#).

Recently, Scharf has shown in [\[22\]](#) that it is possible to get a very general result on pointwise multipliers using atomic decomposition but now with the non-smooth atoms.

Our aim in this section is to extend this result for variable exponent spaces $B_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$ and $F_{p(\cdot),q(\cdot)}^w(\mathbb{R}^n)$. Following [\[22\]](#), we take an atomic decomposition of f as in [Theorem 3.14](#), multiply it by φ and prove that the resulting sum is still a sum of non-smooth atoms. We start by referring two helpful results.

Lemma 4.1. (See [Lemma 4.2](#) in [\[22\]](#).) Let $s \geq 0$. There exists a constant $c > 0$ such that for all $f, g \in \mathcal{C}^s(\mathbb{R}^n)$, the product $f \cdot g$ belongs to $\mathcal{C}^s(\mathbb{R}^n)$ and it holds

$$\|f \cdot g \mid \mathcal{C}^s(\mathbb{R}^n)\| \leq c \|f \mid \mathcal{C}^s(\mathbb{R}^n)\| \cdot \|g \mid \mathcal{C}^s(\mathbb{R}^n)\|.$$

The next result shows that the product of a non-smooth $[K, L]$ -atom with a function $\varphi \in \mathcal{C}^\rho(\mathbb{R}^n)$ is still a non-smooth $[K, L]$ -atom. It is a slight normalization of [Lemma 4.3](#) in [\[22\]](#) and so the proof will not be presented here.

Lemma 4.2. There exists a constant c with the following property: for all $\nu \in \mathbb{N}_0$, $m \in \mathbb{Z}^n$, all non-smooth $[K, L]$ -atoms $a_{\nu,m}$ with support in $dQ_{\nu,m}$ and all $\varphi \in \mathcal{C}^\rho(\mathbb{R}^n)$ with $\rho \geq \max(K, L)$, the product

$$c \|\varphi \mid \mathcal{C}^\rho(\mathbb{R}^n)\|^{-1} \cdot \varphi \cdot a_{\nu,m}$$

is a non-smooth $[K, L]$ -atom with support in $dQ_{\nu,m}$.

We are now in the position of proving the main result.

Theorem 4.3. Let $\mathbf{w} = (w_j)_{j \in \mathbb{N}_0} \in \mathcal{W}_{\alpha_1, \alpha_2}^\alpha(\mathbb{R}^n)$ and $p, q \in \mathcal{P}^{log}(\mathbb{R}^n)$.

(i) Let $\rho > \max(\alpha_2, \sigma_p - \alpha_1 + c_{log}(1/q))$. Then there exists a positive number c such that

$$\|\varphi \cdot f\|_{B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)} \leq c \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \cdot \|f\|_{B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)}$$

for all $\varphi \in \mathcal{C}^\rho(\mathbb{R}^n)$ and all $f \in B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$.

(ii) Let $p^+, q^+ < \infty$ and $\rho > \max(\alpha_2, \sigma_{p, q} - \alpha_1)$. Then there exists a positive number c such that

$$\|\varphi \cdot f\|_{F_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)} \leq c \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \cdot \|f\|_{F_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)}$$

for all $\varphi \in \mathcal{C}^\rho(\mathbb{R}^n)$ and all $f \in F_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$.

Proof. We will prove (i), since (ii) follows similarly. Let $f \in B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$. By Theorem 3.14, there exist $\lambda \in b_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$ and non-smooth $[K, L]$ -atoms $(a_{\nu, m})_{\nu \in \mathbb{N}_0, m \in \mathbb{Z}^n}$ such that f can be represented as

$$f = \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu, m} a_{\nu, m}$$

and

$$\|\lambda\|_{b_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)} \leq c \|f\|_{B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)},$$

where the constant c is independent of $f \in B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$. By Lemma 4.2 we know that

$$\begin{aligned} \varphi \cdot f &= \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \lambda_{\nu, m} (\|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)}^{-1} \cdot \varphi \cdot a_{\nu, m}) \\ &= \sum_{\nu=0}^{\infty} \sum_{m \in \mathbb{Z}^n} \mu_{\nu, m} (\|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)}^{-1} \cdot \varphi \cdot a_{\nu, m}) \end{aligned}$$

is still a linear combination of non-smooth $[K, L]$ -atoms with coefficients $\mu_{\nu, m} := \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \lambda_{\nu, m}$, which means that we obtained an atomic decomposition for the product $\varphi \cdot f$. Using now the other direction of the Theorem 3.14, we know that $\varphi \cdot f$ belongs to $B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)$ and it holds

$$\|\varphi \cdot f\|_{B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)} \leq c \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \cdot \|\lambda\|_{b_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)} \leq c' \|\varphi\|_{\mathcal{C}^\rho(\mathbb{R}^n)} \cdot \|f\|_{B_{p(\cdot), q(\cdot)}^\mathbf{w}(\mathbb{R}^n)},$$

which completes the proof. \square

References

- [1] A. Almeida, P. Hästö, Besov spaces with variable smoothness and integrability, J. Funct. Anal. 258 (5) (2010) 1628–1655.
- [2] J.M. Bony, Second microlocalization and propagation of singularities for semilinear hyperbolic equations, in: Hyperbolic Equations and Related Topics, Katata/Kyoto, 1984, Academic Press, Boston, MA, 1986, pp. 11–46.
- [3] L. Diening, Maximal function on generalized Lebesgue spaces $L^{p(\cdot)}$, Math. Inequal. Appl. 7 (2) (2004) 245–253.
- [4] L. Diening, P. Harjuletho, P. Hästö, Y. Mizuta, T. Shimomura, Maximal functions in variable exponent spaces: limiting cases of the exponent, Ann. Acad. Sci. Fenn. Math. 34 (2) (2009) 503–522.
- [5] L. Diening, P. Harjuletho, P. Hästö, M. Růžička, Lebesgue and Sobolev Spaces with Variable Exponents, Springer-Verlag, 2011.
- [6] L. Diening, P. Hästö, S. Roudenko, Function spaces of variable smoothness and integrability, J. Funct. Anal. 256 (6) (2009) 1731–1768.

- [7] D. Drihem, Atomic decomposition of Besov spaces with variable smoothness and integrability, *J. Math. Anal. Appl.* 389 (2012) 15–31.
- [8] W. Farkas, H.-G. Leopold, Characterizations of function spaces of generalised smoothness, *Ann. Mat.* 185 (2006) 1–62.
- [9] S. Jaffard, Pointwise smoothness, two-microlocalization and wavelet coefficients, in: *Conference on Mathematical Analysis*, El Escorial, 1989, *Publ. Mat.* 35 (1) (1991) 155–168.
- [10] S. Jaffard, Y. Meyer, *Wavelet Methods for Pointwise Regularity and Local Oscillations of Functions*, *Memoirs of the AMS*, vol. 123, 1996.
- [11] H. Kempka, Generalized 2-microlocal Besov spaces, PhD thesis, University of Jena, Germany, 2008.
- [12] H. Kempka, 2-microlocal Besov and Triebel–Lizorkin spaces of variable integrability, *Rev. Mat. Complut.* 22 (1) (2009) 227–251.
- [13] H. Kempka, Atomic, molecular and wavelet decomposition of 2-microlocal Besov and Triebel–Lizorkin spaces with variable integrability, *Funct. Approx. Comment. Math.* 43 (2) (2010) 171–208.
- [14] H. Kempka, J. Vybíral, Spaces of variable smoothness and integrability: characterizations by local means and ball means of differences, *J. Fourier Anal. Appl.* 18 (4) (2012) 852–891.
- [15] O. Kováčik, J. Rákosník, On spaces $L^{p(x)}$ and $W^{k,p(x)}$, *Czechoslovak Math. J.* 40(116) (4) (1991) 592–618.
- [16] J. Lévy Véhel, S. Seuret, A time domain characterization of 2-microlocal spaces, *J. Fourier Anal. Appl.* 9 (5) (2003) 473–495.
- [17] J. Lévy Véhel, S. Seuret, The 2-microlocal formalism, in: *Fractal Geometry and Applications, A Jubilee of Benoit Mandelbrot*, in: *Proceedings of Symposia in Pure Mathematics*, vol. 72, 2004, pp. 153–215 (Part 2).
- [18] S.D. Moura, Function spaces of generalised smoothness, entropy numbers, applications, *Dissertationes Math.* 398 (2001).
- [19] W. Orlicz, Über konjugierte Exponentenfolgen, *Studia Math.* 3 (1931) 200–212.
- [20] J. Peetre, *New Thoughts on Besov Spaces*, *Duke University Mathematics Series*, vol. 1, Mathematics Department, Duke University, Durham, N.C., 1976.
- [21] T. Runst, W. Sickel, *Sobolev Spaces of Fractional Order, Nemytskij Operators and Nonlinear Partial Differential Equations*, de Gruyter, Berlin, 1996.
- [22] B. Scharf, Atomic representations in function spaces and applications to pointwise multipliers and diffeomorphisms, a new approach, *Math. Nachr.* 286 (2–3) (2013) 283–305.
- [23] C. Schneider, J. Vybíral, Non-smooth atomic decompositions, traces on Lipschitz domains and pointwise multipliers in function spaces, *J. Funct. Anal.* 264 (5) (2013) 1197–1237.
- [24] L. Skrzypczak, Atomic decompositions on manifolds with boundary geometry, *Forum Math.* 10 (1) (1998) 19–38.
- [25] H. Triebel, *Theory of Function Spaces II*, *Monographs in Mathematics*, vol. 84, Birkhäuser Verlag, Basel, 1992.
- [26] H. Triebel, H. Winkelhoß, Intrinsic atomic characterization of function spaces on domains, *Math. Z.* 221 (1) (1996) 647–673.
- [27] J. Vybíral, Sobolev and Jawerth embeddings for spaces with variable smoothness and integrability, *Ann. Acad. Sci. Fenn. Math.* 34 (2) (2009) 529–544.