

The Uniform Hopf Inequality for discontinuous coefficients and optimal regularity in BMO for singular problems



Nada El Berdan^a, Jesús Ildelfonso Díaz^b, Jean Michel Rakotoson^{a,*}

^a Université de Poitiers, Laboratoire de Mathématiques et Applications, UMR CNRS 7348 – SP2MI, Boulevard Marie et Pierre Curie – Téléport 2, F-86962 Chasseneuil Futuroscope Cedex, France

^b Instituto de Matematica Interdisciplinar, Departamento de Matemática Aplicada, Universidad Complutense de Madrid, Plaza de Ciencias, 3, 28040 Madrid, Spain

ARTICLE INFO

Article history:

Received 29 May 2015

Available online 29 December 2015

Submitted by V. Radulescu

Keywords:

Uniform Hopf Inequality

BMO and Campanato spaces

Elliptic operator with discontinuous coefficients

Singular problems

ABSTRACT

We consider some singular second order semilinear problems which include, among many other special cases, the boundary layer equations such as they were treated by O.A. Oleinik in her pioneering works. We consider diffusion linear operator with possible discontinuous coefficients and prove an optimal criterion to get a quantitative strong maximum principle that we call Uniform Hopf Inequality UHI. Since the solutions of the singular semilinear problems under consideration are not Lipschitz continuous we carry out a careful study of the regularity of solutions when the coefficients of the diffusion matrix are merely in the vmo space and bounded. We prove that the gradient of the solution is still p-integrable, in absence of any continuity assumption on the spatial potential coefficient in the singular term. To this end, the UHI property is used several times. We also apply and improve previous a priori estimates due to S. Campanato in 1965.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

The paper deals with the study of the singular semilinear problem

$$(\mathcal{SP}) = \begin{cases} Lu = \frac{a(x)}{u^m} & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a regular open domain set of \mathbb{R}^n , $a(x)$ satisfies, $m > 0$

$$a \in L^\infty(\Omega), \quad a \geq 0, \quad a \neq 0, \quad (1.1)$$

* Corresponding author.

E-mail addresses: nada.el.berdan@math.univ-poitiers.fr (N. El Berdan), jidiaz@ucm.es (J.I. Díaz), Jean-Michel.Rakotoson@math.univ-poitiers.fr (J.M. Rakotoson).

and L denotes an elliptic second order operator with non-necessarily $C^1(\overline{\Omega})$ coefficients. We shall pay attention mainly to the “case of pure diffusion” in which

$$Lu = -\operatorname{div}(A(x)\nabla u) \quad (1.2)$$

with the coefficients of the “coercive” matrix $A(x) = (a_{ij}(x))_{i,j=1}^n$ such that

$$(H_1) \quad \begin{cases} a_{ij} \in L^\infty(\Omega) \quad \forall i, j = 1, \dots, n \\ \exists \alpha > 0, \forall \xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n, \sum_{i,j} a_{ij}(x) \xi_i \xi_j \geq \alpha |\xi|^2 \quad \text{a.e. in } \Omega. \end{cases}$$

Problem (\mathcal{SP}) arises in many applied contexts, as for instance, in the study of the boundary layer in Fluid Mechanics. One of the pioneering works of this type of equations was the 1977 paper by M.G. Crandall, P.H. Rabinowitz and L. Tartar [13] in which it was mentioned as applied motivations the case of suitable chemical kinetics (Fulks and Maybee [17]) and the consideration of some similar problems in the context of signal transmission ([33] and [29]). But there are many other contexts in which problem (\mathcal{SP}) turns to be relevant. To indicate some other different motivation, we mention here the study of the “boundary layer” flow past a flat plate such as it was considered by H. Blasius [2], in 1908, after the seminal study by L. Prandtl [36] in 1905. It is well-known that the so-called “Crocco transformations” [14] lead the problem to the consideration of a suitable class of singular parabolic equations which by different arguments are reduced to the study of equations of the type

$$(P_w) \quad \begin{cases} -w_{\eta\eta} + B(\eta)w_\eta + C(\eta)w = \frac{a(\eta)}{w} & \text{in } (0, 1) \\ w(1) = 0 \\ w'(0) = c_0, \end{cases}$$

for suitable coefficients B and C , for some a satisfying (1.1) and $C_0 \in \mathbb{R}$ (see e.g. the expositions made in the monograph Oleinik and Samoklin [35] or the paper Vajravelu et al. [43]).

In spite of the great relevance of the study of the boundary layer in many problems of engineering, meteorology, oceanography etc., the intensive mathematical treatment was only successive after a series of papers by O.A. Oleinik starting in 1952 (see again the exposition made in Oleinik and Samoklin [35]). In her 1968 paper ([34], see also [25]) she derived the a priori estimate

$$\underline{C}(1-\eta)\sqrt{-Ln\mu(1-\eta)} \leq w(\eta) \leq \overline{C}(1-\eta)\sqrt{-Ln\mu(1-\eta)} \quad \text{for any } \eta \in (0, 1)$$

for suitable positive constants \underline{C} , \overline{C} , μ which shows that

$$w \notin C^{0,1}([0, 1]).$$

Since the solution of the problem (\mathcal{SP}) is not Lipschitz continuous, it is important to show that its gradient is in some functional space smaller than $L^2(\Omega)$. A related boundary estimate was proved in Gui and Hua Lin [21] and in [25] for solutions of (\mathcal{SP}) when $L = -\Delta$. We will show that the situation is completely different when we consider an operator L with variable coefficients. We shall show (see Theorem 3) that $u \in W^{1,p}(\Omega)$, for any $p \in [1, +\infty)$ assuming furthermore that the coefficients satisfy

$$a_{ij} \in vmo(\Omega), \quad \forall i, j = 1, \dots, n$$

(see below a full definition).

The regularity $u \in W^{1,p}(\Omega)$ for any $p \in [1, +\infty)$ already improves some previous results in the literature (see e.g. Díaz, Hernández and Rakotoson [16], Bougherara, Giacomoni and Hernández [7], Díaz and Rakotoson [15], and their references). In fact, under the additional regularity $a_{ij} \in C^{0,1}(\overline{\Omega})$, $\forall i, j = 1, \dots, n$ we shall show (Theorem 4.3) that $\nabla u \in bmo_r(\Omega)^n$, a functional space already used by Campanato [8] under the name $\mathcal{L}^{2,n}(\Omega)$ and which can be obtained through some variations from the BMO space of John and Nirenberg [27] (see also Chang [10], Chang, Dafni and Stein [11]).

One of our key arguments, in this paper, is a suitable application of the so-called “Uniform Hopf Inequality” which says that if v is a very weak solution of the linear problem

$$\begin{cases} Lv = f(x) & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

then there exists a positive constant $C_{\Omega,L}$, depending only on Ω and the coefficients of L , such that

$$v(x) \geq C_{\Omega,L} \delta(x) \int_{\Omega} f(y) \delta(y) dy \quad \text{a.e. } x \in \Omega. \quad (1.4)$$

Here, $\delta(x)$ stands for the distance of $x \in \Omega$ to the boundary $\partial\Omega$.

This inequality was used by first time in the paper Díaz, Morel and Oswald [32] for the study of a singular semilinear problem (with $L = -\Delta$) in which the singular term arises in the right hand side of the equality. In fact, the detailed proof of the inequality was announced as a separated independent work by Morel and Oswald [32] but it was unpublished. A proof of it (always with $L = -\Delta$) was offered in the paper Brezis and Cabré [4]. The proof is still valid when $L = -\operatorname{div}(A(x)\nabla \cdot)$ and the coefficients a_{ij} of A are in $C^{0,1}(\overline{\Omega})$. Here, we shall show that the Uniform Hopf Inequality holds even for the case in which L has discontinuous coefficients satisfying $a_{ij} \in C^{0,1}(\overline{\Omega}_b)$, $a_{ij} \in L^\infty(\Omega_l)$, where we assume that Ω admits a partition i.e., $\Omega = \overline{\Omega}_l \cup \Omega_b$, with $\Omega_l \subset \subset \Omega$ and Ω_b contained in a neighborhood of $\partial\Omega$. We shall also prove that the condition $a_{ij} \in C^{0,1}(\overline{\Omega}_b)$, $a_{ij} \in L^\infty(\Omega_l)$ is sharp by giving a counter example of it for the case in which a_{ij} are not continuous in some neighborhood of the boundary.

Sections 4 and 5 will deal with the main regularity results of this paper (first concerning with u^{-1} as nonlinear term and then, in Section 5, with u^{-m} as a general case).

2. Notations and preliminaries

We shall consider Ω an open bounded smooth (say $C^{0,1}$ at least) of \mathbb{R}^n . We recall some spaces (namely the bounded mean oscillation functions (bmo)) that we shall use later (see e.g. [27,10,11,22,28,38,42], ...).

Definition 1 (Of $bmo(\mathbb{R}^n)$). A locally integrable function f on \mathbb{R} is said to be in $bmo(\mathbb{R}^n)$ if

$$\sup_{0 < \operatorname{diam}(Q) < 1} \frac{1}{|Q|} \int_Q |f(x) - f_Q| dx + \sup_{\operatorname{diam}(Q) \geq 1} \frac{1}{|Q|} \int_Q |f(x)| dx \equiv \|f\|_{bmo(\mathbb{R}^n)} < +\infty,$$

where the supremum is taken over all cube $Q \subset \mathbb{R}^n$ whose sides are parallel to the coordinate axis. $\operatorname{diam}(Q)$ stands for the diameter of Q , $|Q|$ the measure of the cube and f_Q the average of f over the cube Q . The cube can be replaced by a ball.

Definition 2 (Of $bmo_r(\Omega)$ and main property). A locally integrable function f on a Lipschitz bounded domain Ω is said to be in $bmo_r(\Omega)$ (r stands for restriction) if

$$\sup_{0 < \text{diam}(Q) < 1} \frac{1}{|Q|} \int_Q |f(x) - f_Q| dx + \int_{\Omega} |f(x)| dx \equiv \|f\|_{bmo_r(\Omega)} < +\infty,$$

where the supremum is taken over all cubes $Q \subset \Omega$ whose sides are parallel to the coordinate axis.

In this case, there exists a function $\tilde{f} \in bmo(\mathbb{R}^n)$ such that

$$\tilde{f}|_{\Omega} = f \text{ and } \|\tilde{f}\|_{bmo(\mathbb{R}^n)} \leq c_{\Omega} \cdot \|f\|_{bmo_r(\Omega)}.$$

Definition 3 (Of the Campanato space $\mathcal{L}^{2,n}(\Omega)$). A function $u \in \mathcal{L}^{2,n}(\Omega)$ if

$$\|u\|_{L^2(\Omega)} + \sup_{x_0 \in \Omega, r > 0} \left[r^{-n} \int_{Q(x_0, r) \cap \Omega} |u - u_r|^2 dx \right]^{\frac{1}{2}} = \|u\|_{\mathcal{L}^{2,n}(\Omega)} < +\infty.$$

Here

$$u_r = \frac{1}{|Q(x_0, r) \cap \Omega|} \int_{Q(x_0, r) \cap \Omega} u(x) dx.$$

$Q(x_0; r)$ (resp $B(x_0; r)$) is the cube (resp the ball) of center at x_0 of side (resp radius) r_0 .

Lemma 2.1 (Equivalence of the two definitions). For a Lipschitz bounded domain Ω , one has:

$$\mathcal{L}^{2,n}(\Omega) = bmo_r(\Omega),$$

with equivalent norms.

This theorem is not essential for our purpose (we refer to [39] for its proof).

We shall also use the associated Sobolev space

$$W_0^1 bmo_r(\Omega) := \left\{ u : \Omega \rightarrow \mathbb{R}; u \in W_0^{1,1}(\Omega) \text{ and } \nabla u \in bmo_r(\Omega)^n \right\}.$$

As in [42], we also introduce the space.

$$vmo(\Omega) := \left\{ f \in bmo_r(\Omega) \text{ and } \lim_{R \rightarrow 0} \sup_{\substack{r \leq R \\ x_0 \in \Omega}} \frac{1}{r^n} \int_{B(x_0, r) \cap \Omega} |f - f_r| dx \rightarrow 0 \right\}.$$

We recall that the Sobolev–Poincaré inequality implies that $W^{1,n}(\Omega) \hookrightarrow vmo_{loc}(\Omega)$. This gives how we can construct elements vmo .

For a measurable set E in \mathbb{R}^n we denote by $|E|$ its Lebesgue measure, and for a measurable function u from the open bounded set Ω into \mathbb{R}^n we define the following auxiliary functions:

1. The distribution function of u . It is a $m : \mathbb{R} \rightarrow]0, |\Omega|]$, such that

$$m(t) = \text{meas} \{x \in \Omega : u(x) > t\} = |u > t|.$$

2. The monotone rearrangement of u (denoted by u_*), is the generalized inverse of m , i.e.

$$u_*(s) = \inf\{t \in \mathbb{R} : |u| > t\} \leq s, \quad s \in]0, |\Omega|[,$$

$$u_*(0) = \operatorname{ess\,sup}_{\Omega} u.$$

We also define $|u|_{**} = \frac{1}{t} \int_0^t |u|_*(s) ds$ for $t > 0$.

The Lorentz spaces $L^{p,q}_0(\Omega)$ are defined, for $1 \leq p < +\infty$, $1 \leq q < +\infty$, as

$$L^{p,q}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable} : \int_0^{|\Omega|} \left[t^{\frac{1}{p}} |u|_{**}(t) \right]^q \frac{dt}{t} < +\infty \right\},$$

$$L^{p,\infty}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable} : \sup_{t \leq |\Omega|} t^{\frac{1}{p}} |u|_{**}(t) < +\infty \right\},$$

$$W_0^1 L^{p,q}(\Omega) = \left\{ u \in W_0^{1,1}(\Omega) : |\nabla u| \in L^{p,q}(\Omega) \right\}.$$

We shall also use the usual notation

$$C^{0,1}(\overline{\Omega}) := \left\{ u : \Omega \rightarrow \mathbb{R} \text{ measurable}, \exists K > 0; |u(x) - u(y)| \leq K|x - y|, \text{ for any } x, y \in \overline{\Omega} \right\}$$

$$:= \left\{ u : \Omega \rightarrow \mathbb{R} : \exists \tilde{u} \in C^{0,1}(\mathbb{R}^n), \tilde{u} \text{ restricted to } \Omega \text{ is } u : \tilde{u}|_{\Omega} = u \right\}.$$

We define the following operator

$$L^* = -\operatorname{div}(A^* \nabla \cdot); \quad A^* \text{ is the adjoint matrix of } A.$$

It is well-known that we can define the Green function associated to those operators and Dirichlet boundary conditions:

Theorem 2.1 (Green function for L^* and L). (See [41,23,44].) There exists a unique function $G_{L^*} : \Omega \times \Omega \rightarrow \overline{\mathbb{R}}$ such that

1) $\forall y \in \Omega$, $G_{L^*}(\cdot, y) \in W_0^1 L^{n',\infty}(\Omega)$ and $\sup_y \|G_{L^*}(\cdot, y)\|_{W_0^1 L^{n',\infty}(\Omega)} \leq C(\Omega)$, satisfying

$$\int_{\Omega} A(x) \nabla G_{L^*}(x, y) \nabla \varphi \, dx = \varphi(y), \quad \forall \varphi \in W_0^1 L^{n,1}(\Omega).$$

2) $G_{L^*}(\cdot, y) \in C(\overline{\Omega} \setminus \{y\}) \cap H^1(\Omega \setminus B(y, r))$, $\forall r > 0$.

3) $\forall \varphi \in C(\overline{\Omega}) \cap H_0^1(\Omega)$ such that $L^* \varphi \in C(\overline{\Omega})$, we have

$$\int_{\Omega} G_{L^*}(x, y) L^* \varphi(x) \, dx = \varphi(y).$$

4) $G_L(x, y) = G_{L^*}(y, x)$, $\forall (x, y) \in \Omega^2$.

5) Given $f \in L^2(\Omega)$, if $u \in H_0^1(\Omega)$ verifies $Lu = f$ then

$$u(x) = \int_{\Omega} G_L(x, y) f(y) dy.$$

G_L (resp G_{L^*}) is called the Green kernel associated to L (resp L^*) for Dirichlet conditions.

Remark 1. Statement 3) in [Theorem 2.1](#) is due to G. Stampacchia [\[41\]](#). Here the definition of the Green kernel is given according to Stampacchia. But, such as it is pointed by this author, this definition is stable by approximation (his proof relies on the approximation of A). Nevertheless, it is already known that it holds for measure data problems (here the measure is e.g. the Dirac measure). In this case the problem stated in 1) has a solution whenever $A(x) \in L^\infty(\Omega)^{n^2}$ (see [\[37,40,41,44\]](#)).

[Theorem 2.2](#) is then a combination of all those properties.

Theorem 2.2 (Comparison of Green kernel). (See [\[41\]](#).) Suppose we have the following operator defined by $L^1\varphi = -\operatorname{div}(A^1(x)\nabla\varphi)$ such that

$$A^1(x) = (a_{ij}^1(x))_{i,j}, \quad a_{ij}^1 \in C^{0,1}(\overline{\Omega}),$$

with the coercivity condition in all the domain Ω ,

$$\sum_{i,j=1}^n a_{ij}^1(x) \zeta_i \zeta_j \geq \alpha |\zeta|^2, \quad \forall \zeta \in \mathbb{R}^n, \alpha > 0$$

and let G_{L^1} be the Green function associated to L^1 . Then for any relatively open compact set $\Omega'_{\ell,0}$ of Ω there exists a constant $K_1 = K_1(\Omega'_{\ell,0}) > 0$ such that

$$K_1^{-1} G_{-\Delta}(x, y) \leq G_{L^1}(x, y) \leq K_1 G_{L^*}(x, y), \quad \forall (x, y) \in \overline{\Omega'}_{\ell,0} \times \overline{\Omega'}_{\ell,0}.$$

In all this paper, we shall use the notation $\delta(x) = \operatorname{dist}(x, \partial\Omega)$. We shall also use the following,

Lemma 2.2 (Hardy inequality). (See [\[15,30\]](#).) Let Ω be of class $C^{0,1}$. Then, $\exists c > 0$ such that $\forall u \in C_c^1(\Omega)$

$$\int_{\Omega} \left(\frac{|u(x)|}{\delta(x)} \right)^2 dx \leq c \int_{\Omega} |\nabla u|^2 dx.$$

Moreover, for $a > 1$, $\exists C_a(\Omega) > 0$, such that $\forall u \in C_c^1(\Omega)$.

$$\int_{\Omega} \frac{|u(x)|}{\delta^a} dx \leq C_a(\Omega) \int_{\Omega} |\nabla u| \delta^{1-a} dx.$$

Lemma 2.3 (Iteration). (See [\[20,8,12\]](#).) Let $\Phi(\rho)$ be a non-negative and non-decreasing function. Assume that for some non-negative constants A, α, β, r_0, B with $\beta < \alpha$, we have $\forall r \in]0, r_0[, \forall \rho \in]0, r[$

$$\Phi(\rho) \leq A \left(\frac{\rho}{r} \right)^\alpha \Phi(r) + Br^\beta.$$

Then, there exists $c > 0$ such that

$$\Phi(\rho) \leq c \left(\frac{\rho}{r} \right)^\beta \Phi(r) + B\rho^\beta, \quad \forall 0 < \rho \leq r \leq r_0.$$

3. On the Uniform Hopf Inequality

To solve (\mathcal{SP}) , our approach will not be based on the notion of sub-solution and super-solution such as it is done in [24,25], when $A(x) = (a_{ij}(x))$, $a_{ij} \in C^{1,\alpha}(\overline{\Omega})$ (smooth). We shall apply the following inequality:

Definition 4 (*Uniform Hopf Inequality*). We say that the operator L satisfies the Uniform Hopf Inequality if there exists a constant $C_{\Omega,L} > 0$ such that for all $f \in L^{\infty}_+(\Omega)$, the unique solution $v \in H^1_0(\Omega)$ of $-\operatorname{div}(A\nabla v) = f$ in $\mathcal{D}'(\Omega)$ satisfies

$$v(x) \geq C_{\Omega,L} \delta(x) \int_{\Omega} f(y) \delta(y) dy, \quad \text{a.e. } x \in \Omega. \quad (3.1)$$

The inequality (3.1) still holds for (v, f) which can be approximate pointwise almost everywhere by a sequence of $(v_n, f_n) \in H^1_0(\Omega) \times L^{\infty}(\Omega)$ with the same matrix A .

Inequality (3.1) holds true if the coefficients of the matrix A are Lipschitz as it is shown in

Theorem 3.1. (See [32,6].) Suppose that $f \in L^{\infty}_+(\Omega)$, and consider L^1 the operator given in Theorem 2.2. Let v be a solution of

$$\begin{cases} L^1 v = f & \text{in } \Omega, \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

Then

$$v(x) \geq C \delta(x) \int_{\Omega} f(y) \delta(y) dy, \quad \text{a.e. } x \in \Omega,$$

where $C > 0$ is a constant depending on Ω and $\delta(x) = \operatorname{dist}(x, \partial\Omega)$.

Remark 2 (On the proof of Theorem 3.1). In the mentioned references, the proofs are given for the Laplacian operator they can be modified to hold for the case where $A = A^1$. An alternative proof can be given using the equivalence of Green functions (see [26]).

Our first result wants to point out that if A is only bounded near the boundary but not Lipschitz continuous then (3.1) may fail to be true.

Theorem 3.2. There exist a smooth open set $\Omega \subset \mathbb{R}^2$, a matrix A with bounded coefficients, $f \in L^{\frac{3}{2}}(\Omega)$ and $u \in H^1_0(\Omega)$ solution of $\int_{\Omega} A(x) \nabla u \nabla \varphi dx = \int_{\Omega} f \varphi dx$ ($\forall \varphi \in H^1_0(\Omega)$), such that the Uniform Hopf Inequality fails to be true.

Proof. Consider $\Omega := \{(x, y) \in \mathbb{R}^2 \text{ such that } x > 0 \text{ and } x^2 + y^2 < 1\}$.

Define the function

$$v(x, y) = \frac{x}{(x^2 + y^2)^{\frac{1}{4}}} - x := v_1(x, y) + v_2(x, y),$$

and the following matrix (already used in Meyers [31]).

$$A(x, y) := \frac{1}{4(x^2 + y^2)} \begin{pmatrix} 4x^2 + y^2 & 3xy \\ 3xy & x^2 + 4y^2 \end{pmatrix}.$$

We have $v \in H_0^1(\Omega)$. Now, we claim that $-\operatorname{div}(A\nabla v) = f \geq 0$ on Ω . Indeed, since

$$\nabla v_1 = \begin{pmatrix} \frac{x^2 + 2y^2}{2(x^2 + y^2)^{\frac{5}{4}}} \\ \frac{-xy}{2(x^2 + y^2)^{\frac{5}{4}}} \end{pmatrix},$$

then

$$A\nabla v_1 = \begin{pmatrix} \frac{2x^2 + y^2}{2(x^2 + y^2)^{\frac{5}{4}}} \\ \frac{xy}{4(x^2 + y^2)^{\frac{5}{4}}} \end{pmatrix}.$$

Moreover

$$\begin{aligned} \frac{\partial}{\partial x} \left(\frac{2x^2 + y^2}{2(x^2 + y^2)^{\frac{5}{4}}} \right) &= \frac{-2x^3 + 3xy^2}{4(x^2 + y^2)^{\frac{9}{4}}}, \\ \frac{\partial}{\partial y} \left(\frac{xy}{4(x^2 + y^2)^{\frac{5}{4}}} \right) &= \frac{2x^3 - 3xy^2}{4(x^2 + y^2)^{\frac{9}{4}}}. \end{aligned}$$

Consequently, we have

$$-\operatorname{div}(A\nabla v_1) = 0 \text{ in } \Omega. \quad (3.2)$$

On the other hand, for $v_2(x, y) = -x$, we have

$$\nabla v_2 = \begin{pmatrix} -1 \\ 0 \end{pmatrix},$$

and then

$$A\nabla v_2 = \begin{pmatrix} -\frac{4x^2 + y^2}{4(x^2 + y^2)} \\ \frac{-3xy}{4(x^2 + y^2)} \end{pmatrix}.$$

In a similar way, we have

$$-\operatorname{div}(A\nabla v_2) = \frac{3x}{4(x^2 + y^2)} \text{ in } \Omega. \quad (3.3)$$

Thus, by (3.2)–(3.3) we conclude that

$$-\operatorname{div}(A\nabla v) = \frac{3x}{4(x^2 + y^2)} := f \geq 0, \text{ in } \Omega.$$

We have $f \in L^{\frac{3}{2}}(\Omega)$ since, by using polar coordinates, we have that

$$\int_{\Omega} |f(x, y)|^{\frac{3}{2}} dx dy \leq c \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \int_0^1 \left| \frac{3r \cos \theta}{4r^2} \right|^{\frac{3}{2}} r dr d\theta$$

$$\leq c \int_0^1 r^{-\frac{1}{2}} dr < +\infty.$$

Let us calculate $\inf_{(x,y) \in \Omega} \frac{v(x,y)}{\delta(x,y)}$. We observe that since $v(x,y) \geq 0$, then

$$\frac{v(x,y)}{\delta(x,y)} \geq 0 \text{ a.e. on } \Omega.$$

By using polar coordinates again, we get

$$v(x,y) = r \cos \theta \left(\frac{1}{\sqrt{r}} - 1 \right).$$

Then,

$$\frac{v(r,\theta)}{\delta(r,\theta)} \leq \sqrt{r} \cos \theta \rightarrow 0 \text{ as } r \searrow 0.$$

Therefore,

$$\inf_{(x,y) \in \Omega} \frac{v(x,y)}{\delta(x,y)} = 0. \quad (3.4)$$

Arguing by contradiction, (3.4) infers that the Uniform Hopf Inequality cannot hold in this case. \square

We shall assume that

$$(H_2) \quad \begin{cases} \text{there exists a matrix } A^1(x) = (a_{ij}^1(x))_{i,j}, \ x \in \overline{\Omega}, \text{ with} \\ \left\{ \begin{array}{l} a_{ij}^1 \in C^{0,1}(\overline{\Omega}), \ \alpha\text{-coercive i.e. } \forall \xi \in \mathbb{R}^n \ (A^1(x)\xi, \xi) \geq \alpha|\xi|^2, \ \forall x \in \overline{\Omega}, \\ a_{ij}^1 \text{ restricted to } \Omega_b \text{ coincides with } a_{ij}, \ \forall i, j : a_{ij}|_{\Omega_b} = a_{ij}^1. \end{array} \right. \end{cases}$$

Here $\Omega = \overline{\Omega}_\ell \cup \Omega_b$, $\Omega_\ell \subset\subset \Omega$.

In Ω , we shall associate to A^1 the operator $L^1 = -\operatorname{div} (A^1(x)\nabla \cdot)$. The main result of this section is the following:

Theorem 3.3. *Under the above assumptions (H_1) , and (H_2) there exists $C_{\Omega,L} > 0$, such for any $f \in L_+^\infty(\Omega)$, the solution $u \in H_0^1(\Omega)$ of (1.3) satisfies for a.e. $\forall y \in \Omega$*

$$u(y) \geq C_{\Omega,L} \delta(y) \int_{\Omega} f(x) \delta(x) dx. \quad (3.5)$$

For its proof, for $\Omega_\ell \subset\subset \Omega$, we shall consider the open set $\Omega'_{\ell,0} \subset\subset \Omega$ such that $\overline{\Omega}_\ell \subset \Omega'_{\ell,0}$. In addition, for $\Omega_b = \Omega \setminus \overline{\Omega}_\ell$ we consider its subset $\Omega'_{b,0} = \Omega \setminus \overline{\Omega'}_{\ell,0}$. We shall need the following lemmas to prove the inequality (3.5).

Lemma 3.1. *Under the same assumptions as in Theorem 3.3, and if $\Omega'_{\ell,0}$ is given as above, the constant K_1 given in Theorem 2.2, is such that $K_1 = K_1(\Omega'_{\ell,0}) > 0$ and*

$$K_1 G_{L^*}(x,y) \geq G_{L^{1*}}(x,y), \ \forall x \in \Omega'_{b,0}, \ \forall y \in \overline{\Omega'}_{\ell,0}.$$

Proof. Let $\varphi \in W_0^1 L^{n,1}(\Omega'_{b,0})$ and let $\tilde{\varphi}$ its extension to Ω by zero.

Then, $\tilde{\varphi} \in W_0^1 L^{n,1}(\Omega)$. For $y \in \overline{\Omega}_l$ fixed, let $w(x) = K_1 G_{L^*}(x, y) - G_{L^{1*}}(x, y)$, $x \in \Omega'_{b,0}$. Then $w \in H^1(\Omega'_{b,0})$ and

$$\int_{\Omega} A(x) \nabla w \cdot \nabla \tilde{\varphi} dx = K_1 \int_{\Omega} A(x) \nabla G_{L^*}(x, y) \nabla \tilde{\varphi} dx - \int_{\Omega} A(x) \nabla G_{L^{1*}}(x, y) \nabla \tilde{\varphi} dx.$$

Since $A(x) = A^1(x)$ on $\Omega'_{b,0}$, thanks to [Theorem 2.1](#), we obtain

$$\int_{\Omega} A(x) \nabla w \cdot \nabla \tilde{\varphi} dx = K_1 \tilde{\varphi}(y) - \int_{\Omega'_{b,0}} A^1(x) \nabla G_{L^{1*}}(x, y) \nabla \varphi dx, \quad \text{for any } y \in \overline{\Omega'}_{\ell,0}$$

Using again $\tilde{\varphi}$ in the last term and [Theorem 2.1](#)

$$\begin{aligned} \int_{\Omega} A(x) \nabla w \cdot \nabla \tilde{\varphi} dx &= K_1 \tilde{\varphi}(y) - \int_{\Omega} A^1(x) \nabla_x G_{L^{1*}}(x, y) \nabla \tilde{\varphi} dx \\ &= K_1 \tilde{\varphi}(y) - \tilde{\varphi}(y) = 0. \end{aligned}$$

Thus,

$$\int_{\Omega'_{b,0}} A(x) \nabla w \cdot \nabla \varphi dx = 0. \quad (3.6)$$

Moreover, its trace verifies that, $\gamma_0 w(x) = 0$ on $\partial\Omega$, and from [Theorem 2.2](#)

$$w(x) = K_1 G_{L^*}(x, y) - G_{L^{1*}}(x, y) \geq 0 \text{ for } x \in \partial\Omega'_{\ell,0}.$$

Consequently, by the maximum principle, $w \geq 0$ on $\Omega'_{b,0}$. \square

Corollary 3.1 (Of [Theorem 2.2](#) and [Lemma 3.1](#)). Under the same assumptions as in [Theorem 3.3](#), the constant $K_1 > 0$ found in [Lemma 3.1](#) satisfies

$$K_1 G_{L^*}(x, y) \geq G_{L^{1*}}(x, y), \quad \forall x \in \Omega, \quad \forall y \in \overline{\Omega'}_{\ell,0}.$$

Proof. Since $\Omega = \Omega'_{b,0} \cup \overline{\Omega'}_{\ell,0}$ and we have shown that $\forall y \in \overline{\Omega'}_{\ell,0}$,

$$w(x) = K_1 G_{L^*}(x, y) - G_{L^{1*}}(x, y) \geq 0, \quad \forall x \in \Omega'_{b,0},$$

and $w(x) \geq 0$, $\forall x \in \overline{\Omega'}_{\ell,0}$, (from [Theorem 2.2](#)), thus, $w(x) \geq 0$ on $\Omega'_{b,0} \cup \overline{\Omega'}_{\ell,0} = \Omega$. \square

Corollary 3.2 (Lower estimates on the subset Ω_l). Let $f \in L_+^\infty(\Omega)$, and let u and v in $H_0^1(\Omega)$ satisfy $Lu = f$ and $L^1 v = f$ respectively.

Then, a.e. $y \in \overline{\Omega}_\ell$

- 1) $u(y) \geq \frac{1}{K_1} \int_{\Omega} G_{L^1}(y, x) f(x) dx$,
- 2) there exists $C_{\Omega, L^1} > 0$ such that

$$v(y) = \int_{\Omega} G_{L^1}(y, x) f(x) dx \geq C_{\Omega, L^1} \delta(y) \int_{\Omega} f(x) \delta(x) dx.$$

In particular,

$$u(y) \geq C_{\Omega'_\ell, 0} \int_{\Omega} f(x) \delta(x) dx > 0. \quad (3.7)$$

Proof. Let $f \in L^{\infty}_+(\Omega)$. Then from [Corollary 3.2](#), for a.e. $y \in \overline{\Omega}_\ell$, after integrating over Ω , we have

$$u(y) = \int_{\Omega} G_{L^*}(x, y) f(x) dx \geq \frac{1}{K_1} \int_{\Omega} G_{L^1}(x, y) f(x) dx = v(y). \quad (3.8)$$

Recalling that the coefficients of L^1 are Lipschitz continuous, we can apply [Theorem 3.1](#) to obtain that $\exists C_{\Omega, L^1} > 0$ such that, for a.e. $y \in \overline{\Omega}_\ell$,

$$u(y) \geq v(y) \geq C_{\Omega, L^1} \delta(y) \int_{\Omega} f(x) \delta(x) dx. \quad \square \quad (3.9)$$

Lemma 3.2 (Lower estimates on Ω_b near the boundary). Under the same assumptions as in [Theorem 3.3](#), there exists $C_{\Omega, L} > 0$ such that

$$u(y) \geq C_{\Omega, L} \delta(y) \int_{\Omega} f(x) \delta(x) dx, \quad \text{a.e. } y \in \Omega_b = \Omega \setminus \overline{\Omega}_\ell.$$

Proof. This procedure is inspired by the method of proof used by Brezis–Cabré [\[7\]](#).

Let $\Gamma_b := \partial\Omega \cup \partial\Omega_\ell = \partial\Omega_b$ and introduce the function $w \in H^1(\Omega)$ solution of:

$$\begin{cases} L^1 w = -\operatorname{div}(A^1(x) \nabla w) = 0 & \text{in } \Omega_b, \\ w = 0 & \text{on } \partial\Omega, \\ w = 1 & \text{on } \partial\Omega_\ell. \end{cases}$$

Since the coefficients of $A = A^1$ are Lipschitz continuous on $\overline{\Omega}_b$, then by the Hopf strong maximum principle, there exists $C'_{\Omega_b} > 0$ such that

$$w(y) \geq C'_{\Omega_b} \delta(y), \quad \forall y \in \Omega_b. \quad (3.10)$$

Now, let us set $\underline{w}(y) = [C'_{\Omega_b} \int_{\Omega} f(x) \delta(x) dx]^{-1} u(y)$. By the linearity of operator L^1 and since $f \geq 0$:

$$\begin{cases} L^1 \underline{w} = \left[C'_{\Omega_b} \int_{\Omega} f(x) \delta(x) dx \right]^{-1} L^1 u \geq 0 = L^1 w & \text{in } \Omega_b, \\ \underline{w}|_{\partial\Omega_b} \geq w|_{\partial\Omega_b}. \end{cases}$$

Thus, thanks to the maximum principle, we obtain $\underline{w} \geq w$ on Ω_b . This means that for all $y \in \Omega_b$

$$u(y) \geq C'_{\Omega_\ell} C'_{\Omega_b} \delta(y) \int_{\Omega} f(x) \delta(x) dx. \quad (3.11)$$

Finally, combining (3.11) with relation (3.9), we get

$$u(y) \geq C \delta(y) \int_{\Omega} f(x) \delta(x) dx, \quad \text{a.e. } y \in \Omega. \quad \square$$

4. On the singular problem: the case of u^{-1} as nonlinear term

The Uniform Hopf Inequality (UHI) is very useful to derive regularity results for the singular semilinear problem

$$(\mathcal{P}) = \begin{cases} -\operatorname{div}(A(x)\nabla u) = \frac{a(x)}{u} & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

An existence and uniqueness results are also proved in [3,9] for linear operators and in [18,19] for non-linear operators. The main difference with our results is double. Indeed, firstly our method to prove the positivity of the solution is different than the above mentioned papers. For instance, in [3] it is obtained by a monotonicity result, nevertheless our method of proof can be extended to a general operator as $Lu = -\operatorname{div}(A\nabla u) + B\nabla u + \operatorname{div}(Cu) + a_0u$ (as it will be presented in the Nada El Berdan's thesis). Secondly the additional regularity that we shall obtain in the following theorem (the term $\frac{a}{u} \in H^{-1}(\Omega)$ or the results given in Theorem 5.2 below) is not mentioned in the above papers.

Theorem 4.1. *Let $a \in L^{\infty}_+(\Omega)$, $a \neq 0$. Then, there exists a unique solution $u \in H^1_0(\Omega)$ of (\mathcal{P}) , such that*

$$\begin{aligned} i) \quad & \frac{a}{u} \in L^1_{loc}(\Omega) \cap H^{-1}(\Omega), \quad u > 0 \text{ in } \Omega, \\ ii) \quad & \int_{\Omega} A(x)\nabla u \cdot \nabla \varphi dx = \int_{\Omega} \frac{a(x)\varphi}{u} dx, \quad \forall \varphi \in H^1_0(\Omega). \end{aligned}$$

Proof. Let us start with the uniqueness of u . If u, \bar{u} satisfy ii) then by the coercivity condition on A and choosing $\varphi = u - \bar{u}$;

$$\alpha \int_{\Omega} |\nabla(u - \bar{u})|^2 dx \leq \int_{\Omega} a(x) \left[\frac{1}{u} - \frac{1}{\bar{u}} \right] (u - \bar{u}) dx \leq 0.$$

This implies that, necessarily, $u = \bar{u}$. For the existence part, we introduce the following regularized problem:

$$(\mathcal{P}_{\varepsilon}) = \begin{cases} Lu_{\varepsilon} = -\operatorname{div}(A(x)\nabla u_{\varepsilon}) = \frac{a}{|u_{\varepsilon}| + \varepsilon} & \text{in } \Omega, \\ u_{\varepsilon} = 0 & \text{on } \partial\Omega, \end{cases}$$

with, $\varepsilon > 0$. The weak (variational) formulation for $(\mathcal{P}_{\varepsilon})$ reads

$$a_L(u_{\varepsilon}, \varphi) := \int_{\Omega} A(x)\nabla u_{\varepsilon} \cdot \nabla \varphi dx = \int_{\Omega} \frac{a(x)\varphi}{|u_{\varepsilon}| + \varepsilon} dx, \quad \forall \varphi \in H^1_0(\Omega). \quad (4.1)$$

Using the Schauder fixed point theorem (see e.g. [21]), we get the existence of u_ε . In addition, if we apply the weak maximum principle [21], we obtain that $u_\varepsilon \geq 0$. The same argument as the used for u ensures that u_ε is unique.

Estimate on u_ε : Taking $\varphi = u_\varepsilon$ as test function in (4.1), we can write

$$a_L(u_\varepsilon, u_\varepsilon) = \int_{\Omega} \frac{au_\varepsilon}{u_\varepsilon + \varepsilon} dx,$$

and then

$$\alpha \|\nabla u_\varepsilon\|_{L^2(\Omega)}^2 \leq \|a\|_{L^\infty} \int_{\Omega} \frac{u_\varepsilon}{u_\varepsilon + \varepsilon} dx \leq C_\Omega \|a\|_{L^\infty}. \quad (4.2)$$

Therefore, u_ε is uniformly bounded in $H_0^1(\Omega)$, and then there exists $u \in H_0^1(\Omega)$ such that (for a subsequence) u_ε converges to u a.e.

Now, we shall prove that $\frac{a}{u_\varepsilon}$ remains in a bounded set of $L_{loc}^1(\Omega) \cap H^{-1}(\Omega)$. By Theorem 2.1 (see also Theorem 9.3 of [41]), we have

$$u_\varepsilon(x) = \int_{\Omega} G_L(x, y) \frac{a(y)}{(u_\varepsilon(y) + \varepsilon)} dy \quad \text{a.e. } x \in \Omega, \quad (4.3)$$

where G_L is the Green function associated to L as it is defined in Section 1.

Let Ω' be a relatively compact open set in Ω . Let $f_0(y) = \frac{a}{u_\varepsilon + \varepsilon} \chi_{\Omega'}(y)$ and consider the following problem

$$\begin{cases} -\Delta w(x) = f_0 & \text{in } \Omega \\ w = 0 & \text{on } \partial\Omega. \end{cases}$$

This problem has a unique solution w in $H_0^1(\Omega) \cap H^2(\Omega)$. According to the Uniform Hopf Inequality (see Theorem 3.1), there exist $C_{\Omega, \Delta} > 0$ such that

$$w(x) \geq C_{\Omega, \Delta} \delta(x) \int_{\Omega'} \frac{a}{u_\varepsilon + \varepsilon} \delta(y) dy, \quad \text{a.e. } x \in \Omega.$$

Returning to Theorem 2.1 and to the inequality in Theorem 2.2, then for $K = K(\Omega') > 0$

$$\begin{aligned} K^{-1}w(x) &= K^{-1} \int_{\Omega} G_{-\Delta}(x, y) f_0(y) dy \leq \int_{\Omega} G_L(x, y) f_0(y) dy \\ &\leq \int_{\Omega} G_L(x, y) \frac{a(y)}{u_\varepsilon(y) + \varepsilon} dy = u_\varepsilon(x), \quad \text{a.e. } x \in \Omega. \end{aligned}$$

Consequently,

$$+\infty > u_\varepsilon(x) \geq K^{-1}w(x) \geq C_{\Omega, \Delta} \delta(x) \int_{\Omega'} \frac{a}{u_\varepsilon + \varepsilon} \delta(y) dy, \quad \text{a.e. } x \in \Omega', \quad (4.4)$$

which yields that $\frac{a}{u_\varepsilon} \in L^1_{loc}(\Omega)$ and $u_\varepsilon > 0$ a.e. in Ω . To prove that $\frac{a}{u} \in L^1_{loc}(\Omega)$, it is enough to pass to the limit at (4.4), since from Fatou's Lemma, we get

$$+\infty > u(x) \geq C_{\Omega,\Delta} \delta(x) \int_{\Omega'} \frac{a(y)}{u(y)} dy, \quad \text{a.e. } x \in \Omega'.$$

In particular, this implies that $\inf_{0 < \varepsilon < \varepsilon_1} \int_{\Omega'} \frac{a}{u_\varepsilon + \varepsilon} \delta(y) dy > 0$. We now have

$$\frac{a}{u_\varepsilon + \varepsilon} \xrightarrow{\varepsilon \rightarrow 0} \frac{a}{u} \quad \text{a.e., on } \Omega,$$

and for all Ω' relatively compact in Ω , since $\inf_{y \in \Omega'} \delta(y) > 0$, then (4.4) yields for a.e. $x \in \Omega'$

$$0 \leq \frac{a}{u_\varepsilon + \varepsilon}(x) \leq \frac{a}{u_\varepsilon}(x) \leq C_{\Omega'} \frac{a}{\int_{\Omega'} \frac{a(x)}{u_\varepsilon(x) + \varepsilon}(y) dy} \leq C_{\Omega'} a(x) < +\infty.$$

Then, by using the Lebesgue dominated convergence theorem, we obtain

$$\int_{\Omega} \frac{a}{u_\varepsilon + \varepsilon} \psi(x) dx \xrightarrow{\varepsilon \rightarrow 0} \int_{\Omega} \frac{a}{u} \psi(x) dx, \quad \forall \psi \in \mathcal{D}(\Omega).$$

Now, we want to show that

$$\int_{\Omega} A(x) \nabla u \cdot \nabla \psi = \int_{\Omega} \frac{a}{u} \psi, \quad \forall \psi \in H_0^1(\Omega).$$

For this we observe that $\frac{a}{u_\varepsilon + \varepsilon}$ belongs to a bounded subset of $H^{-1}(\Omega) \cap L^1_{loc}(\Omega)$ since, $\forall \psi \geq 0$, $\psi \in H_0^1(\Omega)$ we have

$$\int_{\Omega} \frac{a\psi}{u_\varepsilon + \varepsilon} dx = \left| \int_{\Omega} A(x) \nabla u_\varepsilon \cdot \nabla \psi dx \right| \leq C \|\nabla u_\varepsilon\|_{L^2(\Omega)} \|\nabla \psi\|_{L^2(\Omega)} \leq C \|\nabla \psi\|_{L^2(\Omega)}.$$

But, $\frac{a}{u_\varepsilon + \varepsilon}$ converges to $\frac{a}{u}$ a.e. Thus by Fatou's lemma, we deduce (knowing that $|\psi| \in H_0^1(\Omega)$ once that $\psi \in H_0^1(\Omega)$)

$$\sup_{\substack{\psi \in H_0^1 \\ \|\nabla \psi\|_{L^2(\Omega)} = 1}} \left| \int_{\Omega} \frac{a\psi}{u} dx \right| \leq C_\Omega < +\infty.$$

This shows that $\frac{a}{u} \in H^{-1}(\Omega)$ and following the property defined by Brezis–Browder [5], $\frac{a\psi}{u} \in L^1(\Omega)$ for $\psi \in H_0^1(\Omega)$ and also

$$\langle \frac{a}{u}; \psi \rangle_{H^{-1}, H^1} = \int_{\Omega} \frac{a\psi}{u} dx, \quad \forall \psi \in H_0^1(\Omega).$$

But $\forall \psi \in \mathcal{D}(\Omega)$,

$$\int_{\Omega} A(x) \nabla u \cdot \nabla \psi \, dx = \int_{\Omega} \frac{a}{u} \psi \, dx.$$

So, by density, we have

$$\lim_{\substack{\psi_n \in \mathcal{D} \\ \psi_n \rightarrow \psi}} \left\langle \frac{a}{u}; \psi_n \right\rangle = \left\langle \frac{a}{u}; \psi \right\rangle, \quad \forall \psi \in H_0^1(\Omega).$$

This ends the proof. \square

Our next results show that the gradient of the solution is more regular. We start with the study in the $L^p(\Omega)$ spaces.

Theorem 4.2. Assume furthermore that operator L satisfies the Uniform Hopf Inequality, $\partial\Omega$ is C^1 and that $a_{ij} \in vmo(\Omega)$. Then, $\forall p \in [1, \infty[$, $u \in W^{1,p}(\Omega)$, and

$$\|u\|_{W_0^{1,p}(\Omega)} \leq C_{\Omega}^1(A, p) \frac{\|a\|_{\infty}}{\int_{\Omega} \frac{a(x)\delta(x)}{u(x)} \, dx}, \quad (4.5)$$

for some positive constant $C_{\Omega}^1(A, p)$.

Proof. Since L satisfies the Uniform Hopf Inequality, there exists a constant $C_L > 0$ such that we have

$$0 \leq \frac{a(x)}{u_{\varepsilon}(x)} \leq C_L \frac{\|a\|_{\infty}}{\delta(x) \int_{\Omega} \frac{a(y)\delta(y)}{(u_{\varepsilon} + \varepsilon)(y)} \, dy}, \quad \text{a.e. } x \in \Omega.$$

Then, if $\psi \in W_0^{1,p'}(\Omega)$ with $\frac{1}{p'} = 1 - \frac{1}{p}$, $\psi \geq 0$, we then have,

$$\begin{aligned} 0 &\leq \int_{\Omega} \frac{a(x)}{u_{\varepsilon}(x)} \psi(x) \, dx \leq C_L \int_{\Omega} \frac{a(x)\psi(x)}{\delta(x) \int_{\Omega} \frac{a(y)}{(u_{\varepsilon} + \varepsilon)(y)} \, dy} \, dx \\ &\leq C_L \frac{\|a\|_{\infty}}{\int_{\Omega} \frac{a(y)}{(u_{\varepsilon} + \varepsilon)(y)} \, dy} \int_{\Omega} \frac{\psi(x)}{\delta(x)} \, dx \\ &\leq C_{\Omega,L}(a) \int_{\Omega} \frac{\psi(x)}{\delta(x)} \, dx. \end{aligned}$$

By using the Hardy inequality, we obtain:

$$0 \leq \int_{\Omega} \frac{a(x)}{u_{\varepsilon}(x)} \psi(x) \, dx \leq C_{\Omega,L}(a) \|\nabla \psi\|_{L^{p'}(\Omega)}, \quad (4.6)$$

for some positive constant $C_{\Omega,L}(a)$. Therefore, we deduce

$$\frac{a}{u_\varepsilon + \varepsilon} \text{ belongs to a bounded set of } W^{-1,p}(\Omega).$$

From the regularity result applied to the associated linear equation, the unique solution u_ε of $(\mathcal{P}_\varepsilon)$ is in $W^{1,p}(\Omega)$ (see Simader [40], Auscher–Quafsaoui [1] and Byun [4]) and

$$\|\nabla u_\varepsilon\|_{L^p(\Omega)} \leq C_\Omega \left\| \frac{a}{u_\varepsilon + \varepsilon} \right\|_{W^{-1,p}(\Omega)} \leq C_\Omega \frac{\|a\|_\infty}{\int_\Omega \frac{a(y)\delta(y)}{u_\varepsilon + \varepsilon} dy} \leq C(\Omega, a). \quad (4.7)$$

(Notice that

$$\lim_{\varepsilon_1 \rightarrow 0} \left[\inf_{0 < \varepsilon < \varepsilon_1} \int_\Omega \frac{a(y)\delta(y)}{u_\varepsilon + \varepsilon} dy \right] \geq \int_\Omega \frac{a(y)\delta(y)}{u(y)} dy > 0$$

which implies the uniform estimates in (4.7).)

Therefore, ∇u_ε is bounded in $L^p(\Omega)^n$, and since u_ε converges to u weakly in $H_0^1(\Omega)$, thus ∇u_ε converges weakly to ∇u in $L^p(\Omega)^n$. Then

$$\begin{aligned} \|\nabla u\|_{L^p(\Omega)} &\leq \liminf_{\varepsilon \rightarrow 0} \|\nabla u_\varepsilon\|_{L^p(\Omega)} \\ &\leq \liminf_{\varepsilon \rightarrow 0} \frac{C_\Omega \|a\|_\infty}{\int_\Omega \frac{a(y)\delta(y)}{u_\varepsilon + \varepsilon} dy} = C_\Omega(a). \end{aligned}$$

This gives relation (4.5). \square

Theorem 4.3. Assume that $a_{ij} \in C^{0,1}(\overline{\Omega})$, $\forall i, j$ and $\partial\Omega$ is $C^{1,1}$. Then the solution satisfying ii) of Theorem 4.1, belongs to $W_0^1 bmo_r(\Omega)$.

Proof. First let us notice that according to Theorem 3.1, the operator L satisfies the Uniform Hopf Inequality. Therefore the uniform estimates given by (4.6) and Theorem 4.2 hold true. We will prove that $\nabla u \in bmo_r(\Omega)^n$. For this statement, we will partly use some arguments from Campanato [8]. We shall establish two new *a priori* estimates:

i) **Interior local estimate:**

Lemma 4.1. For any open smooth sets $\Omega_0, \widetilde{\Omega}_0$ with $\overline{\Omega}_0 \subset \widetilde{\Omega}_0$ and $\widetilde{\Omega}_0 \subset \Omega$, for all $1 \leq p < \infty$, there exists a constant $C(p; \widetilde{\Omega}_0)$ such that:

$$\|D^2 u_\varepsilon\|_{L^p(\Omega_0)} \leq C(p; \widetilde{\Omega}_0).$$

Proof. Consider $\widetilde{\delta} = \text{dist}(\widetilde{\Omega}_0; \partial\Omega)$ and introduce function θ_0 be such that:

$$\begin{cases} \theta_0 \in C_c^\infty(\widetilde{\Omega}_0), & 0 \leq \theta_0 \leq 1 \text{ and } \text{supp } \theta_0 \subset \widetilde{\Omega}_0 \subset \Omega, \\ \theta_0 = 1 \text{ on } \Omega_0, & |D^\alpha \theta_0| \leq \frac{M}{\widetilde{\delta}^{|\alpha|}}, \quad \alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n, |\alpha| = \alpha_1 + \dots + \alpha_n. \end{cases}$$

Let $v_\varepsilon = u_\varepsilon \theta_0 \in H_0^1(\widetilde{\Omega}_0)$. Then v_ε verifies the local problem:

$$(\mathcal{P}_l) : \begin{cases} -\operatorname{div}(A(x)\nabla v_\varepsilon) = \frac{a\theta_0}{u_\varepsilon + \varepsilon} - \operatorname{div}(A(x)u_\varepsilon\nabla\theta_0) & \text{in } \widetilde{\Omega}_0, \\ v_\varepsilon = 0 & \text{on } \partial\widetilde{\Omega}_0. \end{cases}$$

Let $F_\varepsilon := \frac{a\theta_0}{u_\varepsilon + \varepsilon} - \operatorname{div}(A(x)u_\varepsilon\nabla\theta_0)$, and let $U_0 = A(x)\nabla\theta_0$. Then $F_\varepsilon = \frac{a\theta_0}{u_\varepsilon + \varepsilon} + U_0\nabla u_\varepsilon + u_\varepsilon\operatorname{div}(U_0)$, support $U_0 \subset \widetilde{\Omega}_0 \setminus \Omega_0$ and

$$\|\operatorname{div}(U_0)\|_{L^\infty} + \|U_0\|_{L^\infty} \leq \frac{M}{\delta^2} = M(\widetilde{\Omega}_0).$$

Therefore, using estimates (4.7)

$$\|F_\varepsilon\|_{L^p(\widetilde{\Omega}_0)} \leq C_a(\widetilde{\Omega}_0) \left(\left\| \frac{\theta_0}{\delta} \right\|_{L^p(\Omega)} + C(\Omega, a) \right). \quad (4.8)$$

By the well known Agmon–Douglis–Nirenberg regularity results we have,

$$\|v_\varepsilon\|_{W^{2,p}(\widetilde{\Omega}_0)} \leq C_{\Omega}(p) \|F_\varepsilon\|_{L^p(\widetilde{\Omega}_0)}. \quad (4.9)$$

Since $D^\alpha v_\varepsilon = D^\alpha u_\varepsilon$ on Ω_0 , then relation (4.9) leads finally to

$$\|u_\varepsilon\|_{W^{2,p}(\Omega_0)} \leq C \|v_\varepsilon\|_{W^{2,p}(\widetilde{\Omega}_0)} \leq C_p(\widetilde{\Omega}_0). \quad \square$$

As a consequence of Lemma 4.1 one has,

Lemma 4.2. *For all $p \geq 1$, for all open smooth set Ω_0 relatively compact in Ω , the sequence u_ε remains in a bounded set of $W^{2,p}(\Omega_0)$. Moreover, the sequence remains in a bounded set of $C^1(\overline{\Omega}_0)$.*

Proof. It is a consequence of Lemma 4.1 and the Sobolev embedding,

$$W^{2,p}(\Omega_0) \hookrightarrow C^1(\overline{\Omega}_0), \quad p > n. \quad \square$$

ii) **Estimates in a neighborhood of the boundary:** Since we assume that Ω is of class $C^{1,1}$, for every $x \in \partial\Omega$, we can find, an open neighborhood of x denoted by $\Omega_{0,1}(x)$ and a bijection

$$\tau : \Omega_{0,1} = \Omega_{0,1} \longrightarrow I^+(1),$$


such that

$$\tau \in C^{1,1}(\Omega_{0,1}(x))^n, \quad \tau^{-1} \in C^{1,1}(\overline{I^+(1)})^n \quad \text{and} \quad \tau(\partial\Omega \cap \Omega_{0,1}) = \Gamma_1,$$

where

$$I^+(1) := \left\{ x = (x', x_n) \in \mathbb{R}^n \text{ such that } |x| < 1 \text{ and } x_n > 0 \right\}$$

and



This means that we can continuously deform the boundary to a hyperplane and that this transformation is regular. After this transformation the problem $(\mathcal{P}_\varepsilon)$ reads

$$\tau(\mathcal{P}_\varepsilon) : \begin{cases} -\operatorname{div}(B(y)\nabla w_\varepsilon) = \frac{\tilde{a}(y)}{w_\varepsilon + \varepsilon} & \text{in } I^+(1), \\ w_\varepsilon = 0 & \text{on } \Gamma_1, \end{cases}$$

with $B(y) \in C^{0,1}(\overline{I^+(1)})$, $\tilde{a}(y) \in L^\infty(I^+(1))$ and $\forall \zeta = (\zeta_1, \dots, \zeta_n) \in \mathbb{R}^n, \exists \nu > 0$ such that $\sum_{i,j} b_{ij} \zeta_i \zeta_j \geq \nu |\zeta|^2$. On each ball $I^+(0, r) = I^+(r)$, more generally, for $x_0 \in \overline{I^+(1)}$, we set

$$I^+(x_0, r) := \left\{ x \in I^+(1) : |x - x_0| < r \right\}, \quad \Gamma_r = \left\{ x \in \overline{I^+(r)} : x_n = 0 \right\},$$

$I(x_0, r)$ is the ball of radius r centered at x_0 .

We will construct two Dirichlet problems with constant coefficients such that the sum of the two solutions of these two problems coincides with w_ε .

Let us fix $0 < R < 1$, $x_0 \in \overline{I^+(R)}$, R closed to 1. The first problem will be defined without the right-hand side of $\tau(\mathcal{P}_\varepsilon)$, and having the same trace of w_ε , i.e.,

$$\tau(\mathcal{P}_\varepsilon)_1 : \begin{cases} -\operatorname{div}(B(x_0)\nabla w_\varepsilon^1) = 0 & \text{in } I^+(x_0, r), \\ w_\varepsilon^1 = w_\varepsilon & \text{on } \partial I^+(x_0, r). \end{cases}$$

Here, $0 < r \leq \frac{1-R}{2}$.

Theorem 4.4. (See Campanato [pages 338, 352].) $\tau(\mathcal{P}_\varepsilon)_1$ admits a positive solution w_ε^1 with

$$\int_{I^+(x_0; r)} |\nabla w_\varepsilon^1|^2 dx \leq c_B \int_{I^+(x_0; r)} |\nabla w_\varepsilon|^2 dx \leq c(\Omega, a) \text{ (independent of } x_0, \text{ and } r).$$

There exists $c(\nu) > 0$, $\forall 0 < \rho < r$ such that

1. If $\Gamma_R \cap I(x_0, r) = \emptyset$ then

$$\|\nabla w_\varepsilon^1 - \{\nabla w_\varepsilon^1\}_\rho\|_{L^2(I^+(x_0, \rho))}^2 \leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \|\nabla w_\varepsilon^1 - \{\nabla w_\varepsilon^1\}_r\|_{L^2(I^+(x_0, r))}^2. \quad (4.10)$$

2. If $\Gamma_R \cap I(x_0, r) \neq \emptyset$ then

$$\int_{I^+(x_0; \rho)} |D_j w_\varepsilon^1|^2 dx \leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; r)} |D_j w_\varepsilon^1|^2 dx, \quad j = 1, \dots, n-1, \quad (4.11)$$

$$\int_{I^+(x_0; \rho)} |D_n w_\varepsilon^1 - \{D_n w_\varepsilon^1\}_\rho|^2 \leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; \rho)} |D_n w_\varepsilon^1 - \{D_n w_\varepsilon^1\}_r|^2 dx. \quad (4.12)$$

Here D_j denotes the partial derivative in the x_j -direction, $j = 1, \dots, n$ and $\{\cdot\}_\rho$ is the average over $I^+(x_0, \rho)$.

Proof. The problem $\tau(\mathcal{P}_\varepsilon)_1$ is identical to the one considered by Campanato [8]. Therefore, his proof can be reproduced line by line to get (4.10), (this estimate is proven in p. 338 by Campanato [8], see relation (8.12) for the local estimate, observing in that case $I(x_0, r) = B(x_0; r) \subset I^+(1)$).

The second set of relations (4.11) and (4.12), are given in page 352 (Corollary I.11 and Lemma II.11) of Campanato [8]. \square

Now, we construct the second problem as follows:

$$\tau(\mathcal{P}_\varepsilon)_2 := \begin{cases} -\operatorname{div}(B(x_0)\nabla w_\varepsilon^2) = \frac{\tilde{a}(y)}{w_\varepsilon + \varepsilon} + \operatorname{div}((B(y) - B(x_0))\nabla w_\varepsilon) & \text{in } I^+(x_0, r), \\ w_\varepsilon^2 = 0 & \text{on } \partial I^+(x_0, r). \end{cases}$$

Theorem 4.5. $\tau(\mathcal{P}_\varepsilon)_2$ admits a unique solution $w_\varepsilon^2 \in H_0^1(I^+(x_0; r))$ and for all $\lambda \in [0, n]$,

$$\|\nabla w_\varepsilon^2\|_{L^2(I^+(x_0; r))}^2 \leq c \int_{I^+(x_0; r)} \tilde{a}(y) dy + r^{\lambda+2} C_\Omega(\lambda, a),$$

for some $c > 0$, which depends only on Ω .

Proof. This problem is well-posed since the right hand side is in $H^{-1}(I^+(x_0, r))$ and admits a unique solution $w_\varepsilon^2 \in H_0^1(I^+(x_0, r))$ (by using Lax–Milgram Theorem). Note that we have $w_\varepsilon = w_\varepsilon^1 + w_\varepsilon^2$.

Estimate on ∇w_ε^2 : By multiplying $\tau(\mathcal{P}_\varepsilon)_2$ by w_ε^2 , we get

$$\nu \int_{I^+(x_0, r)} |\nabla w_\varepsilon^2|^2 dy \leq \int_{I^+(x_0, r)} \frac{\tilde{a}(y)w_\varepsilon^2(y)}{w_\varepsilon + \varepsilon} dy + \int_{I^+(x_0, r)} (B(y) - B(x_0))\nabla w_\varepsilon \nabla w_\varepsilon^2 dy.$$

Since $w_\varepsilon^2 = w_\varepsilon - w_\varepsilon^1$ and $w_\varepsilon^1 \geq 0$ then $w_\varepsilon^2 \leq w_\varepsilon$ the Lipschitz continuity condition on B , Cauchy–Schwartz inequality and Young’s inequality, yield

$$\begin{aligned} \int_{I^+(x_0, r)} |\nabla w_\varepsilon^2|^2 dy &\leq c \int_{I^+(x_0, r)} \frac{\tilde{a}(y) w_\varepsilon^2}{w_\varepsilon + \varepsilon} dy + \int_{I^+(x_0, r)} cr |\nabla w_\varepsilon \cdot \nabla w_\varepsilon^2| dy, \\ &\leq c \int_{I^+(x_0, r)} \tilde{a}(y) dy + \frac{c^2 r^2}{2} \int_{I^+(x_0, r)} |\nabla w_\varepsilon|^2 dy + \frac{1}{2} \int_{I^+(x_0, r)} |\nabla w_\varepsilon^2|^2 dy. \end{aligned}$$

Thus

$$\int_{I^+(x_0; r)} |\nabla w_\varepsilon^2|^2 dy \leq c \int_{I^+(x_0; r)} \tilde{a}(y) dy + cr^2 \int_{I^+(x_0; r)} |\nabla w_\varepsilon|^2 dy.$$

Next, we want to show that

$$\int_{I^+(x_0; r)} |\nabla w_\varepsilon|^2 dx \leq r^\lambda c_\Omega(\lambda, a). \quad (4.13)$$

From relation (4.6) of the proof of Theorem 4.2, we have proved that

$$\|\nabla u_\varepsilon\|_{L^p(\Omega)} \leq C_\Omega(p, a), \quad \forall p \in [1, +\infty[.$$

Lemma 4.3. *Let $\lambda \in [0, n[$. Then, $L^p(\Omega) \hookrightarrow L^{2, \lambda}(\Omega)$ provided for $p \geq \frac{2n}{n - \lambda}$. Moreover, there exists $C(\Omega) > 0$ such that*

$$\sup_{\substack{x \in \Omega \\ r > 0}} [r^{-\lambda} |v|_{L^2(B(x, r) \cap \Omega)}] \leq C(\Omega) |v|_{L^p(\Omega)}.$$

Proof. Setting $\Omega_{0, r} = B(x, r) \cap \Omega$, we have

$$r^{-\lambda} \int_{\Omega \cap B(x, r)} |v|^2 dx = r^{-\lambda} \int_{\Omega} |v|^2 \chi_{\Omega_{0, r}} dx \leq r^{-\lambda} \|v\|_{L^p(\Omega)}^2 |\Omega_{0, r}|^{\frac{p-2}{p}} \leq c(\Omega) \|v\|_{L^p(\Omega)}^2.$$

Thus,

$$\sup_{\substack{r > 0 \\ x_0 \in \Omega}} (r^{-\lambda} \int_{\Omega \cap B(x, r)} |v|^2 dx) \leq c(\Omega) \|v\|_{L^p(\Omega)}^2. \quad \square$$

Using Lemma 4.3 and relation (4.6), we obtain that

$$\|\nabla u_\varepsilon\|_{L^2(\Omega \cap B(x, r))}^2 \leq r^\lambda C_\Omega(n, \lambda, a). \quad (4.14)$$

By applying the homeomorphism function τ , we obtain

$$\|\nabla w_\varepsilon\|_{L^2(I^+(x_0; r))}^2 \leq r^\lambda C_\Omega(\lambda, a). \quad (4.15)$$


Therefore,

$$\|\nabla w_\varepsilon^2\|_{L^2(I^+(x_0;r))}^2 \leq c \int_{I^+(x_0;r)} \tilde{a}(y) dy + r^{\lambda+2} C_\Omega(\lambda, a). \quad \square \quad (4.16)$$

Next, we want to prove that for all $R < 1$

$$\sup_{\substack{\forall x_0 \in \overline{I^+(R)} \\ \forall \rho > 0}} \rho^{-n} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0;\rho))}^2 \leq C(R) < \infty. \quad (4.17)$$

Let $x_0 \in \overline{I^+(R)}$, we set $\delta_0 = \frac{1-R}{2}$ with $0 < R < 1$. We have two cases to be analyzed:



1st case: $\rho \geq \frac{1-R}{2}$

In this case, we have for any x_0

$$\rho^{-n} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0;\rho))}^2 \leq \left(\frac{2}{1-R}\right)^n \|\nabla w_\varepsilon\|_{L^2(I^+(1))}^2 \leq c_\Omega \frac{\|a\|_\infty}{(1-R)^n}. \quad (4.18)$$

2nd case: $0 < \rho < \frac{1-R}{2}$. Let r be such $0 < \rho < r \leq \frac{1-R}{2}$. We have two subcases: either $\Gamma_R \cap I(x_0;r) = \emptyset$ or $\Gamma_R \cap I(x_0;r) \neq \emptyset$.

(a) For the first case, $\Gamma_R \cap I(x_0;r) = \emptyset$, we first write $w_\varepsilon = w_\varepsilon^1 + w_\varepsilon^2$ and apply estimate (4.10) of Theorem 4.4 to derive

$$\begin{aligned} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0;\rho))}^2 &\leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \|\nabla w_\varepsilon^1 - \{\nabla w_\varepsilon^1\}_r\|_{L^2(I(x_0;r))}^2 \\ &\quad + \|\nabla w_\varepsilon^2 - \{\nabla w_\varepsilon^2\}_\rho\|_{L^2(I(x_0;\rho))}^2. \end{aligned}$$

Applying Theorem 4.5 and the decomposition $w_\varepsilon^1 = w_\varepsilon - w_\varepsilon^2$,

$$\begin{aligned} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0;\rho))}^2 &\leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_r\|_{L^2(I^+(x_0;r))}^2 \\ &\quad + r^{\lambda+2} C_\Omega(\lambda, a) + 2 \int_{I^+(x_0;r)} \tilde{a}(y) dy. \end{aligned}$$

Choosing $\lambda = n - 1$, one has for all $0 < \rho < r$

$$\|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0, \rho))}^2 \leq c(\nu) \left(\frac{\rho}{r}\right)^{n+2} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_r\|_{L^2(I^+(x_0, r))}^2 + c(\Omega)r^n. \quad (4.19)$$

Applying the iteration [Lemma 2.3](#) (see [\[8,12\]](#)) on [\(4.13\)](#) with

$$\Phi(\rho) = \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0, \rho))}^2,$$

we get

$$\|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0, \rho))}^2 \leq c \left(\frac{\rho}{r}\right)^n \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_r\|_{L^2(I^+(x_0, r))}^2 + c(\Omega)\rho^n.$$

Dividing by ρ^n , we obtain

$$\frac{1}{\rho^n} \|\nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho\|_{L^2(I^+(x_0, \rho))}^2 \leq \frac{c}{r^n} \|\nabla w_\varepsilon\|_{L^2(I^+(1))}^2 + c(\Omega). \quad (4.20)$$

(b) For the second case, $\Gamma_R \cap I(x_0; r) \neq \emptyset$, we need to use relation [\(4.11\)](#) of [Theorem 4.4](#) by distinguishing the x_j -direction, $j \leq n - 1$ and x_n -direction.

For $j = 1, \dots, n - 2$, since $w_\varepsilon = w_\varepsilon^1 + w_\varepsilon^2$, we have

$$\int_{I^+(x_0; \rho)} |D_j w_\varepsilon|^2 dx \leq 2 \int_{I^+(x_0; \rho)} |D_j w_\varepsilon^1|^2 dx + 2 \int_{I^+(x_0; \rho)} |D_j w_\varepsilon^2|^2 dx \quad (4.21)$$

Using [Theorem 4.4](#) relation [\(4.11\)](#) and [Theorem 4.5](#), we derive from this last relation that

$$\begin{aligned} \int_{I^+(x_0; \rho)} |D_j w_\varepsilon|^2 &\leq c \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; r)} |D_j w_\varepsilon^1|^2 dx + c\rho^n \\ &\leq c \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; r)} |D_j w_\varepsilon|^2 dx + c \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; r)} |D_j w_\varepsilon^2|^2 dx + c\rho^n. \end{aligned} \quad (4.22)$$

Using again [Theorem 4.5](#) with relation [\(4.21\)](#), we deduce

$$\int_{I^+(x_0; \rho)} |D_j w_\varepsilon|^2 dx \leq c \left(\frac{\rho}{r}\right)^{n+2} \int_{I^+(x_0; r)} |D_j w_\varepsilon|^2 dx + c\rho^n. \quad (4.23)$$

This last relation is valid for all $0 < \rho < r$ we may appeal the iteration lemma (see [\[12,39\]](#)), to derive

$$\int_{I^+(x_0; \rho)} |D_j w_\varepsilon|^2 dx \leq c \left(\frac{\rho}{r}\right)^n \int_{I^+(x_0; r)} |D_j w_\varepsilon|^2 dx + c\rho^n \quad (4.24)$$

Thus

$$\rho^{-n} \int_{I^+(x_0; \rho)} \left| D_j w_\varepsilon - \{D_j w_\varepsilon\}_\rho \right|^2 dx \leq \rho^{-n} \int_{I^+(x_0; \rho)} |D_j w_\varepsilon|^2 \leq c \frac{1}{r^n} \int_{I^+(1)} |D_j w_\varepsilon|^2 dx + c. \quad (4.25)$$

In the x_n -direction, we have from [Theorem 4.4](#), relation [\(4.12\)](#) and [Theorem 4.5](#) with $\lambda = n - 1$, for all $0 < \rho < r$

$$\begin{aligned}
\int_{I^+(x_0;\rho)} \left| D_n w_\varepsilon - \{D_n w_\varepsilon\}_\rho \right|^2 &\leq \int_{I^+(x_0;\rho)} \left| D_n w_\varepsilon^1 - \{D_n w_\varepsilon^1\}_\rho \right|^2 dx + \int_{I^+(x_0;\rho)} \left| D_n w_\varepsilon^2 \right|^2 dx \\
&\leq c \left(\frac{\rho}{r} \right)^{n+2} \int_{I^+(x_0;r)} \left| D_n w_\varepsilon^1 - \{D_n w_\varepsilon^1\}_r \right|^2 dx + c r^n \\
&\leq c \left(\frac{\rho}{r} \right)^{n+2} \int_{I^+(x_0;r)} \left| D_n w_\varepsilon - \{D_n w_\varepsilon\}_r \right|^2 dx.
\end{aligned} \tag{4.26}$$

Thus, we may appeal the iteration [Lemma 2.3](#) with $\Phi(\rho) = \int_{I^+(x_0;\rho)} \left| D_n w_\varepsilon - \{D_n w_\varepsilon\}_\rho \right|^2 dx$ to derive

$$\rho^{-n} \int_{I^+(x_0;\rho)} \left| D_n w_\varepsilon - \{D_n w_\varepsilon\}_\rho \right|^2 dx \leq \frac{c}{r^n} \int_{I^+(1)} |D_n w_\varepsilon|^2 dx + c. \tag{4.27}$$

In all the cases, from relations [\(4.20\)](#), [\(4.25\)](#), [\(4.27\)](#), there exists a constant c , $\forall 0 < \rho < r$, for all $x_0 \in \overline{I^+(R)}$

$$\rho^{-n} \left\| \nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho \right\|_{L^2(I^+(x_0;\rho))}^2 \leq c \frac{1}{r^n} \left\| \nabla w_\varepsilon \right\|_{L^2(I^+(1))}^2 + c, \tag{4.28}$$

which infer that

$$\sup_{\rho > 0, x_0 \in \overline{I^+(R)}} \rho^{-n} \left\| \nabla w_\varepsilon - \{\nabla w_\varepsilon\}_\rho \right\|_{L^2(I^+(x_0;\rho))}^2 \leq \frac{c}{(1-R)^n} < \infty.$$

This ends the proof of [\(4.17\)](#). \square

Applying τ^{-1} on relation [\(4.17\)](#), we derive that

$$u_\varepsilon \text{ remains in a bounded set of } W^1 bmo_r \left(\tau^{-1}(I^+(R)) \right) \text{ for all } R < 1.$$

In the local estimate, we have proved that $\nabla u_\varepsilon \in bmo_r(\Omega_0)^n$, $\forall \Omega_0 \subset\subset \Omega$, and in the estimate in a neighborhood of the boundary $\tau^{-1}(I^+(R)) = \Omega_{0,R}(x)$ for all $R < 1$, we proved that $\nabla u_\varepsilon \in bmo_r(\Omega_{0,R}(x))^n$ with $x \in \partial\Omega$. Collecting both results on local estimates and boundary estimates, we can conclude as in Campanato [\[8\]](#) that ∇u_ε remains in a bounded set of $bmo_r(\Omega)^n$. This implies that u_ε belongs to a bounded set of $W_0^1 bmo_r(\Omega)$. This shows that $u \in W_0^1 bmo_r(\Omega)$. \square

5. Case where the right hand side is $a(x)u^{-m}(x)$, $m > 0$

In this paragraph, we want to discuss the existence and the regularity of solution for the following problem

$$\begin{cases} Lu = -\operatorname{div}(A(x)\nabla u) = a(x)u^{-m}(x) & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

As we shall see the regularity of the solution relies not only on the value of m but also on the regularity of the coefficients of $A(x) = (a_{ij}(x))_{i,j}$ the domain is still a Lipschitz one. For an alternative proof see also [\[3\]](#). More precisely, we want to show:

Theorem 5.1 (Existence). Let $a \in L_+^\infty(\Omega)$, $m > 0$, and let $A(x) = (a_{ij}(x))_{i,j}$ be an α -coercive matrix with bounded coefficients. Then there exists a positive function $u \in H_{loc}^1(\Omega)$ such that

1) $u \in L^q(\Omega)$, $q = \frac{2^*}{2}(m+1)$, [2^* is the Sobolev exponent], $\frac{a}{u^m} \in L_{loc}^1(\Omega)$, and

$$\int_{\Omega} |u(x)|^q dx \leq C_{\Omega} \left(\frac{\|a\|_{\infty}}{\alpha} \right)^{\frac{2^*}{2}}.$$

2) $\forall \psi \in \mathcal{D}(\Omega)$, $\int_{\Omega} A(x) \nabla u \cdot \nabla \psi dx = \int_{\Omega} \frac{a(x)\psi(x)}{u^m(x)} dx$, and $u^{\frac{m+1}{2}} \in H_0^1(\Omega)$. Moreover, $\int_{\Omega} |\nabla u|^2 u^{m-1} dx \leq \frac{\|a\|_{\infty}}{\alpha m} \left(\frac{m+1}{2} \right)^2$.

In the special case of $0 < m \leq 1$, then $\frac{a}{u} \in L^1(\Omega, \delta)$ and $u \in H_0^1(\Omega)$.

Proof. Let $\varepsilon > 0$. Then, there exists a non-negative function $u_{\varepsilon} \in H_0^1(\Omega) \cap L^\infty(\Omega)$ satisfying

$$\int_{\Omega} A(x) \nabla u_{\varepsilon} \cdot \nabla \varphi dx = \int_{\Omega} \frac{a(x)\varphi(x)}{u_{\varepsilon}^m + \varepsilon} dx, \quad \forall \varphi \in H_0^1(\Omega). \quad (5.1)$$

Then, one has the following *a priori* estimates

$$\int_{\Omega} |\nabla(u_{\varepsilon}^{\frac{m+1}{2}})|^2 dx \leq \frac{\|a\|_{\infty}}{\alpha m} \left(\frac{m+1}{2} \right)^2, \quad (5.2)$$

and there exists $C_{\Omega} > 0$, such that

$$\int_{\Omega} |u_{\varepsilon}|^q dx \leq C_{\Omega} \left(\frac{\|a\|_{\infty}}{\alpha} \right)^{\frac{2^*}{2}} \quad (5.3)$$

with $q = \frac{2^*}{2}(m+1)$, again [2^* is the Sobolev exponent].

Indeed, for the first inequality, we choose $\varphi = u_{\varepsilon}^m$ as a test function and we use the coercivity condition on A to derive that

$$m\alpha \int_{\Omega} \nabla u_{\varepsilon} \cdot \nabla u_{\varepsilon}^m dx \leq \int_{\Omega} A(x) \nabla u_{\varepsilon} \cdot \nabla u_{\varepsilon}^m dx = \int_{\Omega} \frac{a(x)u_{\varepsilon}^m}{u_{\varepsilon}^m + \varepsilon} \leq \|a\|_{\infty}.$$

Therefore,

$$\int_{\Omega} |\nabla u_{\varepsilon}|^2 u_{\varepsilon}^{m-1} dx \leq \frac{\|a\|_{\infty}}{\alpha m}.$$

Consequently,

$$\int_{\Omega} |\nabla(u_{\varepsilon}^{\frac{m+1}{2}})|^2 dx = \left(\frac{m+1}{2} \right)^2 \int_{\Omega} |\nabla u_{\varepsilon}|^2 u_{\varepsilon}^{m-1} dx \leq \left(\frac{m+1}{2} \right)^2 \frac{\|a\|_{\infty}}{\alpha m}. \quad (5.4)$$

While for the second inequality, we shall set $v_\varepsilon = u_\varepsilon^{\frac{m+1}{2}}$ and we have

$$\int_{\Omega} |u_\varepsilon|^q dx = \int_{\Omega} |v_\varepsilon|^{2^*} dx \leq C_{\Omega} \left(\int_{\Omega} |\nabla v_\varepsilon|^2 dx \right)^{\frac{2^*}{2}} \leq C_{\Omega} \left(\frac{\|a\|_{\infty}}{\alpha m} \right)^{\frac{2^*}{2}} \left(\frac{m+1}{2} \right)^2.$$

Using the Rellich–Kondrachov compactness, we may assume that there exists $v \in H_0^1(\Omega)$ such that $v \geq 0$ and $v_\varepsilon(x) \rightarrow v(x)$ a.e. in Ω . Then,

$$v_\varepsilon^{\frac{2}{m+1}} = u_\varepsilon \rightarrow v(x)^{\frac{2}{m+1}} = u(x), \quad \text{a.e. and } v_\varepsilon \rightarrow v \text{ weakly in } H_0^1(\Omega).$$

So,

$$\int_{\Omega} |\nabla v(x)|^2 dx \leq \frac{\|a\|_{\infty}}{\alpha m} \left(\frac{m+1}{2} \right)^2.$$

By Fatou's lemma, we have from the estimate (5.5)

$$\int_{\Omega} |u|^q dx \leq C_{\Omega} \left(\frac{\|a\|_{\infty}}{\alpha m} \right)^{\frac{2^*}{2}} \left(\frac{m+1}{2} \right)^2. \quad (5.5)$$

Then we deduce from the Theorem 2.2 that for all open set Ω_0 relatively compact in Ω , we have for a.e. $y \in \overline{\Omega}_0$

$$u_\varepsilon(y) \geq C'_{\Omega_0} \int_{\Omega_0} \frac{a(x)\delta(x)}{u_\varepsilon^m + \varepsilon} dx > 0. \quad (5.6)$$

We know from the Egorov theorem that there is a set B in Ω_0 of positive measure on which $\sup_{y \in B} \sup_{\varepsilon > 0} u_\varepsilon(y) = M$ is finite. Thus, there is a constant $C(\Omega_0) > 0$ such that $\forall \varepsilon > 0$

$$\int_{\Omega_0} \frac{a(x)\delta(x)}{u_\varepsilon^m + \varepsilon} dx \leq C(\Omega_0). \quad (5.7)$$

By Fatou's lemma, we deduce

$$0 < \int_{\Omega_0} \frac{a(x)}{u^m(x)} \delta(x) dx \leq C(\Omega_0) < +\infty. \quad (5.8)$$

• If $0 < m \leq 1$, we can choose $\varphi = u_\varepsilon$ as a test function and get

$$\alpha \int_{\Omega} |\nabla u_\varepsilon|^2 dx \leq \int_{\Omega} A(x)(\nabla u_\varepsilon)^2 dx = \int_{\Omega} \frac{a(x)u_\varepsilon}{u_\varepsilon^m + \varepsilon} \leq \|a\|_{\infty} \int_{\Omega} u_\varepsilon^{1-m} dx.$$

This implies that

$$\int_{\Omega} |\nabla u_\varepsilon|^2 dx \leq \frac{\|a\|_{\infty}}{\alpha} \int_{\Omega} u_\varepsilon^{1-m} dx \leq C(\Omega, a, \alpha), \quad \left(\text{since } 0 \leq 1-m < \frac{2^*}{2}(m+1) \right).$$

This and (5.7) yield that $u \in H_0^1(\Omega)$ and that u_ε converges weakly to u in $H_0^1(\Omega)$.

• If $m > 1$, then from (5.7) and (5.8), we have

$$\lambda_0 = \inf_{\varepsilon < \varepsilon_1} \int_{\Omega_0} \frac{a(x)\delta(x)}{u_\varepsilon^m(x) + \varepsilon} dx > 0, \quad \text{for some } \varepsilon_1 > 0.$$

From (5.4) and (5.6), we have

$$\frac{\|a\|_\infty}{\alpha} \geq \int_{\Omega_0} |\nabla u_\varepsilon|^2 u_\varepsilon^{m-1} dx \geq C_{\Omega_0}'^{m-1} \lambda_0^{m-1} \int_{\Omega_0} |\nabla u_\varepsilon|^2 dx. \quad (5.9)$$

We then deduce that

$$\int_{\Omega_0} |\nabla u_\varepsilon|^2 dx \leq C_{\Omega_0} < +\infty,$$

and jointly with the estimate on the L^q norm of u_ε , we deduce that u_ε remains in a bounded set of $H_{loc}^1(\Omega)$. Therefore, since for all $\psi \in \mathcal{D}(\Omega)$, we have

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} \frac{a(x)\psi(x)}{u_\varepsilon^m(x) + \varepsilon} dx = \int_{\Omega} \frac{a(x)\psi(x)}{u^m(x)} dx$$

$\forall \Omega_0 \subset\subset \Omega$, recalling that $u_\varepsilon(y) \geq C_{\Omega_0}' \lambda_0 > 0$, and u_ε converges weakly to u in $H^1(\Omega_0)$ for all $\Omega_0 \subset\subset \Omega$, we get the statement 2) of the theorem. \square

Remark 3 (The uniqueness of u_ε and u). The function u_ε is unique since the mapping $t \rightarrow \frac{a(x)}{t^m + \varepsilon}$ is decreasing and the regularity of u_ε allows at to choose $\varphi = u_\varepsilon - \overline{u_\varepsilon}$ as a test function whenever u_ε and $\overline{u_\varepsilon}$ are two different solutions. The same remarks holds for u when $0 < m \leq 1$.

Next, we want to study the regularity of the function $u = \lim u_\varepsilon$ constructed in Theorem 5.1.

Theorem 5.2 (Regularity). Assume that the operator L satisfies the Uniform Hopf Inequality. Then, the function $u = \lim u_\varepsilon$ satisfies

$$1) \quad \frac{a}{u^m} \in L^1(\Omega; \delta) \text{ and } u(y) \geq C_\Omega \delta(y) \int_{\Omega} \frac{a\delta}{u^m}(x) dx \text{ for a.e. } y \in \Omega.$$

2) For all $m \geq 1$,

$$\int_{\Omega} |\nabla u|^2 \delta^{m-1} dx \leq C_\Omega' \frac{\|a\|_\infty}{m\alpha \left(\int_{\Omega} \frac{a\delta}{u^m}(x) dx \right)^{m-1}}.$$

In particular, $u \in W^{1,2}(\Omega; \delta^{m-1})$.

3) Assume that $a_{ij} \in vmo(\Omega) \cap L^\infty(\Omega)$ and $\partial\Omega$ is C^1 , then

a) if $0 < m \leq 1$, then $u \in W_0^{1,p}(\Omega)$ for all $1 \leq p < +\infty$,

b) if $1 < m < 2$, then $u \in W_0^{1, \frac{1}{m-1}, \infty}(\Omega)$.

4) Assume that $a_{ij} \in C^{0,1}(\overline{\Omega})$ and $\partial\Omega$ is $C^{1,1}$. Then, if $0 < m \leq 1$, we have

$$u \in W_0^1 bmo_r(\Omega).$$

Proof. By Fatou's lemma, we have

$$\liminf_{\varepsilon} \int_{\Omega} \frac{a\delta}{u_{\varepsilon}^m + \varepsilon}(x) dx \geq \int_{\Omega} \frac{a\delta}{u^m}(x) dx > 0, \quad \text{since } a \neq 0, \quad (5.10)$$

and $u(x) < +\infty$ a.e. On the other hand, since L satisfies the Uniform Hopf Inequality, we get

$$u_{\varepsilon}(x) \geq C_{\Omega} \delta(x) \int_{\Omega} \frac{a\delta}{u_{\varepsilon}^m + \varepsilon}(y) dy. \quad (5.11)$$

Relations (5.10) and (5.11) with the fact that $u_{\varepsilon}(x)$ converges to $u(x)$ a.e., infers statement 1).

Now combining this last inequality with (5.4), one has, for $m \geq 1$,

$$m\alpha C_{\Omega}^{m-1} \left(\int_{\Omega} \frac{a\delta}{u_{\varepsilon}^m + \varepsilon}(y) dy \right)^{m-1} \int_{\Omega} |\nabla u_{\varepsilon}|^2 \delta^{m-1} dx \leq \|a\|_{\infty}. \quad (5.12)$$

Thanks to (5.10) and (5.12), we deduce that u_{ε} remains in a bounded set of $W^{1,2}(\Omega; \delta^{m-1}) = \{\varphi \in L^2(\Omega; \delta^{m-1}) : \int_{\Omega} |\nabla \varphi|^2 \delta^{m-1} dx < +\infty\}$. Therefore, u_{ε} converges weakly to u in $W^{1,2}(\Omega; \delta^{m-1})$ and from relation (5.12), we then have

$$m\alpha C_{\Omega}^{m-1} \left(\int_{\Omega} \frac{a\delta}{u^m}(y) dy \right)^{m-1} \int_{\Omega} |\nabla u|^2 \delta^{m-1} dx \leq \|a\|_{\infty}. \quad (5.13)$$

This proves the second statement.

To prove 3.a), let us show that $\frac{a}{u_{\varepsilon}^m + \varepsilon} \in W^{-1,p}(\Omega)$ with $1 < p < +\infty$. If $0 < m \leq 1$, one has from (5.11), by taking $\varphi \in W_0^{1,p'}(\Omega)$, $\frac{1}{p} + \frac{1}{p'} = 1$, $\varphi \geq 0$,

$$0 \leq \int_{\Omega} \frac{a}{u_{\varepsilon}^m + \varepsilon} \varphi dx \leq \frac{C_{\Omega}^0 \|a\|_{\infty}}{\left(\int_{\Omega} \frac{a\delta}{u_{\varepsilon}^m + \varepsilon} dy \right)^m} \int_{\Omega} \frac{\varphi}{\delta^m} dx \leq C_{\Omega}^1 \left(\int_{\Omega} \frac{\varphi}{\delta^m} dx \right). \quad (5.14)$$

If $0 < m \leq 1$, we have $\frac{1}{\delta^m} = \frac{1}{\delta} \delta^{1-m} \leq C_{\Omega} \frac{1}{\delta}$, so the Hardy inequality leads to

$$0 \leq \int_{\Omega} \frac{a}{u_{\varepsilon}^m + \varepsilon} \varphi dx \leq C_{\Omega}^1 \int_{\Omega} \frac{\varphi}{\delta} dx \leq C_{\Omega}^2 \left\| \frac{\varphi}{\delta} \right\|_{L^{p'}(\Omega)} \leq C \|\nabla \varphi\|_{W^{1,p'}(\Omega)}, \quad p' > 1,$$

which implies that

$$\left\| \frac{a}{u_{\varepsilon}^m + \varepsilon} \right\|_{W^{-1,p}(\Omega)} \leq C(p) \quad \text{uniformly in } \varepsilon.$$

Now, to prove 3.b), let us show that $\frac{a}{u_\varepsilon^m + \varepsilon} \in W^{-1}L^{\frac{1}{m-1}, +\infty}(\Omega)$, with $W^{-1}L^{\frac{1}{m-1}, +\infty}(\Omega)$ being the dual of $W_0^1L^{\frac{1}{m-1}, +\infty}(\Omega)$.

If $1 < m < 2$, we deduce from relation (5.11) and Hardy inequality with weights (see statement 2, section 1) $\forall \varphi \in C_c^1(\Omega)$

$$\begin{aligned} 0 &\leq \int_{\Omega} \frac{a}{u_\varepsilon^m + \varepsilon} |\varphi| dx \leq C'_\Omega \int_{\Omega} |\varphi| \delta^{-m} dx \leq C'_\Omega \int_{\Omega} |\nabla \varphi| \delta^{1-m} dx \\ &\leq C'_\Omega \|\nabla \varphi\|_{L^{n_m, 1}(\Omega)} \|\delta^{1-m}\|_{L^{\frac{1}{m+1}, \infty}(\Omega)}, \quad n_m = \frac{1}{2-m}, \end{aligned}$$

according to Díaz–Rakotoson ([15], pp. 53–54), $\delta^{1-m} \in L^{\frac{1}{m-1}, +\infty}(\Omega)$. Thus,

$$\frac{a}{u_\varepsilon^m + \varepsilon} \in W^{-1}L^{\frac{1}{m-1}, \infty}(\Omega).$$

Now, we apply the regularity result to u_ε satisfying $-\operatorname{div}(A(x)\nabla u_\varepsilon) = \frac{a}{u_\varepsilon^m + \varepsilon}$.

If $1 < m < 2$,

$$\|\nabla u_\varepsilon\|_{L^{\frac{1}{m-1}, \infty}(\Omega)} \leq C_\Omega \|\delta^{1-m}\|_{L^{\frac{1}{m-1}, \infty}(\Omega)} < +\infty.$$

If $0 < m \leq 1$, for $1 < p < +\infty$,

$$\|\nabla u_\varepsilon\|_{L^p(\Omega)} \leq C_\Omega(p) < +\infty.$$

Finally, for statement 4), if $a_{ij} \in C^{0,1}(\overline{\Omega})$, then u_ε remains in a bounded set of $W^1bmo_r(\Omega)$. Indeed, we write $\frac{a}{u_\varepsilon^m + \varepsilon} = \frac{a}{u_\varepsilon + \varepsilon} \cdot \frac{u_\varepsilon + \varepsilon}{u_\varepsilon^m + \varepsilon} = \frac{a_\varepsilon}{u_\varepsilon + \varepsilon}$. We have

$$0 \leq a_\varepsilon = a \cdot \frac{u_\varepsilon + \varepsilon}{u_\varepsilon^m + \varepsilon} \leq \|a\|_\infty u_\varepsilon^{1-m} \leq C_\Omega.$$

If $0 < m \leq 1$, since u_ε remains in a bounded set of $W_0^{1,p}(\Omega)$, $p > n$ according to statement 3.a), thus $u_\varepsilon \in W_0^{1,p}(\Omega)$, $\forall p < +\infty$ and satisfies

$$-\operatorname{div}(A(x)\nabla u_\varepsilon) = \frac{a_\varepsilon}{u_\varepsilon + \varepsilon}, \quad (5.15)$$

with a_ε remaining in a bounded set of $L^\infty(\Omega)$. Thus, we conclude (following the $bmo_r(\Omega)$ result previously proved) that the solution u_ε of (5.15) is in a bounded set of $W_0^1bmo_r(\Omega)$. \square

Acknowledgments

The research of J.I. Díaz was partially supported by the projects Ref. MTM2011-26119 and MTM2014-57113 of the DGISPI (Spain) and Research Group MOMAT (Ref. 910180) of the UCM.

References

- [1] P. Ausher, M. Qafsaoui, Observations on $W^{1,p}$ estimates for divergence elliptic equations with vmo coefficients, *Boll. Unione Mat. Ital.* 8 5-B (2002) 487–509.
- [2] H. Blasius, Grenzschichten in Flüssigkeiten mit Kleiner Reibung, *Z. Math. Phys.* 56 (1908) 1.
- [3] L. Boccardo, L. Orsina, Semilinear elliptic equations with singular nonlinearities, *Calc. Var. Partial Differential Equations* 37 (2010) 363–370.
- [4] B. Bougherara, J. Giacomoni, J. Hernández, Some remarks on the regularity of solutions to a singular elliptic problem, DCDS, in preparation.
- [5] H. Brézis, F. Browder, A property of Sobolev spaces, *Comm. Partial Differential Equations* 4 (1979) 1077–1083.
- [6] H. Brezis, X. Cabré, Some simple nonlinear PDE's without solutions, *Boll. Unione Mat. Ital.* B (1998) 223–262.
- [7] S. Byun, Elliptic equations with BMO coefficients in Lipschitz domains, *Trans. Amer. Math. Soc.* 375 (2005) 1025–1046.
- [8] S. Campanato, Equazioni ellittiche del secondo ordine e spazi $L^{2,\lambda}(\Omega)$, *Ann. Mat. Pura Appl.* (4) LXIX (1965) 321–381.
- [9] A. Canino, M. Degiovanni, A variational approach to a class of singular semilinear elliptic equations, *J. Convex Anal.* 11 (1) (2004) 147–162.
- [10] D.C. Chang, The dual of Hardy spaces on a bounded domain in \mathbb{R}^n , *Forum Math.* 6 (1) (1994) 65–81.
- [11] D.R. Chang, G. Dafni, E. Stein, Hardy spaces, BMO, and boundary value problems for the Laplacian on a smooth domain in \mathbb{R}^N , *Trans. Amer. Math. Soc.* 351 (4) (1999) 1605–1661.
- [12] Y.-Z. Chen, L.-C. Wu, Second Order Elliptic Equations and Elliptic Systems, vol. 174, Amer. Mathematical Society, Philadelphia, 1998.
- [13] M.G. Crandall, P.H. Rabinowitz, L. Tartar, On a Dirichlet problem with singular nonlinearity, *Comm. Partial Differential Equations* 2 (1977) 193–222.
- [14] L. Crocco, *Monografie Scientifiche: Aeronautica*, vol. 3, 1946 (translated from North American Aviation, Report CF-1038).
- [15] J.I. Díaz, J.M. Rakotoson, Elliptic problems on the space of weighted with the distance to the boundary integrable functions revisited, *Electron. J. Differ. Equ. Conf.* 21 (2014) 45–59.
- [16] J.I. Díaz, J. Hernández, J.M. Rakotoson, On very weak positive solutions to some semilinear elliptic problems with simultaneous singular nonlinear and spatial dependence terms, *Milan J. Math.* 79 (2011) 233–245.
- [17] W. Fulks, J.S. Maybee, A singular non-linear equation, *Osaka J. Math.* 12 (1960) 1–19.
- [18] J. Giacomoni, I. Schindler, P. Takac, Sobolev versus Holder local minimizers and global multiplicity for singular and quasilinear equations, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (5) 6 (1) (2007) 117–158.
- [19] J. Giacomoni, I. Schindler, P. Takac, $C^{0,\beta}$ -regularity and singular quasilinear elliptic systems, *Adv. Differential Equations* 20 (3–4) (2015) 259–298.
- [20] M. Giaquinta, Multiple integrals in the calculus of variations of a quasi conformal mapping, in: *Elliptic Systems*, Princeton Univ. Press, Princeton, NJ, 1983.
- [21] D. Gilbarg, S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, Berlin, 2001.
- [22] D. Goldberg, A local version of real Hardy spaces, *Duke Math. Notes* 46 (1979) 27–42.
- [23] M. Grüter, K.O. Widman, The Green function for uniformly elliptic equations, *Manuscripta Math.* 37 (3) (1982) 303–342.
- [24] C. Gui, F. Hua Lin, Regularity of an elliptic problem with a singular nonlinearity, *Proc. Roy. Soc. Edinburgh Sect. A* 123 (1993) 1021–1029.
- [25] J. Hernández, F.J. Mancebo, Singular elliptic and parabolic equations, in: M. Chipot, P. Quittner (Eds.), *Handbook of Differential Equations: Stationary Partial Differential Equations*, vol. 3, Elsevier BV, 2006, pp. 319–395.
- [26] H. Huber, M. Sieveking, Uniform bounds for quotients of Green functions on $C^{1,1}$ -domains, *Ann. Inst. Fourier* 32 (1) (1982) 105–117.
- [27] F. John, L. Nirenberg, On function of bounded mean oscillation, *Comm. Pure Appl. Math.* 14 (1961) 415–426.
- [28] P.W. Jones, Extension theorems for BMO, *Indiana Univ. Math. Notes* 29 (1) (1980) 41–66.
- [29] S. Karlin, Nirenberg, On a theorem of P. Nowosad, *J. Math. Anal. Appl.* 17 (1967) 61–67.
- [30] A. Kufner, *Weighted Sobolev Spaces*, Teuber Verlagsgesellschaft, Prague, 1980.
- [31] N. Meyers, An L^p -estimate for the gradient of solutions of second order elliptic divergence equations, *Ann. Sc. Norm. Super. Pisa* 17 (1980) 409–442.
- [32] J.M. Morel, L. Oswald, A uniform formulation for Hopf Maximum Principle, 1985, unpublished preprint.
- [33] P. Nowosad, On the integral equation $Kf = \frac{1}{f}$ arising in a problem in communication, *J. Math. Appl.* 14 (1966) 484–492.
- [34] O.A. Oleinik, Mathematical problem in the theory of the boundary layer, *Uspekhi Mat. Nauk* 23 (3) (1968) 3–65.
- [35] O.A. Oleinik, V.N. Samoklin, *Mathematical Models in Boundary Layer Theory*, Chapman & Hall/CRC, Boca Raton, USA, 1999.
- [36] L. Prandtl, Über Flüssigkeitsbewegung bei sehr kleiner Reibung, in: *Verhandlungen des III. Internationalen Mathematiker-kongresses*, Heidelberg, 1905, pp. 484–491, *LPGA* 2, 575–584.
- [37] J.M. Rakotoson, Quasilinear elliptic problems with measures as data, *Differential Integral Equations* 4 (1991) 449–457.
- [38] J.M. Rakotoson, Linear equation with data in non-standard spaces, *Atti Accad. Naz. Lincei, Math. Appl.* 26 (2015) 241–262, <http://dx.doi.org/10.4171/RLM/705>.
- [39] J.M. Rakotoson, Linear equations with variable coefficient and Banach functions spaces, forthcoming.
- [40] C. Simader, *On Dirichlet's Boundary Value Problem*, Springer, Berlin, 1972.
- [41] G. Stampacchia, Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus, *Ann. Inst. Fourier* 15 (1) (1965) 189–257.
- [42] E.M. Stein, *Harmonic Analysis Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, New Jersey, 1995.

- [43] K. Vajravelu, E. Soewono, R.N. Mohapatra, On solutions of some singular non linear differential equations arising in the theory of boundary layers, *J. Math. Anal. Appl.* 155 (2) (1991) 499–512.
- [44] L. Véron, Elliptic equations involving measures, in: M. Chipot, P. Quittner (Eds.), *Handbook of Differential Equations: Stationary Partial Differential Equations*, vol. 1, Elsevier BV, 2004, pp. 593–712.