



Optimal control and the value of information for a stochastic epidemiological SIS-model [☆]



Peter Grandits, Raimund M. Kovacevic ^{*}, Vladimir M. Veliov

*Institute of Statistics and Mathematical Methods in Economics, Vienna University of Technology,
Wiedner Hauptstraße 8/E105-4, A-1040 Vienna, Austria*

ARTICLE INFO

Article history:

Received 28 April 2017
Available online 6 April 2019
Submitted by M. Quincampoix

Keywords:

Stochastic differential equations
Optimal control
Epidemiology
SIS models
Hamilton-Jacobi-Bellman equation
Fokker-Planck equation

ABSTRACT

We analyze a continuous time stochastic SIS epidemiological model, describing the transmission dynamics of infectious diseases where infected individuals can become susceptible again after recovery. The infectivity period of infected individuals can be influenced dynamically by a decision maker, with the aim to minimize the expected aggregated economic costs due to illness and drug treatment. This stochastic control problem is investigated under two alternative assumptions about the information available. If a complete and exact measurement of the size of the infected population is available at any time, then the optimal control is sought in a state-feedback form and the related Hamilton-Jacobi-Bellman (HJB) equation can be used. If no state measurement is available, then the optimal control is sought in an open-loop form and the problem can be reformulated as an optimal control problem for the Kolmogorov forward equation. In both cases, deriving optimality conditions requires non-standard arguments due to the degeneracy of the involved HJB and Kolmogorov equations. Based on the obtained theoretical results, the role of the information pattern is numerically investigated.

© 2019 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

SIS-models form a class of simple prototypical epidemiological models. They assume that individuals can be infected multiple times throughout their lives with no immunity after each infection. Typical examples mentioned in literature are rota-viruses, some sexually transmitted as well as bacterial infections, see e.g. [17,5].

[☆] This work was supported by the Austrian Science Foundation (FWF) under grants P24125-N13 and P30864-N35.

^{*} Corresponding author.

E-mail addresses: peter.grandits@tuwien.ac.at (P. Grandits), raimund.kovacevic@tuwien.ac.at (R.M. Kovacevic), veliov@tuwien.ac.at (V.M. Veliov).

In the following we denote the number of infected individuals¹ at time $t \geq 0$ by $x(t)$ and the number of susceptible individuals by $y(t)$. We will use capital letters, when referring to stochastic entities. These notations differ from the usual notations in epidemiology ($I(t)$ for infected and $S(t)$ for susceptible) but are more in line with the conventions in stochastic calculus. Throughout this paper we restrict the discussion to the simplest case of a stationary population. Normalizing the population size to one,

$$x(t) + y(t) = 1,$$

means that $x(t)$ and $y(t)$ represent the fractions of infected and susceptible individuals. In this setup it suffices to regard only the number of infected, replacing $y(t)$ by $1 - x(t)$ wherever it appears.

The basic SIS model involves two parameters: the disease transmission coefficient (or strength of infection) $\beta > 0$, leading to a force of infection (or incidence rate) of $\beta x(t)$, and the recovery rate $\gamma > 0$. The force of infection models the rate at which susceptible individuals become infected, while the recovery rate is the rate at which infected individuals become susceptible again. One can interpret the reciprocal $1/\gamma$ as the average duration of infection.

Within this setup, the deterministic SIS model is described by the ordinary differential equation

$$\begin{aligned} \frac{dx(t)}{dt} &= \beta x(t)(1 - x(t)) - \gamma x(t), \\ x(0) &= x_0 \in [0, 1], \end{aligned} \quad (1)$$

which has an explicit solution whose properties are well known, see e.g. [27]. In particular the equation can be solved explicitly. Moreover, if the basic reproduction number $R_0 = \frac{\beta}{\gamma}$ is larger than 1, then the number of infected converges to a positive stable equilibrium (the disease becomes endemic), while otherwise it converges to zero (the disease dies out).

While many different approaches for stochastic epidemiological models exist in literature, see e.g. [11] or Chapter 6 of [20], in the present paper we analyze a diffusion-type version of the above SIS model: the deterministic equation (1) is replaced by a stochastic (Ito) differential equation (SDE)

$$\begin{aligned} dX(t) &= [\beta(1 - X(t))X(t) - \gamma X(t)] dt + \sigma[(1 - X(t))X(t)] d\mathcal{B}(t), \\ X(0) &= x_0 \in (0, 1). \end{aligned} \quad (2)$$

Here σ is a positive real number, $\mathcal{B}(\cdot)$ represents a standard Brownian motion in \mathbb{R} and the solution $X(\cdot)$ of (2) is a one dimensional Ito diffusion process (in contrast to $x(\cdot)$, which is a deterministic function).

Such a formulation can be obtained, if one e.g. assumes that the parameter β in (1) is not constant and deterministically known but random, when βdt is replaced with the specification

$$d\hat{\beta}(t) = \beta dt + \sigma d\mathcal{B}(t).$$

In this way it is possible to model random fluctuations of the parameter around the value β . Here the volatility parameter $\sigma > 0$ models the uncertainty of the true parameter process $\hat{\beta}$.

Keeling and Rohani [19] call the specification of noise used in (2) *external parameter noise* (in contrast to constant noise, scaled additive noise and heterogeneous parameter noise). While in other noise specifications the number of infected is directly subject to random fluctuations, here the parameters of an originally deterministic epidemiological model are disturbed by random noise due to external unpredictable forces. See also [1,2,10] for possible underlying mechanisms.

¹ Strictly speaking, “number of susceptible/infected individuals” means “size of the susceptible/infected sub-population”, which is a real number.

So far, the literature on stochastic epidemiological SDE-models mainly focuses on descriptive models, where the parameters – in our case β, σ and in particular γ are static and fixed. The simple stochastic SIS model (2) e.g., has been analyzed in [16], see also [12,6] and [23]. The diffusion term of (2) may become zero, which complicates the analysis considerably. While it is not possible to solve this equation analytically, the four cited papers analyze the existence of stationary distributions. [23] shows their existence if the modified basic reproduction number $R_0 - \frac{1}{2} \frac{\sigma^2}{\gamma}$ is larger than one and also gives an explicit representation of the stationary density.

In the present paper we focus on finite time horizons and make a step from descriptive to optimization models in the stochastic context. To this end, we assume that the “natural” development of the disease can be influenced by a decision maker. Various specifications are possible, but here we focus on the case where the decision maker is able to control the recovery rate γ that reflects the duration of the infectivity period of the infected individual, leading to faster recovery. The recovery rate can be controlled dynamically over time, which leads to a stochastic optimal control problem. Such a control can be implemented e.g. by increasing the treatment capacity or the efficiency of medication, dependent on time or the observed circumstances.

Using (deterministic) optimal control theory in epidemiological models dates from the early days of this theory, see e.g. [26]. The literature is vast and varies from simple compartmental models, as the SIS one, to models involving distributed dynamics (heterogeneity in space, age, contact rates, etc.). We mention [4], where treatment and prevention (vaccination) are used as controls in a SIR model, [7] and [8] where control theory with treatment and vaccination is applied to a SIRS model in a network. An extended SIR-type model with the same controls is investigated in the recent paper [9], which also contains more detailed bibliography on the mathematical tools involved. Besides optimal control theory, also alternative approaches to optimization and control of deterministic epidemiological models have been proposed in literature, see e.g. the broad survey [28].

There is also some work on (static) optimization for stochastic epidemiological models in literature. Several papers analyze vaccination in a stochastic environment, see e.g. [3]. [23] analyzes cost optimal long term treatment for two kinds of stochastic SIS models, aiming at their stationary distribution. The recent work [24] numerically compares several (fixed) control strategies for SIS processes on networks. However, optimality of stochastic control problems for epidemiological models seems to be a largely neglected topic so far. As we will see later, one reason may be that for epidemiological SDE-models usually there are implied boundaries due to the fact that the diffusion term is not bounded away from zero. This issue requires separate analysis depending on the concrete model specification, and does not allow to use the “usual” standard results on optimal control.

From a mathematical point of view, the main contribution of this work consists in dealing with this kind of degeneracy in the analysis of the underlying partial differential equations. We do this for a specification with fully observable states and for a specification, where it is not possible to observe the states over the planning horizon.

In order to clarify the issue we first consider the deterministic SIS model (1). Over the whole paper we will use the notations

$$a(x, u) = \beta x(1 - x) - (\gamma + u)x, \quad b(x) = \sigma^2 x^2(1 - x)^2 \tag{3}$$

(the notation b will come into play a bit later).

As before, we assume that γ is the recovery rate without any action by the decision maker, while $\gamma + u(t)$ is the recovery rate at time t if a control $u(t) \geq 0$ is applied at t . Thus the control variable $u(t)$ can be interpreted as treatment, accelerating recovery.

Then, building on the model (1) it is reasonable to consider the following standard optimal control problem, further referred to as the *Deterministic Problem* (DP):

$$\min_u \int_0^T e^{-rt} [F(x(t), u(t))] dt + c_3 x(T),$$

subject to

$$\begin{aligned} \frac{dx(t)}{dt} &= a(x(t), u(t)), & x(0) &= x_0 \in [0, 1], \\ u(t) &\geq 0, \end{aligned} \quad (4)$$

where $[0, T]$ is the given planning horizon, $r \geq 0$ is an interest rate (which can be taken equal to zero). Moreover, $F(x, u)$ is some cost function, depending on the number of infected and on the control and $c_3 x(T)$ is the expected accumulated future cost of the individuals that remain infected at T (can be taken equal to zero if the disease is expected to nearly die out till time T).

We mention that DP is formulated as an open-loop problem (the control function u is sought as a function of time only), but even if a feedback control is allowed, in absence of disturbances it would generate the same optimal trajectory $x(t)$ as the optimal open-loop control.

Using a cost function F , we assume that the control is costly, especially in case it is decided that the administration of the drug has to take place widely at the population level during an epidemic. Over the whole paper we give results for one specific cost model, that can be motivated as follows. We assume that all infected individuals are equally treated, that the cost is proportional to the number of treated individuals, $x(t)$, and the cost per a treated individual depends quadratically on the treatment effort $u(t)$, which means that it is marginally increasing in $u(t)$. Therefore, we specify the overall control costs per unit time as $c_2 x(t)u(t)^2$, where c_2 is a positive constant. In addition, taking an economic point of view, we assume that any infected individual creates for the society cost $c_1 > 0$ per unit of time, due to lost working hours and standard medical care, not including the treatment u . This leads to the cost function

$$F(x, u) = c_1 x + c_2 x u^2. \quad (5)$$

For one of the stochastic counterparts of the DP (the case of exact measurement and feedback control, see the two model cases described below) exclusively this cost specification will be used. Any alternative cost specification would afford a compatible verification theorem. In the open-loop formulation of the stochastic problem, a more general form of the cost function F is considered. The more general results are used to derive results for the original specification (5) for completeness.

Now we consider stochastic counterparts of DP. Since the Ito equation (2) generates a stochastic process, rather than a single trajectory, the information pattern becomes of crucial importance. Note that throughout this paper we will write $\mathbf{E}[\cdot]$ for the expectation and – if $X(t)$ denotes a stochastic process – $\mathbf{E}_{t,x}[\cdot]$ or even $\mathbf{E}_t[\cdot]$ for the conditional expectation given $X(t) = x$. We consider the following two “extreme” cases.

1. *Full information case* (Section 2). Here we assume that at any time t the exact current number of infected individuals becomes known to the decision maker. Correspondingly, the optimal control (which is a stochastic process) can be sought in feedback form: $u = u(X(s), s)$. Then the following optimal control problem (called further *Stochastic Feedback Problem* – SFP) becomes relevant:

$$\min_u \mathbf{E} \int_0^T e^{-rs} [F(X(s), u(X(s), s))] ds + c_3 \mathbf{E}[X(T)]. \quad (6)$$

subject to

$$dX(s) = a(X(s), u(X(s), s)) ds + \sqrt{b(X(s))} dB(s), \quad X(0) = x_0 \in (0, 1], \tag{7}$$

$$u(X(s), s) \geq 0. \tag{8}$$

Here the methods of stochastic control can be applied, in particular the Hamilton-Jacobi-Bellman (HJB) equation, which is employed for the analysis in Section 2. We show that the constraint $u(x, t) \geq 0$ is not necessary at the optimum and use the related necessary condition for the HJB equation to derive a reduced partial differential equation for the value function of the SFP. Strict arguments (including a regularity result and a verification theorem) are given for showing that under certain standing assumptions the unique solution of the reduced equation in fact is the value function of the SFP.

2. *No information case* (Section 3). Exact information about the current state of the disease is usually not available, especially in less developed countries. Therefore, we consider a second “extreme” scenario, where the probability density of $X(0)$ is given, but no information about the current state becomes available over the planning horizon $[0, T]$. In this case the feedback control cannot be implemented, therefore the optimal control is sought as function only of time: $u = u(t)$. Then a *Stochastic Open-loop Problem* (SOP) is considered, which has a similar formulation as SFP, but with $u(X(t), t)$ replaced by $u(t)$, a deterministic function of time. The HJB equation cannot be applied directly in this case. Still it is possible to use the Ito-Lemma in order to formulate a verification theorem that comes close to a HJB-formulation. However, we will see that this needs quite strong restrictions on the control function. Therefore we take a different approach. Namely, we employ the Kolmogorov-forward (or Fokker-Planck) equation related to (7), which describes the evolution of the density of $X(t)$ over time. Then SOP can be reformulated as an optimal control problem for a second order parabolic partial differential equation (the Fokker-Planck equation). In Section 3 we present and strictly prove Pontryagin type optimality conditions, which provide also a base for a numerical procedure.

The results for both model versions are used for an extensive illustrative example in Section 4, where both, the SFP and the SOP are solved numerically. While the reduced partial differential equation for the value function analyzed in Section 3 is used for solving the SFP, we use the steepest descent method for PDE-constraint optimal control problems in order to solve the SOP. In addition, we also solve the deterministic problem in a standard way. In particular we use the numerical results for comparing optimal solutions under differing information patterns: Assuming that reality is represented by the dynamics of the SOP, we analyze the gain (in terms of the objective function of SOP) from using the SOP instead of using the deterministic model DP. This quantifies the value of using the stochastic approach SOP. Moreover we also analyze the situation where the same dynamics still represents the reality for a decision maker, but another agent is able to observe the states. The expected gain that the agent may achieve (using SFP) compared to the decision maker (SOP) gives us an estimate for the amount that the decision maker is willing to pay for the information related to the observed states.

In the short Section 5 we discuss the possibility to include vaccination in the stochastic SIS model.

Acknowledgment

We wish to thank three anonymous referees for their valuable suggestions.

2. The stochastic closed loop optimal control problem

In the following we consider a decision maker who is able to observe the realized states $X(t)$ at any time $t \in [0, T]$ and who therefore is also able to react to these observations by choosing a feedback (that is, a state and time dependent) control $u(x, t)$. In this section we discuss properties of the optimization problem SFP (see (6)–(8)) in the HJB framework, including a regularity result and a verification theorem for the

associated value function. This is done for the cost specification (5) with the following additional assumption needed for technical reasons in the proof of the verification result.

Assumption 1. The constant c_3 in our model fulfills $c_3 < \tilde{c}_3$, where \tilde{c}_3 is the positive solution of the equation

$$-\gamma c_3 + e^{-rT} c_1 - \frac{e^{rT}}{4c_2} c_3^2 = 0.$$

2.1. Stochastic control for the SIS model with SDE constraints

With $X(t)$ observable at time t , the relevant σ -field is $\mathcal{F} = \{\mathcal{F}_t = \sigma(X(t))\}$, i.e. the σ -field generated by the process X . The control $u(\cdot, \cdot)$, is a function $(0, 1) \times [0, T] \rightarrow \mathbb{R}$ of observed states and time. The set of admissible feedback control functions, \mathcal{A} , is given by

$$\mathcal{A} := \{u(x, t) | u \text{ is Lipschitz w.r.t. } t \in [0, T] \text{ and locally Lipschitz w.r.t. } x \in [0, 1), u \geq 0\}. \tag{9}$$

In order to apply the dynamic programming approach we embed SFB into a family of problems parametrized by the initial time t and initial state x :

$$V(x, t) = \min_u \mathbf{E}_{t,x} \left[\int_t^T e^{-rs} [F(X(s), u(X(s), s))] ds + c_3 X(T) \right] \tag{10}$$

subject to

$$dX(s) = a(X(s), u(X(s), s)) ds + \sqrt{b(X(s))} dB(s) \quad X(t) = x, \tag{11}$$

$$u(X(s), s) \geq 0. \tag{12}$$

Remark 1. It is important to keep in mind the fact that, provided $u(x, t) \geq 0$ as assumed in our problem specification, the process $X(t)$, described by (11), stays in $I = (0, 1)$ forever, when started within this interval. The diffusion term $b(x) = \sigma^2 x^2 (1 - x)^2$ equals zero at $x = 0$ and $x = 1$. Moreover, the derivative of $b(x)$ also is equal to zero at these points, which in consequence leads to prescribed boundaries: at $x = 1$ the drift is negative, hence there is an entrance boundary, whereas at $x = 0$ we have $a(0, t) = 0$, which indicates a natural boundary (see e.g. [15] p. 119). As a natural boundary, $x = 0$ is absorbing but is never reached if the process starts in $I = (0, 1)$. In similar manner the entrance boundary $x = 1$ is never reached.

For a fixed function u , the backward operator \mathcal{L} related to (11) is given by

$$\mathcal{L}V(x, t) = V_t(x, t) + a(x, u(x, t))V_x(x, t) + \frac{1}{2}b(x)V_{xx}(x, t),$$

and we denote

$$\mathcal{L}^u V(x, t) = V_t(x, t) + a(x, u)V_x(x, t) + \frac{1}{2}b(x)V_{xx}(x, t). \tag{13}$$

Moreover, the related Hamilton-Jacobi-Bellman (HJB) equation is given by

$$\inf_{u \geq 0} \{ \mathcal{L}^u W(x, t) + e^{-rt} F(x, u) \} = 0 \tag{14}$$

$$W(x, T) = c_3 x,$$

which can be written as

$$W_t(x, t) + a(x, 0)W_x(x, t) + \frac{1}{2}b(x)W_{xx}(x, t) + e^{-rt}c_1x + \inf_{u \geq 0} \{e^{-rt}c_2xu^2 - uxW_x(x, t)\} = 0 \quad (15)$$

$$W(x, T) = c_3x.$$

In the following subsection we will show a regularity result and a verification theorem, in particular it is shown that under certain regularity conditions the HJB equation has a solution, which is the optimal value function of the SFB problem (10). Here, we start with some simple facts about the optimal value function V and the optimal control u , namely that V is monotonically increasing in its first argument and that the optimal u is nonnegative, even when the condition $u(X(s), s) \geq 0$ is omitted in SFB (10).

To this end, consider the optimization problem

$$\mathcal{V}(x, t) = \min_u \mathbf{E}_{t,x} \left[\int_t^T e^{-rs} [F(X(s), u(X(s), s))] ds + c_3X(T) \right] \quad (16)$$

subject to

$$dX(s) = a(X(s), u(X(s), s)) ds + \sqrt{b(X(s))} d\mathcal{B}(s) \quad X(t) = x, \quad (17)$$

which is in fact (10)-(12) without the restriction on u .

Lemma 1. *The value function $\mathcal{V}(x, t)$ of problem (16)-(17) is strictly increasing in x .*

Proof. Consider two states x_0, x_1 with $x_0 < x_1$. For $i \in \{0, 1\}$, denote an optimal strategy for a process starting at x_i at time t by $u_i^*(x, t)$ and the corresponding state process by $X_i^*(t)$. Using some $\varepsilon > 0$ define the following (non-optimal) strategy for a controlled process $\tilde{X}_0(t)$, starting at state x_0 at time t :

$$\tilde{u}_0(x, t) = \begin{cases} u_1^*(x, t) - \varepsilon & \text{if } x < X_1^*(t) \\ u_1^*(x, t) & \text{if } x = X_1^*(t). \end{cases} \quad (18)$$

In addition, define the stopping time $\tau = \inf \{t : \tilde{X}_0(t) = X_1^*(t)\}$ with $\inf \emptyset = T$ and denote $\tilde{\mathcal{V}}(x_1, t) = E \left[\int_t^T e^{-rs} (c_1\tilde{X}_0(s) + c_2\tilde{X}_0(s)\tilde{u}_1(\tilde{X}_0(s), s)^2) ds + c_3\tilde{X}_0(T) \right]$

Then we have $\tilde{X}_0(s) \leq X_1^*(s)$ almost surely at each time $t \in [0, T]$ and (due to continuity of the sample paths) $\tau > 0$ almost surely. For $t \in [0, \tau \wedge T)$ we have $\tilde{X}_0(s) < X_1^*(s)$. Furthermore $\tilde{u}_0(s, x) \leq u_1^*(s, x)$ is ensured by (18). This implies

$$\begin{aligned} \mathcal{V}(x_1, t) - \tilde{\mathcal{V}}(x_0, t) &= c_1 \mathbf{E} \left[\int_t^{\tau \wedge T} e^{-rs} (X_1^*(s) - \tilde{X}_0(s)) ds + \int_{\tau \wedge T}^T e^{-rs} (X_1^*(s) - \tilde{X}_0(s)) ds \right] \\ &+ c_2 \mathbf{E} \left[\int_0^T e^{-rs} (X_1^*(s)u_1^*(X_1^*(s), s)^2 - \tilde{X}_0(s)\tilde{u}_0(\tilde{X}_0(s), s)^2) ds \right] \\ &+ c_3 [X_1^*(T) - \tilde{X}_0(T)] > 0, \end{aligned}$$

because the first integral is positive ($\tau > 0$), whereas all other integrals as well as the last line are nonnegative.

Finally, because \tilde{u}_0 is not the optimal strategy for a controlled process starting at x_0 , we have

$$\mathcal{V}(x_0, t) \leq \tilde{\mathcal{V}}(x_0, t) < \mathcal{V}(x_1, t). \quad \square$$

It seems that the constraint $u(x, t) \geq 0$ may complicate the situation. However, based on Lemma (1) it can be shown that this constraint is irrelevant for the optimal solution.

Lemma 2. *The optimal control u^* of (16)-(17) is nonnegative. It follows that it also solves (10)-(12) and therefore that $V(x, t)$ is strictly monotone increasing in x .*

Proof. Let $u^*(x, t)$ be an optimal control for (16)-(17) and denote the related state process by $X^*(t)$.

Suppose that $u^*(t) < 0$ with positive probability on a set S of points in time, where S has positive Lebesgue-measure and choose a point in time τ_1 such that $\mathbb{P}(u(x, \tau_1) < 0) > 0$ and $\mathbb{P}(u(x, t) < 0) > 0$ for t in a set $S_1 \subseteq S$ with $\lambda(S_1) > 0$. Define now the control law

$$u'(x, t) = \begin{cases} u^*(x, t) & \text{for } t < \tau_1 \\ |u^*(x, t)| & \text{for } t \geq \tau_1. \end{cases}$$

Write $X'(t)$ for the related process of states. Because $X^*(\tau_1) = X'(\tau_1)$ we have almost surely $d(X^*(\tau_1) - X'(\tau_1)) \geq 0$ and $d(X^*(\tau_1) - X'(\tau_1)) = -2 \cdot u^*(\tau_1)X^*(\tau_1) > 0$ with positive probability. Therefore it can be inferred that $X^*(t) \geq X'(t)$ almost surely and $X^*(t) > X'(t)$ with positive probability for t in some (maybe small) interval $[\tau_1, \tau_2)$. In addition, $u^*(x, t)^2 = u'(x, t)^2$ holds almost surely. These facts imply

$$\mathbf{E}_{\tau_1} \int_{\tau_1}^{\tau_2} e^{-rs} (c_1 X^*(s) + c_2 X^*(s) u^*(X^*(s), s)^2) ds > \mathbf{E}_{\tau_1} \int_{\tau_1}^{\tau_2} e^{-rs} (c_1 X'(s) + c_2 X'(s) u'(X'(s), s)^2) ds. \quad (19)$$

Moreover, at time τ_2 the inequality $X^*(\tau_2) > X'(\tau_2)$ implies

$$\mathcal{V}(X'(\tau_2), \tau_2) < \mathcal{V}(X^*(\tau_2), \tau) \quad (20)$$

by Lemma 1.

It follows that a strategy u'' with related process X'' that switches from u^* to u' at time τ_1 and switches back to u^* at time τ_2 is a better strategy than keeping the optimal strategy u^* over the whole planning horizon. This contradicts the assumption that u^* is optimal, if $\tau_1 < T$. Hence, $u^*(x, t) \geq 0$ for problem (16)-(17), the condition $u^*(x, t) \geq 0$ is redundant in (10)-(12), and $V(x, t) = \mathcal{V}(x, t)$ holds. By applying Lemma 1, the function $V(x, t)$ must then be strictly monotone increasing. \square

The minimization in the HJB equation (14) leads to

$$u^*(t) = \frac{e^{rt}}{2c_2} W_x(x, t), \quad (21)$$

which is true only if the process starts in the open interval $(0, 1)$ – when $x = 0$ is almost surely never reached. Applying this result, reduces the HJB equation to the nonlinear equation

$$W_t(c, t) + a(x, 0)W_x(x, t) + \frac{1}{2}b(x)W_{xx}(x, t) + e^{-rt}c_1x - \frac{e^{rt}}{4c_2}x[W_x(x, t)]^2 = 0. \quad (22)$$

In fact we will show below (see Corollary 1) that there exists a smooth enough solution W of the equation (22) which fulfills $W_x(x, t) \geq 0$. This readily implies that W is also a solution of the HJB-equation (14) with optimal control (21).

2.2. A regularity result and the verification theorem

In order to show that the solution of the HJB equation (14), resp. (22), is indeed the value function of our problem, we will have to show a so called verification result. To prove this, one needs some regularity for the solution of the HJB equation. We provide this in the following theorem. Note that here and in the following $C^{2+\beta,1+\frac{\beta}{2}}$ denotes the space of functions, such that second derivative with respect to x is Hölder continuous with index β and the first derivative with respect to t is Hölder continuous with index $\frac{\beta}{2}$.

Theorem 1. Equation (22) has at least one solution $W(x, t)$, which is bounded by a positive constant M and which is an element of $C^{2+\beta,1+\beta/2}(Q_t)$, with $0 < \beta < 1$ and $Q_T := \{(x, t) | \epsilon \leq x \leq 1 - \epsilon, 0 \leq t \leq T\}$, for arbitrary small positive ϵ . Moreover, $W(x, t)$ is uniformly continuous at $x = 0$.

Proof. We want to apply in this proof standard results for parabolic Cauchy problems, see [25], hence we will first transform our x -variable to a z -variable, varying on $] - \infty, \infty[$. So let $z = \phi(x)$, with $\phi \in C^\infty$ and $\phi : (0, 1) \rightarrow] - \infty, \infty[$. We will need to have control of the transformation near the boundary, so we prescribe its value there explicitly, whereas we just impose ϕ to be strictly monotone increasing in the middle part, i.e.

$$\phi(x) := \begin{cases} \ln x, & x \in]0, \frac{1}{10}], \\ \text{strict. mon. increasing,} & x \in]\frac{1}{10}, \frac{9}{10}[, \\ -\ln(1-x), & x \in [\frac{9}{10}, 1[. \end{cases} \tag{23}$$

Hence, we also have

$$\begin{aligned} 0 < \text{const} \leq \phi'(x) \leq \text{Const} \\ |\phi''(x)| \leq \text{Const} \end{aligned} \tag{24}$$

for $x \in [\frac{1}{10}, \frac{9}{10}]$. Performing the transformation, we find

$$W_x = W_z z_x = \begin{cases} W_z e^{-z}, & x \in]0, \frac{1}{10}], \\ W_z \phi'(\phi^{(-1)}(z)), & x \in]\frac{1}{10}, \frac{9}{10}[, \\ W_z e^z, & x \in [\frac{9}{10}, 1[. \end{cases} \tag{25}$$

and

$$W_{xx} = \begin{cases} W_{zz} e^{-2z} - W_z e^{-2z}, & x \in]0, \frac{1}{10}], \\ W_{zz} \phi'(\phi^{(-1)}(z))^2 + W_z \phi''(\phi^{(-1)}(z)), & x \in]\frac{1}{10}, \frac{9}{10}[, \\ W_{zz} e^{2z} + W_z e^{2z}, & x \in [\frac{9}{10}, 1[. \end{cases} \tag{26}$$

It is a lengthy, but elementary calculation to plug this transformation into the equation (22). The result is

$$W_t + \kappa_1(z)W_z + \kappa_2(z)W_{zz} + \frac{e^{rt}}{4c_2} \kappa_3(z)W_z^2 + \kappa_4(z) = 0. \tag{27}$$

We do not provide the functions κ_i explicitly, but we give their properties, which we will need in the sequel, i.e.

$$\begin{aligned} \kappa_i(z) &\in C^\infty(-\infty, \infty), & i &= 1, \dots, 4 \\ |\kappa_i(z)| &\leq \text{const.}, & z &\in [-\ln 10, \ln 10], i = 1, \dots, 4 \\ \kappa_i(z) &\geq \text{const.} > 0, & z &\in [-\ln 10, \ln 10], i = 2, \dots, 4. \end{aligned} \tag{28}$$

Finally, the boundary condition transforms to

$$W(z, T) = c_3 \phi^{(-1)}(z) = c_3 \begin{cases} e^z, & z \leq -\ln 10, \\ \phi^{(-1)}(z), & z \in]-\ln 10, \ln 10[, \\ 1 - e^{-z}, & z \geq \ln 10. \end{cases} \tag{29}$$

We want to apply Theorem V.8.1 in [25], and in order to get useful information from this theorem, it is helpful to transform also the dependent variable via $\tilde{W}(z, t) := \frac{W(z, t)}{h(z)}$, with $h(z) \in C^\infty(\mathbf{R})$ and

$$h(z) := \begin{cases} e^z, & z \leq -\ln 10, \\ \text{strict. mon. increasing,} & z \in]-\ln 10, \ln 10[, \\ 1, & z \geq \ln 10. \end{cases} \tag{30}$$

Moreover, we choose $h(z)$ s.t. it fulfills

$$\begin{aligned} |h'(z)| &\leq \text{const.}, \forall z \in \mathbf{R}, \\ |h''(z)| &\leq \text{const.}, \forall z \in \mathbf{R}, \\ h(z) &\in [1/10, 1], z \in [-\ln 10, \ln 10]. \end{aligned} \tag{31}$$

We also introduce the “backwards time” $\tau := T - t$. Performing all these transformations gives after some elementary calculations instead of (27)

$$\begin{aligned} &\tilde{W}_\tau - \tilde{W}_{zz} \cdot \begin{cases} \sigma^2(1 - e^z)^2 \\ \kappa_2(z) \\ \sigma^2(1 - e^{-z})^2 \end{cases} + \tilde{W}_z^2 \frac{e^{r(T-\tau)}}{4c_2} \cdot \begin{cases} 1 \\ h(z)\kappa_3(z) \\ (1 - e^{-z})e^{2z} \end{cases} + \tilde{W}\tilde{W}_z \frac{e^{r(T-\tau)}}{4c_2} \cdot \begin{cases} 2 \\ h'(z)\kappa_3(z) \\ 0 \end{cases} \\ &+ \tilde{W}_z \cdot \begin{cases} -\beta(1 - e^z) + \gamma + \sigma^2(1 - e^z)^2 - 2\sigma^2(1 - e^z)^2 \\ -\kappa_1(z) - 2(h'(z)/h(z))\kappa_2(z) \\ -\beta(1 - e^{-z}) + \gamma(e^z - 1) - \sigma^2(1 - e^{-z})^2 \end{cases} + \tilde{W}^2 \frac{e^{r(T-\tau)}}{4c_2} \cdot \begin{cases} 1 \\ ((h'(z))^2/h(z))\kappa_3(z) \\ 0 \end{cases} \\ &+ \tilde{W} \cdot \begin{cases} -\beta(1 - e^z) + \gamma + \sigma^2(1 - e^z)^2 - \sigma^2(1 - e^z)^2 \\ -\kappa_1(z)(h'(z)/h(z)) - \kappa_2(z)(h''(z)/h(z)) \\ 0 \end{cases} - e^{-(T-\tau)}c_1 \cdot \begin{cases} 1 \\ \phi^{(-1)}(z)/h(z) \\ 1 - e^{-z} \end{cases} = 0 \end{aligned} \tag{32}$$

with the initial condition

$$\tilde{W}(z, 0) = \frac{c_3 \phi^{(-1)}(z)}{h(z)} = c_3 \begin{cases} 1, & z \leq -\ln 10, \\ \phi^{(-1)}(z)/h(z), & -\ln 10 < z < \ln 10, \\ 1 - e^{-z}, & z \geq \ln 10. \end{cases} \tag{33}$$

In this equation and in some of the following ones we adopt the convention that the first line holds always for $z \leq -\ln 10$, the third line for $z \geq \ln 10$ and the second one for the area in-between

As announced earlier, we now apply Theorem V.8.1 in [25]. In order to do this, we match first some notation and find using V.(6.4) there

$$\frac{\partial a_1}{\partial \tilde{W}_z}(z, \tau, \tilde{W}, \tilde{W}_z) = \begin{cases} \sigma^2(1 - e^z)^2, \\ \kappa_2(z), \\ \sigma^2(1 - e^{-z})^2. \end{cases} \tag{34}$$

Moreover, $A(z, \tau, \tilde{W}, \tilde{W}_z)$ denotes in our case all the terms of the l.h.s. of equation (32), but the \tilde{W}_τ and the \tilde{W}_{zz} -term.

We now start to check the assumptions of Theorem V.8.1. Concerning assumption a), it is clear that our initial condition, defined by (33), is bounded and an element of $C^{2+\beta}(\Omega)$, for all bounded domains $\Omega \subset \mathbf{R}$.

We proceed with condition b) : Equation V.(8.6) there is satisfied by our (34) and the third line of (28). Concerning the condition on A , we find that in our case

$$A(z, \tau, \tilde{W}, 0)\tilde{W} \geq -b_1|\tilde{W}|^3 - b_2$$

holds for some positive constants b_1, b_2 , because all the coefficients of \tilde{W}^2, \tilde{W} and “1” in (32) are uniformly bounded. This means that the second condition of b) with $\Phi(\tau) = b_1\tau^2$ is fulfilled.

Coming to condition c), we first find

$$a_1(z, \tau, \tilde{W}, \tilde{W}_z) = \tilde{W}_z \begin{cases} \sigma^2(1 - e^z)^2, \\ \kappa_2(z), \\ \sigma^2(1 - e^{-z})^2, \end{cases} \tag{35}$$

and, using V.(6.4) of [25], one gets

$$a(z, \tau, \tilde{W}, \tilde{W}_z) = A(z, \tau, \tilde{W}, \tilde{W}_z) + \frac{\partial a_1}{\partial \tilde{W}}\tilde{W}_z + \frac{\partial a_1}{\partial z} = A(z, \tau, \tilde{W}, \tilde{W}_z) + \tilde{W}_z \begin{cases} -2\sigma^2(1 - e^z)e^z, \\ \kappa'_2(z), \\ 2\sigma^2(1 - e^{-z})^2e^{-z}. \end{cases} \tag{36}$$

We have to check the conditions b) and c) of Theorem V.6.1 in [25] now. One easily sees that a_1 and a are continuous with respect to its variables and that a_1 is also differentiable.

Let now $Q_T := [-N_1, N_2] \times [0, T]$, for some (large) positive constants N_1, N_2 . One first finds that the constants of inequality V.(6.9/1) are independent of Q_T in our case.

Concerning the inequality V.(6.9/2), we realize for later use the

Fact 1. *The constants in inequality V.(6.9/2) depend on N_2 but not on N_1 .*

The reason for this are the e^{2z} , resp. the $(e^z - 1)$ terms in the coefficient of \tilde{W}_z^2 , resp. \tilde{W}_z of (32).

Finally, the Hölder continuity conditions of Theorem V.6.1c are easily seen to be satisfied. All in all, we can now apply Theorem V.8.1, giving us the following.

Assertion 1. *There exists at least one solution $\tilde{W}(z, \tau)$ of the Cauchy problem (32)+(33) in $[0, T] \times \mathbf{R}$, which does not exceed a positive constant M in modulus and belongs to $C^{2+\beta, 1+\beta/2}(Q_T)$ with $Q_T := [-N_1, N_2] \times [0, T]$ for some (large) positive constants N_1, N_2 . Moreover, because of Fact 1 above, it is also an element of $C^{2+\beta, 1+\beta/2}([-\infty, N_2] \times [0, T])$.*

Now, transforming back to the original variables, Assertion 1 provides almost Theorem 1, we just have to show continuity at $x = 0$. But this is easy: By Assertion 1, we have $|\tilde{W}(z, \tau)| \leq \text{const.}$ for all $(z, t) \in [-\infty, N_2] \times [0, T]$, giving $|W(z, \tau)| \leq \text{const.}h(z)$, or $|W(z, \tau)| \leq \text{const.}e^z$, for $z \leq -\ln 10$. Hence, finally $|W(x, \tau)| \leq \text{const.}x$, for $x \leq 1/10$. □

In a next step we will have to show that a solution $W(x, t)$ of (22), provided by Theorem 1, fulfills $W_x(x, t) \geq 0$ (we know this already for the value function, see Lemma 2). Once we know this, one gets that the optimizer u^* appearing in the HJB equation (14) is also nonnegative and given by the formula $u^* = \frac{e^{rt}}{2c_2}W_x$. This allows us in the proof of the verification theorem to restrict to nonnegative controls.

The method of our proof will be the following. We first consider an approximating optimization problem, analogous to the original one, but where we prescribe an artificial boundary condition at $x = 1 - \epsilon$. For this problem we do not have to show that the upper boundary is an entrance boundary. So a standard verification theorem will show that the solution of the HJB equation $W^{(\epsilon)}$ is the value function $V^{(\epsilon)}$ of our approximating problem. If we choose the artificial boundary condition properly, a simple modification of Lemma 2 will show $V_x^{(\epsilon)} > 0$, hence $W_x^{(\epsilon)} > 0$. Going with epsilon to zero, will yield the desired result.

So let us start with the definition of the approximating problem

$$V^{(\epsilon)}(x, t) = \min_{u \in \mathcal{A}^\epsilon} \mathbf{E}_{t,x} \left[\int_t^{\tau^{(\epsilon)}} e^{-rs} [F(X(s), u(X(s), s))] ds + c_3 X(\tau^{(\epsilon)}) \mathbf{1}_{\{X(\tau^{(\epsilon)}) < 1-\epsilon\}} + \Psi^{(\epsilon)}(X(\tau^{(\epsilon)})) \mathbf{1}_{\{X(\tau^{(\epsilon)}) = 1-\epsilon\}} \right], \tag{37}$$

where $\tau^{(\epsilon)}$ is the stopping time $\tau^{(\epsilon)} := \inf\{s \geq t | X(s) = 0\} \wedge \inf\{s \geq t | X(s) = 1 - \epsilon\} = \inf\{s \geq t | X(s) = 1 - \epsilon\}$, with the convention $\inf \emptyset = T$ and accounting for the fact that $x = 0$ is a natural boundary (see Remark 1). Moreover, $\mathcal{A}^\epsilon := \{u(x, t) | u \text{ is Lipschitz w.r.t. } t \in [0, T] \text{ and } x \in [0, 1 - \epsilon]\}$ are the admissible strategies. The state process is as in Lemma 1. For the artificial boundary condition we assume

$$\Psi^{(\epsilon)}(t) \in C^\infty([0, T]), \quad (\Psi^{(\epsilon)})'(t) \leq 0. \tag{38}$$

We now consider the boundary value problem

$$\begin{aligned} W_t^{(\epsilon)} + [\beta x(1-x) - \gamma x] W_x^{(\epsilon)} + \sigma^2 x^2(1-x)^2 W_{xx}^{(\epsilon)} + e^{-rt} c_1 x - \frac{e^{rt}}{4c_2} x(W_x^{(\epsilon)})^2 &= 0, \\ W^{(\epsilon)}(x, T) &= c_3 x, \\ W^{(\epsilon)}(0, t) &= 0, \\ W^{(\epsilon)}(1-\epsilon, t) &= \Psi^{(\epsilon)}(t). \end{aligned} \tag{39}$$

If we impose a compatibility condition on our boundary data at the point $(1 - \epsilon, T)$, i.e., if we assume that the first line of (39) holds in this point, we get

$$(\Psi^{(\epsilon)})'(T) + O(\epsilon) - \gamma c_3 + e^{-rT} c_1 - \frac{e^{rT}}{4c_2} c_3^2 = 0$$

Now, Assumption 1 guarantees that one can find a function $\Psi^{(\epsilon)}$ fulfilling (38), as well as the compatibility condition, for ϵ small enough. As the compatibility condition is fulfilled now, one finds, using the beginning of chapter V.8 of [25], the following lemma.

Lemma 3. *The boundary value problem (39) has a solution in $C^{2+\beta, 1+\beta/2}([0, 1 - \epsilon] \times [0, T])$, which converges – as $\epsilon \rightarrow 0$, after possibly extracting a subsequence – together with its derivatives on $]0, 1[$ to $W(x, t)$, a solution provided by Theorem 1.*

We proceed with the announced verification result for the approximating problem.

Proposition 1. *A $C^{2+\beta, 1+\beta/2}([0, 1 - \epsilon] \times [0, T])$ solution $W^{(\epsilon)}$ of the boundary value problem (BVP)*

$$\inf_{u \in \mathbf{R}} \left\{ \mathcal{L}^u W^{(\epsilon)} + F(x, u) \right\} = 0,$$

$$\begin{aligned} W^{(\epsilon)}(x, T) &= c_3 x, \\ W^{(\epsilon)}(0, t) &= 0, \\ W^{(\epsilon)}(1 - \epsilon, t) &= \Psi^{(\epsilon)}(t), \end{aligned}$$

is the value function of problem (37). (Hence, there is only one such solution.)

Proof. Note that the BVP above is just another way of writing the system (39) (\mathcal{L}^u stands for the generator of our state process).

We first remark that our assumption $u \in \mathcal{A}^\epsilon$ guarantees a unique strong solution of the underlying SDE. Applying Ito’s formula to $W^{(\epsilon)}$, which is regular enough, gives

$$W^{(\epsilon)}(X(\tau^{(\epsilon)}), \tau^{(\epsilon)}) = W^{(\epsilon)}(x, t) + \int_t^{\tau^{(\epsilon)}} \mathcal{L}^u W^{(\epsilon)}(X(s), s) \, ds + \int_t^{\tau^{(\epsilon)}} W_x^{(\epsilon)}(X(s), s) \sqrt{b(X(s))} \, dB(s).$$

Since σ and $W_x^{(\epsilon)}$ are bounded, the last integral is a true martingale and vanishes in the expectation. This gives

$$\begin{aligned} \mathbf{E}_{t,x} \left[\int_t^{\tau^{(\epsilon)}} F(X(s), u(s)) \, ds + W^{(\epsilon)}(X(\tau^{(\epsilon)}), \tau^{(\epsilon)}) \right] &= W^{(\epsilon)}(x, t) + \mathbf{E}_{t,x} \left[\int_t^{\tau^{(\epsilon)}} F(X(s), u(s)) \right. \\ &\quad \left. + \mathcal{L}^u W^{(\epsilon)}(X(s), s) \, ds \right]. \end{aligned}$$

By the HJB equation in the formulation of the proposition, the expectational value in the r.h.s. of the last equation is greater or equal to 0 for a general admissible strategy u , whereas it is equal to zero for the optimizer u^* . This concludes our proof. \square

Checking the proof of Lemma (2), we note that it holds as well for our approximating optimization problem, because of the second assumption in (38). This gives $W_x^{(\epsilon)}(x, t) = V_x^{(\epsilon)}(x, t) > 0$ and, together with Lemma 3, we end up with

Corollary 1. A solution $W(x, t)$ provided by Theorem 1 fulfills $W_x(x, t) \geq 0$ for $(x, t) \in [0, 1) \times [0, T]$.

Because of (21) this corollary implies that the restriction $u(t) \geq 0$ in the definition (9) of the admissible set is fulfilled automatically.

Finally we come to the main result of this section

Theorem 2. A solution W given by Theorem 1 is the value function of problem (10)-(12) for $0 \leq x < 1$ under Assumptions 1. (Hence, there is only one such solution.)

Proof. Let $\tau^{(\epsilon)}$ be as it is defined after equation (37) and let $u \in \mathcal{A}$, then we get, as in the proof of Proposition 1,

$$\mathbf{E}_{t,x} \left[\int_t^{\tau^{(\epsilon)}} F(X(s), u(s)) \, ds + W(X(\tau^{(\epsilon)}), \tau^{(\epsilon)}) \right] \geq W(x, t).$$

Since $x = 1$ is entrance (see the discussion in Remark 1), we can define the stopping time τ as $\tau := \inf\{s \geq t | X(s) = 0\}$, with the convention $\inf \emptyset = T$. There is a natural boundary at $x = 0$ (see Remark 1), hence we get a.s. $\tau^{(\epsilon)} \rightarrow \tau = T$, for $\epsilon \rightarrow 0$. Since the state process $X(t)$ is continuous and $W(x, t)$ is continuous on $[0, 1] \times [0, T]$, we conclude $W(X(\tau^{(\epsilon)}), \tau^{(\epsilon)}) \rightarrow W(X(\tau), \tau)$ a.s. By bounded convergence (W is bounded) and monotone convergence (ν is positive), we end up with

$$\mathbf{E}_{t,x} \left[\int_t^T F(X(s), u(s)) ds + W(X(T), T, \cdot) \right] \geq W(x, t).$$

If we take instead of an arbitrary u the optimizer u^* , we get instead of the inequality an equality above, which concludes the proof. \square

Remark 2. Let us finally remark that the existence of a strong solution of the underlying SDE (11) for our admissible strategies u_t follows by Exercise IX.2.10 of the monograph [32].

3. The stochastic open-loop optimal control problem

In this section we investigate the stochastic optimal control problem SOP without state observation, in which only (deterministic) open-loop controls, $u = u(t)$ are implementable. The only state-information available is the probability distribution of $X(0)$, represented by its density function $p_0(\cdot) : [0, 1] \rightarrow \mathbb{R}_0^+$, so that $\mathbb{P}(a \leq X(0) \leq b) = \int_a^b p_0(x) dx$ for any interval $0 \leq a \leq b \leq 1$. This initial distribution can be based on statistical estimation or on subjective probabilistic views.

The stochastic optimal open-loop control problem, SOP, reads as

$$\mathcal{U}(p_0(\cdot)) = \min_u \mathbf{E} \left[\int_0^T e^{-rs} [F(X(s), u(s))] ds + c_3 X(T) \right] \tag{40}$$

subject to

$$dX(s) = a(X(s), u(s)) ds + \sqrt{b(X(s))} dB(s),$$

$X(0)$ is a random variable with probability density $p_0(x)$.

We embed SOP into a family of optimization problems

$$U(t) = \min_u \mathbf{E} \left[\int_t^T e^{-rs} [F(X(s), u(s))] ds + c_3 X(T) \right] \tag{41}$$

subject to

$$dX(s) = a(X(s), u(s)) ds + \sqrt{b(X(s))} dB(s), \tag{42}$$

$X(t)$ is a random variable with (unconditional) probability density $p(x, t)$, $\tag{43}$

parametrized by time $t \in [0, T]$. $U(t)$ is the optimal rest value of SOP on $[t, T]$. With $p(x, 0) = p_0(x)$ we have $\mathcal{U}(p_0(\cdot)) = U(0)$. Moreover, if u^* is the optimal control function of SOP, we define the conditional value function as the expectation

$$U(x, t) = \mathbf{E}_{t,x} \left[\int_t^T e^{-rs} [F(X(s), u^*(s))] ds + c_3 X(T) \right] \tag{44}$$

We consider two approaches with different admissible control set: in Subsection 3.1 the HJB conditions are modified in order to deal with the no information case. Here the cost specification (5) is used. In Subsection 3.2 we formulate a control problem based on the related Fokker-Planck equation, which allows for much more general cost models (and also more general stochastic dynamics). Applied to the cost model (5), both approaches lead to comparable results.

3.1. Analyzing the open-loop problem with stochastic calculus

In this section we use stochastic calculus for deriving a sufficient condition for optimal solutions of the open-loop problem. As we will see, this condition is in some sense very similar to the HJB-condition but accounts for the fact that the states cannot be observed.

For technical reasons, in this subsection we restrict the set of admissible controls to functions $u : [0, T] \rightarrow [0, \infty)$ with bounded derivatives up to order 3.

Nonnegativity again is a natural assumption for the control in our context. Moreover it ensures that there is an entrance boundary at $x = 1$ (which can be ensured also by the weaker assumption $u(t) > -\gamma$). This fact is used in the proof of the following Theorem. Note however, that the arguments in the proof of Lemma 2 – showing the nonnegativity is automatically fulfilled in the optimum – can be applied also to the open loop problem.

Theorem 3. *For any admissible control $u(\cdot)$ (in the sense of this subsection) the PDE*

$$\begin{aligned} \mathcal{L}^{u(s)}Q(x, s) + e^{-rs}F(x, u(s)) &= 0, \\ Q(x, T) &= c_3x \text{ for } x \in (0, 1), \end{aligned} \tag{45}$$

with \mathcal{L}^u defined in (13), has a unique $C^2([0, 1] \times [0, T])$ -solution Q . Moreover, if an admissible control \hat{u} and the related solution \hat{Q} of (45) satisfy

$$\hat{u}(s) = \underset{u}{\operatorname{argmin}} \mathbf{E} \left[\mathcal{L}^u \hat{Q}(X(s), s) + e^{-rs}F(X(s), u) \right] \tag{46}$$

for all $s \in [t, T]$, then \hat{u} is an optimal control of SOP (41) and

$$\hat{Q}(x, s) = U(x, s). \tag{47}$$

This also leads to

$$\mathbf{E} \hat{Q}(X(s), s) = U(s). \tag{48}$$

Proof. Equation (45) is a linear second order equation with non-negative characteristic form ($b(x) \cdot \xi^2$ is obviously nonnegative for any $x \in [0, 1]$ and $\xi \in \mathbb{R}$). Applying the Fichera-classification (see e.g. [13]) to the piecewise smooth boundary of our equation, we get $\Sigma_0 = \{0\} \times (0, T)$, $\Sigma_1 = (0, 1) \times \{0\} \cup \{1\} \times (0, T)$, $\Sigma_2 = (0, 1) \times \{T\}$ and $\Sigma_3 = \emptyset$. This shows that (45) is a well formulated first boundary-value problem (i.e. the boundary condition is formulated on $\Sigma_2 \cup \Sigma_3$). Observe now that due to the assumed properties of u the coefficients of (45) as well as the function F have bounded derivatives up to order 3 on the whole domain $[0, 1] \times [0, T]$. Moreover the coefficients can be extended outside the domain with the same smoothness and keeping nonnegativity of the characteristic form (e.g., set $b(x) = \sigma^2 x^2(1 - 2x)$ for small negative x). The function $x \mapsto c_3x$ is linear. Furthermore the intersection of any two of the sets $\Sigma_3, \Sigma_2, \Sigma_0 \cup \Sigma_1$ is nonempty.

Using the result in [29] p. 775, we get that there exists a solution² with bounded derivatives up to order 3. Altogether, this implies that the solution is in $W^{3,p}((0, 1) \times (0, T))$ for any $p > 0$.

The rectangle $[0, 1] \times [0, T]$ has a Lipschitz boundary and we can apply Sobolev’s embedding results (see e.g. [18], Theorem 1.14) to infer that Q can be represented (almost everywhere) by a $C^2([0, 1] \times [0, T])$ -function. This gives enough regularity for applying Ito’s formula to the solution Q . Moreover, we have a natural boundary at $x = 0$ and an entry boundary at $x = 1$ (recall that $u(t) \geq 0$ by assumption) – both points are never reached if the process starts in the interior. Based on the dynamics (42) and using some starting point $X(t) = x_0$, we therefore get

$$Q(X(T), T) = Q(x_0, t) + \int_t^T \mathcal{L}^u Q(X(s), s) \, ds + \int_t^T Q_x(X(s), s) \sqrt{b(X(s))} \, d\mathcal{B}(s).$$

Since b and Q_x are bounded on $(0, 1)$, the last integral is a true martingale and vanishes in expectation. This gives

$$\mathbf{E}_{t,x} \left[\int_t^T e^{-rs} F(X(s), u(s)) \, ds + Q(X(T), T) \right] = Q(x_0, t) + \mathbf{E}_{t,x} \left[\int_t^T e^{-rs} F(X(s), u(s)) \right. \\ \left. + \mathcal{L}^u Q(X(s), s) \, ds \right]. \tag{49}$$

If (45) holds, the expectation on the right-hand side vanishes, i.e.

$$\mathbf{E}_{t,x} \left[\int_t^T e^{-rs} F(X(s), u(s)) \, ds + c_3 X(T) \right] = Q(x_0, t). \tag{50}$$

Moreover, the start value x_0 was chosen arbitrarily so far. Applying now the assumption that $X(t)$ is distributed with density function $p(x, t)$ and taking expectation at both sides of (49), we get

$$\mathbf{E} \left[\int_t^T e^{-rs} F(X(s), u(s)) \, ds + Q(X(T), T) \right] = \mathbf{E} [Q(X(t), t)] + \int_t^T \mathbf{E} \left[e^{-rs} F(X(s), u(s)) \right. \\ \left. + \mathcal{L}^{u(t)} Q(X(s), s) \right] \, ds. \tag{51}$$

If now $u^*(t)$ is the minimizer of (46), it minimizes the right-hand and hence also the left hand side of (51). Again the integral at the right hand side vanishes by (45) and in consequence

$$\mathbf{E}Q(X(t), t) = U(t). \tag{52}$$

Plugging u^* into (50), we also get

$$Q(x, t) = U(x, t). \quad \square \tag{53}$$

² Strictly speaking, the result is valid for equations that involve a further term $c(x)Q(x, t)$ in the left hand side, such that $c(x)$ is negative and bounded by a constant that is small enough. For parabolic PDEs however, this always can be achieved by a simple transformation of Q .

We mention that restricting the set of admissible control to 3 times differentiable functions, as in this subsection, may be too strong. Moreover, while it is clear that condition (46) is equivalent to the optimality condition

$$u(t) = \frac{e^{rt}}{2c_2} \frac{\mathbf{E}[X(t)Q_x(X(t), t)]}{\mathbf{E}X(t)}, \tag{54}$$

it is not clear at this point, how the conditions given in Theorem 1 can be used for calculations, in particular how the expectations in (54) could be calculated. For this reason, in the next subsection, we employ a more general approach, which uses arguments from PDE-constraint optimization.

3.2. The optimal control problem for the related Fokker-Planck equation

Denote $D := [0, 1] \times [0, T]$. Because $X(t)$ is a (Markovian) diffusion process, the objective function in (40) can be rewritten in terms of the transition density function $p : D \rightarrow [0, \infty)$. This leads to the equivalent formulation

$$\mathcal{U} = \min_u \iint_D e^{-rt} F(x, u(t)) p(x, t) \, dx \, dt + \int_0^1 c_3 x p(x, T) \, dx \tag{55}$$

subject to

$$p_t(x, t) = -[a(x, u(t))p(x, t)]_x + \frac{1}{2}[b(x)p(x, t)]_{xx}, \tag{56}$$

$$p(x, 0) = p_0(x), \quad p(1, t) = 0, \tag{57}$$

where the subscripts t and x mean partial differentiation. This is the Kolmogorov-forward (or Fokker-Planck) equation of the diffusion process (see e.g. [14] p. 132).

We remind that according to the specification of the functions a and b in (3), we have

$$a(0, u) = 0, \quad a(1, u) = -(\gamma + u) < 0, \quad b(0) = b'(0) = b(1) = b'(1) = 0. \tag{58}$$

Then the boundary condition (57) is equivalent to the zero-flux condition at $x = 1$:

$$0 = a(1, u(t))p(1, t) - \frac{1}{2} \frac{\partial}{\partial x} [b(x)p(x, t)]_{x=1} = a(1, u(t))p(1, t).$$

This condition ensures conservation of the probability mass at the “entrance boundary” $x = 1$. A similar boundary condition is not needed at the “natural boundary” (in the Feller terminology, see e.g. [15], p. 119) $x = 0$, since the flux there is automatically equal to zero due to (58).

Introducing the notations

$$A(x) = \frac{1}{2}b(x), \quad B(x, u) = a(x, u) - \frac{1}{2}b_x(x), \quad C(x, u) = a_x(x, u) - \frac{1}{2}b_{xx}(x), \quad f(x, t) = 0, \tag{59}$$

we can rewrite (56) (for a given measurable control $u(t) \geq 0$ inserted in B and C) in the symmetric form

$$p_t - (Ap_x)_x + Bp_x + Cp = f. \tag{60}$$

Thus problem (55) is a special case of the following problem:

$$\text{minimize } \left\{ J(u) := \iint_D h(x, t, u(t)) p(x, t) \, dx \, dt + \int_0^1 g(x) p(x, T) \, dx \right\}, \tag{61}$$

where in our particular case $h(x, t, u) = e^{-rt}F(x, u)$ and $g(x) = c_3x$.

In fact, in the analysis below we do not use the particular forms of A, B, C , and f . We consider the more general problem (61) subject to (60) and (57). Admissible controls are all measurable and bounded functions $u : [0, T] \rightarrow [0, \bar{u}]$, where $\bar{u} \geq 0$ is an upper bound for the control.

For this more general problem we assume that the functions A, B, C, h , and f (where A depends only on x , B and C depend on (x, u) , f depends on (x, t) , h depends on (x, t, u)) and the derivatives $A_x, A_{xx}, B_x, B_u, C_u$, and h_u are Lipschitz continuous; there are constants \hat{c}_1 and \hat{c}_2 such that the following inequalities are fulfilled for every $x \in [0, 1]$ and $u \in [0, \bar{u}]$:

$$\begin{aligned} A(0) = A(1) = A'(0) = A'(1) = 0, \quad B(1, u) < 0, \\ A(x) \geq \hat{c}_1x^2, \quad |B(x, u)| + |B_u(x, u)| + |B_{uu}(x, u)| \leq \hat{c}_2x. \end{aligned} \tag{62}$$

Clearly, the last conditions are fulfilled for our original SIS problem due to (59) and (3). Some of them may be weakened, but we do not aim for generality in this paper.

Below we briefly introduce some preliminary material, which is not quite standard and seems not to be available in the literature in the form that is needed in the optimal control context. The reason is that the function A in the elliptic term of (60) degenerates (even together with its first derivative) on the boundary $x \in \{0, 1\}$, which requires special care.

Denote

$$V := \{v : (0, 1] \rightarrow \mathbb{R} : v \text{ is locally absolutely continuous, } v \in L_2(0, 1), x v_x \in L_2(0, 1)\}.$$

Here v_x is the derivative of v , which exists almost everywhere. With the scalar product

$$\langle v, q \rangle := \int_0^1 [v(x)q(x) + x^2v_x(x)q_x(x)] \, dx + v(1)q(1), \quad v, q \in V,$$

V is a Hilbert space. Denote by V^* its dual space, and by $\langle v^*, v \rangle_{V^*, V}$ – the pairing of V^* and V .

For a Banach space Y we denote by $L_2(0, T; Y)$ the space of all functions $y : (0, T) \rightarrow Y$ such that

$$\|y\|_{L_2(0, T; Y)} := \left(\int_0^T \|y(t)\|^2 \, dt \right)^{1/2} < \infty.$$

It is said that $y \in L_2(0, T; Y)$ has a weak derivative z , written later as $\dot{y} = z$, if

$$\int_0^T \dot{\varphi}(t)y(t) \, dt = - \int_0^T \varphi(t)z(t) \, dt \quad \forall \varphi \in C_c^\infty(0, T),$$

where the integrals are in the sense of Bochner. In particular, we will use the spaces $L_2(0, T; V)$, $L_2(0, T; V^*)$ and

$$W := \{p \in L_2(0, T; V) : \dot{p} \in L_2(0, T; V^*)\}.$$

It is known (see e.g. [35, Theorem 3.10]) that every element of W actually belongs to $C([0, T]; V)$ (defined similarly as $L_2(0, T; V)$ but using the uniform norm), and for every two elements $p, \lambda \in W$ the formula for integration by parts is valid:

$$\int_0^T \langle \dot{p}(t), \lambda(t) \rangle_{V^*, V} dt = \langle p(T), \lambda(T) \rangle_{L_2(0,1)} - \langle p(0), \lambda(0) \rangle_{L_2(0,1)} - \int_0^T \langle \dot{\lambda}(t), p(t) \rangle_{V^*, V} dt. \tag{63}$$

In the space $V \times V$ we define the following bi-linear functional depending on $u \geq 0$:

$$\mathbf{B}(q, v; u) := \int_0^1 [A(x)q_x v_x + B(x, u)q_x v + C(x, u) q v] dx - B(1, u) q(1)v(1).$$

From the definition of V and the properties of A and B it is easy to verify that \mathbf{B} is well defined. Indeed, using (62) we have

$$\begin{aligned} \|Aq_x v_x\|_{L_1} &\leq \frac{1}{2} \|A_{xx}\|_{L_\infty} \|xq_x\|_{L_2} \|xv_x\|_{L_2} \leq \frac{1}{2} \|A_{xx}\|_{L_\infty} \|q\|_V \|v\|_V, \\ \|B(x, u)q_x v\|_{L_1} &\leq \|B/x\|_{L_\infty} \|xq_x\|_{L_2} \|v\|_{L_2} \leq \hat{c}_2 \|q\|_V \|v\|_V. \end{aligned}$$

Definition. A function $p : D \rightarrow \mathbb{R}$ is a *weak solution* of equation (60) with side conditions (57) (for a fixed admissible control $u(\cdot)$) if $p \in W$,

$$\langle \dot{p}(t), v \rangle_{V^*, V} + \mathbf{B}(p(t), v; u(t)) = \langle f(t), v \rangle_{L_2(0,1)} \quad \text{for every } v \in V \text{ and a.e. } t \in (0, T),$$

and $p(\cdot, 0) = p_0(\cdot)$.

We mention that, thanks to the inequality $B(1, u) \neq 0$, the last term of \mathbf{B} ensures satisfaction of the boundary condition $p(1, t) = 0$.

Now, let us fix a reference admissible control u , and introduce the so-called adjoint equation:

$$\dot{\lambda} + (A\lambda_x)_x + (B\lambda)_x - C\lambda = -h, \tag{64}$$

with the end-condition

$$\lambda(x, T) = g(x). \tag{65}$$

Notice that by changing the time variable $t \rightarrow T - t$ we obtain the following equation for $\mu(x, t) = \lambda(x, T - t)$:

$$\dot{\mu} - (A(x)\mu_x)_x - (B(x, u(T - t))\mu)_x + C(x, u(T - t))\mu = h(x, T - t, u(T - t)).$$

This equation has a similar form as (60) with the difference that the transport term is now $-B\mu$, and since $-B(1, u(t)) > 0$ and $A(1) = 0$, there is no need of a boundary condition at $x = 1$ (“natural boundary” in the Feller terminology, see e.g. [15], p. 119).

Correspondingly, the definition of a solution is a slight modification of the previous one. Namely, a solution of (64), (65) is any element $\lambda \in W$ such that

$$\langle \dot{\lambda}(t), v \rangle_{V^*, V} + \mathbf{D}(\lambda(t), v; u(t)) = -\langle h(\cdot, t, u(t)), v \rangle_{L_2(0,1)} \quad \forall v \in V \text{ and a.e. } t \in (0, T),$$

and $\lambda(T) = g$, where

$$\mathbf{D}(q, v; u) := \int_0^1 [-A(x)q_x v_x + (B(x, u)q)_x v - C(x, u)qv] dx, \quad q, v \in V,$$

Proposition 2. *Given any measurable selection $u(\cdot)$ of the interval $[0, \bar{u}]$, both systems (60), (57) and (64), (65) have unique solutions, p and λ , respectively, in the space W . Moreover, there exists a constant c (depending on the functions A , B , and C , and the numbers \bar{u} and T , but not on f , p_0 , and the particular choice of the measurable selection u of $[0, \bar{u}]$), such that*

$$\max_{t \in [0, 1]} \|p(\cdot, t)\|_{L_2(0, 1)} + \|p\|_{L_2(0, T; V)} \leq c (\|p_0\|_{L_2(0, 1)} + \|f\|_{L_2(D)}).$$

The proof of this proposition adopts known approaches combining [30][Chapter 1] and [35, Chapter 3]. It is long but does not involve new ideas, therefore we do not present it here. Issues of regularity of the solution of problem (60), (57) will not be discussed (and used) in order to keep the presentation as focused as possible.

We introduce the notation

$$H(x, t, p, q, u, \mu) := h(x, t, u)p - (B(x, u)q + C(x, u)p) \mu.$$

The following is the main result of this section.

Theorem 4. *Let \hat{u} be an admissible control and let \hat{p} and λ be the corresponding solutions of the primal equation (60), (57) and the adjoint equation (64), (65), respectively.*

(i) *Then for every number $M \geq 0$ there exists a constant $c(M)$ such that the inequality*

$$\left| J(u) - J(\hat{u}) - \int_0^T J'(\hat{u})(t) (u(t) - \hat{u}(t)) dt \right| \leq c(M) \|u - \hat{u}\|_{L_2(0, T)}^2$$

is fulfilled for every measurable selection $u(\cdot)$ of $[0, \bar{u} + M]$. Here $J'(\hat{u}) \in L_\infty(0, T)$ is defined as

$$J'(\hat{u})(t) = \int_0^1 H_u(x, t, \hat{p}(x, t), \hat{p}_x(x, t), \hat{u}(t), \lambda(x, t)) dx.$$

(ii) *If \hat{u} is an optimal control, then the following maximum condition holds:*

$$\int_0^1 H(x, t, \hat{p}(x, t), \hat{p}_x(x, t), \hat{u}(t), \lambda(x, t)) dx = \min_{u \in [0, \bar{u}]} \int_0^1 H(x, t, \hat{p}(x, t), \hat{p}_x(x, t), u, \lambda(x, t)) dx.$$

We mention that property (i) directly implies Fréchet differentiability of the mapping $L_\infty(0, T) \ni u \mapsto J(u)$ with the L_∞ -representative $J'(u)(\cdot)$ of the derivative. However, property (i) is stronger (as resembling some properties of an L_2 -derivative), and this is essential for application of the gradient projection method for solving the optimal control problem at hand.

Proof. Let M be arbitrarily fixed and u be an arbitrary measurable selection of $[0, \bar{u} + M]$. Denote $\varepsilon := \|u - \hat{u}\|_{L_2(0, T)}$. Let p be the corresponding solution of equations (60), (57). Further, we abbreviate

$$\begin{aligned} \hat{B}(x, t) &:= B(x, \hat{u}(t)), & \hat{C}(x, t) &:= C(x, \hat{u}(t)), & \hat{h}(x, t) &:= h(x, t, \hat{u}(t)), \\ \Delta p &= p - \hat{p}, & \Delta B(x, t) &:= B(x, u(t)) - B(x, \hat{u}(t)), \\ \Delta C(x, t) &:= C(x, u(t)) - C(x, \hat{u}(t)), & \Delta h(x, t) &:= h(x, t, u(t)) - h(x, t, \hat{u}(t)). \end{aligned}$$

Then the following equations are satisfied:

$$\begin{aligned} \Delta p_t - (A \Delta p_x)_x + B \Delta p_x + C \Delta p &= -\Delta B(x, t) \hat{p}_x(x, t) - \Delta C(x, t) \hat{p}(x, t), \\ \Delta p(x, 0) = 0, \quad \Delta p(1, x) &= 0. \end{aligned} \tag{66}$$

We shall apply to the above equation Proposition 2 with $p_0 = 0$ and $f = -\Delta B(x, t) \hat{p}_x(x, t) - \Delta C(x, t) \hat{p}(x, t)$. For this we need the following estimations that use the assumptions about A and B , the definition of the space V and the fact that $p \in C([0, T]; V)$. First,

$$\begin{aligned} \iint_D |\Delta C(x, t) \hat{p}(x, t)|^2 dx dt &\leq \int_0^T \hat{c}_4 |u(t) - \hat{u}(t)|^2 \int_0^1 |\hat{p}(x, t)|^2 dx dt \\ &= \hat{c}_4 \int_0^T |u(t) - \hat{u}(t)|^2 \|\hat{p}(\cdot, t)\|_{L_2(0,1)}^2 dt \leq \hat{c}_4 \|\hat{p}\|_{C([0,T];V)}^2 \varepsilon^2, \end{aligned}$$

where \hat{c}_4 , and also \hat{c}_5 , etc., below, are constants (independent of ε). Second, for some measurable function $B^u(x, t) \in B_u(x, [u(t), \hat{u}(t)])$ we have

$$\begin{aligned} \iint_D |\Delta B(x, t) \hat{p}_x(x, t)|^2 dx dt &= \int_0^T \int_0^1 |B^u(x, t)(u(t) - \hat{u}(t))|^2 |\hat{p}_x(x, t)|^2 dx dt \\ &\leq \int_0^T \int_0^1 |B^u(x, t)/x|^2 |u(t) - \hat{u}(t)|^2 \|\hat{p}(\cdot, t)\|_V^2 dx dt \leq \hat{c}_3^2 \|\hat{p}\|_{C([0,T];V)}^2 \varepsilon^2. \end{aligned}$$

The obtained estimation of $\|f\|_{L_2(D)}$, together with Proposition 2, implies existence of a constant \hat{c}_5 which is independent of ε and such that

$$\|\Delta p\|_{L_2(0,T;V)} \leq \hat{c}_5 \varepsilon. \tag{67}$$

Now, we rewrite equation (66) in the more detailed form

$$\Delta p_t - (A \Delta p_x)_x + \hat{B} \Delta p_x + \hat{C} \Delta p = -\Delta B \hat{p}_x - \Delta C \hat{p} + \gamma,$$

where $\gamma = -\Delta B \Delta p_x - \Delta C \Delta p$. We notice that

$$\hat{c}_5 \varepsilon \geq \|\Delta p\|_{L_2(0,T;V)} \geq \|\Delta p\|_{L_2(D)} + \|x \Delta p_x\|_{L_2(D)}.$$

Thus

$$\|\gamma\|_{L_2(D)} \leq \|\Delta B \Delta p_x\|_{L_2(D)} + \|\Delta C \Delta p\|_{L_2(D)} \leq \|(\Delta B^u/x)(u - \hat{u})(x \Delta p_x)\|_{L_2(D)} + \hat{c}_6 \varepsilon^2 \leq \hat{c}_7 \varepsilon^2.$$

Let λ be the solution of the adjoint equation (64), (65) corresponding to \hat{u} and \hat{p} . Since $\lambda(t) \in V$ for a.e. t , we have

$$\langle \hat{\Delta} p(t), \lambda(t) \rangle_{V, V^*} + \mathbf{B}(\Delta p(t), \lambda(t); \hat{u}(t)) = -\langle \Delta B(t) \hat{p}_x(t) + \Delta C(t) \hat{p}(t), \lambda(t) \rangle_{L_2(0,1)} + \langle \gamma(t), \lambda(t) \rangle_{L_2(0,1)}.$$

Using the integration by part formula (63) and the relations $\Delta p(x, 0) = 0$, $\lambda(x, T) = g(x)$ we obtain that

$$\begin{aligned} & - \int_0^T \langle \dot{\lambda}(t), \Delta p(t) \rangle_{V^*, V} dt + \langle g, \Delta p(T) \rangle_{L_2(0,1)} + \int_0^T \mathbf{B}(\Delta p(t), \lambda(t); \hat{u}(t)) dt \\ & = - \int_0^T \langle (\Delta B(t)\hat{p}_x(t) + \Delta C(t)\hat{p}(t)), \lambda(t) \rangle_{L_2(0,1)} dt + \zeta_1, \end{aligned} \quad (68)$$

where $|\zeta_1| \leq \hat{c}_8 \varepsilon^2$ and \hat{c}_8 is independent of ε and the particular choice of the selection u of $[0, \bar{u} + M]$.

The term $\int_0^T \mathbf{B}(\Delta p(t), \lambda(t); \hat{u}(t)) dt$ contains the integral $\iint_D \hat{B} \lambda \Delta p_x dx dt$, in which we want to shift the differentiation from Δp to λ . Applying integration by parts needs a special care, due to the (a priori) possible unboundedness of Δp and λ at $x = 0$. For this reason we first show that for every $t \in [0, T]$ there exists a sequence $x_k \in (0, 1]$, $x_k \rightarrow 0$, such that $x_k \Delta p(x_k, t) \lambda(x_k, t)$ converges to zero. Indeed, if for some t such a sequence does not exist then there exist $\alpha > 0$ and $\beta > 0$ such that $x |\Delta p(x, t) \lambda(x, t)| \geq \alpha$ for any $x \in (0, \beta]$. Since $\Delta p(\cdot, t), \lambda(\cdot, t) \in L_2(0, 1)$, the function $|\Delta p(\cdot, t) \lambda(\cdot, t)|$ is integrable, which contradicts the inequality $|\Delta p(\cdot, t) \lambda(\cdot, t)| \geq \alpha/x$ on $(0, \beta]$.

Then, due to the absolute integrability of $\hat{B} \lambda \Delta p_x$, $\hat{B} \lambda_x \Delta p$ (already shown) and $\hat{B}_x \lambda \Delta p$ with respect to x , we have (skipping the fixed arguments t and $\hat{u}(t)$ in the notations)

$$\begin{aligned} & \int_0^1 \hat{B}(x) \lambda(x) \Delta p_x(x) dx = \lim_k \int_{x_k}^1 \hat{B}(x) \lambda(x) \Delta p_x(x) dx \\ & = \lim_k \left[\hat{B}(1) \lambda(1) \Delta p(1) - \hat{B}(x_k) \lambda(x_k) \Delta p(x_k) - \int_{x_k}^1 [B_x(x) \lambda(x) \Delta p(x) + [B(x) \lambda_x(x) \Delta p(x)] dx \right] \\ & = \hat{B}(1) \lambda(1) \Delta p(1) - \int_0^1 [B_x(x) \lambda(x) \Delta p(x) + [B(x) \lambda_x(x) \Delta p(x)] dx, \end{aligned}$$

where we also use the inequality $|B(x, u)| \leq \hat{c}_2 x$ in (62). Substituting the obtained expression in (68) we obtain the equality

$$\begin{aligned} & - \int_0^T \langle \dot{\lambda}(t), \Delta p(t) \rangle_{V^*, V} dt + \int_0^1 \lambda^T(x) \Delta p(x, T) dx + \int_0^T \mathbf{D}(\lambda(t), \Delta p(t); t, \hat{u}(t)) dt \\ & = - \iint_D (\Delta B(x, t)\hat{p}_x(x, t) + \Delta C(x, t)\hat{p}(x, t)) \lambda(x, t) dx dt + \zeta_1. \end{aligned}$$

Having in mind the definition of solution of the adjoint equation and the fact that $\Delta p(t) \in V$, we obtain that

$$\begin{aligned} \int_0^1 g(x) \Delta p(x, T) dx & = - \iint_D \hat{h}(x, t) \Delta p(x, t) dx dt \\ & \quad - \iint_D (\Delta B(x, t)\hat{p}_x(x, t) + \Delta C(x, t)\hat{p}(x, t)) \lambda(x, t) dx dt + \zeta_1. \end{aligned} \quad (69)$$

Now consider the difference

$$J(u) - J(\hat{u}) = \iint_D [h(x, t, u(t))p(x, t) - h(x, t, \hat{u}(t))\hat{p}(x, t)] dx dt + \int_0^1 g(x) \Delta p(x, T) dx,$$

which, using the properties of h and (67), can be represented (skipping the arguments (x, t)) as

$$\begin{aligned} J(u) - J(\hat{u}) &= \iint_D [\hat{h} \Delta p + \Delta h \hat{p} + \Delta h \Delta p] dx dt + \int_0^1 g(x) \Delta p(x, T) dx \\ &= \iint_D [\hat{h} \Delta p + \Delta h \hat{p}] dx dt + \int_0^1 g(x) \Delta p(x, T) dx + \zeta_2, \end{aligned}$$

where $|\zeta_2| \leq \hat{c}_9 \|u - \hat{u}\|_{L_2(0,T)}^2 = \hat{c}_9 \varepsilon^2$. Using (69), we obtain that

$$\begin{aligned} J(u) - J(\hat{u}) &= \iint_D [\hat{h} \Delta p + \Delta h \hat{p}] dx dt - \iint_D \hat{h} \Delta p dx dt \\ &\quad - \iint_D (\Delta B(x, t)\hat{p}_x(x, t) + \Delta C(x, t)\hat{p}(x, t)) \lambda(x, t) dx dt + \zeta_1 + \zeta_2 \\ &= \iint_D [H(x, t, \hat{p}(x, t), \hat{p}_x(x, t), u(t), \lambda(x, t)) \\ &\quad - H(x, t, \hat{p}(x, t), \hat{p}_x(x, t), \hat{u}(t), \lambda(x, t))] dx dt + \zeta_1 + \zeta_2. \end{aligned} \tag{70}$$

Using the abbreviation $\hat{H}(x, t, u) := H(x, t, \hat{p}(x, t), \hat{p}_x(x, t), u, \lambda(x, t))$ we represent in a standard way

$$\begin{aligned} &\hat{H}(x, t, u(t)) - \hat{H}(x, a, \hat{u}(t)) \\ &= \hat{H}_u(x, t, \hat{u}(t))(u(t) - \hat{u}(t)) + \int_0^1 [\hat{H}_u(x, t, \hat{u}(t) + s(u(t) - \hat{u}(t))) - \hat{H}_u(x, t, \hat{u}(t))] ds (u(t) - \hat{u}(t)). \end{aligned}$$

Combining this with (70) and using the particular form of the function H and again the properties (62) in a similar way as above, we obtain the representation in part (i) of the theorem and the boundedness of the function $J'(\hat{u})(t)$.

Now let \hat{u} be an optimal control. Then applying (70) to controls of the form

$$u(t) = \begin{cases} v & \text{if } t \in [\tau, \tau + \varepsilon'], \\ \hat{u}(t) & \text{if } t \notin [\tau, \tau + \varepsilon'], \end{cases}$$

where $v \in [0, \bar{u}]$, $\tau \in (0, T)$, and $\varepsilon' > 0$ is so small that $\tau + \varepsilon' \leq T$ and $4\bar{u}^2\varepsilon' \leq \varepsilon$ (so that $\int_\tau^{\tau+\varepsilon'} |v - \hat{u}(t)| dt \leq \varepsilon$), one can prove the second claim of the theorem in a standard way. \square

The second part of Theorem 4 gives a Pontryagin type necessary optimality condition for the problem (61), (60), (57). The first part provides a formula for calculation of the derivative of the objective functional with respect to the control function in the space $L_2(0, T)$, in directions that are bounded. This derivative can be used for computing an approximate solution by using the gradient projection method, as we do it

in our numerical experiments in Section 4 (although the justification needs a lot more work that we do not present in this paper).

We mention that with the particular specifications for our original SIS model the adjoint equation (64) (by substituting the definitions (59)) takes the form

$$-\lambda_t = a(x, u)\lambda_x + \frac{1}{2}b(x)\lambda_{xx} + e^{-rt}F(x, t, u). \tag{71}$$

If we apply Theorem 4 to the original problem (55)-(57), respectively (40), we can see a very close relation to Theorem 3.

Corollary 2. *Let the assumptions of Theorem 4 be fulfilled with optimal $\hat{p}, \hat{u}, \hat{\lambda}$. Moreover let $\hat{X}(s)$ denote the process defined by*

$$d\hat{X}(t) = a(\hat{X}(t), \hat{u}(t)) dt + \sqrt{b(\hat{X}(t))} d\mathcal{B}(t). \tag{72}$$

Then the equations

$$\begin{aligned} \mathcal{L}^{\hat{u}(s)}\hat{\lambda}(x, s) + e^{-rs}F(x, \hat{u}(s)) &= 0, \\ \hat{\lambda}(x, T) &= c_3x \text{ for } x \in (0, 1), \end{aligned} \tag{73}$$

are fulfilled and

$$\hat{u}(s) = \underset{u}{\operatorname{argmin}} \mathbf{E} \left[\mathcal{L}^u \hat{\lambda}(\hat{X}(s), s) + e^{-rs}F(\hat{X}(s), u) \right] \tag{74}$$

holds for all $s \in [t, T]$. (We remind that the notation \mathcal{L}^u is introduced in (13).)

Proof. Clearly, the adjoint equation (71) for $\hat{u}, \hat{\lambda}$ can be reformulated as (72). Observe now that (using the definitions (59) and applying integration by parts) we have (without denoting the dependencies on (x, s))

$$\begin{aligned} &\int_0^1 H(x, s, \hat{p}(x, s), \hat{p}_x(x, s), \hat{u}(s), \hat{\lambda}(x, s)) dt \\ &= \int_0^1 \left[h(x, s, \hat{u}(s)) + \left(a(x, \hat{u}(s)) - \frac{1}{2}b_x(x) \right) \hat{\lambda}_x(x, s) \right] \hat{p}(x, s) ds \\ &= \mathbf{E} \left[h(\hat{X}(s), s, \hat{u}(s)) + \left(a(\hat{X}(s), s, \hat{u}(s)) - \frac{1}{2}b_x(\hat{X}(s)) \right) \hat{\lambda}_x(\hat{X}(s), s) \right]. \end{aligned} \tag{75}$$

Because $b(\cdot)$ and $\hat{\lambda}_t$ do not depend on u , minimizing (75) is equivalent to minimizing (74). \square

This shows that the results in Theorems 4 and 3 are fully consistent with each other: in fact, (73) and (74) are the main conditions in Theorem 3.

4. The role of information in the stochastic SIS optimal control problem: a numerical study

Based on an illustrative example, in this section we investigate what improvement in the decision performance is brought by the availability of exact measurements of the current state. This numerical analysis uses the results in sections 2 and 3, where state-feedback (for the SFP problem) or open-loop controls (SOP)

are involved, respectively. In addition, we compare these performance assessments with the ones obtained by using the deterministic model (DP), assuming that the stochastic model represents the reality.

In order to ensure comparability we take a viewpoint immediately before $X(0) = x_0$ is realized and assume that $X(0)$ is distributed according to the density $p_0(x)$ as assumed in Section 3. The expectation

$$\mathcal{W}(p_0(\cdot)) = \mathbf{E}[V(X(0), 0)] = \int_0^1 V(x)p_0(x) dx,$$

where V was defined for SFP in (10), then gives the minimal expected cost for a decision maker, who is able to observe all values $X(t)$ and to calculate state-dependent strategies. This expectation can be compared with the optimal value $\mathcal{U}(p_0(\cdot))$, i.e. the minimal expected cost for a decision maker who is not able to observe any values of the process $X(t)$, see (40).

The *value of observation*

$$VO = \mathcal{U}(p_0(\cdot)) - \mathcal{W}(p_0(\cdot))$$

expresses the value of the additional information, achieved by observing the state values over time. Clearly, VO is positive, because restricting $u(x, t)$ to $u(t)$ puts an additional constraint on the original optimization problem. In our case the difference can be interpreted as the (maximal) amount of money, a decision maker without the possibility to observe the process $X(t)$ is willing to pay in order to acquire full information about $X(t)$ at any time t . In the context of disease modelling, VO helps to assess the meaningfulness of improved reporting schemes: a small value speaks against improved efforts, whereas large values suggest that investment in information will pay off later. The idea of VO is closely related to the notion of *value of information*, where $\mathcal{W}(p_0(\cdot))$ is calculated as the expectation of a clairvoyant’s optimal value. The value of information, introduced in [31], has a long history in stochastic optimization, see e.g. also [34].

When considering optimal control in a stochastic context, one may ask the question, whether the substantial extra effort for using SDE and/or PDE constraints is worthwhile. In our case, the deterministic SIS model (1) could be easily reduced to the deterministic control problem (1), where the starting value is chosen as $x_0 = \mathbf{E}[X(0)] = \int_0^1 xp_0(x) dx$. In this way one replaces distributions and stochastic processes by (instantaneous) expectations and gets a simpler, ODE constraint problem, which can be treated by classical methods. The question then is, how much (in terms of objective value) we lose if the system is in fact stochastic (according to either (10) or (55)), but decisions are based on the deterministic model (1). We therefore calculate the optimal control $u_d(t)$ for (1) and plug it into the Fokker-Planck equation of the stochastic system, which leads to a transition density $p_d(x, t)$ for a decision maker who faces the stochastic system but nevertheless uses the deterministic model:

$$\begin{aligned} \frac{\partial p_d(x, t)}{\partial t} &= -\frac{\partial}{\partial x} a(x, u_d(t))p_d(x, t) + \frac{1}{2} \frac{\partial^2}{\partial x^2} [b(x)p_d(x, t)], \\ p_d(1, t) &= 0, \\ p_d(x, 0) &= p_0(x). \end{aligned}$$

The cost for this strategy is given by

$$\mathcal{R}(p_0(\cdot)) = \int_t^T \int_0^1 e^{-rs} F(X(s), u(s))p_d(x, s) dx ds + \int_0^1 c_3 x p_d(x, T) dx.$$

It is then possible to compare \mathcal{R}, \mathcal{U} and \mathcal{W} . Clearly we have $\mathcal{R}(p_0(\cdot)) \geq \mathcal{U}(p_0(\cdot)) \geq \mathcal{W}(p_0(\cdot))$. In particular the difference

$$VSOL = \mathcal{R}(p_0(\cdot)) - \mathcal{U}(p_0(\cdot))$$

expresses the value of the stochastic open loop solution (55), whereas

$$VSCS = \mathcal{R}(p_0(\cdot)) - \mathcal{W}(p_0(\cdot))$$

is the value of the stochastic closed loop solution (10).

In the following example the value function $V(x, t) = W(x, t)$ of the stochastic control problem (10) was calculated by solving the HJB equation (14) numerically. The related optimal control $u(x, t)$ is then calculated from (21).

For finding the optimal solution of the Fokker-Planck system (55) we used the Fréchet derivative J' given in Theorem 4 and applied the gradient projection method described e.g. in [35], Section 5.9.1 (see also [21] for convergence results). All partial differential equations (the HJB equation as well as all PDEs used within the gradient projection descent algorithm) were solved by the numerical method of lines (see e.g. [33]). Finally, we solved the deterministic problem (1) by the steepest descent method for ODE constraint control problems, see e.g. [22] p. 334. All the algorithms were implemented in Wolfram Mathematica™.

As a numerical example we compare now the effects of available information and of a low versus a high volatility parameter value. This example is purely illustrative with no specific disease in view. We choose $\gamma = 1$ and $\beta = 4.5$. This setup may be interpreted as a disease where the mean length of infection is one time unit, e.g. a week, whereas each infected person infects 4.5 persons in the mean over this time unit. In the deterministic model without control (1) such a setup describes a disease which does not die out (the basic reproduction number is $R_0 = 4.5$) and has a stationary point (endemic equilibrium) at

$$\bar{x} = 1 - \frac{\gamma}{\beta} = 0.7.$$

For the volatility parameter σ we consider two cases: low volatility with $\sigma = 0.5$ and high volatility with $\sigma = 2.7$. In all cases we plan over a horizon of $T = 8$ time units and the cost parameters are chosen as $c_1 = 15.0$, $c_2 = 80.0$, $c_3 = 0.0$. The discount rate is set to $r = 0.00057$, which corresponds to an interest rate of 3% p.a. if time units are weeks. Finally, we assume that distribution of the initial value $X(0)$ was estimated as a uniform distribution with density.

$$p_0(x) = \begin{cases} 20 & \text{if } x \in [0.03, 0.08], \\ 0 & \text{else.} \end{cases}$$

This expresses the view that the initial fraction of infected individuals lies between 0.03 and 0.08, but each fraction in this range is equally likely.

Fig. 1 shows the transition density function $p(x, t)$ for the optimal solution of the Fokker-Planck constraint problem (55) when $\sigma = 0.5$ (low volatility case). The optimal value is (within the accuracy of three digits) equal to the optimal value of the deterministic solution, i.e.

$$\mathcal{U}(p_0(\cdot)) \approx \mathcal{R}(p_0(\cdot)) \approx 73.816.$$

This shows that the value of the open loop solution $VSOL$ is vanishingly small in this case, which can be interpreted as a sort of justification for using deterministic control in case of low volatility. Visually, the transition density in Fig. 1 shows time dependent densities (cuts parallel to the x-axis) with small variation.

We consider now the case with larger volatility parameter $\sigma = 2.7$. Fig. 2 shows a comparison between the deterministic optimal control and the stochastic optimal open loop control. While both strategies use more control at the beginning (which is due to the fact that we chose $c_3 = 0$ in this example), the stochastic control uses higher values over the whole planning period.




Fig. 1. Low volatility case $\sigma = 0.5$: Transition density $p(x, t)$ for the optimal solution of the Fokker-Planck constraint problem (55).




Fig. 2. High volatility case $\sigma = 2.7$: Comparison between the optimal control for the deterministic problem (dashed line) and the optimal open loop control for the stochastic model (solid line).

The related transition density when applying the optimal open loop control is shown in Fig. 3. Here the control is able to bring down the number of infected near to zero and hold them there with high probability. Still there is a substantial probability that the disease grasps a high fraction of the population.

The stochastic open loop solution leads to a substantial reduction in expected costs: we have $\mathcal{U}(p_0(\cdot)) \approx 36.167$ for the optimal value of SOP and $\mathcal{R}(p_0(\cdot)) \approx 38.147$ for the expected cost of the deterministic solution, hence the value of the stochastic open loop problem is given by $VSOL \approx 1.98$. Note that already using the deterministic solution leads to a slight reduction of costs compared to the case without control ($u(x, t) = 0$), where the expected cost is equal to 39.669.

Finally we consider the full information case SFP, i.e. the solution of problem (10). The optimal transition density and the optimal control function are shown in Fig. 4.

The transition density shows that the probabilities of staying near zero can be substantially improved, when being able to observe the states. The likelihoods of high fractions of infected then can be reduced a lot, also compared with the optimal open loop solution. The optimal control function again shows some decrease over time. More important are the differences in treatment intensities when compared over possible states. It shows that when it is possible to observe the states, one would put high effort per case if the fraction of infected is small, whereas the effort per case has to be reduced a lot if the number of cases becomes




Fig. 3. High volatility case $\sigma = 2.7$: Transition density $p(x, t)$ for the optimal solution of the Fokker-Planck constraint problem (55). The third axis (density $p(x, t)$) is cut at $p = 2.5$, so the small values contain substantially more probability mass than shown in the picture.




Fig. 4. High volatility case: (a) transition density and (b) optimal feedback control $u(x, t)$ for the stochastic control problem (10). The third axis (density $p(x, t)$) is cut at $p = 2.5$, so the small values contain substantially more probability mass than shown in the picture. Note also that the x -axis is cut at $x = 0.5$ as the surface becomes very flat for larger values of x .

larger. Being able to observe the values of the process $X(t)$ is very valuable: the optimal value of the control problem is $\mathcal{W}(p_0(\cdot)) = 24.507$ and the value of observation therefore is $VO \approx 11.66$.

For both stochastic optimization model SOP and SFP and also for the deterministic approach, with $\sigma = 2.7$ the respective transition densities shows a similar twin-peaked visual pattern. There is a region near zero and a region with higher fraction of infected where the system tends to be concentrated. Because the pictures would look quite similar to Fig. 3 at first glance, we do not show the transition densities for all the models. Instead we summarize the differences between the cases in Table 1, which shows the values of the related cumulative distribution functions of $X(T)$ at the end time $T = 8$ for selected fractions of infected. Not only SFP dominates SOP, SOP dominates the deterministic and the deterministic solution dominates the model without control in terms of the expected costs. In addition the same order of dominance can be observed at all considered fractions of infected: the probability that the fraction of infected is smaller or equal to the value given in the first column is larger for SFP than for SOP, larger for SOP than for the deterministic solution and larger for the deterministic solution than without any control.

5. A remark about vaccination

As one of the anonymous referees suggested, one can incorporate vaccination (prior to the beginning of the disease) into basically the same stochastic model as considered in this paper. We shortly describe the idea.

Table 1

Expected costs and cumulative distribution function for the analyzed approaches (no control with $u(x, t) = 0$, deterministic DP, no observation SOP and full observation SFP).

Infected x	No control	Deterministic	No observation (SOP)	Full observation (SFP)
	39.669	38.147	36.167	24.507
0.001	0.24535	0.25891	0.31358	0.85869
0.010	0.31168	0.32767	0.38703	0.86568
0.100	0.38831	0.40631	0.46629	0.87559
0.200	0.41923	0.43767	0.49630	0.88084
0.500	0.49929	0.51762	0.57010	0.89649

The second line shows the expected costs of the respective (optimal) strategies. The rest of the table shows the values of the cumulative distribution function $P_m(X(T) \leq x)$ at time $T = 8$ for the respective models at the fraction x of infected individuals specified in the first column.

Let $v \in [0, 1)$ be the fraction of the population vaccinated before the outburst of the epidemic disease, and let $\kappa \in (0, 1]$ be the fraction of the vaccinated individuals who become immune. Then we can modify the equation (2) as

$$dX(t) = [\beta(1 - \kappa v - X(t))X(t) - \gamma X(t)] dt + \sigma[(1 - \kappa v - X(t))X(t)] dB(t),$$

$$X(0) = x_0 \in (0, 1).$$

If $c(v)$ is the cost of vaccinating fraction v of the population, we obtain the following extension of problem (6):

$$\min_{v, u(\cdot)} \mathbf{E} \int_0^T e^{-rs} [F(X(s), u(X(s), s))] ds + c_3 \mathbf{E}[X(T)] + c(v).$$

Here the decision parameter $v \in [0, 1)$ is a scalar, while $u(\cdot)$ is either a feedback or open-loop control, depending on the information pattern as above in the paper. Observe that for a fixed v the problem is completely analogous to (2), (6), with the only difference that now the range of X is $[0, 1 - \kappa v]$ instead of $[0, 1]$. Thus the problem for fixed v can be solved in the same way as in the case without vaccination. The scalar minimization is then a matter of repeated calculations.

We also mention that including vaccination during the disease may lead to complications in the stochastic case, since the control variable representing vaccination may appear (depending on the particular model) in the diffusion term of the relevant equations.

6. Conclusions

In this paper we analyze the stochastic SIS model both in the full information case (SFP) and in the no information case (SOP). In particular, we give (under some technical assumptions)

- a sufficient optimality condition (the HJB equation) together with a verification theorem for the optimal solution of the SFP.
- a sufficient optimality condition with verification theorem for the SOP under quite strong assumptions about the control function, using stochastic calculus as the basis.
- a necessary optimality condition for the SOP under weaker assumptions, applying the idea of optimization under PDE-constraints.

It turns out that the two approaches for the SOP lead to formally identical optimality conditions, but the sufficiency requires more regularity of the optimal control. In a numerical part we compare the solutions (the control functions) of SFP and SOP with each other and with the solution of the deterministic problem.

Plugging these solutions into the same stochastic dynamics leads to the insight that using stochastic optimization has an advantage over using just (instantaneous) expectation for defining the dynamics. Moreover it also leads to decreasing expected costs if one is able to observe the number of infected individuals over time, thus the measurement information has a quantitative value for the decision maker.

Clearly several open questions have to be left for further research: The SIS model is one of the simplest epidemiological models, so it is natural to apply the basic ideas of stochastic SFP and SOP to more elaborated models (e.g. more compartments or heterogeneity). In the optimization model so far we restricted the consideration to controlling the recovery rate (which only influences the drift), while it might also be important to influence the infection process (e.g. by vaccination). Moreover, one may also consider different forms of the cost function.

References

- [1] F.X. Abad, R.M. Pinto, A. Bosch, Survival of enteric viruses on environmental fomites, *Appl. Environ. Microbiol.* 60 (1994) 3704–3710.
- [2] E. Albina, Epidemiology of porcine reproductive and respiratory syndrome (PRRS): an overview, *Vet. Microbiol.* 55 (1997) 309–316.
- [3] F. Ball, T. Britton, O. Lyne, Stochastic multitype epidemics in a community of households: estimation and form of optimal vaccination schemes, *Math. Biosci.* (2004).
- [4] H. Behnke, Optimal control of deterministic epidemics, *Optimal Control Appl. Methods* 21 (2000) 269–285.
- [5] F. Brauer, L.J.S. Allen, P. Van den Driessche, J. Wu, *Mathematical Epidemiology*, Lecture Notes in Math. 1945, Math. Biosci. Subser., Springer, 2008.
- [6] C. Chen, Y. Kang, Dynamics of stochastic SIS epidemic model with saturated incidence, *Abstr. Appl. Anal.* (2014).
- [7] L. Chen, J. Sun, Optimal vaccination and treatment of an epidemic network model, *Phys. Lett. A* 378 (41) (2014) 3028–3036.
- [8] L. Chen, J. Sun, Global stability and optimal control of an sirs epidemic model on heterogeneous networks, *Phys. A* 410 (2014) 196–204.
- [9] M. de Pinho, H. Maurer, H. Zidani, Optimal control of normalized simr models with vaccination and treatment, *Discrete Contin. Dyn. Syst. Ser. B* 23 (2018) 7–99.
- [10] N. Dexter, Stochastic models of foot and mouth disease in feral pigs in the Australian semi-arid rangelands, *J. Appl. Ecol.* 40 (2003) 293–306.
- [11] O. Diekmann, H. Heesterbeek, T. Britton, *Mathematical Tools for Understanding Infectious Disease Dynamics*, Princeton Series in Theoretical and Computational Biology, Princeton University Press, Princeton, 2013.
- [12] J. Dureau, K. Kalogeropoulos, M. Baquelin, Capturing the time-varying drivers of an epidemic using stochastic dynamical systems, *Biostatistics* 14 (3) (2013) 541–555.
- [13] G. Fichera, *On a Unified Theory of Boundary Value Problems for Elliptic-Parabolic Equations of Second Order*, University of Wisconsin Press, Madison/Wisconsin, 1960, pp. 97–120.
- [14] W.H. Fleming, R.W. Rishel, *Deterministic and Stochastic Optimal Control*, Springer-Verlag, New York, Heidelberg, Berlin, 1975.
- [15] C. Gardiner, *Stochastic Methods — A Handbook for the Natural and Social Sciences*, Springer, Berlin, Heidelberg, New York, 2009.
- [16] A. Gray, D. Greenhalgh, L. Hu, X. Mao, J. Pan, A stochastic differential equation SIS epidemic model, *SIAM J. Appl. Math.* 71 (3) (2011) 870–902.
- [17] H.W. Hethcote, J.A. Yorke, *Gonorrhoea Transmission Dynamics and Control*, Lecture Notes in Biomath., vol. 56, Springer, 1984.
- [18] M. Hinze, R. Pinnau, M. Ulbrich, S. Ulbrich, *Optimization with PDE Constraints*, Mathematical Modelling: Theory and Applications, vol. 23, Springer, Heidelberg, New York, 2009.
- [19] M.J. Keeling, P. Rohani, *Modeling Infectious Disease in Humans and Animals*, Princeton University Press, Princeton and Oxford, 2008.
- [20] M.J. Keeling, J.V. Ross, On methods for studying stochastic disease dynamics, *J. R. Soc. Interface* 5 (2008) 171–181.
- [21] C.T. Kelley, *Iterative Methods for Optimization*, SIAM, Philadelphia, 1999.
- [22] D.E. Kirk, *Optimal Control Theory — An Introduction*, Dover, Mineola, New York, 2004.
- [23] R.M. Kovacevic, Stochastic contagion models without immunity: their long term behaviour and the optimal level of treatment, *CEJOR Cent. Eur. J. Oper. Res.* 26 (2) (2018) 395–421.
- [24] A.L. Krause, L. Kurowski, K. Yawar, R.A. Van Gorder, Stochastic epidemic metapopulation models on networks: Sis dynamics and control strategies, *J. Theoret. Biol.* 449 (2018) 35–52.
- [25] V.A. Ladyženskaja, O.A. Solonnikov, N.N. Uralceva, *Linear and Quasilinear Equations of Parabolic Type*, American Mathematical Society, Providence, Rhode Island, 1968.
- [26] R. Marton, K. Wickwire, On the optimal control of a deterministic epidemic, *Adv. in Appl. Probab.* 6 (1974) 622–635.
- [27] J.D. Murray, *Mathematical Biology*, Springer, New York, 1989.
- [28] C. Nowzari, V.M. Preciado, G.J. Pappas, Analysis and control of epidemics: a survey of spreading processes on complex networks, *IEEE Control Syst.* 36 (1) (2016) 26–46.

- [29] O.A. Oleinik, On linear second order equations with non-negative characteristic form, *Actes, Congres Intern. Math.* 2 (1970) 771–777.
- [30] O.A. Oleinik Oleinik, E.V. Radkevich, *Second Order Equations with Nonnegative Characteristic Form*, Plenum Press, New York, London, 1973.
- [31] H. Raiffa, R. Schlaifer, *Applied Statistical Decision Theory*, Harvard University Press, 1961.
- [32] D. Revuz, M. Yor, *Continuous Martingales and Brownian Motion*, 3rd edition, Springer, Berlin, 1998.
- [33] W.E. Schiesser, *The Numerical Method of Lines*, Academic Press Inc., San Diego, New York, Boston, 1991.
- [34] W.A. Spurr, C. Bonini, *Statistical Analysis of Business Decisions*, Richard D. Irwin Publishing House, 1973.
- [35] F. Tröltzsch, *Optimal Control of Partial Differential Equations: Theory, Methods and Applications*, Graduate Studies in Mathematics, vol. 112, American Mathematical Society, Providence, Rhode Island, 2010.