

Unusual Role of a Cysteine Residue in Substrate Binding and Activity of Human AP-Endonuclease 1

Anil K. Mantha^{1†}, Numan Oezguen^{1,2†}, Kishor K. Bhakat¹,
Tadahide Izumi³, Werner Braun^{1,2} and Sankar Mitra^{1*}

¹Department of Biochemistry and Molecular Biology, University of Texas Medical Branch, Galveston, TX 77555, USA

²Sealy Center for Structural Biology and Molecular Biophysics, University of Texas Medical Branch, Galveston, TX 77555, USA

³Stanley S. Scott Cancer Center and Department of Otolaryngology, 533 Bolivar St., 5th Floor, Louisiana State University Health Sciences Center, New Orleans, LA 70112, USA

Received 20 December 2007;
received in revised form
10 March 2008;
accepted 25 March 2008
Available online
3 April 2008

The mammalian AP-endonuclease (APE1) repairs apurinic/aprimidinic (AP) sites and strand breaks with 3' blocks in the genome that are formed both endogenously and as intermediates during base excision repair. APE1 has an unrelated activity as a redox activator (and named Ref-1) for several trans-acting factors. In order to identify whether any of the seven cysteine residues in human APE1 affects its enzymatic function, we substituted these singly or multiply with serine. The repair activity is not affected in any of the mutants except those with C99S mutation. The Ser99-containing mutant lost affinity for DNA and its activity was inhibited by 10 mM Mg²⁺. However, the Ser99 mutant has normal activity in 2 mM Mg²⁺. Using crystallographic data and molecular dynamics simulation, we have provided a mechanistic basis for the altered properties of the C99S mutant. We earlier predicted that Mg²⁺, with potential binding sites A and B, binds at the B site of wild-type APE1–substrate complex and moves to the A site after cleavage occurs, as observed in the crystal structure. The APE1–substrate complex is stabilized by a H bond between His309 and the AP site. We now show that this bond is broken to destabilize the complex in the absence of the Mg²⁺. This effect due to the mutation of Cys99, ~16 Å from the active site, on the DNA binding and activity is surprising. Mg²⁺ at the B site promotes stabilization of the C99S mutant complex. At higher Mg²⁺ concentration the A site is also filled, causing the B-site Mg²⁺ to shift together with the AP site. At the same time, the H bond between His309 and the AP site shifts toward the 5' site of DNA. These shifts could explain the lower activity of the C99S mutant at higher [Mg²⁺]. The unexpected involvement of Cys99 in APE1's substrate binding and catalysis provides an example of involvement of a residue far from the active site.

© 2008 Elsevier Ltd. All rights reserved.

Keywords: human APE1; Ref-1; AP-endonuclease; DNA base excision repair; DNA binding

Edited by J. Karn

Introduction

The genomes of all organisms are being continuously damaged due to both spontaneous chemical

reactions and exposure to endogenous genotoxic agents, in particular, reactive oxygen species that are generated as by-products of respiration. A total of 10³ to 10⁵ DNA lesions are estimated to be induced in a genome per day including oxidized bases, apurinic/aprimidinic (AP) sites and DNA single-strand breaks.^{1,2} Unrepaired AP sites and their cleavage products are cytotoxic and mutagenic.³ AP-endonucleases (APEs), responsible for repair of the AP sites and 3' end cleaning at single-strand breaks, are highly conserved among various species. APEs belong to two families, Xth and Nfo, whose prototypes are exonuclease III (Xth) and endonuclease IV (Nfo) in *Escherichia coli*.^{4–6} Mammalian cells express

*Corresponding author. E-mail address: samitra@utmb.edu.

† A.K.M. and N.O. contributed equally to this study.

Abbreviations used: APE, AP-endonuclease; AP site, apurinic/aprimidinic site; BER, base excision repair; Ref-1, redox factor 1; hAPE1, human APE1; WT, wild type; MD, molecular dynamics; THF, tetrahydrofuran; EDTA, ethylenediaminetetraacetic acid.

only one active APE, APE1, an ortholog of *E. coli* Xth with an unconserved N-terminal extension.^{4,7} All APEs have two intrinsic activities in DNA repair. They act as an endonuclease in cleaving AP sites to generate 3' OH and 5' phosphodeoxyribose termini. They also act as a 3' phosphodiesterase/exonuclease to remove 3' blocking phosphodeoxyribose or its fragments generated during strand breaks, and by reactive oxygen species or DNA glycosylases that excise oxidized bases in the first step of base excision repair (BER).⁴ Thus, APE's key function is to produce a 3' OH terminus that serves as a primer for repair synthesis.⁸

The mammalian APE1 was independently discovered as redox factor 1 (Ref-1), involved in activation shown to many critical trans-acting proteins including C-Jun and p53.^{9–12} Cys65 (Cys64 in mouse APE1), initially identified as the reductant for such activation, was subsequently shown to be dispensable for this function.^{13,14} APE1/Ref-1 was also identified as a component of the nCaRE-bound protein complexes.^{15,16} We have recently shown that Lys6 or Lys7 of APE1 is acetylated by the histone acetyltransferase p300 that enhances APE1's binding to nCaREs.¹⁷ APE1 was earlier shown to be essential for embryo development and no APE1-null mammalian cell could be established.¹³ Our subsequent studies established the essentiality of APE1 in somatic cells, which is due to both acetylation-dependent functions and DNA repair activity.⁹

The X-ray crystallographic structure of mammalian/human APE1 (hAPE1) has provided significant insight into the molecular mechanism of AP-site recognition and cleavage by APE1.^{18,19} The phosphodiester bond cleavage by APE1 requires Mg^{2+} , similar to other DNase I family members. We proposed earlier that this divalent metal ion shuttles between two metal-binding sites, A and B, in hAPE1.²⁰ Kinetic and binding studies of wild-type (WT) and mutant APE1 show that product dissociation is dependent on Mg^{2+} concentration.²¹ Arg156 and Tyr128 are the predominant residues bound to the minor groove of AP-site-containing DNA for stabilizing the enzyme–DNA complex.²² Several amino acid residues in hAPE1 have been identified by biochemical and structural studies that are required for phosphodiester cleavage at the AP site.^{4,18,21,23–26} The critical residues required for phosphodiester bond cleavage and Mg^{2+} coordination are Glu96, Asp210 and His309, all of which are conserved in the Xth family members. His309 forms an H bond with the AP phosphate and is thus critical for substrate/product binding.^{20,21,24,27} APE1 contains several highly conserved sequence motifs, including 94LQETK98, in which Glu96 is involved in coordinating Mg^{2+} .^{23,28,35}

Redox regulation by APE1 via one or more of seven Cys residues has been investigated for more than a decade. Cys65, Cys93 and Cys310 have been implicated in its redox activity.^{13,29} Interestingly, some Cys residues are located close to the catalytic core domain for APE activity. These include C93 and C99, which are adjacent to the metal-coordinating

domain, and C310, which is adjacent to the active-site H309. However, there have been no systematic studies to elucidate the possible role of the Cys residues in catalysis. This study is an effort to identify the active site for redox activation; we individually mutated all seven Cys residues in hAPE1 to Ser and tested for a possible linkage between the redox function and AP-site cleavage activity. Our results indicate that all Cys residues are dispensable for the AP-endonuclease or phosphodiesterase activity. However, unexpectedly, the C99S mutant lost intrinsic affinity for the AP-DNA substrate in the absence of Mg^{2+} , and its AP-endonuclease activity, comparable to that of the WT APE1 at low Mg^{2+} concentration, is strongly inhibited at high Mg^{2+} concentration. This unusual behaviour of the Cys99 mutant can be explained by molecular dynamics (MD) simulation analysis based on our previously proposed moving-metal mechanism.²⁰

Results

Cys residues in APE1 are dispensable for AP-endonuclease activity

WT APE1 and its Cys mutants were expressed as His-tag fusion polypeptides as described in Materials and Methods. After affinity purification and cleavage of the His-tag, we purified the untagged proteins to near homogeneity by fast protein liquid chromatography using heparin–Sepharose (Fig. 1). To test whether the mutation affected APE1 activity,




Fig. 1. Purification of WT and mutant (Cys→Ser) APE1 proteins. Coomassie Blue staining of purified WT APE1 and Ser mutants (0.5 μ g protein) after SDS-PAGE (10% polyacrylamide). Lane 1, protein marker; lane 2, WT APE1; lane 3, C65S; lane 4, C93S; lane 5, C99S; lane 6, C138S; lane 7, C208S; lane 8, C296S; and lane 9, C310S.




Fig. 2. Mg²⁺-dependent AP-endonuclease activity of the WT and mutant (Cys→Ser, Ala and Thr) APE1 proteins. The 43-mer, THF-containing oligonucleotide was incubated with 33 pM APEs for 3 min at 37 °C. AP-endonuclease activity was measured using a buffer containing 10 mM Mg²⁺. (a) Lane 1, no protein; lane 2, WT APE1; lanes 3–5, three different batches of C99S mutant; lane 6, C99A mutant; lane 7, C99T mutant; lane 8, C99S/C138S double mutant;

lane 9, C99S/C65S/C138S triple mutant. (b) Lane 1, WT APE1; lane 2, C65S mutant; lane 3, C93S mutant; lanes 4–6, three different batches of C99S mutant; lane 7, C138S mutant; lane 8, C208S mutant; lane 9, C296S mutant; lane 10, C310S mutant.

we used a 43-mer duplex oligo containing the stable AP-site analog tetrahydrofuran (THF) at position 31.³⁰ Our standard assay buffer contained 10 mM Mg²⁺, and we ensured a linear dependence of activity with respect to the amount of enzyme and time. We observed that all Cys mutants except those with C99S mutation were at least as active as the WT APE1 (Fig. 2b). We concluded that the Cys residues, including those reported to be involved in Ref-1 function, are dispensable for APE1 repair activity. However, the AP-endonuclease activity of the C99S mutant as well as of the double mutants C138S/C99S and C65S/C99S was strongly inhibited in the presence of 10 mM Mg²⁺. Furthermore, the loss of activity due to Cys99 mutation was independent of Cys substitution with Ser, Ala or Thr (Fig. 2a). Finally, in order to confirm that the activity loss due to Cys99 was not an artefact, we produced several independent batches of the C99S mutant, all of which behaved identically.

The Cys99 APE1 mutant is inhibited by high Mg²⁺

Further studies revealed that the C99S mutant was active in the presence of low Mg²⁺ concentration. In fact, the activity of Cys99 mutants was comparable to that of the WT and other Cys mutants in 0.5 or 2 mM Mg²⁺ (Supplementary Fig. 1), while only these

Cys99 mutants were strongly inhibited at 5 mM and higher Mg²⁺ concentration.

Loss of substrate DNA binding due to C99S mutation

APE1 has significant affinity for the AP-DNA substrate in the absence of Mg²⁺, and the structure of APE1–substrate oligo complex without a divalent metal ion was solved earlier by X-ray crystallography.¹⁹ We used electrophoretic mobility shift analysis (EMSA) to confirm binding of WT APE1 to the AP site in the absence of Mg²⁺ (to prevent cleavage of the bound substrate). However, unlike WT and other Ser mutant polypeptides, the C99S mutant did not bind to the AP site (Fig. 3). This suggested that, unlike other Cys residues, Cys99 in hAPE1 is directly or indirectly involved in contact with the substrate AP site.

Fluorescence emission spectra of WT and C99S mutant APE1

Intrinsic tryptophan (Trp) fluorescence emission spectra were generated for the WT and C99S APE1. The fluorescence output for the mutant was less than that of the WT protein without any change in the emission spectra (Supplementary Fig. 2). Similar decrease in emission intensity without a change in




Fig. 3. EMSA of THF–DNA binding by WT APE1 and mutants. Lane 1, no protein; lane 2, WT APE1; lane 3, C65S mutant; lane 4, C93S mutant; lane 5, C99S mutant; lane 6, C138S mutant; lane 7, C208S mutant; lane 8, C296S mutant; and lane 9, C310S mutant. The experiments were repeated at least four times. Other details are given in Materials and Methods.

the spectra was earlier observed for the K98R mutant APE1, which we had interpreted to be due to an increase in collisional quenching.²³

Kinetic parameters for WT versus C99S APE1

To further probe the effect of the C99S mutation on APE1 enzymatic activity, we determined the kinetic parameters of purified WT and mutant enzymes. Increasing Mg^{2+} concentration to 10 mM inhibited product formation by 5.4-fold in the mutant as compared to the WT APE1 (Fig. 4c). At 20 mM Mg^{2+} , the product formation with WT APE1 was inhibited 4.2-fold and with the C99S mutant ~14-fold relative to the activity of the WT protein in 5 mM Mg^{2+} (Fig. 4c). Figure 5 shows that the K_m of the C99S mutant in 2 mM Mg^{2+} was slightly higher (53.6 nM) than that of the WT enzyme (36.8 nM), which is comparable to the published values.^{23,31} However, the k_{cat} values of the mutant and WT proteins in 2 mM Mg^{2+} were comparable (Table 1).

Molecular dynamics simulation

The impact of the C99S mutation on the DNA binding and endonuclease activity of APE1 was quite surprising in view of the fact that Cys99 is ~16 Å away from the active site. We then performed




Fig. 4. Effect of Mg^{2+} on AP-endonuclease activity of (a) WT and (b) C99S mutant APE1. Lane 1, control; lanes 2–11, 0, 0.1, 0.5, 1, 2, 3, 5, 10, 20, and 40 mM Mg^{2+} , respectively. Lane 2 also contains 0.1 mM EDTA. (c) Effect of Mg^{2+} on the activity of WT and Ser99 mutants.




Fig. 5. Reaction kinetics of WT and C99S mutant APE1 (33 pM) in the presence of 2 mM $[Mg^{2+}]$. (a) WT APE1 and (b) C99S mutant proteins were incubated with 0–160 nM of THF-containing oligonucleotide at 37 °C for 3 min. Other details are described in Materials and Methods. Data represent mean values of four independent experiments.

MD simulation analysis to examine the effect of this mutation on the structure and dynamics of hAPE1, as our previous work on simulation for WT hAPE1 led us to postulate a “moving-ion mechanism”.²⁰ We predicted that in the precleavage complex of APE1 and DNA, the divalent metal ion is present at the buried B site, where it is coordinated by Asp210, Glu96, Asn212 and AP phosphate. We also showed that two H bonds are stably formed between Asp283 and His309 and between His309 and the AP phosphate oxygen. Here we compare the presence of the H bond between His309 and the AP site in the

Table 1. Kinetic parameters for AP-endonuclease activity of wild-type and C99S mutant APE1 proteins in the presence of 2 mM Mg^{2+}

	K_m (nM)	k_{cat} (s^{-1})	k_{cat}/K_m ($nM^{-1} s^{-1}$)
WT APE1	36.8±7.6	3.36±0.23	0.091
C99S APE1	53.6±13.1	2.91±0.28	0.054

WT and C99S APE1 during a 2-ns dynamics simulation. Figure 6 shows whether this H bond is formed (1) or broken (0), while Fig. 7 shows the distances between the atoms NE2 of His309 and P of the AP site. The H bond in the WT enzyme is stable and independent of the presence of Mg^{2+} (Figs. 6a and 7a). For the C99S mutant this H bond between His309 and AP site is metastable in the presence of Mg^{2+} (Fig. 6b), and in the absence of Mg^{2+} this bond is completely broken at ~ 1800 ps (Figs. 6c and 7a and b). At this time, the AP site moves away, increasing the distance from ~ 4.2 to 8 \AA between the atoms at NE2 of His309 and the P on the AP site (Fig. 7). At the same time, the binding to DNA weakens as the distance to the CA atom of Asp210 increases (Fig. 8b). Adding Mg^{2+} to the C99S mutant changes the dynamics of the AP site along with the rest of the DNA. As shown in Fig. 8a, Mg^{2+} pulls the AP site down by $\sim 1 \text{ \AA}$ into the active-site pocket towards Asp210.

Cys99 in WT APE1 does not interact with DNA at all, as we had shown earlier in several independent simulations.²⁰ The shortest H bonds to the complementary strand in DNA are formed with the Arg73 side chain. However, in several independent simulations with C99S mutation, Arg73 forms more H bonds to DNA phosphate compared to WT APE1, and Ser99 directly forms an H bond to the DNA or to Arg73. These additional H bonds pull the complementary strand closer to Arg73 and Ser99. The change in the relative position of the complementary strand is transmitted via the DNA backbone and weakens the H bond between the AP site and His309. Figure 9 shows two snapshots of simulation




Fig. 7. Time course of the distance between atoms NE2 on the His309 and P on the AP site for (a) WT APE1, C99S and C99S+ Mg^{2+} simulations and (b) C99S APE1 mutant simulation.

with the C99S mutant at 1 ps and at 22 ns. At the beginning, there are seven H bonds between Arg73 and the DNA, and one H bond is also formed between Arg73 and Ser99; towards the end, only




Fig. 6. Kinetics of H-bond formation/breakage between NE2 of H309 and O1P of the AP-site phosphate for (a) WT APE1, (b) C99S mutant with Mg^{2+} at the B site, and (c) C99S APE1 mutant. The value is 1 when a H bond is formed and 0 when not formed.




Fig. 8. Typical time course for the distance between atoms CA of Asp210 and P of the AP site for (a) WT APE1, C99S and C99S+Mg²⁺ simulations and (b) C99S APE1 mutant simulation.

three H bonds are formed between Arg73 and DNA, which is similar to the situation with WT APE1.

Our earlier study with WT APE1 showed that Mg²⁺ located at the A or B site in WT APE1 were 5.7 Å apart.²⁰ However, at higher [Mg²⁺] when both sites are filled, two Mg²⁺ ions move apart and the distance increases to 7.7 Å after 500 ps. Analogous simulation with the C99S mutant shows further increase of the intermetal distance to 9.6 Å (Fig. 10). The final

positions of Mg²⁺ are shown by red spheres, and the unperturbed single ion positions of the A and B sites are shown by blue spheres. In this study and our previous simulations, Mg²⁺ at the B site was coordinated by the phosphate oxygen of the AP site and Asp210. Accordingly, the AP site and the P are also displaced by 2.6 Å in the C99S simulation with 2 Mg²⁺ ions. The AP site is still bound at the active site as indicated by its distance of 5.7 Å from the CA atom of Asp210. However, the substrate-stabilizing H bond is now between His309 and O3'. It is shifted towards the 5' end of the DNA. At the end of the simulation, the relative positions of the active-site constituents were significantly disturbed.

Taken together, our simulations show that the Ser99 mutation introduces stronger interaction between the DNA, Arg73 and Ser99 sites. This tension propagates via H bonds between the DNA strands along the backbone to the AP site. In the absence of Mg²⁺, this weakens and finally breaks the AP-site-stabilizing H bond between the H309 to an AP-site phosphate oxygen. The Cys99Ser mutation thus has a similar effect, as the His309 to Asn309 mutation also breaks the H bond to the phosphate oxygen and releases the substrate at a higher rate than the WT enzyme.²¹ In the presence of Mg²⁺, the tension, which originates from the pull of the Arg73 and Ser99 site, is partially compensated for by the interaction with Mg²⁺. Thus, the H bond between His309 and an AP-site phosphate oxygen is weakened; consequently, the internuclear distance fluctuates. Nevertheless, at higher Mg²⁺ concentration, when both metal-binding sites are expected to be filled, the enzymatic activity decreases. This is consistent with our earlier finding that having two metal ions in the active site caused the metal at the B site to move about 2.6 Å and the internuclear distance to increase from 5.7 Å (unperturbed A to B site distance) to 7.7 Å in WT APE1.²⁰ In simulation with the Ser99 mutant and two Mg²⁺ ions at the




Fig. 9. Snapshots of the C99S mutant simulation at the beginning after 1 ps (a) and at the end after 22,500 ps (b). Only the DNA and key residues that are important for this study in the C99S mutant protein are shown. The AP-site-containing DNA strand is shown in cyan and the complementary strand in magenta. The side chains of key residues of the C99S mutant are shown in yellow. Dotted green and blue lines represent H bonds and the distance is indicated.




Fig. 10. Snapshots of simulation of the C99S mutant after 500 ps with two Mg^{2+} ions in the active site. The positions of key DNA residues and Mg^{2+} are shown. The final Mg^{2+} positions are shown in red; for comparison the starting positions, which are identical to the unperturbed precleavage A and B site, are shown in blue. Important H bonds are shown as dotted green lines.

active site, this distance increased even further to 9.6 Å. This alters the active-site geometry to the point where the phosphodiester bond cleavage activity is strongly suppressed while the substrate is still bound in the active site.

Discussion

The mammalian APE1 is an extremely unusual DNA repair protein because of its second unrelated activity in transcriptional regulation. The situation turned out to be more complex with the discovery of two regulatory activities in APE1/Ref-1, one in reductive activation of some transcription factors^{11,32} and the other in acetylation-dependent modulation of other trans-acting factors.¹⁷ We have shown that the second regulatory function, in addition to the DNA repair activity, is essential for survival.⁹ However, essentiality of the repair activity was unexpected because the lack of APEs including APE1 orthologs in *E. coli* and yeast does not cause cell death.^{33,34}

The Cys residues in mammalian APE1 became important after the discovery of the redox activation (Ref-1) function of this protein. It is interesting to note that among the seven Cys residues in hAPE1, Cys208 and Cys310 are completely conserved among all APEs, while Cys99, Cys93 and Cys296 are conserved only among metazoan APEs. It is also intriguing that Cys208, Cys310 and Cys99 are close to the key residues Asp210, His309 and Glu96, respectively, which are involved in catalysis and metal binding for the AP-endonuclease activity. Furthermore, an earlier study suggested direct involvement of Cys310 in the reaction chemistry of hAPE1.²⁹ Although Cys residues have not been implicated in the catalytic steps of other APEs, we decided to directly assess their role in hAPE1 by characterizing site-specific mutations. Our results

clearly show that at low $[Mg^{2+}]$, substitution of individual Cys with Ser for all seven Cys residues did not affect enzyme turnover in regard to either endonuclease or phosphodiesterase activity. In fact, Ser138 and Ser208 mutants may have slightly higher specific activity than the WT enzyme. However, the aberrant behaviour of the Ser99 mutant in regard to its affinity loss for the substrate in the absence of Mg^{2+} was surprising. A small reduction in quantum yield without a change in the emission spectra of intrinsic Trp fluorescence suggested only a subtle change in the tertiary structure of hAPE1 due to the C99S mutation. To gain more definitive insight into the structural perturbation that caused this unusual behaviour, we exploited MD modeling and extended our previous simulation studies.

Earlier we have shown that APE1's binding to the substrate/product DNA via H bonds is necessary for efficient functioning of the BER pathway.³⁵ The present study shows the importance of the AP-site-stabilizing H bond from His309 to the AP-site phosphate oxygen. Weakening of this H bond in the C99S mutant reduces the affinity of the AP-site DNA in the absence of Mg^{2+} . At low Mg^{2+} concentration, this interaction is partially restored. The reduced affinity of the C99S mutant for AP-site-containing DNA was specifically addressed in MD simulations. Being so far from the active site (~ 16 Å), Cys99 indirectly affects the relative position of the AP site that is bound in the active site to H309. We postulate that in the Ser99 mutant, Mg^{2+} at high concentration occupies both A and B sites resulting in inhibition of the endonuclease activity as we had proposed earlier.²⁰ MD simulation gave further indication that the H bond between the active-site H309 and the AP site is metastable in the C99S mutant in the absence of Mg^{2+} as compared to the WT enzyme. In the absence of the metal ion, once the H bond is broken, the AP-site-containing DNA moves away resulting in increase in distance between NE2 of H309 and P on the AP site. This also leads to an increase in distance between the CA atom of Asp210, and DNA binding weakens. The presence of Mg^{2+} at the B site changes the dynamics of the AP site, forcing it downward to the active site towards Asp210.

In summary, we have identified Cys99 in hAPE1 as a critical residue modulating APE's function for ensuring optimum activity. Substitution of this residue alters the enzyme's ability to bind the AP site substrate in the absence of Mg^{2+} . Further studies with this mutant protein may shed light on metal-dependent substrate recognition and processing in the BER pathway.

Materials and Methods

Purification of WT and mutant APE1s

The coding sequence for full-length APE1 was inserted in the pET15b vector (Novagen) at Nde I/Xho I sites for expression of APE1 in *E. coli*. Other single site-directed mutations at all seven Cys amino acid positions including

Cys65Ser, Cys93Ser, Cys99Ser, Cys138Ser, Cys208Ser, Cys296Ser and Cys310Ser were created in the APE1 cDNA with a site-directed mutagenesis kit (QuickChange II XL, Stratagene) by following the manufacturer's guidelines. The DNA sequence of the WT APE1 and mutant cDNAs were confirmed by the University of Texas Medical Branch recombinant DNA laboratory (R.D.L.).

WT APE1 and the Ser mutants were purified as previously described²³ with slight modifications. After transformation with the pET15b-based APE1 expression plasmids, *E. coli* BL21 (DE3) cultures (1 L in L-broth) were grown to 0.6 OD at 600 nm. APE1 expression was then induced with 0.5 mM IPTG at 16 °C for 14–16 h.²³ Cells were collected in suspension buffer containing 20 mM Tris (pH 8.0) and 0.5 M NaCl, sonicated and centrifuged. The supernatant was loaded onto a Ni-NTA (Qiagen) column (3 mL), which was first washed with 18 mL sonication buffer containing 20 mM imidazole. WT and mutant APE1s were then eluted with 10 mL sonication buffer containing 200 mM imidazole, and the eluate was dialyzed *versus* 20 mM Tris–Cl (pH 8.0), 100 mM NaCl, 1 mM ethylenediaminetetraacetic acid (EDTA), 1 mM DTT and 10% glycerol. The poly His-tags in the proteins were cleaved by overnight incubation at 4 °C with thrombin. The APE1 polypeptides were finally purified to near-homogeneity by fast protein liquid chromatography in a 1-mL SP-Sepharose column (LCC-500 PLUS; Pharmacia) and the final preparations were dialyzed *versus* 20 mM Tris (pH 8.0), 300 mM NaCl, 0.1 mM EDTA, 1 mM DTT and 50% glycerol and stored at –20 °C.

AP-endonuclease assay

A 43-mer oligonucleotide containing the AP-site analog THF at nt 31 (Midland Corp), was prepared as described previously.^{23,36,37} The oligo was 5'-end-labeled with [γ -³²P]ATP using T4 polynucleotide kinase. After annealing to the complementary strand with A opposite THF, the duplex oligo was purified from the free nucleotides by gel filtration column (Chroma Spin TE 10, Clontech).

The duplex THF-containing oligo was incubated with recombinant WT APE1 or Cys mutants at 37 °C for 3 min during which the reaction rate was linear in a 15- μ L reaction mixture containing 50 mM Tris–HCl (pH 8.5), 50 mM KCl, 1 mM DTT, 0.1 mM EDTA and 100 μ g/mL bovine serum albumin and various concentrations of MgCl₂. The reaction was stopped with 10 μ L 80% formamide/40 mM NaOH containing 0.05% xylene cyanol, followed by heating at 95 °C for 5 min; the samples were kept on ice until denaturing gel electrophoresis in 20% polyacrylamide containing 8 M urea to separate the substrate oligo from the cleaved product. The gels were dried and the radioactivity was quantitated by phosphorimager analysis in a Storm system (Molecular Dynamics). The enzyme kinetics data were fitted by nonlinear least-squares regression to obtain V_{\max} and K_m with the Michaelis–Menten equation using Sigma plot software. The kinetic parameters K_m and k_{cat} were calculated after incubating 33 pM enzyme at 37 °C for 3 min with substrate at various concentrations.

Electrophoretic mobility shift analysis

The 5'- γ -³²P-labeled 43-mer THF-oligo (about 25 pM) was incubated with 5 ng of WT or mutant APE1 in the EMSA-binding buffer containing 40 mM HEPES, 50 mM KCl, 1 mM DTT and 2 mM EDTA for 15 min at 25 °C

followed by electrophoresis in nondenaturing 6% polyacrylamide gel containing 7 mM Tris–HCl (pH 8), 3.3 mM sodium acetate and 1 mM EDTA at 4 °C for 1 h at 150 V. The gels were processed as described earlier.³⁵

Fluorescence spectroscopy

The fluorescence emission spectra with 0.5 μ M of WT and C99S mutant APE1 proteins in phosphate-buffered saline were acquired using an excitation wavelength of 295 nm in a Shimadzu RF-5301PC fluorimeter. An average of four repetitive scans were considered.³⁵

Molecular dynamic simulation

MD simulations were done with the AMBER9 program suite³⁸ with complexes of WT APE1, uncleaved AP-site-containing DNA substrate and the cofactor Mg²⁺. Models for the WT APE1, Cys99→Ser99 mutant and the DNA were generated from the crystal structure 1DE8.¹⁹ The 11-bp DNA oligo has flipped out and the AP site bound to the WT APE1. Active-site His309 was protonated in all models and succeeding simulation to account for its role in stabilizing the substrate/product. It was shown experimentally that H309N mutation abolishes endonuclease activity and increases substrate release rate.^{21,24} In previous work we showed that His309 only forms the stabilizing H bond to the AP-site oxygen when it is protonated.²⁰ For simulations with the cofactor Mg²⁺ we placed the Mg²⁺ ions at the A- and B-site positions.^{20,25}

Na⁺ ions were added to all complexes to achieve electrostatic neutrality and then soaked in a TIP3 water box.³⁹ The dimensions of the water box were chosen to be 10 Å beyond any complex atom. These models were first energy-minimized over 2000 steps using a standard procedure with the Sander program of the AMBER9 distribution.³⁸ Dynamic simulations over several nanoseconds followed the minimization. The parameters for the MD simulations were constant pressure and temperature, periodic boundary conditions and the ff03 force field.⁴⁰ Finally, analysis of the trajectories and visualizations were done with the programs CARNAL of the AMBER6 distribution and MOLMOL.⁴¹

Acknowledgements

This research was supported by USPHS grants RO1 CA53791 and RO1 ES08457 and grant P30 ES006676 (to S.M.), RO1 CA98664 (to T.I.) and US DOE grant DE-FG02-04ER63826 (to W.E.B.). We thank Drs. T. Hazra and M. Hegde at the University of Texas Medical Branch (UTMB), and Dr. G. Tell at University of Udine for helpful discussions during the study. We thank Dr. R. Chattopadhyay, University of Virginia, for his technical guidance. We acknowledge Dr. R. Roy of Georgetown University for critically reviewing the manuscript. We also thank Drs. T. Wood of the Recombinant DNA Laboratory for DNA sequencing and J. Navarro for providing fluorescence spectroscopy facility at UTMB. Ms. W. Smith's secretarial assistance is gratefully acknowledged.

Supplementary Data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.jmb.2008.03.052](https://doi.org/10.1016/j.jmb.2008.03.052)

References

- Barnes, D. E. & Lindahl, T. (2004). Repair and genetic consequences of endogenous DNA base damage in mammalian cells. *Annu. Rev. Genet.* **38**, 445–476.
- Mitra, S., Izumi, T., Boldogh, I., Bhakat, K. K., Chattopadhyay, R. & Szczesny, B. (2007). Intracellular trafficking and regulation of mammalian AP-endonuclease 1 (APE1), an essential DNA repair protein. *DNA Repair (Amst)*, **6**, 461–469.
- Doetsch, P. W. & Cunningham, R. P. (1990). The enzymology of apurinic/aprimidinic endonucleases. *Mutat. Res.* **236**, 173–201.
- Demple, B. & Harrison, L. (1994). Repair of oxidative damage to DNA: enzymology and biology. *Annu. Rev. Biochem.* **63**, 915–948.
- Johnson, R. E., Torres-Ramos, C. A., Izumi, T., Mitra, S., Prakash, S. & Prakash, L. (1998). Identification of APN2, the *Saccharomyces cerevisiae* homolog of the major human AP endonuclease HAP1, and its role in the repair of abasic sites. *Genes Dev.* **12**, 3137–3143.
- Ribar, B., Izumi, T. & Mitra, S. (2004). The major role of human AP-endonuclease homolog Apn2 in repair of abasic sites in *Schizosaccharomyces pombe*. *Nucleic Acids Res.* **32**, 115–126.
- Mitra, S., Boldogh, I., Izumi, T. & Hazra, T. K. (2001). Complexities of the DNA base excision repair pathway for repair of oxidative DNA damage. *Environ. Mol. Mutagen.* **38**, 180–190.
- Wilson, S. H. (1998). Mammalian base excision repair and DNA polymerase beta. *Mutat. Res.* **407**, 203–215.
- Izumi, T., Brown, D. B., Naidu, C. V., Bhakat, K. K., Macinnes, M. A., Saito, H. *et al.* (2005). Two essential but distinct functions of the mammalian abasic endonuclease. *Proc. Natl. Acad. Sci. USA*, **102**, 5739–5743.
- Evans, A. R., Limp-Foster, M. & Kelley, M. R. (2000). Going APE over ref-1. *Mutat. Res.* **461**, 83–108.
- Xanthoudakis, S., Miao, G., Wang, F., Pan, Y. C. & Curran, T. (1992). Redox activation of Fos-Jun DNA binding activity is mediated by a DNA repair enzyme. *EMBO J.* **11**, 3323–3335.
- Tell, G., Damante, G., Caldwell, D. & Kelley, M. R. (2005). The intracellular localization of APE1/Ref-1: more than a passive phenomenon? *Antioxid. Redox Signal.* **7**, 367–384.
- Walker, L. J., Robson, C. N., Black, E., Gillespie, D. & Hickson, I. D. (1993). Identification of residues in the human DNA repair enzyme HAP1 (Ref-1) that are essential for redox regulation of Jun DNA binding. *Mol. Cell. Biol.* **13**, 5370–5376.
- Ordway, J. M., Eberhart, D. & Curran, T. (2003). Cysteine 64 of Ref-1 is not essential for redox regulation of AP-1 DNA binding. *Mol. Cell. Biol.* **23**, 4257–4266.
- Okazaki, T., Chung, U., Nishishita, T., Ebisu, S., Usuda, S., Mishiro, S. *et al.* (1994). A redox factor protein, ref1, is involved in negative gene regulation by extracellular calcium. *J. Biol. Chem.* **269**, 27855–27862.
- Fuchs, S., Philippe, J., Corvol, P. & Pinet, F. (2003). Implication of Ref-1 in the repression of renin gene transcription by intracellular calcium. *J. Hypertens.* **21**, 327–335.
- Bhakat, K. K., Izumi, T., Yang, S. H., Hazra, T. K. & Mitra, S. (2003). Role of acetylated human AP-endonuclease (APE1/Ref-1) in regulation of the parathyroid hormone gene. *EMBO J.* **22**, 6299–6309.
- Gorman, M. A., Morera, S., Rothwell, D. G., de La Fortelle, E., Mol, C. D., Tainer, J. A. *et al.* (1997). The crystal structure of the human DNA repair endonuclease HAP1 suggests the recognition of extra-helical deoxyribose at DNA abasic sites. *EMBO J.* **16**, 6548–6558.
- Mol, C. D., Izumi, T., Mitra, S. & Tainer, J. A. (2000). DNA-bound structures and mutants reveal abasic DNA binding by APE1 and DNA repair coordination [corrected]. *Nature*, **403**, 451–456.
- Oezguen, N., Schein, C. H., Peddi, S. R., Power, T. D., Izumi, T. & Braun, W. (2007). A “moving metal mechanism” for substrate cleavage by the DNA repair endonuclease APE-1. *Proteins*, **68**, 313–323.
- Masuda, Y., Bennett, R. A. O. & Demple, B. (1998). Rapid dissociation of human apurinic endonuclease (Ape1) from incised DNA induced by magnesium. *J. Biol. Chem.* **273**, 30360–30365.
- Nguyen, L. H., Barsky, D., Erzberger, J. P. & Wilson, D. M., III (2000). Mapping the protein–DNA interface and the metal-binding site of the major human apurinic/aprimidinic endonuclease. *J. Mol. Biol.* **298**, 447–459.
- Izumi, T., Malecki, J., Chaudhry, M. A., Weinfeld, M., Hill, J. H., Lee, J. C. & Mitra, S. (1999). Intragenic suppression of an active site mutation in the human apurinic/aprimidinic endonuclease. *J. Mol. Biol.* **287**, 47–57.
- Masuda, Y., Bennett, R. A. O. & Demple, B. (1998). Dynamics of the interaction of human apurinic endonuclease (Ape1) with its substrate and product. *J. Biol. Chem.* **273**, 30352–30359.
- Beermink, P. T., Segelke, B. W., Hadi, M. Z., Erzberger, J. P., Wilson, D. M., III & Rupp, B. (2001). Two divalent metal ions in the active site of a new crystal form of human apurinic/aprimidinic endonuclease, Ape1: implications for the catalytic mechanism. *J. Mol. Biol.* **307**, 1023–1034.
- Mol, C. D., Hosfield, D. J. & Tainer, J. A. (2000). Abasic site recognition by two apurinic/aprimidinic endonuclease families in DNA base excision repair: the 3' ends justify the means. *Mutat. Res.* **460**, 211–229.
- Rothwell, D. G., Hang, B., Gorman, M. A., Freemont, P. S., Singer, B. & Hickson, I. D. (2000). Substitution of Asp-210 in HAP1 (APE/Ref-1) eliminates endonuclease activity but stabilises substrate binding. *Nucleic Acids Res.* **28**, 2207–2213.
- Schein, C. H., Ozgun, N., Izumi, T. & Braun, W. (2002). Total sequence decomposition distinguishes functional modules, “molegos” in apurinic/aprimidinic endonucleases. *BMC Bioinf.* **3**, 37.
- Kelley, M. R. & Parsons, S. H. (2001). Redox regulation of the DNA repair function of the human AP endonuclease Ape1/ref-1. *Antioxid. Redox Signal.* **3**, 671–683.
- Barzilay, G., Mol, C. D., Robson, C. N., Walker, L. J., Cunningham, R. P., Tainer, J. A. & Hickson, I. D. (1995). Identification of critical active-site residues in the multifunctional human DNA repair enzyme HAP1. *Nat. Struct. Biol.* **2**, 561–568.
- Wilson, D. M., III, Takeshita, M., Grollman, A. P. & Demple, B. (1995). Incision activity of human apurinic endonuclease (Ape) at abasic site analogs in DNA. *J. Biol. Chem.* **270**, 16002–16007.

32. Xanthoudakis, S. & Curran, T. (1992). Identification and characterization of Ref-1, a nuclear protein that facilitates AP-1 DNA-binding activity. *EMBO J.* **11**, 653–665.
33. Guillet, M. & Boiteux, S. (2003). Origin of endogenous DNA abasic sites in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* **23**, 8386–8394.
34. Saporito, S. M., Gedenk, M. & Cunningham, R. P. (1989). Role of exonuclease III and endonuclease IV in repair of pyrimidine dimers initiated by bacteriophage T4 pyrimidine dimer–DNA glycosylase. *J. Bacteriol.* **171**, 2542–2546.
35. Izumi, T., Schein, C. H., Oezguen, N., Feng, Y. & Braun, W. (2004). Effects of backbone contacts 3' to the abasic site on the cleavage and the product binding by human apurinic/aprimidinic endonuclease (APE1). *Biochemistry*, **43**, 684–689.
36. Chattopadhyay, R., Wiederhold, L., Szczesny, B., Boldogh, I., Hazra, T. K., Izumi, T. & Mitra, S. (2006). Identification and characterization of mitochondrial abasic (AP)-endonuclease in mammalian cells. *Nucleic Acids Res.* **34**, 2067–2076.
37. Ramana, C. V., Boldogh, I., Izumi, T. & Mitra, S. (1998). Activation of apurinic/aprimidinic endonuclease in human cells by reactive oxygen species and its correlation with their adaptive response to genotoxicity of free radicals. *Proc. Natl. Acad. Sci. USA*, **95**, 5061–5066.
38. Case, D. A., Cheatham, T. E., Darden, T., Gohlke, H., Luo, R., Merz, K. M. *et al.* (2005). The Amber biomolecular simulation programs. *J. Comput. Chem.* **26**, 1668–1688.
39. Jorgensen, W. L., Chandrasekhar, J. & Madura, J. D. (1983). Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926–935.
40. Ponder, J. W. & Case, D. A. (2003). Force fields for protein simulations. *Adv. Protein Chem.* **66**, 27–85.
41. Koradi, R., Billeter, M. & Wuthrich, K. (1996). MOLMOL: a program for display and analysis of macromolecular structures. *J. Mol. Graphics*, **14**, 51–55.