


Accepted Manuscript

Olive Biophenol Integral Extraction at a Two-Phase Olive Mill

Riccardo Delisi, Rosaria Ciriminna, Stefano Arvati, Francesco Meneguzzo, Mario Pagliaro



PII: S0959-6526(17)32583-0
DOI: 10.1016/j.jclepro.2017.10.278
Reference: JCLP 11070
To appear in: *Journal of Cleaner Production*

Received Date: 30 April 2017
Revised Date: 24 October 2017
Accepted Date: 25 October 2017

Please cite this article as: Riccardo Delisi, Rosaria Ciriminna, Stefano Arvati, Francesco Meneguzzo, Mario Pagliaro, Olive Biophenol Integral Extraction at a Two-Phase Olive Mill, *Journal of Cleaner Production* (2017), doi: 10.1016/j.jclepro.2017.10.278

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Olive Biophenol Integral Extraction at a Two-Phase Olive Mill**Riccardo Delisi,^{*a,b} Rosaria Ciriminna,^b Stefano Arvati,^a Francesco Meneguzzo,^c Mario Pagliaro^{*b}**

^aRenovo Bioindustry, via P. Verri 1, 46100 Mantova, Italy; ^bIstituto per lo Studio dei Materiali Nanostrutturati, CNR, via U. La Malfa 153, 90146 Palermo, Italy; ^cIstituto di Biometeorologia, CNR, via G. Caproni 8, 50145 Firenze, Italy

Abstract

The biophenol integral extraction protocol from vegetation water developed in the early 2000s for the three-phase olive mill was adapted to a large two-phase mill operating in Sicily during the 2016/2017 season. The new set-up allows extensive recovery of olive phenolics, affording after nine months natural extracts with hydroxytyrosol concentration exceeding 1500 ppm. Previous toxic waste is transformed into a source of additional revenues for the milling company, and of valued bioproducts for the bioeconomy partner, eliminating altogether a source of potential pollution.

Keywords: biophenols; olive; polyphenols; hydroxytyrosol; tyrosol

**Corresponding authors*

Dr M. Pagliaro

E-mail: mario.pagliaro@cnr.it

Dr R. Delisi

E-mail: riccardo.delisi@renovospa.it

1. Introduction

The production of large amounts (10-30 million m³) of wastewater at olive mills in the short period of time of olive harvesting and milling is a well known environmental and economic hurdle in Mediterranean countries where most (about 75%) olive oil production takes place every year.¹ Called also vegetation water, olive mill wastewater (OMWW) originates from water contained in the olive fruits plus added water (0.3-0.8 m³ per 1000 kg of olives, depending on the process) to the milled paste in order to extract the highest quantity of oil from the paste via centrifugation in the three-phase milling process. For decades, the latter method has been the process of choice to produce extra virgin olive oil (EVOO). The oil yield increases proportionally to the amount of added water, but external processing water further increases dissolution in the aqueous phase of valued biophenols, showing strong antioxidant, anti-inflammatory, nutrigenomic, chemoprotective and anti-atherosclerotic activities,² widely associated to health benefits of EVOO.³

In general, OMWW cannot be used for irrigation and need to be disposed of as a pollutant at significant economic cost for olive mills.⁴ Several treatment methods have been proposed to improve the quality of OMWW and reduce its phytotoxicity prior to disposal.¹ The simplest disposal practices are storage/evaporation in lagoons and controlled dispersion in soil. Emission limits, however, are generally surpassed and soil and water contamination easily occurs with serious environmental consequences (and closure of olive mills not conforming to environmental laws).⁵

Starting in the early 1990s, a new continuous centrifuge two-phase milling process in which very little water is added, started to be commercially adopted,⁶ first in Spain and then in Greece and in Italy. In place of OMWW, the new milling process affords high quality EVOO along with a wet pomace consisting of olive pulp (ca. 35%) and seed residues dispersed in water released by the olive fruit and about 2.5% of

processing water. Composted with agricultural wastes, this two-phase olive mill waste (TPOMW) becomes an excellent fertilizer ideally recycled in land growing olive trees,⁷ though its significant phytotoxicity still requires its disposal as an industrial waste, whereas the large amount of solid matter can lead to fermentative phenomena complicating its processing.

We argued elsewhere⁸ that olive biophenols extracted from OMWW will shortly find large scale utilization, in a variety of nutraceutical,⁹ food, beverage,¹⁰ and cosmetic formulations,¹¹ in which they exert multiple functions and provide health benefits including powerful anti-inflammatory, antioxidant, antimicrobial, anti carcinogenic, skin and bone regenerative action. Furthermore, recent research suggests that olive phenolics could play a significant role in prevention and treatment of inflammation and of free-radicals mediated chronic diseases.¹² In the following, we show how the biophenol integral extraction protocol from vegetation water developed by Crea in the early 2000s for the three-phase olive mill was successfully adapted to a large two-phase mill operating in Sicily during the 2016/2017 season.

2. Integral Biophenol Extraction from Wet Pomace

Among several biophenol extraction techniques from olive mill waste developed in the last two decades,¹ the process invented by Crea in 2001 allows to obtain an integral *bouquet* of olive biophenols particularly rich in hydroxytyrosol (3,4-dihydroxyphenylethanol) starting from OMWW obtained in the three-phase milling process.¹³ In the latter process, the vegetation waters are centrifuged using a three-phase spin-dryer in order to remove the solid residuals and oil traces. The resulting aqueous phase, called “olive juice”, is chemically stabilized by the addition of citric acid, and then left to age in closed vessels for several months in order to allow for the full hydrolysis of the main glycosides (oleuropein and verbascoside).

Once aging is complete (full hydrolysis of oleuropein), the resulting biophenol extract possessing negligible COD and BOD₅ values (tradenamed *Hidrox*)¹⁴ can be processed in different ways to obtain nutraceutical or cosmetic formulations. This process has been in use in California now for more than a decade, resulting in manifold applications of the aqueous extract. Aiming at adapting the Crea’s protocol to the two-phase milling process, we focused on the use of TPOMW as a source of biophenols, as well as trying to recover the emulsified olive oil usually lost in the wet pomace. The technology was developed at a large olive mill based in Chiaramonte Gulfi, Sicily, a renown Italian production centre of high quality extra virgin olive oil (EVOO) during the 2016 olive milling season.

The simple processing stages for the recovery and concentration of olive biophenol are briefly described below. No fractionation such as partitioning with an organic solvent, chromatographic methods, supercritical fluids, or membranes are employed. Two different modified industrial centrifuges were instead placed downstream the EVOO production line. In the first step, the TPOMW was forced in a two-phase horizontal spin-dryer in order to separate the solid particles from the liquid

(oil+water) phase. The emulsified liquid phase thus underwent centrifugation in a vertical three-phase centrifuge, namely a 3-phase separating decanter in which two liquids of different densities are separated from each other, and from the residual solid particles ($d > 1 \text{ mm}$).



Figure 1. Water phase output of the three-phase vertical centrifugation.



Figure 2. Solid-cake resulting from the three-phase vertical centrifugation removed using a press filter.

The water phase output of the vertical centrifugation, containing approximately 1-2% of micrometer solid residue comprised of olive pulp, peel and wax (Figure 1), was clarified using an industrial press filter (Figure 2). Following the Crea's protocol, the resulting olive juice was stabilized with citric acid. In this olive juice biophenols are more concentrated than in the corresponding juice obtained from the three-phase OMWW since very little water is added during the two-phase milling process.

3. Enhanced EVOO Production

Along with the olive juice, the new process recovers from the wet pomace a significant yield of extra virgin olive oil, namely 90% of the oil not extracted during the conventional milling, without the need to use *n*-hexane. The EVOO obtained from this second round of extraction had significant economic value, 4-5 times higher than the value of hexane-extracted olive oil. It is important to notice, here, that the latter EVOO never comes in contact with citric acid, as the acid is only used in a subsequent step only to stabilize the olive juice and promote the hydrolysis of its glycoside components.

No oxidative or other degenerative phenomena of this EVOO fraction could take place, as the new extractor was placed directly downstream the mill's wet pomace output, with the newly recovered oil maintaining retaining excellent flavour with a greenish colour was, more eye-catching than first extraction oil, due to additional chlorophyll extraction from the TPOMW.

4. Additional Pomace

Another valued output of the new extraction process was a dried pomace with 60% less humidity compared to that of conventional wet pomace which could indeed be sold to a local factory which further commercializes the dried ‘nocciolino’ (seed) as solid biofuel to power domestic and commercial biomass heaters. No waste was discharged into the environment, whereas the milling company supplying the wet pomace not only got rid of previous significant costs associated to processing the TPOMW as industrial waste, but converted it in a triple source of extra revenues obtained from selling: *i*) the TPOMW used as phenolics source to the extraction company; *ii*) high quality second-extraction EVOO; and *iii*) the dried pomace to a ‘nocciolino’ company.

6. High Quality Biophenol Extracts

Three lots of stabilized biophenol extracts obtained according to the harvesting and milling time (late September-early October; late October; early November) were stored in Sicily in stainless steel silos (Figure 3). Every lot is composed of olive juice coming from different organic Sicilian *Cultivar* (mainly tonda Iblea and Nocellara del Belice but also Bioancollilla, Cerasuola, Moresca and Nocellara Etnea). Regardless of the poor season (year-to-year decrease in production of olives approaching 70%) about 800 tonnes of olive juice could be obtained.



Figure 3. Silos stocking the olive juice obtained in Sicily in 2016 from a two-phase olive mill.

The juice from different lots was left to undergo hydrolysis according to Crea's original protocol for several months at buffered pH. The HPLC analyses aimed to assess the total biophenol, hydroxytyrosol (HT) and tyrosol (T; 4-hydroxyphenethyl alcohol) concentration were carried out in Italy by an accredited laboratory (Centro Analisi Biochimiche, Rizziconi) further registered for olive oil analysis by Italy's Agriculture Ministry.

All juice samples contain a large amount of total phenols, given in ppm (mg kg^{-1}) in Table 1, and in the following text and Tables. The first lot, coming from the early harvested drupes, contained the highest biophenol amount, in agreement with the known higher phenolics content in greener olive drupes.

Table 1. Total phenolics, tyrosol and hydroxytyrosol in in three lots of olive juice as obtained from early to late harvesting^a

Entry	Total biophenol (ppm)	Hydroxytyrosol (ppm)	Tyrosol (ppm)	Hydroxytyrosol (wt%)	Tyrosol (wt%)
Lot 1	7288	545.1	79	7.48	1.42
Lot 2	5318	444	68.8	8.40	1.41
Lot 3	5810	355.3	53.7	6.11	0.98

^aHPLC analytical method 0178-00 with syringic acid as internal standard; data from the accredited Laboratory Centro Analisi Biochimiche (Rizziconi, Italy).

Data in Table 2 show that after one month of aging, the pH buffered at 3.2 and the activity of the olives yeast,¹⁵ promoted the hydrolysis of the biophenol glycosides leading to increasing concentration of both HT and T.

Table 2. Total phenolics, tyrosol and hydroxytyrosol in in three lots of olive juice obtained from early to late harvesting, after one month^a

Entry	Total biophenol (ppm)	Hydroxytyrosol (ppm)	Tyrosol (ppm)	Hydroxytyrosol (wt%)	Tyrosol (wt%)
Lot 1	5542	448	102.9	8.08	1.4
Lot 2	5144	486	74.7	9.45	1.4
Lot 3	4623	412.6	57	8.93	1.2

^aHPLC analytical method 0178-00 with syringic acid as internal standard; data from the accredited Laboratory Centro Analisi Biochimiche (Rizziconi, Italy).

Remarkably, the trend observed in Sicily with olive juice originating from the two-phase mill was considerably slower when compared with the hydrolytic profile observed in California for juice coming from a three-phase mill, likely due to the higher phenol concentration, as well as to the initial lower outdoor temperature during December 2017 and January 2017 in the area hosting the silos (Caltagirone, 610 m above sea level), both responsible for partial inhibition of the *Saccharomyces* yeast

strain. This hypothesis was confirmed by the data obtained after four months aging (Table 3).

Table 3. Total phenolics, tyrosol and hydroxytyrosol in three lots of olive juice obtained from early to late harvesting, after 4 months^a

Entry	Total biophenol (ppm)	Hydroxytyrosol (ppm)	Tyrosol (ppm)	Hydroxytyrosol (wt%)	Tyrosol (wt%)
Lot 1	5045	743	126	14.73	2.5
Lot 2	3530	1004	127	28.44	3.6
Lot 3	1387	711	79	51.26	5.7

^aHPLC analytical method 0178-00 with syringic acid as internal standard; data from the accredited Laboratory Centro Analisi Biochimiche (Rizziconi, Italy).

The considerably higher temperatures recorded in February and March 2017 (+2.69 °C for February and +2.52°C for March, in Italy, in comparison to the 1971-2000 mean)¹⁶ promoted the glycoside hydrolysis to such an extent that in the aqueous extract from Lot 2, hydroxytyrosol represented 51.26 wt. % of the olive polyphenol composition, namely in the very high concentration range (40-90 wt. %) required for advanced medical applications such as the treatment of early-stage Parkinson disease.¹⁷ In 2001, Crea was the first scholar to associate a powerful anti-inflammatory activity to hydroxytyrosol.¹⁸


Table 4. Total phenolics, tyrosol and hydroxytyrosol in three lots of olive juice obtained from early to late harvesting, after 9 months^a

Entry	Total biophenol (ppm)	Hydroxytyrosol (ppm)	Tyrosol (ppm)	Hydroxytyrosol (wt%)	Tyrosol (wt%)
Lot 1	2766	1491	252	53.90	9.11
Lot 2	3690	1531	221	41.49	5.98
Lot 3	2107	1153	190	51.77	8.53


^aHPLC analytical method 0178-00 with syringic acid as internal standard; data from the accredited Laboratory Centro Analisi Biochimiche (Rizziconi, Italy).

Finally, data in Table 4 show that after nine months of aging with the temperature further raising due to Sicily's hot climate, the hydrolysis of the biophenol glycosides led to record concentration of both HT and T: 53.9% for hydroxytyrosol in the first lot and, in general, concentration approaching or even exceeding 1500 ppm which is, to the best


of our knowledge, the highest concentration of hydroxytyrosol in an olive biophenol extract ever reported.



Silos 1



Silos 2



Silos 3

Figure 4. Evolution of hydroxytyrosol and polyphenol total concentration in the silos of Figure 3.

7. Conclusions

We have adapted to the two-phase olive mill the integral biophenol extraction protocol originally developed by Crea for the three-phase olive mill in 2001. The extraction, now, is from the wet pomace (“alperujo” in Spanish), and not from the vegetation waters. The new set-up allowed extensive recovery of olive phenolics, transforming previous toxic waste into a source of revenues for the milling company, and of valued bioproducts for its bioeconomy partner.

The olives were specific cultivars grown in southern Sicily where the unique climate and geographical conditions afford particularly high amounts of biophenols.¹⁹ This translates into significant higher amounts of total biophenols in the aqueous extract, eventually affording an extract with hydroxytyrosol concentration exceeding 1500 ppm and 53.0 wt.% of the olive polyphenol composition.

Natural extracts with high concentration of hydroxytyrosol, the oleuropein metabolite, possessing potent pharmacological activities *in vivo*, are now required for human intervention studies² that could lead to the first biophenol-based drugs, ending what Quideau has called the century-long neglect of plant polyphenols in the development of new drugs.²⁰

In brief, the integral extraction process originally developed in 2001 for the three-phase olive mill, has kept its promises to make the olive bioeconomy possible at affordable costs. The olive mill company transformed waste formerly processed at high cost in a triple source of revenues (waste disposal costs eliminated, extra EVOO, extra dried pomace); the bioeconomy company obtained a biophenol-rich extract ready for multiple health beneficial applications; and the environment benefited from not receiving pollutants. On November 18 2016, Italy’s Labour Minister and the State Secretary for Research visited the new Sicily-based biophenol extraction plant,²¹ as a prominent example of the circular bioeconomy now actively supported in Europe as a

tool to promote economic growth and to address environmental and societal challenges.

22

Conflict of interest disclosure

Renovo Biondustry, a company, has an interest in the biophenol market. M.P., F.M. and R.C., affiliate with Italy's Research Council, have no competing financial interest.

List of abbreviations

EVOO = Extra Virgin Olive Oil

HT = Hydroxytyrosol

T = Tyrosol

OMWW = Olive Mill Waste Water

TPOMW = Two-Phase Olive Mill Waste

COD = Chemical Oxygen Demand

BOD₅ = Biological Oxygen Demand

Acknowledgements

This article is dedicated to Professor Roberto Crea, prominent biochemist, for his pioneering work on olive biophenol biochemistry and extraction methodologies. R.D. is grateful to Salvatore Cutrera, Giacomo Ardagna and their wives at Frantoi Cutrera (Chiamonte Gulfi), for they have done during his stay in the 2016 milling season. Thanks are due to Mr Cesare Pinotti and Dr Alfredo Tamburino for valued collaboration.

References

1. C. M. Galanakis, K. Kotsiou, Recovery of Bioactive Compounds from Olive Mill Waste In *Olive Mill Waste: Recent Advances for Sustainable Management*, C. M. Galanakis (Ed.), Academic Press, Amsterdam: 2017; pp. 205-230.
2. L. Parkinson, S. Cicerale, The Health Benefiting Mechanisms of Virgin Olive Oil Phenolic Compounds, *Molecules* 21 (2016) 1734.
3. R. Valls, M. Farràs, M. Suárez, S. Fernández-Castillejo, M. Fitó, V. Konstantinidou, F. Fuentes, J. López-Miranda, M. Giralt, M. Covas, M. Motilva, R. Solà, Effects of functional olive oil enriched with its own phenolic compounds on endothelial function in hypertensive patients. A randomised controlled trial, *Food Chem.* 167 (2015), 30-35.
4. E. Tsagaraki, H. N. Lazarides, K. B. Petrotos, Olive Mill Wastewater Treatment In *Utilization of By-Products and Treatment of Waste in the Food Industry*, V. Oreopoulou, W. Russ (Ed.s) Springer, Boston: 2007; 133-157.
5. G. Golia, I danni ambientali dell'olio di oliva, mediaset.it, 9 November 2015. See at the URL: <https://goo.gl/d9Oh9I>
6. A. Roig, M. L. Cayuela, M. A. Sánchez-Monedero, An overview on olive mill wastes and their valorisation methods, *Waste Manag.* 26 (2006) 960-969.
7. A. Fernández-Hernández, A. Roig, N. Serramiá, C. G. Civantos, M. A. Sánchez-Monedero, Application of compost of two-phase olive mill waste on olive grove: effects on soil, olive fruit and olive oil quality, *Waste Manag.* 34 (2014) 1139-1147.
8. R. Ciriminna, A. Fidalgo, F. Meneguzzo, L. M. Ilharco, M. Pagliaro, Extraction, Benefits and Valorization of Olive Polyphenols, *Eur. J. Lipid Sci. Technol.* 118 (2016) 503–511.
9. S. Rigacci, M. Stefani, Nutraceutical Properties of Olive Oil Polyphenols. An Itinerary from Cultured Cells through Animal Models to Humans, *Int. J. Mol. Sci.* 17 (2016) 843.
10. R. Ciriminna, F. Meneguzzo, R. Delisi, M. Pagliaro, Olive Biophenols as New Antioxidant Additives in Food and Beverage, *ChemistrySelect* 2 (2017) 1360-1365.
11. M. Lueder, The Active Ingredients Mixture of Olives Provides Skin Whitening and Age Spot Reduction, *Cosmetic Science Technology* (2011) 1-6.
12. F. Casamenti, M. Stefani, Olive polyphenols: new promising agents to combat aging-associated neurodegeneration, *Expert Rev. Neurother.* 17 (2017) 345-358.
13. R. Crea, Method of obtaining a hydroxytyrosol-rich composition from vegetation water, US20020058078 (2001).
14. A tradename of CreAgri, Inc. (Hayward, CA).
15. G. Ciafardini, B. A. Zullo, *Int. J. Food Microbiol.* 275 (2002), 111-118]
16. Data from Italy's Research Council ISAC Institute. See at the URL: www.isac.cnr.it/climstor/climate_news.html (last time accessed, July 12, 2017).
17. R. Crea, Treatment of early stage Parkinson's disease with a hydroxytyrosol-containing polyphenol formulation, US20160374959A1 (2016).
18. R. Crea, Method for treatment of inflammation, US8216599B2 (2002).
19. R. Delisi, M. Pagliaro, F. Saiano, R. Ciriminna, C18 Alkyl-Modified Silica: A Suitable Tool for Olive Biophenol Green Extraction, *Chemical Data Collections* 7-8 (2017) 102-106.
20. S. Quideau, D. Deffieux, C. Douat-Casassus, L. Pouységu, *Angew. Chem. Int. Ed.* 50 (2011) 586-621.
21. M. Guccione, Da scarto a risorsa, il sole regala un tesoro dalle acque dei frantoi, *La Sicilia*, 18 November 2016.
22. R. Peters, Societal Challenges and Bioeconomy in Europe, *Bioeconomy Impact 2016*, Dublin, 10-11 February 2016.

Olive Biophenol Integral Extraction at a Two-Phase Olive Mill



**Highly valued
biophenol extract**

**Waste disposal costs
eliminated**

**Additional highly
valued EVOO**

**Additional valued
dried pomace**