



A determining form for the damped driven nonlinear Schrödinger equation—Fourier modes case

Michael S. Jolly^{a,*}, Tural Sadigov^a, Edriss S. Titi^{b,c}

^a Department of Mathematics, Indiana University, Bloomington, Bloomington, IN 47405, USA

^b The Department of Computer Science and Applied Mathematics, The Weizmann Institute of Science, Rehovot 76100, Israel

^c Department of Mathematics, Texas A&M University, 3368 TAMU, College Station, TX 77843-3368, USA

Received 10 June 2014; revised 17 December 2014

Available online 19 January 2015

Abstract

In this paper we show that the global attractor of the 1D damped, driven, nonlinear Schrödinger equation (NLS) is embedded in the long-time dynamics of a determining form. The determining form is an ordinary differential equation in a space of trajectories $X = C_b^1(\mathbb{R}, P_m H^2)$ where P_m is the L^2 -projector onto the span of the first m Fourier modes. There is a one-to-one identification with the trajectories in the global attractor of the NLS and the steady states of the determining form. We also give an improved estimate for the number of the determining modes.

© 2014 Elsevier Inc. All rights reserved.

MSC: 35Q55; 34G20; 37L05; 37L25

Keywords: Nonlinear Schrödinger equation; Determining forms; Determining modes; Determining nodes; Inertial manifolds

* Corresponding author.

E-mail addresses: msjolly@indiana.edu (M.S. Jolly), tsadigov@indiana.edu (T. Sadigov), ctiti@math.uci.edu, edriss.titi@weizmann.ac.il (E.S. Titi).

1. Introduction

The damped, driven, nonlinear Schrödinger equation (NLS), (2.1), has been derived in various areas of physics, and widely investigated (see e.g. [4] and references therein). In plasma physics, the NLS is a model for the propagation of an intense laser beam through a nonlinear medium (see e.g. [7]). In this model the unknown function $u(x, t)$ is the electrical field amplitude, t is the distance in the direction of the propagation and x is a transverse spatial variable. Absorption of the electromagnetic wave by the medium is accounted for by linear damping. Some resonant forcing of small amplitude (for example, a traveling wave) is used to compensate weak dissipative losses (i.e., absorption). The NLS also describes the single particle properties of Bose–Einstein condensate (BEC) (see e.g. [3]), in which a gas of bosons is cooled to very low temperatures. In this case, known as the Gross–Pitaevski equation, $u(x, t)$ describes the macroscopic wave function of the condensate; t is time and x is a spatial variable. A constant damping rate (absorption) γ describes inelastic collisions with the background gas which occur when the particle density is very large. The forcing term represents the interatomic forces of the condensate. We note that the NLS is also investigated in deep-water phenomena and in the collapse of Langmuir waves (see e.g. [7]). In this paper, we consider a force general enough to include the above applications where it can be periodic in space and independent of time.

The undamped, unforced case has been extensively studied in modern mathematical physics (see e.g. [5]). Well-posedness of (2.1), for nonzero forcing and $\gamma > 0$ is established by Ghidaglia in [12], where, under the assumption that the force is either time independent or time periodic, it is also proved that there exists a weak attractor in the Sobolev spaces H^1 and H^2 . Later, it is proved in [18] that this weak attractor is in fact a global attractor in H^2 in the strong sense. In [13], assuming the force is smooth enough and periodic in spatial variable, Goubet proved that the global attractor \mathcal{A} is smooth, meaning it is included and bounded in H^k , for any $k \geq 1$. This implies that \mathcal{A} is in C^∞ due to classical Sobolev embeddings theorems. Finally in [16], it is proved that \mathcal{A} is in fact contained in a subclass of the space of real analytical functions provided that the forcing term is real analytic. The long-time dynamics of the damped, driven NLS is entirely contained in the *global attractor* \mathcal{A} , a compact finite-dimensional set within the infinite-dimensional phase space H^k for any $k \geq 1$ (see [13]). It is shown in [16], for real analytical forcing, that the solutions on the attractor of the NLS are determined uniquely by their nodal values on only two sufficiently close nodes.

The finite dimensionality for the NLS can be stated more explicitly. It is also known that solutions of the NLS in \mathcal{A} are determined by the asymptotic behavior of a sufficient finite number of Fourier modes (see [13,14]). To be precise, this means that if two complete trajectories in the global attractor coincide under the projection P_m onto a sufficiently large number, m , of low modes, then they are the same trajectory. These m -modes are called *determining modes* (see [11]). This notion of determining modes was used in [8] to find a *determining form* for the 2D Navier–Stokes equations (NSE). In [8], the determining form is an ordinary differential equation in an *infinite dimensional* Banach space $X = C_b(\mathbb{R}, P_m H)$, governing the evolution of trajectories. Here H is a Hilbert space which is a natural phase space for the 2D NSE (see [6,17]). The trajectories in the attractor of the 2D NSE are identified with traveling wave solutions of the determining form in [8].

A determining form of a different sort was found in [9] for the 2D NSE. It is based on *data assimilation by feedback control* through a general interpolant operator. It is general in the sense that it can be induced by a variety of determining parameters such as determining modes, nodal values and finite volumes (see [10], [15] and references therein). The steady states of this deter-

mining form are precisely the trajectories in the global attractor of the 2D NSE. Thus, in general, a determining form for a dissipative system can be defined as an ODE in a phase space of trajectories which characterizes those in the global attractor. The second type of determining form may ultimately prove useful in data assimilation; if a time series for the projection of a solution is corrupted by noise, then the true solution could be recovered by evolving the form toward a steady state.

Motivation for the determining form comes from the notion of an inertial form. An inertial form for a partial differential equation is an ordinary differential equation restricted to a *finite* dimensional manifold called an *inertial manifold*. We note that it is not known if there is an inertial manifold for 2D NSE. Nor is it known whether there is such a manifold for the NLS. In this paper we adapt the approach in [9] for the NLS. While the feedback control approach potentially allows for a variety of interpolant operators, our analysis for the NLS is restricted to the case of Fourier modes. This is done in order to close the *a priori* estimates needed in L^2 , H^1 , H^2 , even though there is no dissipative term to absorb the highest derivative. The key step to get a determining form is defining and extending the map W which recovers the high frequency components of a trajectory on the global attractor from the low frequency components. This is done by adding a feedback control term to the NLS (see [1,2] for feedback controls). The determining form in [8] has the map W inserted in the bilinear term of the NSE. The feedback control approach allows us to avoid doing this for the cubic nonlinear term in the NLS. The idea of the feedback control approach is that if we know the P_m projection of the solution of the damped driven NLS on the attractor, we can feed this information into the system to construct the complete solution. It is worth pointing out that this equation is dispersive and merely damped, not strongly dissipative. The analysis used here involves compound functionals motivated by the Hamiltonian structure of the Schrödinger equation.

Section 2 introduces the NLS and special notation. The statements of the main results are mentioned in Section 3. A priori estimates are done in Section 4. Section 5 contains the main results that we need to obtain a determining form. Section 6 introduces the determining form. Finally, in Section 7, we give a different proof of the determining modes property of the NLS through a ‘reverse’ Poincaré type inequality. This approach produces an improved estimate for the number of the determining modes for the NLS.

2. Preliminaries

We consider the 1D damped, driven, nonlinear Schrödinger equation subject to periodic boundary conditions

$$\begin{aligned} iu_t + u_{xx} + |u|^2 u + i\gamma u &= f, \\ u(t, x) &= u(t, x + L), \quad \forall (t, x) \in \mathbb{R} \times \mathbb{R}, \\ u(0, x) &= u_0(x), \end{aligned} \tag{2.1}$$

where $0 < L < \infty$, $0 < \gamma$ and $f \neq 0$. We assume that f is time independent, and $f \in L^2_{per}$. Let $0 \leq k < \infty$. We denote by $H^k[0, L]$ (or simply H^k) the Sobolev space of order k ,

$$H^k[0, L] := \{f \in L^2[0, L]: \alpha \leq k, D^\alpha f \text{ exists and } D^\alpha f \in L^2[0, L]\},$$

and by H_{per}^k , the subspace of H^k consisting of functions which is periodic in x , with period L . Note that $H_{per}^0[0, L] = L_{per}^2[0, L]$. We assume that $u_0(x) \in H_{per}^2$. It has been proven in [12] that (2.1) has a unique solution $u(x, t)$ such that the mapping

$$u_0 \rightarrow u(t)$$

is continuous on H^1 , with $u \in L^\infty(\mathbb{R}; H^1)$. The global attractor is the maximal compact invariant set under the solution operator $S(t, \cdot)$. Alternatively, it can be defined as $\mathcal{A} = \bigcap_{u_0 \in \mathcal{B}} S(t, u_0)$ where \mathcal{B} is an absorbing ball (see e.g. [17]). Throughout the paper, we will use the notation

$$\begin{aligned}\|u\|^2 &:= \|u\|_{L^2}^2, \\ \|u\|_{H^1}^2 &:= \|u\|_{L^2}^2 + \|u_x\|_{L^2}^2,\end{aligned}$$

and

$$\|u\|_{H^2}^2 := \|u\|_{L^2}^2 + \|u_{xx}\|_{L^2}^2.$$

To make the flow of the analysis more transparent, we adopt some specialized notation for certain bounding expressions. Bounding expressions that depend on γ , f (and μ , see (3.4)) will be denoted by capital letters \mathcal{R} and K with specific indices. The bounding expressions \mathcal{R} with indices 0, 1 and 2 are L^2 , H^1 and H^2 bounds, respectively, for the solution of (3.4). Those bounding expressions accented with $\tilde{\cdot}$ and $\tilde{\cdot}$ will be subsequently improved. As they are improved once, we remove a $\tilde{\cdot}$. For example, \tilde{K}_2 will be improved once, and we use \tilde{K}_2 for the improvement. Then we improve \tilde{K}_2 again to get K_2 which is the final improvement. Universal constants will be denoted by c and updated throughout the paper. We denote by P_m the L^2 -projection onto the space H_m , where

$$H_m := \text{span}\{e^{ikx \frac{2\pi}{L}} : |k| \leq m\}. \quad (2.2)$$

3. The statements of the main results

We define the following norms:

$$\begin{aligned}|v|_X &= \sup_{s \in \mathbb{R}} \|v(s)\| + \sup_{s \in \mathbb{R}} \|v_s(s)\|, \\ |v|_{X,0} &= \sup_{s \in \mathbb{R}} \|v(s)\|, \\ |w|_Y &= \sup_{s \in \mathbb{R}} \|w(s)\|_{H^2},\end{aligned} \quad (3.1)$$

and the following Banach spaces:

$$X = C_b^1(\mathbb{R}, P_m H^2) = \{v : \mathbb{R} \rightarrow P_m H^2 : v(s) \text{ is continuous } \forall s \in \mathbb{R} \text{ and } |v|_X < \infty\}, \quad (3.2)$$

$$Y = C_b(\mathbb{R}, H^2) = \{w : \mathbb{R} \rightarrow H^2 : w(s) \text{ is continuous } \forall s \in \mathbb{R} \text{ and } |w|_Y < \infty\}. \quad (3.3)$$

Let $v \in X$, and consider the equation

$$i w_s + w_{xx} + |w|^2 w + i \gamma w = f - i \mu [P_m(w) - v], \quad (3.4)$$

subject to periodic boundary condition

$$w(s, x) = w(s, x + L), \quad \forall (s, x) \in \mathbb{R} \times \mathbb{R}.$$

We assume that $f \in L^2_{per}$. We first state a new estimate for the number of determining modes.

Theorem 3.1. *Assume*

$$m \geq \frac{L}{2\pi} K_{11} - 1,$$

where K_{11} is defined in (7.1). Then the Fourier projection P_m of L^2 onto the space H_m , where H_m is defined in (2.2), is determining for (2.1) i.e., for all $u_1(\cdot), u_2(\cdot) \subset \mathcal{A}$, $P_m u_1(t) = P_m u_2(t)$, for all $t \in \mathbb{R}$ implies that $u_1(t) = u_2(t)$, for all $t \in \mathbb{R}$.

Remark 3.2. By tracking the $\|f\|$ dependence of the bounds throughout the paper, we will show that a sufficient number of determining modes is of order $O(\|f\|^{10})$ as $\|f\| \rightarrow \infty$ and $O(\gamma^{-12})$ as $\gamma \rightarrow 0$. Following the analysis in Goubet [13], one can show that a sufficient number of determining modes is of order $O(\gamma^{-12.5})$ as $\gamma \rightarrow 0$ and $O(\|f\|^{12})$ as $\|f\| \rightarrow \infty$. Thus the functionals in our analysis in (4.8), (4.16), (5.6) and (5.16), which are naturally motivated by the Hamiltonian structure of the Schrödinger equation, lead to sharper explicit estimates. We also mention that the abstract treatment of determining modes by Hale and Raugel is applied to the damped, driven, nonlinear Schrödinger equation in [14], but that approach does not provide estimates for the number of modes needed.

The proof of Theorem 3.1 is given in Section 7. It is a byproduct of the proof for the following main result.

Theorem 3.3. *Let $v \in X$, and u^* be a steady state of Eq. (2.1). Then we have the following:*

- (1) *There exists a unique bounded solution $w \in Y$ of (3.4), which defines a map $W : X \rightarrow Y$, such that $w = W(v)$.*
- (2) *For sufficiently large m and μ , we have $W(P_m u) = u$, for any trajectory $u(s)$, $s \in \mathbb{R}$, in the global attractor of (2.1).*
- (3) *For sufficiently large m and μ , $P_m W : X \rightarrow X$ is a locally Lipschitz map.*
- (4) *The determining form*

$$\frac{dv}{dt} = F(v) = -|v - P_m W(v)|_{X,0}^2 (v - P_m u^*)$$

is an ordinary differential equation in a forward invariant set

$$\{v \in X : |v - P_m u^*|_X < 3(\mathcal{R}_0^0 + \mathcal{R}'^0)\},$$

and F restricted to that set is globally Lipschitz. Moreover, $P_m u(s)$ is included in that set, for every $u(s) \in \mathcal{A}$. Here $\mathcal{R}_0^0 = \mathcal{R}_0|_{\mu=0}$ and $\mathcal{R}'^0 = \mathcal{R}'|_{\mu=0}$ are defined in (5.8) and (4.26), respectively.

Theorem 3.3 is a combination of results to follow. Item (1) corresponds to [Proposition 4.1](#) and the first part of [Theorem 5.3](#), item (2) is equivalent to [Proposition 5.1](#), item (3) is analogous to the second part of [Theorem 5.3](#), and finally item (4) is the summary of [Theorem 6.1](#). The basic idea is to use the Galerkin method to establish a unique bounded solution to (3.4), which defines the map W . This involves a series of a priori estimates undertaken in the next section.

Let $n > m$. Note that $P_n v = v$, for every $v \in X$, and consider a Galerkin approximation of (3.4),

$$i \partial_s w_n + \partial_x^2 w_n + P_n(|w_n|^2 w_n) + i \gamma w_n = P_n f - i \mu (P_m w - v), \quad (3.5)$$

subject to periodic boundary condition, and with the initial data

$$w_n(-k, x) = 0, \quad (3.6)$$

where $w_n \in H_n$, for some $k \in \mathbb{N}$. For simplicity we will drop the subscript n . Since (3.5) is an ordinary differential equation with locally Lipschitz nonlinearity, it has a unique, bounded solution w_n on a small interval $[-k, S^*)$, for some $S^* > -k$. We will show that w_n exists globally on the interval $[-k, \infty)$ and is uniformly bounded with respect to $s \in [-k, \infty)$, n and k , in the norms of the spaces L^2 , H^1 and H^2 .

4. A priori estimates

4.1. L^2 bound

Let $[-k, S^*)$ be the maximal interval of existence for (3.5). We will establish here global (in time) uniform in n bounds which will imply, among other things, that $S^* = \infty$. Let us focus below on the interval $[-k, S^*)$. Multiply (3.5) by \bar{w} , and integrate

$$i \int_0^L w_s \bar{w} + \int_0^L w_{xx} \bar{w} + \int_0^L |w|^4 + i \int_0^L \gamma |w|^2 + i \mu \int_0^L P_m(w) \bar{w} = \int_0^L f \bar{w} + i \mu \int_0^L v \bar{w}. \quad (4.1)$$

Take the imaginary parts of both sides, and use the fact that P_m is an orthogonal projection and $v \in X$ to get

$$\frac{1}{2} \frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 + \mu \|P_m(w)\|^2 = \operatorname{Im} \int_0^L f \bar{w} + \mu \operatorname{Re} \int_0^L v P_m w \bar{w}.$$

By using Hölder's and Young's inequalities, we have

$$\begin{aligned} \frac{1}{2} \frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 + \mu \|P_m w\|^2 &\leq \|f\| \|w\| + \mu \|v\| \|P_m w\| \\ &\leq \frac{\|f\|^2}{2\gamma} + \frac{\gamma \|w\|^2}{2} + \frac{\mu \|v\|^2}{2} + \frac{\mu \|P_m w\|^2}{2}, \end{aligned}$$

and hence,

$$\frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 + \mu \|P_m w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu \|v\|^2,$$

for all $s \in [-k, S^*)$. Since $v \in X$, we get

$$\frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 + \mu \|P_m w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu |v|_X^2. \quad (4.2)$$

Thus,

$$\frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu |v|_X^2.$$

Since, $w(-k, x) = 0$, we deduce by Gronwall's lemma that,

$$\|w(s)\|^2 \leq \frac{\|f\|^2}{\gamma^2} + \frac{\mu}{\gamma} |v|_X^2,$$

for all $s \in [-k, S^*)$. Since the right-hand side is constant, we conclude that $S^* = \infty$, and therefore

$$\|w(s)\| \leq \frac{\|f\|}{\gamma} + \frac{\mu^{\frac{1}{2}}}{\gamma^{\frac{1}{2}}} |v|_X =: \tilde{\mathcal{R}}_0,$$

for all $s \in [-k, \infty)$, and as a result

$$\sup_{s \geq -k} \|w(s)\| \leq \tilde{\mathcal{R}}_0. \quad (4.3)$$

Note that the constant $\tilde{\mathcal{R}}_0$ satisfies $\tilde{\mathcal{R}}_0 = O(\mu^{\frac{1}{2}})$ as $\mu \rightarrow \infty$, and is independent of k and n .

4.2. H^1 bound

Use again the fact that P_m is an orthogonal projection in (4.1), and take the real parts of Eq. (4.1):

$$\operatorname{Im} \int_0^L w \bar{w}_s = \|w_x\|^2 - \|w\|_{L^4}^4 + \operatorname{Re} \int_0^L f \bar{w} - \mu \operatorname{Im} \int_0^L v \bar{w}, \quad (4.4)$$

for all $s \in [-k, \infty)$. Now, multiply (3.5) by \bar{w}_s and integrate with respect to x over $[0, L]$ to obtain

$$i \|w_s\|^2 - \int_0^L w_x \bar{w}_{xs} + \int_0^L |w|^2 w \bar{w}_s + i \gamma \int_0^L w \bar{w}_s + i \mu \int_0^L P_m(w) \bar{w}_s = \int_0^L f \bar{w}_s + i \mu \int_0^L v \bar{w}_s.$$

Take the real part of the above equation to obtain

$$\begin{aligned} \frac{d}{ds} \|w_x\|^2 - \frac{1}{2} \frac{d}{ds} \|w\|_{L^4}^4 + 2\gamma \operatorname{Im} \int_0^L w \bar{w}_s + 2\mu \operatorname{Im} \int_0^L P_m w P_m \bar{w}_s \\ = -2 \operatorname{Re} \int_0^L f \bar{w}_s + 2\mu \operatorname{Im} \int_0^L v \bar{w}_s. \end{aligned} \quad (4.5)$$

To eliminate the third term in (4.5), we will use (4.4). For the fourth term, we take the P_m projection of Eq. (3.4), then multiply by $P_m \bar{w}$, integrate, and take the real parts to find

$$\begin{aligned} \operatorname{Im} \int_0^L P_m w P_m \bar{w}_s - \|P_m w_x\|^2 + \operatorname{Re} \int_0^L P_m (|w|^2 w) P_m \bar{w} \\ = \operatorname{Re} \int_0^L P_m f P_m \bar{w} - \mu \operatorname{Im} \int_0^L v P_m \bar{w}. \end{aligned} \quad (4.6)$$

Now combine as follows: $(-2\gamma) \times (4.4) + (-2\mu) \times (4.6) + (4.5)$ to get

$$\begin{aligned} \frac{d\phi}{ds} + 4\gamma\phi = 2\gamma \|w_x\|^2 + 6\gamma \operatorname{Re} \int_0^L f \bar{w} - 6\mu\gamma \operatorname{Im} \int_0^L v \bar{w} \\ + 2\mu \operatorname{Re} \int_0^L P_m (|w|^2 w) P_m \bar{w} - 2\mu \|P_m w_x\|^2 \\ - 2\mu \operatorname{Re} \int_0^L P_m f P_m \bar{w} + 2\mu^2 \operatorname{Im} \int_0^L v P_m \bar{w} \\ - 2\mu \operatorname{Im} \int_0^L v_s \bar{w}, \end{aligned} \quad (4.7)$$

where

$$\phi(s) = \|w_x\|^2 - \frac{1}{2} \|w\|_{L^4}^4 + 2 \operatorname{Re} \int_0^L f \bar{w} - 2\mu \operatorname{Im} \int_0^L v \bar{w}. \quad (4.8)$$

Since $w(-k, x) = 0$ for all x , we have that $w_x(-k, x) = 0$ for all x , and thus $\phi(-k) = 0$.

We estimate the right-hand side of (4.7) using Hölder's, Young's and Agmon's inequalities and (4.3) as follows

$$\begin{aligned}
2\gamma \|w_x\|^2 &\leq 2\gamma \|w\|_{H^1}^2, \\
6\gamma \operatorname{Re} \int_0^L f \bar{w} &\leq 6\gamma \|f\| \|w\| \leq 6\gamma \|f\| \tilde{\mathcal{R}}_0, \\
-6\mu\gamma \operatorname{Im} \int_0^L v \bar{w} &\leq 6\mu\gamma \|v\| \|w\| \leq 6\mu\gamma |v|_X \tilde{\mathcal{R}}_0, \\
2\mu \operatorname{Re} \int_0^L P_m(|w|^2 w) P_m \bar{w} &= 2\mu \operatorname{Re} \int_0^L |w|^2 w P_m \bar{w} \\
&\leq 2\mu \|w\|_\infty^2 \|w\| \|P_m w\| \\
&\leq 2\mu (c \|w\| \|w\|_{H^1}) \|w\| \|P_m w\| \\
&\leq 2\mu c \|w\|^3 \|w\|_{H^1} \\
&\leq \frac{\mu^2 c^2 \|w\|^6}{\gamma} + \gamma \|w\|_{H^1}^2 \\
&\leq \frac{\mu^2 c^2 \tilde{\mathcal{R}}_0^6}{\gamma} + \gamma \|w\|_{H^1}^2.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
-2\mu \operatorname{Re} \int_0^L P_m f P_m \bar{w} &\leq 2\mu \|P_m f\| \|P_m w\| \leq 2\mu \|f\| \|w\| \leq 2\mu \|f\| \tilde{\mathcal{R}}_0, \\
2\mu^2 \operatorname{Im} \int_0^L v P_m \bar{w} &\leq 2\mu^2 \|v\| \|P_m w\| \leq 2\mu^2 |v|_X \tilde{\mathcal{R}}_0, \\
-2\mu \operatorname{Im} \int_0^L v_s \bar{w} &\leq 2\mu \|v_s\| \|w\| \leq 2\mu |v|_X \tilde{\mathcal{R}}_0.
\end{aligned}$$

Putting together the above estimates, we obtain

$$\frac{d\phi}{ds} + 4\gamma\phi \leq 3\gamma \|w\|_{H^1}^2 + \tilde{K}_1, \quad (4.9)$$

where

$$\tilde{K}_1 := 6\gamma \|f\| \tilde{\mathcal{R}}_0 + 6\mu\gamma |v|_X \tilde{\mathcal{R}}_0 + \frac{\mu^2 c^2 \tilde{\mathcal{R}}_0^6}{\gamma} + 2\mu \|f\| \tilde{\mathcal{R}}_0 + 2\mu^2 |v|_X \tilde{\mathcal{R}}_0 + 2\mu |v|_X \tilde{\mathcal{R}}_0.$$

Then by using Agmon's, Hölder's, and Young's inequalities in (4.8), we get

$$\begin{aligned}
 \phi(s) &\geq \|w_x\|^2 - \frac{1}{2}c\|w\|^3\|w\|_{H^1} - 2\|f\|\|w\| - 2\mu\|v\|\|w\| \\
 &= \|w_x\|^2 - \frac{1}{2}c\|w\|^3\|w_x\| - \frac{1}{2}c\|w\|^4 - 2\|f\|\|w\| - 2\mu\|v\|\|w\| \\
 &\geq \|w_x\|^2 - \frac{c^2\|w\|^6}{16\xi} - \xi\|w_x\|^2 - \frac{1}{2}c\|w\|^4 - 2\|f\|\tilde{\mathcal{R}}_0 - 2\mu|v|_X\tilde{\mathcal{R}}_0 \\
 &\geq (1-\xi)\|w_x\|^2 + (1-\xi)\|w\|^2 \\
 &\quad - \left[\frac{c^2\tilde{\mathcal{R}}_0^6}{16\xi} + \frac{1}{2}c\tilde{\mathcal{R}}_0^4 + 2\|f\|\tilde{\mathcal{R}}_0 + 2\mu|v|_X\tilde{\mathcal{R}}_0 + (1-\xi)\tilde{\mathcal{R}}_0^2 \right], \tag{4.10}
 \end{aligned}$$

where $0 < \xi < 1$, ξ to be chosen later. Thus, we have

$$\phi \geq (1-\xi)\|w\|_{H^1}^2 - \tilde{K}_2,$$

where

$$\tilde{K}_2 := \frac{c^2\tilde{\mathcal{R}}_0^6}{16\xi} + \frac{1}{2}c\tilde{\mathcal{R}}_0^4 + 2\|f\|\tilde{\mathcal{R}}_0 + 2\mu|v|_X\tilde{\mathcal{R}}_0 + (1-\xi)\tilde{\mathcal{R}}_0^2,$$

and hence

$$\|w\|_{H^1}^2 \leq \frac{1}{1-\xi}\phi + \frac{\tilde{K}_2}{1-\xi}. \tag{4.11}$$

Use (4.11) in (4.9) to obtain

$$\frac{d\phi}{ds} + \frac{1-4\xi}{1-\xi}\gamma\phi \leq \tilde{K}_3,$$

where

$$\tilde{K}_3 := \frac{3\gamma\tilde{K}_2}{1-\xi} + \tilde{K}_1.$$

Choose $\xi = \frac{1}{7}$, so that

$$\frac{1-4\xi}{1-\xi} = \frac{1}{2}.$$

So we have

$$\frac{d\phi}{ds} + \frac{\gamma}{2}\phi \leq \tilde{K}_3.$$

Since $\phi(-k) = 0$, applying the Gronwall lemma, we have

$$\phi(w(s)) \leq \tilde{K}_4,$$

for all $s \geq -k$, where $\tilde{K}_4 := 2\tilde{K}_3/\gamma$. From (4.11), we obtain

$$\|w(s)\|_{H^1}^2 \leq \frac{7}{6}\phi + \frac{7}{6}\tilde{K}_2 \leq \frac{7}{6}(\tilde{K}_4 + \tilde{K}_2),$$

for all $s \geq -k$. Therefore, we have $\tilde{K}_1 = O(\mu^5)$, $\tilde{K}_2 = O(\mu^3)$, $\tilde{K}_3 = O(\mu^5)$, $\tilde{K}_4 = O(\mu^5)$ as $\mu \rightarrow \infty$. Thus

$$\sup_{s \geq -k} \|w(s)\|_{H^1} \leq \tilde{\mathcal{R}}_1 := \sqrt{\frac{7}{6}(\tilde{K}_4 + \tilde{K}_2)} = \sqrt{\frac{28}{3}\tilde{K}_2 + \frac{7}{3\gamma}\tilde{K}_1} = O(\mu^{\frac{5}{2}}), \quad (4.12)$$

as $\mu \rightarrow \infty$. Note that $\tilde{\mathcal{R}}_1$ is independent of k and n .

4.3. Improved L^2 bound

We now use the H^1 -bound in (4.12) to obtain a better L^2 -bound. We rewrite (4.2) as

$$\frac{d}{ds} \|w\|^2 + \gamma \|w\|^2 + \mu \|w\|^2 - \mu \|Q_m w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu |v|_X^2,$$

where $Q_m = I - P_m$. Thus,

$$\frac{d}{ds} \|w\|^2 + (\gamma + \mu) \|w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu |v|_X^2 + \mu \|Q_m w\|^2.$$

By the generalized Poincaré inequality we have

$$\|Q_m w\|^2 \leq \frac{L^2}{4\pi^2} \frac{1}{(m+1)^2} \|w_x\|^2 \leq \frac{L^2}{4\pi^2} \frac{\tilde{\mathcal{R}}_1^2}{(m+1)^2}.$$

If we choose m large enough such that

$$\frac{\tilde{\mathcal{R}}_1^2}{(m+1)^2} \frac{L^2}{4\pi^2} < 1, \quad (4.13)$$

then

$$\frac{d}{ds} \|w\|^2 + (\gamma + \mu) \|w\|^2 \leq \frac{\|f\|^2}{\gamma} + \mu |v|_X^2 + \mu.$$

Now, we apply the Gronwall lemma, using the fact that $\|w(-k)\| = 0$, to obtain

$$\|w(s)\|^2 \leq \frac{\|f\|^2}{\gamma(\gamma + \mu)} + \frac{\mu|v|_X^2}{\gamma + \mu} + \frac{\mu}{\gamma + \mu},$$

for every $s \geq -k$, and hence,

$$\sup_{s \geq -k} \|w(s)\|^2 \leq \frac{\|f\|^2}{\gamma(\gamma + \mu)} + \frac{\mu|v|_X^2}{\gamma + \mu} + \frac{\mu}{\gamma + \mu}.$$

As a result,

$$\sup_{s \geq -k} \|w(s)\| \leq \mathcal{R}_0,$$

where

$$\mathcal{R}_0 := \frac{\|f\|}{\sqrt{\gamma(\gamma + \mu)}} + \sqrt{\frac{\mu}{\gamma + \mu}}|v|_X + \sqrt{\frac{\mu}{\gamma + \mu}} = O(\mu^0), \quad (4.14)$$

as $\mu \rightarrow \infty$. Note that \mathcal{R}_0 depends neither on k , nor on n . So by choosing m large enough satisfying (4.13), we get an L^2 -bound which is uniform in μ . Inserting \mathcal{R}_0 in place of $\tilde{\mathcal{R}}_0$ in the proof of the H^1 -bound, yields new constants

$$\tilde{K}_1 = O(\mu^2), \quad \tilde{K}_2 = O(\mu), \quad \tilde{K}_3 = O(\mu^2), \quad \tilde{K}_4 = O(\mu^2),$$

replacing \tilde{K}_1 , \tilde{K}_2 , \tilde{K}_3 and \tilde{K}_4 , respectively as $\mu \rightarrow \infty$. As a result,

$$\sup_{s \geq -k} \|w(s)\|_{H^1} \leq \tilde{\mathcal{R}}_1 := \sqrt{\frac{28}{3}\tilde{K}_2 + \frac{7}{3\gamma}\tilde{K}_1} = O(\mu),$$

as $\mu \rightarrow \infty$, where

$$\tilde{K}_1 := 6\gamma\|f\|\mathcal{R}_0 + 6\mu\gamma|v|_X\mathcal{R}_0 + \frac{\mu^2 c^2 \mathcal{R}_0^6}{\gamma} + 2\mu\|f\|\mathcal{R}_0 + 2\mu^2|v|_X\mathcal{R}_0 + 2\mu|v|_X\mathcal{R}_0,$$

and

$$\tilde{K}_2 := \frac{c^2 \mathcal{R}_0^6}{16\xi} + \frac{1}{2}c\mathcal{R}_0^4 + 2\|f\|\mathcal{R}_0 + 2\mu|v|_X\mathcal{R}_0 + (1 - \xi)\mathcal{R}_0^2.$$

4.4. H^2 bound

Multiply (3.5) with $\bar{w}_{xxs} + \gamma \bar{w}_{xx}$, integrate, and take the real parts:

$$\begin{aligned}
 & \operatorname{Re} \int_0^L i w_s \bar{w}_{xxs} + \gamma \operatorname{Re} \int_0^L i w_s \bar{w}_{xx} + \operatorname{Re} \int_0^L w_{xx} \bar{w}_{xxs} + \gamma \operatorname{Re} \int_0^L |w_{xx}|^2 \\
 & + \operatorname{Re} \int_0^L |w|^2 w \bar{w}_{xxs} + \gamma \operatorname{Re} \int_0^L |w|^2 w \bar{w}_{xx} + \gamma \operatorname{Re} \int_0^L i w \bar{w}_{xxs} \\
 & + \gamma^2 \operatorname{Re} \int_0^L i w \bar{w}_{xx} + \mu \operatorname{Re} \int_0^L i P_m w P_m \bar{w}_{xxs} + \mu \gamma \operatorname{Re} \int_0^L i P_m w P_m \bar{w}_{xx} \\
 & = \operatorname{Re} \int_0^L f \bar{w}_{xxs} + \gamma \operatorname{Re} \int_0^L f \bar{w}_{xx} + \mu \operatorname{Re} \int_0^L i v \bar{w}_{xxs} + \mu \gamma \operatorname{Re} \int_0^L i v \bar{w}_{xx}.
 \end{aligned}$$

We now estimate term by term, using integration by parts in most cases.

$$\begin{aligned}
 \operatorname{Re} \int_0^L i w_s \bar{w}_{xxs} &= -\operatorname{Re} \int_0^L i |w_{xs}|^2 = 0, \\
 \gamma \operatorname{Re} \int_0^L i w_s \bar{w}_{xx} &= -\gamma \operatorname{Re} \int_0^L i w_{xs} \bar{w}_x = \gamma \operatorname{Im} \int_0^L w_{xs} \bar{w}_x = -\gamma \operatorname{Im} \int_0^L w_x \bar{w}_{xs}, \\
 \operatorname{Re} \int_0^L w_{xx} \bar{w}_{xxs} &= \frac{1}{2} \frac{d}{ds} \|w_{xx}\|^2, \\
 \gamma \operatorname{Re} \int_0^L |w_{xx}|^2 &= \gamma \|w_{xx}\|^2, \\
 \operatorname{Re} \int_0^L |w|^2 w \bar{w}_{xxs} &= -\operatorname{Re} \int_0^L (w^2 \bar{w})_x \bar{w}_{xs} = -\operatorname{Re} \int_0^L 2w w_x \bar{w} \bar{w}_{xs} - \operatorname{Re} \int_0^L w^2 \bar{w}_x \bar{w}_{xs} \\
 &= -\int_0^L |w|^2 \frac{d}{ds} |w_x|^2 - \frac{1}{2} \operatorname{Re} \int_0^L w^2 \frac{d}{ds} \bar{w}_x^2,
 \end{aligned}$$

$$\begin{aligned}
\gamma \operatorname{Re} \int_0^L |w|^2 w \bar{w}_{xx} &= -\gamma \operatorname{Re} \int_0^L (w^2 \bar{w})_x \bar{w}_x = -\gamma \operatorname{Re} \int_0^L 2w w_x \bar{w} \bar{w}_x - \gamma \operatorname{Re} \int_0^L w^2 \bar{w}_x \bar{w}_x \\
&= -2\gamma \int_0^L |w|^2 |w_x|^2 - \gamma \operatorname{Re} \int_0^L w^2 \bar{w}_x^2, \\
\gamma \operatorname{Re} \int_0^L i w \bar{w}_{xxs} &= -\gamma \operatorname{Re} \int_0^L i w_x \bar{w}_{xs} = \gamma \operatorname{Im} \int_0^L w_x \bar{w}_{xs}, \\
\gamma^2 \operatorname{Re} \int_0^L i w \bar{w}_{xx} &= -\gamma^2 \operatorname{Re} \int_0^L i |w_x|^2 = 0, \\
\mu \operatorname{Re} \int_0^L i P_m w P_m \bar{w}_{xss} &= -\mu \operatorname{Re} \int_0^L i P_m w_x P_m \bar{w}_{xs} = \mu \operatorname{Im} \int_0^L P_m w_x P_m \bar{w}_{xs}, \\
\mu \gamma \operatorname{Re} \int_0^L i P_m w P_m \bar{w}_{xx} &= -\mu \gamma \operatorname{Re} \int_0^L i |P_m w_x|^2 = 0.
\end{aligned}$$

Now, we combine the above terms to get

$$\begin{aligned}
\frac{1}{2} \frac{d\varphi}{ds} + \gamma \varphi &= - \int_0^L 2 \operatorname{Re}(w \bar{w}_s |w_x|^2) - \operatorname{Re} \int_0^L w w_s \bar{w}_x^2 \\
&\quad + \mu \operatorname{Im} \int_0^L v_s \bar{w}_{xx} + \gamma \mu \operatorname{Im} \int_0^L v \bar{w}_{xx} \\
&\quad - \gamma \operatorname{Re} \int_0^L f \bar{w}_{xx} - \mu \operatorname{Im} \int_0^L P_m w_x P_m \bar{w}_{xs}, \tag{4.15}
\end{aligned}$$

for all $s \geq -k$, where

$$\begin{aligned}
\varphi(w) &:= \|w_{xx}\|^2 - 2 \int_0^L |w|^2 |w_x|^2 - \operatorname{Re} \int_0^L w^2 \bar{w}_x^2 \\
&\quad - 2 \operatorname{Re} \int_0^L f \bar{w}_{xx} + 2\mu \operatorname{Im} \int_0^L v \bar{w}_{xx}. \tag{4.16}
\end{aligned}$$

Observe again that since $w(-k, x) = w_x(-k, x) = w_{xx}(-k, x) = 0$, for all $x \in [0, L]$, we have $\varphi(-k) = 0$. We write

$$w_s = i w_{xx} + h, \quad (4.17)$$

where $h := i|w|^2 w - \gamma w - \mu P_m w + \mu v - i f$. Observe, thanks to Agmon's inequality, that

$$\begin{aligned} \|h(s)\| &\leq c\|w\|^2\|w\|_{H^1} + (\gamma + \mu)\|w\| + \mu\|v\| + \|f\| \\ &\leq c\mathcal{R}_0^2\tilde{\mathcal{R}}_1 + (\gamma + \mu)\mathcal{R}_0 + \mu\|v\|_X + \|f\|, \end{aligned}$$

for all $s \geq -k$. We estimate each term on the right-hand side of (4.15). We use (4.17), as well as the Young, the Hölder, and the Agmon inequalities to obtain

$$\begin{aligned} -\int_0^L 2 \operatorname{Re}(w \bar{w}_s |w_x|^2) &\leq 2 \int_0^L |w| |w_s| |w_x|^2 = 2 \int_0^L |w| |i w_{xx} + h| |w_x|^2 \\ &\leq 2 \int_0^L |w| |w_{xx}| |w_x|^2 + 2 \int_0^L |w| |h| |w_x|^2 \\ &\leq 2\|w\|_\infty \|w_x\|_\infty \int_0^L |w_{xx}| |w_x| + 2\|w\| \|h\| \|w_x\|_\infty^2 \\ &\leq 2\|w\|_\infty \|w_x\|_\infty \|w_{xx}\| \|w_x\| + 2\|w\| \|h\| \|w_x\|_\infty^2 \\ &\leq c\|w\|_\infty \|w_x\|^\frac{3}{2} \|w_{xx}\|^\frac{3}{2} + c\|w\| \|h\| \|w_x\| \|w_{xx}\| \\ &\leq c(\|w\|^\frac{1}{2} \|w\|_{H^1}^\frac{1}{2}) \|w_{xx}\|^\frac{3}{2} + c(\|w\| \|h\| \|w_x\|) \|w_{xx}\| \\ &\leq c(\mathcal{R}_0^\frac{1}{2} \tilde{\mathcal{R}}_1^\frac{1}{2}) \|w_{xx}\|^\frac{3}{2} \\ &\quad + c\mathcal{R}_0 \tilde{\mathcal{R}}_1 (\mathcal{R}_0^2 \tilde{\mathcal{R}}_1 + (\gamma + \mu)\mathcal{R}_0 + \mu\|v\|_X + \|f\|) \|w_{xx}\| \\ &\leq \frac{\gamma}{20} \|w_{xx}\|^2 + \frac{c(\mathcal{R}_0^\frac{1}{2} \tilde{\mathcal{R}}_1^\frac{1}{2})^4}{\gamma^3} \\ &\quad + \frac{c(\mathcal{R}_0 \tilde{\mathcal{R}}_1 (\mathcal{R}_0^2 \tilde{\mathcal{R}}_1 + (\gamma + \mu)\mathcal{R}_0 + \mu\|v\|_X + \|f\|))^2}{\gamma} \\ &= \frac{\gamma}{20} \|w_{xx}\|^2 + \tilde{K}_5, \end{aligned}$$

where

$$\tilde{K}_5 := \frac{c(\mathcal{R}_0^\frac{1}{2} \tilde{\mathcal{R}}_1^\frac{1}{2})^4}{\gamma^3} + \frac{c\{\mathcal{R}_0 \tilde{\mathcal{R}}_1 [\mathcal{R}_0^2 \tilde{\mathcal{R}}_1 + (\gamma + \mu)\mathcal{R}_0 + \mu\|v\|_X + \|f\|]\}^2}{\gamma}.$$

Similarly, we have

$$-Re \int_0^L w w_s \bar{w}_x^2 \leq \frac{\gamma}{10} \|w_{xx}\|^2 + \tilde{K}_5,$$

and

$$\begin{aligned} \mu Im \int_0^L v_s \bar{w}_{xx} &\leq \mu \|v_s\| \|w_{xx}\| \leq \frac{\gamma}{20} \|w_{xx}\|^2 + \frac{c\mu^2 |v|_X^2}{\gamma}, \\ \gamma \mu Im \int_0^L v \bar{w}_{xx} &\leq \gamma \mu \|v\| \|w_{xx}\| \leq \frac{\gamma}{20} \|w_{xx}\|^2 + c\gamma \mu^2 |v|_X^2, \\ -\gamma Re \int_0^L f \bar{w}_{xx} &\leq \gamma \|f\| \|w_{xx}\| \leq \frac{\gamma}{20} \|w_{xx}\|^2 + c\gamma \|f\|^2. \end{aligned}$$

For the term $-\mu Im \int_0^L P_m w_x P_m \bar{w}_{xs}$, we take the P_m projection of Eq. (3.5), multiply it with $\mu P_m \bar{w}_{xx}$, integrate, and take the real part to get

$$\begin{aligned} -\mu Im \int_0^L P_m w_x P_m \bar{w}_{xs} &= -\mu \|P_m w_{xx}\|^2 - \mu Re \int_0^L P_m (|w|^2 w) P_m \bar{w}_{xx} \\ &\quad + \mu Re \int_0^L f P_m \bar{w}_{xx} - \mu^2 Im \int_0^L v P_m \bar{w}_{xx} \\ &\leq -\mu \|P_m w_{xx}\|^2 + \mu \|w\|_\infty^2 \|w\| \|P_m w_{xx}\| \\ &\quad + \mu \|f\| \|P_m w_{xx}\| + \mu^2 \|v\| \|P_m w_{xx}\|. \end{aligned}$$

Apply Young's inequality to eliminate $\mu \|P_m w_{xx}\|^2$, and then the Agmon inequality, to get

$$-\mu Im \int_0^L P_m w_x P_m \bar{w}_{xs} \leq c\mu (\mathcal{R}_0^2 \tilde{\mathcal{R}}_1)^2 + c\mu \|f\|^2 + c\mu^3 |v|_X^2.$$

Combine the above terms to obtain

$$\frac{1}{2} \frac{d\varphi}{ds} + \gamma \varphi \leq \tilde{K}_6 + \frac{\gamma}{4} \|w_{xx}\|^2, \quad (4.18)$$

where

$$\begin{aligned}\tilde{K}_6 := & 2\tilde{K}_5 + \frac{c\mu^2|v|_X^2}{\gamma} + c\gamma\mu^2|v|_X^2 + c\gamma\|f\|^2 \\ & + c\mu(\mathcal{R}_0^2\tilde{\mathcal{R}}_1)^2 + c\mu\|f\|^2 + c\mu^3|v|_X^2.\end{aligned}$$

Note that $\tilde{K}_6 = O(\mu^8)$ as $\mu \rightarrow \infty$. From (4.16) we obtain

$$\varphi \geq \frac{1}{2}\|w_{xx}\|^2 - \tilde{K}_7, \quad (4.19)$$

where $\tilde{K}_7 := c(\mathcal{R}_0\tilde{\mathcal{R}}_1^3 + \|f\|^2 + \mu^2|v|_X^2)$. Use (4.19) in (4.18) to get

$$\begin{aligned}\frac{1}{2}\frac{d\varphi}{ds} + \gamma\varphi & \leq \tilde{K}_6 + \frac{\gamma}{2}(\varphi + \tilde{K}_7) \\ & = \frac{\gamma}{2}\varphi + \left(\frac{\gamma}{2}\tilde{K}_7 + \tilde{K}_6\right) \\ & = \frac{\gamma}{2}\varphi + \left(\frac{\gamma}{2}\tilde{K}_7 + \tilde{K}_6\right).\end{aligned}$$

Thus we have

$$\frac{d\varphi}{ds} + \gamma\varphi \leq \gamma\tilde{K}_7 + 2\tilde{K}_6.$$

Since $\varphi(-k) = 0$, we have, thanks to Gronwall's lemma, $\varphi(s) \leq \tilde{K}_7 + \frac{2\tilde{K}_6}{\gamma}$, for all $s \geq -k$. From (4.19), we get

$$\|w_{xx}(s)\|^2 \leq 2\varphi + 2\tilde{K}_7 \leq \frac{4\tilde{K}_6}{\gamma} + 4\tilde{K}_7,$$

for all $s \geq -k$. Thus we get

$$\sup_{s \geq -k} \|w(s)\|_{H^2} \leq \tilde{\mathcal{R}}_2,$$

where

$$\tilde{\mathcal{R}}_2 := \sqrt{\frac{4\tilde{K}_6}{\gamma} + 4\tilde{K}_7} + \mathcal{R}_0 = O(\mu^4), \quad (4.20)$$

as $\mu \rightarrow \infty$. Comparing to (4.12), we observe that $\tilde{\mathcal{R}}_2 \gg \tilde{\mathcal{R}}_1$, for large μ .

4.5. Improved H^1 bound

We now use the H^2 -bound to obtain a better H^1 -bound. From (4.7) and (4.8), we realize that

$$\begin{aligned} \frac{d\phi}{ds} + 4\gamma\phi + \mu\phi &= 2\gamma\|w_x\|^2 + 6\gamma \operatorname{Re} \int_0^L f \bar{w} - 6\mu\gamma \operatorname{Im} \int_0^L v \bar{w} \\ &\quad + 2\mu \operatorname{Re} \int_0^L P_m(|w|^2 w) P_m \bar{w} - 2\mu\|P_m w_x\|^2 \\ &\quad - 2\mu \operatorname{Re} \int_0^L P_m f P_m \bar{w} - 2\mu \operatorname{Im} \int_0^L v_s \bar{w} \\ &\quad + \mu\|w_x\|^2 - \frac{\mu}{2}\|w\|_{L^4}^4 + 2\mu \operatorname{Re} \int_0^L f \bar{w}. \end{aligned} \quad (4.21)$$

We estimate the right-hand side of (4.21) as follows

$$\begin{aligned} 2\gamma\|w_x\|^2 &\leq 2\gamma\|w\|_{H^1}^2, \\ 6\gamma \operatorname{Re} \int_0^L f \bar{w} &\leq 6\gamma\|f\|\|w\| \leq 6\gamma\|f\|\mathcal{R}_0, \\ -6\mu\gamma \operatorname{Im} \int_0^L v \bar{w} &\leq 6\mu\gamma\|v\|\|w\| \leq 6\mu\gamma\|v\|_X \mathcal{R}_0, \\ 2\mu \operatorname{Re} \int_0^L P_m(|w|^2 w) P_m \bar{w} &= 2\mu \operatorname{Re} \int_0^L |w|^2 w P_m \bar{w} \\ &\leq 2\mu\|w\|_\infty^2\|w\|\|P_m w\| \\ &\leq 2\mu c(\|w\|\|w\|_{H^1})\|w\|\|P_m w\| \\ &\leq 2\mu c\|w\|^3\|w\|_{H^1} \\ &\leq \mu c^2\|w\|^6 + \mu\|w\|_{H^1}^2 \\ &\leq \mu(c^2\|w\|^6 + \|w\|^2) + \mu\|w_x\|^2 \\ &\leq \mu(c^2\mathcal{R}_0^6 + \mathcal{R}_0^2) + \mu\|w_x\|^2. \end{aligned}$$

Choose m large enough so that

$$\frac{\tilde{\mathcal{R}}_2^2 L^2}{4\pi^2(m+1)^2} \leq 1. \quad (4.22)$$

Then,

$$\begin{aligned} -2\mu \|P_m w_x\|^2 &= -2\mu \|w_x\|^2 + 2\mu \|Q_m w_x\|^2 \\ &\leq -2\mu \|w_x\|^2 + \frac{2\mu}{((m+1)\frac{2\pi}{L})^2} \|w_{xx}\|^2 \\ &\leq -2\mu \|w_x\|^2 + 2\mu, \\ 2\mu \operatorname{Re} \int_0^L f \bar{w} - 2\mu \operatorname{Re} \int_0^L P_m f P_m \bar{w} &\leq 2\mu \|f\| \|Q_m w\| \\ &\leq 2\mu \|f\| \|w\| \\ &\leq 2\mu \|f\| \mathcal{R}_0, \\ -2\mu \operatorname{Im} \int_0^L v_s \bar{w} &\leq 2\mu \|v_s\| \|w\| \leq 2\mu |v|_X \mathcal{R}_0. \end{aligned}$$

Add the above terms to obtain

$$\frac{d\phi}{ds} + 4\gamma\phi + \mu\phi \leq 2\gamma \|w\|_{H^1}^2 + K_1, \quad (4.23)$$

where

$$K_1 := 6\gamma \|f\| \mathcal{R}_0 + 6\mu\gamma |v|_X \mathcal{R}_0 + \mu(c^2 \mathcal{R}_0^6 + \mathcal{R}_0^2) + 2\mu + 2\mu \|f\| \mathcal{R}_0 + 2\mu |v|_X \mathcal{R}_0.$$

Now, as in (4.10)

$$\begin{aligned} \phi(s) &\geq \|w_x\|^2 - \frac{3c^2 \|w\|^6}{16} - \frac{1}{3} \|w_x\|^2 - \frac{1}{2} c \|w\|^4 - 2\|f\| \mathcal{R}_0 - 2\mu |v|_X \mathcal{R}_0 \\ &\geq \frac{2}{3} \|w_x\|^2 + \frac{2}{3} \|w\|^2 - \left[\frac{3c^2 \mathcal{R}_0^6}{16} + \frac{1}{2} c \mathcal{R}_0^4 + 2\|f\| \mathcal{R}_0 + 2\mu |v|_X \mathcal{R}_0 + \frac{2}{3} \mathcal{R}_0^2 \right] \\ &= \frac{2}{3} \|w\|_{H^1}^2 - K_2, \end{aligned}$$

where

$$K_2 := \frac{3c^2 \mathcal{R}_0^6}{16} + \frac{1}{2} c \mathcal{R}_0^4 + 2\|f\| \mathcal{R}_0 + 2\mu |v|_X \mathcal{R}_0 + \frac{2}{3} \mathcal{R}_0^2.$$

Thus $\|w\|_{H^1}^2 \leq \frac{3}{2} \phi(w) + \frac{3}{2} K_2$. Use this in (4.23) to obtain

$$\frac{d\phi}{ds} + \gamma\phi + \mu\phi \leq 3\gamma K_2 + K_1.$$

Thus, since $\phi(-k) = 0$, by virtue of the Gronwall lemma, we have

$$\phi(s) \leq \frac{3\gamma K_2 + K_1}{\gamma + \mu},$$

for all $s \geq -k$. Thus,

$$\sup_{s \geq -k} \|w(s)\|_{H^1} \leq \mathcal{R}_1 := \sqrt{\frac{(\frac{3}{2}\mu + 6\gamma)K_2 + K_1}{\gamma + \mu}} = O(\mu^{\frac{1}{2}}), \quad (4.24)$$

as $\mu \rightarrow \infty$. Inserting \mathcal{R}_1 in place of $\tilde{\mathcal{R}}_1$ in the proof of the H^2 -bound yields new H^2 -bound

$$\sup_{s \geq -k} \|w(s)\|_{H^2} \leq \mathcal{R}_2 = \sqrt{\frac{4K_6}{\gamma} + 4K_7} + \mathcal{R}_0 = O(\mu^2), \quad (4.25)$$

as $\mu \rightarrow \infty$, where

$$K_6 := 2K_5 + \frac{c\mu^2|v|_X^2}{\gamma} + c\gamma\mu^2|v|_X^2 + c\gamma\|f\|^2 + c\mu(\mathcal{R}_0^2\mathcal{R}_1)^2 + c\mu\|f\|^2 + c\mu^3|v|_X^2,$$

$$K_7 := c(\mathcal{R}_0\mathcal{R}_1^3 + \|f\|^2 + \mu^2|v|_X^2),$$

$$K_5 := \frac{c(\mathcal{R}_0^{\frac{1}{2}}\mathcal{R}_1^2)^4}{\gamma^3} + \frac{c\{\mathcal{R}_0\mathcal{R}_1[\mathcal{R}_0^2\mathcal{R}_1 + (\gamma + \mu)\mathcal{R}_0 + \mu|v|_X + \|f\|]\}^2}{\gamma}.$$

4.6. Time derivative bound

We realize from (3.5) that we have

$$\|w'(s)\| \leq \mathcal{R}_2 + c\mathcal{R}_0^2\mathcal{R}_1 + (\gamma + \mu)\mathcal{R}_0 + \|f\| + \mu|v|_X.$$

Thus

$$\sup_{s \geq -k} \|w'(s)\| \leq \mathcal{R}',$$

where

$$\mathcal{R}' := \mathcal{R}_2 + c\mathcal{R}_0^2\mathcal{R}_1 + (\gamma + \mu)\mathcal{R}_0 + \|f\| + \mu|v|_X. \quad (4.26)$$

4.7. Passing to the limit

To summarize our L^2 , H^1 , H^2 and the time derivative bounds, we have

$$\begin{aligned} \sup_{s \geq -k} \|w_n(s)\| &\leq \mathcal{R}_0 = O(\mu^0), \\ \sup_{s \geq -k} \|w_n(s)\|_{H^1} &\leq \mathcal{R}_1 = O(\mu^{\frac{1}{2}}), \\ \sup_{s \geq -k} \|w_n(s)\|_{H^2} &\leq \mathcal{R}_2 = O(\mu^2), \\ \sup_{s \geq -k} \|w'_n(s)\| &\leq \mathcal{R}' = O(\mu^2), \end{aligned} \quad (4.27)$$

as $\mu \rightarrow \infty$, where \mathcal{R}_0 , \mathcal{R}_1 , \mathcal{R}_2 and \mathcal{R}' are independent of k and n , and defined in (4.14), (4.24), (4.25) and (4.26), respectively, and where w_n is the solution of the initial value problem (3.5)–(3.6). Thus we have a bounded solution w_n to the Galerkin approximation (3.5) of Eq. (3.4) with initial condition (3.6), on the interval $[-k, \infty)$, and satisfying (4.27); we will call it $w_{n,k}$ to emphasize the initial time $-k$, and consider

$$w_{n,k} \in C_b([-k, \infty); H^2) \cap C_b^1([-k, \infty); L^2).$$

We now focus on the interval $[-1, 1]$. Since H_n , defined in (2.2), is finite-dimensional, we may invoke the Arzela–Ascoli compactness theorem to extract a subsequence of $w_{n,k}$, denoted by $w_{n,k}^{(1)}$, such that $w_{n,k}^{(1)} \rightarrow w_n^{(1)}$, as $k \rightarrow \infty$, where $w_n^{(1)}$ is a bounded solution of the Galerkin approximation (3.5) on the interval $[-1, 1]$. Let $j \in \mathbb{N}$, we will use an induction iterative procedure to define $w_{n,k}^{(j+1)}$ to be subsequence of $w_{n,k}^{(j)}$, all of which are subsequences of $w_{n,k}$. Indeed, we have already defined $w_{n,k}^{(1)}$. Suppose $w_{n,k}^{(j)}$ is defined, and is a subsequence of $w_{n,k}$. We apply again the Arzela–Ascoli compactness theorem to extract a subsequence of $w_{n,k}^{(j)}$, denoted by $w_{n,k}^{(j+1)}$, such that $w_{n,k}^{(j+1)} \rightarrow w_n^{(j+1)}$, as $k \rightarrow \infty$, uniformly on $[-(j+1), (j+1)]$, where $w_n^{(j+1)}$ is a bounded solution of the Galerkin approximation (3.5) on the interval $[-(j+1), (j+1)]$. Notice that $w_n^{(j)}$ satisfies all the estimates in (4.27) in the interval $[-j, j]$. By the Cantor diagonal process we have that $w_{n,k}^{(k)} \rightarrow w_n$, where w_n is a bounded solution of the Galerkin approximation (3.5) on all of \mathbb{R} satisfying all the estimates above. Thank to the compact embeddings

$$H^2 \hookrightarrow H^1 \hookrightarrow L^2,$$

and

$$\sup_{s \in \mathbb{R}} \|w_n\|_{H^2} \leq \mathcal{R}_2 \quad \text{and} \quad \sup_{s \in \mathbb{R}} \|w'_n\| \leq \mathcal{R}',$$

we can apply Aubin's compactness theorem (see, e.g., [6] and [17]). For every $m \in \mathbb{N}$ there exist a subsequence $w_n^{(m)}$ of w_n such that $w_n^{(m)} \rightarrow w^{(m)}$ on the interval $[-m, m]$, where $w^{(m)}$

is a bounded solution of (3.4) on the interval $[-m, m]$. Also, $w^{(m)}$ and $\frac{d}{ds}w^{(m)}$ satisfy estimates (4.27) on the interval $[-m, m]$. Again by the Cantor diagonal process we have a subsequence $w_n^{(n)} \rightarrow w$ where w is a bounded solution of (3.4) on all of \mathbb{R} . Since w and w' also satisfy (4.27) for all $s \in \mathbb{R}$, we have the following theorem:

Proposition 4.1. *Let $v \in X$, where X is defined as in (3.2). Then there exists a bounded solution $w \in Y$ of (3.4), where Y is defined as in (3.3).*

Note that conditions (4.13) and (4.22) are only needed to get sharper bounds. Even without these conditions, there exists a bounded solution of (3.4). But to have the bounds (4.27), we need conditions (4.13) and (4.22).

Theorem 4.2. *Let w be any bounded solution of (3.4), on \mathbb{R} , for some $v \in X$. Assume that m is large enough such that both conditions (4.13) and (4.22) hold i.e.,*

$$\max \left\{ \frac{\tilde{\mathcal{R}}_1 L}{2\pi(m+1)}, \frac{\tilde{\mathcal{R}}_2 L}{2\pi(m+1)} \right\} \leq 1, \quad (4.28)$$

where $\tilde{\mathcal{R}}_1, \tilde{\mathcal{R}}_2$ are defined as in (4.12), (4.20), respectively. Then w satisfies the bounds in (4.27).

Proof. Given that w is a bounded solution of (3.4), we can mimic Section 4. We integrate the evolution inequalities in that section from s_0 to s , then take s_0 to $-\infty$ to obtain the same bounds. \square

Remark 4.3. Assume that $v \in X$ is given.

- (1) v is independent of μ and γ . All the estimates we have depend on $|v|_X$.
- (2) Observe that $\tilde{\mathcal{R}}_1 = O(\mu^{\frac{5}{2}})$ and $\tilde{\mathcal{R}}_2 = O(\mu^4)$ as $\mu \rightarrow \infty$. Therefore, condition (4.28) implies that $m \geq O(\mu^4)$.
- (3) We note the γ dependence on these constants is of the form $\mathcal{R}_0 = O(\gamma^{-1})$, $\mathcal{R}_1 = O(\gamma^{-\frac{7}{2}})$, $\mathcal{R}_2 = O(\gamma^{-17})$, and $\mathcal{R}' = O(\gamma^{-17})$ as $\gamma \rightarrow 0$.
- (4) Note that since $\tilde{\mathcal{R}}_1 = O(\gamma^{-4})$ and $\tilde{\mathcal{R}}_2 = O(\gamma^{-19.5})$ as $\gamma \rightarrow 0$, then condition (4.28) implies that $m \geq O(\gamma^{-19.5})$.

5. Main results

Proposition 5.1. *Let u be a trajectory on the global attractor of the damped, driven NLS*

$$iu_t + u_{xx} + |u|^2 u + i\gamma u = f, \quad (5.1)$$

and let w be a bounded solution of Eq. (3.4) with $v = P_m u$. Assume that μ is large enough so that

$$\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}} < \mu \quad (5.2)$$

holds, where $\mathcal{R}_\infty = c\mathcal{R}_0\mathcal{R}_1$, and $\mathcal{R}_\infty^0 = \mathcal{R}_\infty|_{\mu=0}$. In addition, assume that m is large enough such that both (4.28) and

$$\frac{cL^2K_9}{\gamma^2(m+1)^2} \leq 1 \quad (5.3)$$

hold, where K_9 is defined in (5.12), and c is a universal constant. Then we have $w \equiv u$.

Remark 5.2. Note that condition (5.2) can be achieved since $\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}} = O(\mu^{\frac{1}{4}})$ as $\mu \rightarrow \infty$. We realize that

$$\frac{cL^2K_9}{2\gamma^2} \leq O(\gamma^{-\frac{383}{12}}), \quad \frac{cL^2K_9}{2\gamma^2} \leq O(\mu^{\frac{35}{12}}),$$

as $\gamma \rightarrow 0$ and $\mu \rightarrow \infty$. Condition (5.3) implies that we need to choose m large enough such that $m \geq O(\gamma^{-\frac{383}{24}})$ and $m \geq O(\mu^{\frac{35}{24}})$. We note that these conditions are already achieved by (5.2), (4.28) and Remark 4.3, up to a constant.

Now we give a proof for Proposition 5.1.

Proof of Proposition 5.1. We first mention that all of the bounds that we obtained in Section 4 also hold for the solution of Eq. (5.1), with $\mu = 0$. Our notation for the time derivative bound and the square of the L^∞ bound for the solution u will be \mathcal{R}'^0 and \mathcal{R}_∞^0 (see (5.8) for \mathcal{R}_∞), respectively. We will use the superscript 0 in K_j^0 to denote the constant K_j , but with $\mu = 0$ in its formula. Taking the difference of the following equations

$$\begin{aligned} iw_s + w_{xx} + |w|^2w + i\gamma w &= f - i\mu[P_m(w) - u], \\ iu_s + u_{xx} + |u|^2u + i\gamma u &= f, \end{aligned}$$

we get

$$i\delta_s + \delta_{xx} + |w|^2w - |u|^2u + i\gamma\delta = -i\mu P_m\delta,$$

where $\delta := w - u$. Note that

$$\begin{aligned} |w|^2w - |u|^2u &= |\delta|^2\delta + w\bar{u}\delta + \bar{w}u\delta + wu\bar{\delta} \\ &= |\delta|^2\delta + 2\operatorname{Re}(w\bar{u})\delta + wu\bar{\delta}, \end{aligned}$$

and hence

$$i\delta_s + \delta_{xx} + |\delta|^2\delta + 2\operatorname{Re}(w\bar{u})\delta + wu\bar{\delta} + i\gamma\delta = -i\mu P_m\delta. \quad (5.4)$$

Multiply (5.4) by $\bar{\delta}$, integrate, and take the real parts to get

$$\operatorname{Im} \int_0^L \delta \bar{\delta}_s = \|\delta_x\|^2 - \int_0^L |\delta|^4 - 2 \int_0^L \operatorname{Re}(w\bar{u})|\delta|^2 - \operatorname{Re} \int_0^L wu\bar{\delta}^2. \quad (5.5)$$

Define $\Phi(s)$ as follows

$$\Phi(s) = \|\delta_x\|^2 - \frac{1}{2} \int_0^L |\delta|^4 - 2 \int_0^L \operatorname{Re}(w\bar{u})|\delta|^2 - \operatorname{Re} \int_0^L wu\bar{\delta}^2. \quad (5.6)$$

Thus from (5.5), we have

$$\operatorname{Im} \int_0^L \delta \bar{\delta}_s = \Phi(s) - \frac{1}{2} \int_0^L |\delta|^4.$$

Now multiply (5.4) by $P_m \bar{\delta}$, integrate, and take the real parts to get

$$\begin{aligned} \operatorname{Im} \int_0^L P_m \delta P_m \bar{\delta}_s &= \|P_m \delta_x\|^2 - \operatorname{Re} \int_0^L |\delta|^2 \delta P_m \bar{\delta} \\ &\quad - 2 \int_0^L \operatorname{Re}(w\bar{u}) \operatorname{Re}(\delta P_m \bar{\delta}) - \operatorname{Re} \int_0^L wu\bar{\delta} P_m \bar{\delta}. \end{aligned} \quad (5.7)$$

Multiply (5.4) by $\bar{\delta}_s$, integrate, and take the real parts to get

$$\begin{aligned} \frac{d}{ds} \|\delta_x\|^2 - \frac{1}{2} \frac{d}{ds} \int_0^L |\delta|^4 &= -2\gamma \operatorname{Im} \int_0^L \delta \bar{\delta}_s - 2\mu \operatorname{Im} \int_0^L P_m \delta P_m \bar{\delta}_s \\ &\quad + 4 \int_0^L \operatorname{Re}(w\bar{u}) \operatorname{Re}(\delta \bar{\delta}_s) + 2 \operatorname{Re} \int_0^L wu\bar{\delta} \bar{\delta}_s. \end{aligned}$$

We then realize that,

$$\begin{aligned} \frac{d}{ds} \Phi(s) + 2\gamma \Phi(s) &= -2\mu \operatorname{Im} \int_0^L P_m \delta P_m \bar{\delta}_s - 2 \int_0^L \operatorname{Re}(w\bar{u}_s)|\delta|^2 \\ &\quad - \operatorname{Re} \int_0^L (wu)_s \bar{\delta}^2 + \gamma \int_0^L |\delta|^4. \end{aligned}$$

Use (5.7) above to get

$$\begin{aligned}
\frac{d}{ds}\Phi(s) + 2\gamma\Phi(s) &= -2\mu\|P_m\delta_x\|^2 + 2\mu\operatorname{Re}\int_0^L |\delta|^2\delta P_m\bar{\delta} \\
&\quad + 4\mu\int_0^L \operatorname{Re}(w\bar{u})\operatorname{Re}(\delta P_m\bar{\delta}) + 2\mu\operatorname{Re}\int_0^L w\bar{u}\bar{\delta}P_m\bar{\delta} \\
&\quad - 2\int_0^L \operatorname{Re}(w\bar{u}_s)|\delta|^2 - \operatorname{Re}\int_0^L (wu)_s\bar{\delta}^2 \\
&\quad + \gamma\int_0^L |\delta|^4.
\end{aligned}$$

Since condition (4.28) is an assumption of the proposition, we may use the bounds we obtained in Section 4. By using Agmon's inequality, along with derivative bound (4.26) and

$$\|w\|_\infty^2 \leq c\|w\|\|w\|_{H^1} \leq c(\mathcal{R}_0\mathcal{R}_1) := \mathcal{R}_\infty = O(\mu^{\frac{1}{2}}), \quad (5.8)$$

as $\mu \rightarrow \infty$, so we have

$$\begin{aligned}
-2\int_0^L \operatorname{Re}(w\bar{u}_s)|\delta|^2 &\leq 2\|w\|_\infty\|\delta\|_\infty\|u_s\|\|\delta\| \\
&\leq c\mathcal{R}_\infty^{\frac{1}{2}}(\|\delta\|^{\frac{1}{2}}\|\delta\|_{H^1}^{\frac{1}{2}})\|u_s\|\|\delta\| \\
&\leq c\mathcal{R}_\infty^{\frac{1}{2}}\mathcal{R}'^0(\|\delta\|^{\frac{3}{2}}\|\delta\|_{H^1}^{\frac{1}{2}}) \\
&\leq c\mathcal{R}_\infty^{\frac{1}{2}}\mathcal{R}'^0\|\delta\|^2 \\
&\quad + c\mathcal{R}_\infty^{\frac{1}{2}}\mathcal{R}'^0\|\delta\|^{\frac{3}{2}}\|\delta_x\|^{\frac{1}{2}} \\
&\leq c\left[\mathcal{R}_\infty^{\frac{1}{2}}\mathcal{R}'^0 + \frac{1}{\gamma^{\frac{1}{3}}}\mathcal{R}_\infty^{\frac{2}{3}}(\mathcal{R}'^0)^{\frac{4}{3}}\right]\|\delta\|^2 + \frac{\gamma}{3}\|\delta_x\|^2.
\end{aligned}$$

Similar analysis can be done for the term $-\operatorname{Re}\int_0^L (wu)_s\bar{\delta}^2$. Since

$$-\operatorname{Re}\int_0^L (wu)_s\bar{\delta}^2 = -\operatorname{Re}\int_0^L w_s u\bar{\delta}^2 - \operatorname{Re}\int_0^L w u_s\bar{\delta}^2,$$

we have

$$\begin{aligned}
 -\operatorname{Re} \int_0^L (wu)_s \bar{\delta}^2 &\leq c \left[(\mathcal{R}_\infty^0)^{\frac{1}{2}} \mathcal{R}' + \frac{1}{\gamma^{\frac{1}{3}}} (\mathcal{R}_\infty^0)^{\frac{2}{3}} (\mathcal{R}')^{\frac{4}{3}} \right] \|\delta\|^2 + \frac{\gamma}{3} \|\delta_x\|^2 \\
 &\quad + c \left[\mathcal{R}_\infty^{\frac{1}{2}} \mathcal{R}'^0 + \frac{1}{\gamma^{\frac{1}{3}}} \mathcal{R}_\infty^{\frac{2}{3}} (\mathcal{R}'^0)^{\frac{4}{3}} \right] \|\delta\|^2 + \frac{\gamma}{3} \|\delta_x\|^2.
 \end{aligned}$$

We also have

$$\begin{aligned}
 2\mu \operatorname{Re} \int_0^L |\delta|^2 \delta P_m \bar{\delta} &\leq 2\mu \|\delta\|_\infty^2 \|\delta\| \|P_m \delta\| \\
 &\leq 4\mu (\mathcal{R}_\infty + \mathcal{R}_\infty^0) \|\delta\| \|P_m \delta\| \\
 &\leq 4\mu (\mathcal{R}_\infty + \mathcal{R}_\infty^0) \|\delta\|^2.
 \end{aligned}$$

After similar treatment of the terms $4\mu \int_0^L \operatorname{Re}(w\bar{u}) \operatorname{Re}(\delta P_m \bar{\delta})$, $2\mu \operatorname{Re} \int_0^L wu \bar{\delta} P_m \bar{\delta}$ and $\gamma \int_0^L |\delta|^4$, we obtain

$$\begin{aligned}
 4\mu \int_0^L \operatorname{Re}(w\bar{u}) \operatorname{Re}(\delta P_m \bar{\delta}) &\leq 4\mu (\mathcal{R}_\infty)^{\frac{1}{2}} (\mathcal{R}_\infty^0)^{\frac{1}{2}} \|\delta\|^2, \\
 2\mu \operatorname{Re} \int_0^L wu \bar{\delta} P_m \bar{\delta} &\leq 2\mu (\mathcal{R}_\infty)^{\frac{1}{2}} (\mathcal{R}_\infty^0)^{\frac{1}{2}} \|\delta\|^2, \\
 \gamma \int_0^L |\delta|^4 &\leq \gamma \|\delta\|_\infty^2 \|\delta\|^2 \leq 2\gamma (\mathcal{R}_\infty + \mathcal{R}_\infty^0) \|\delta\|^2.
 \end{aligned}$$

Combine the above terms to obtain

$$\frac{d}{ds} \Phi(s) + 2\gamma \Phi(s) + 2\mu \|P_m \delta_x\|^2 \leq cK_8 \|\delta\|^2 + \gamma \|\delta_x\|^2, \quad (5.9)$$

where

$$\begin{aligned}
 K_8 &:= \mu (\mathcal{R}_\infty + \mathcal{R}_\infty^0) + \mu (\mathcal{R}_\infty)^{\frac{1}{2}} (\mathcal{R}_\infty^0)^{\frac{1}{2}} + (\mathcal{R}_\infty^0)^{\frac{1}{2}} \mathcal{R}' + \frac{1}{\gamma^{\frac{1}{3}}} (\mathcal{R}_\infty^0)^{\frac{2}{3}} (\mathcal{R}')^{\frac{4}{3}} \\
 &\quad + \mathcal{R}_\infty^{\frac{1}{2}} \mathcal{R}'^0 + \frac{1}{\gamma^{\frac{1}{3}}} \mathcal{R}_\infty^{\frac{2}{3}} (\mathcal{R}'^0)^{\frac{4}{3}}.
 \end{aligned}$$

We realize that $K_8 = O(\mu^{\frac{8}{3}})$ as $\mu \rightarrow \infty$. Also, from (5.6)

$$\Phi(s) \geq \|\delta_x\|^2 - c(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}} (\mathcal{R}_\infty^0)^{\frac{1}{2}}) \|\delta\|^2,$$

and hence

$$\|\delta_x\|^2 \leq \Phi(s) + c(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})\|\delta\|^2. \quad (5.10)$$

Using (5.9) and (5.10), we conclude that

$$\frac{d}{ds}\Phi(s) + 2\gamma\Phi(s) \leq c[K_8 + \gamma(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})]\|\delta\|^2 + \gamma\Phi(s),$$

so

$$\frac{d}{ds}\Phi(s) + \gamma\Phi(s) \leq c[K_8 + \gamma(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})]\|\delta\|^2.$$

So we have

$$\Phi(s) \leq \frac{c[K_8 + \gamma(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})]}{\gamma} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2.$$

Thus,

$$\|\delta_x\|^2 \leq \frac{c[K_8 + \gamma(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})]}{\gamma} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2. \quad (5.11)$$

The inequality (5.11) is a ‘reverse’ Poincaré type inequality. From (4.2),

$$\begin{aligned} \frac{d}{ds}\|\delta\|^2 + 2\gamma\|\delta\|^2 + 2\mu\|P_m\delta\|^2 &\leq 2\|w\|_\infty\|u\|_\infty\|\delta\|^2 \\ &\leq 2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}\|\delta\|^2 \\ &= 2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}\|P_m\delta\|^2 + 2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}\|Q_m\delta\|^2 \\ &\leq 2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}\|P_m\delta\|^2 + \frac{2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}}{(m+1)^{\frac{2\pi}{L}}}\|\delta_x\|^2 \\ &\leq 2\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}\|P_m\delta\|^2 + \frac{cL^2K_9}{\gamma(m+1)^2} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2, \end{aligned}$$

where

$$K_9 := \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}}[K_8 + \gamma(\mathcal{R}_\infty + \mathcal{R}_\infty^0 + \mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}})]. \quad (5.12)$$

Thus, if we choose μ large enough so that $c\mathcal{R}_\infty^{\frac{1}{2}}(\mathcal{R}_\infty^0)^{\frac{1}{2}} = O(\mu^{\frac{1}{4}}) \leq 2\mu$ (which is the condition (5.2)), we get

$$\frac{d}{ds}\|\delta\|^2 + 2\gamma\|\delta\|^2 \leq \frac{cL^2K_9}{\gamma(m+1)^2} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2,$$

and hence

$$\sup_{s \in \mathbb{R}} \|\delta(s)\|^2 \leq \frac{cL^2 K_9}{2\gamma^2(m+1)^2} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2.$$

Since m is chosen to be large enough satisfying condition (5.3), we conclude that $\sup_{s \in \mathbb{R}} \|\delta(s)\|^2 = 0$. Thus $\delta \equiv 0$. This implies that $w \equiv u$. \square

Theorem 5.3. Let $v \in B_\rho := \{v \in X; |v|_X \leq \rho\}$ for some positive ρ . Assume that

$$c(\mathcal{R}_0 \mathcal{R}_1) < \mu \quad (5.13)$$

holds, and for such μ , conditions (4.28) and

$$\frac{cL^2 K_{10}}{\gamma^2(m+1)^2} \leq 1 \quad (5.14)$$

hold, where K_{10} is defined in (5.18). Then the map $W : X \rightarrow Y$, where $W(v) := w$ is a bounded solution of (3.4) provided by Proposition 4.1, is well-defined, and $P_m W : X \rightarrow X$ is a locally Lipschitz function with Lipschitz constant $L_W(\rho)$ given in (5.21).

Remark 5.4.

- (1) Recall that $\mathcal{R}_0, \mathcal{R}_1, \mathcal{R}', \mathcal{R}_\infty$ depend on $|v|_X$ which is controlled by ρ . Also note that condition (5.13) can be achieved since $c(\mathcal{R}_0 \mathcal{R}_1) = O(\mu^{\frac{1}{2}})$ as $\mu \rightarrow \infty$.
- (2) Notice that $L_W(\rho) = O(\mu^{\frac{3}{2}})$ as $\mu \rightarrow \infty$, $L_W(\rho) = O(\gamma^{-\frac{9}{2}})$ as $\gamma \rightarrow 0$ and $L_W(\rho)$ can be bounded independent of m .
- (3) We note that

$$\frac{cL^2 K_{10}}{2\gamma^2} \leq O(\gamma^{-\frac{65}{2}}), \quad \frac{cL^2 K_{10}}{2\gamma^2} \leq O(\mu^{\frac{7}{2}}),$$

as $\gamma \rightarrow 0$ and $\mu \rightarrow \infty$. Condition (5.14) implies that we need to choose m large enough such that $m \geq O(\gamma^{-\frac{65}{4}})$ and $m \geq O(\mu^{\frac{7}{4}})$. These conditions are already achieved by (5.13), (4.28) and Remark 4.3, up to a constant.

Now we give the proof of Theorem 5.3.

Proof of Theorem 5.3. Note that all constants $\mathcal{R}_0, \mathcal{R}_1, \mathcal{R}_2, \mathcal{R}'$ and \mathcal{R}_∞ depend on $|v|_X$. So, since we are in a ball $B_\rho \subset X$, all of these constants will depend on ρ . Let $v, \tilde{v} \in B_\rho$ such that $W(v) = w$ and $W(\tilde{v}) = \tilde{w}$. Since w and \tilde{w} are the solutions of Eq. (3.4) for v and \tilde{v} , respectively, the following hold:

$$\begin{aligned} iw_s + w_{xx} + |w|^2 w + i\gamma w &= f - i\mu[P_m(w) - v], \\ i\tilde{w}_s + \tilde{w}_{xx} + |\tilde{w}|^2 \tilde{w} + i\gamma \tilde{w} &= f - i\mu[P_m(\tilde{w}) - \tilde{v}]. \end{aligned}$$

Subtract, denoting $\delta := w - \tilde{w}$ and $\eta := v - \tilde{v}$, to obtain

$$i\delta_s + \delta_{xx} + |\delta|^2\delta + 2\operatorname{Re}(w\tilde{w})\delta + w\tilde{w}\bar{\delta} + i\gamma\delta + i\mu P_m\delta = i\mu\eta. \quad (5.15)$$

Multiply (5.15) by $\bar{\delta}$, integrate, and take the real parts to get

$$\begin{aligned} \operatorname{Im} \int_0^L \delta \bar{\delta}_s &= \|\delta_x\|^2 - \int_0^L |\delta|^4 - 2 \int_0^L \operatorname{Re}(w\tilde{w})|\delta|^2 \\ &\quad - \operatorname{Re} \int_0^L w\tilde{w}\bar{\delta}^2 - \mu \operatorname{Im} \int_0^L \eta \bar{\delta}. \end{aligned}$$

Define $\Psi(s)$ as follows

$$\begin{aligned} \Psi(s) &= \|\delta_x\|^2 - \frac{1}{2} \int_0^L |\delta|^4 - 2 \int_0^L \operatorname{Re}(w\tilde{w})|\delta|^2 \\ &\quad - \operatorname{Re} \int_0^L w\tilde{w}\bar{\delta}^2 - \mu \operatorname{Im} \int_0^L \eta \bar{\delta}. \end{aligned} \quad (5.16)$$

Now multiply (5.15) by $P_m\bar{\delta}$, integrate, and take the real parts to get

$$\begin{aligned} \operatorname{Im} \int_0^L P_m\delta P_m\bar{\delta}_s &= \|P_m\delta_x\|^2 - \operatorname{Re} \int_0^L |\delta|^2\delta P_m\bar{\delta} \\ &\quad - 2 \int_0^L \operatorname{Re}(w\tilde{w})\operatorname{Re}(\delta P_m\bar{\delta}) - \operatorname{Re} \int_0^L w\tilde{w}\bar{\delta} P_m\bar{\delta} \\ &\quad - \mu \operatorname{Im} \int_0^L \eta P_m\bar{\delta}. \end{aligned} \quad (5.17)$$

Multiply (5.15) by $\bar{\delta}_s$, integrate, and take the real parts to get

$$\begin{aligned} \frac{d}{ds} \|\delta_x\|^2 - \frac{1}{2} \frac{d}{ds} \int_0^L |\delta|^4 &= -2\gamma \operatorname{Im} \int_0^L \delta \bar{\delta}_s - 2\mu \operatorname{Im} \int_0^L P_m\delta P_m\bar{\delta}_s \\ &\quad + 4 \int_0^L \operatorname{Re}(w\tilde{w})\operatorname{Re}(\delta \bar{\delta}_s) + 2 \operatorname{Re} \int_0^L w\tilde{w}\bar{\delta} \bar{\delta}_s \\ &\quad + \mu \operatorname{Im} \int_0^L \eta \bar{\delta}_s. \end{aligned}$$

We then realize that

$$\begin{aligned} \frac{d}{ds}\Psi(s) + 2\gamma\Psi(s) &= -2\mu \operatorname{Im} \int_0^L P_m \delta P_m \bar{\delta}_s - 2 \int_0^L \operatorname{Re}(w\bar{w})_s |\delta|^2 \\ &\quad - \operatorname{Re} \int_0^L (w\bar{w})_s \bar{\delta}^2 + \gamma \int_0^L |\delta|^4 \\ &\quad - \mu \operatorname{Im} \int_0^L \eta_s \bar{\delta} - 2\gamma\mu \operatorname{Im} \int_0^L \eta \bar{\delta}. \end{aligned}$$

Use (5.17) above to get

$$\begin{aligned} \frac{d}{ds}\Psi(s) + 2\gamma\Psi(s) &= -2\mu \|P_m \delta_x\|^2 + 2\mu \operatorname{Re} \int_0^L |\delta|^2 \delta P_m \bar{\delta} \\ &\quad + 4\mu \int_0^L \operatorname{Re}(w\bar{u}) \operatorname{Re}(\delta P_m \bar{\delta}) \\ &\quad + 2\mu \operatorname{Re} \int_0^L wu \bar{\delta} P_m \bar{\delta} - 2 \int_0^L \operatorname{Re}(w\bar{u}_s) |\delta|^2 \\ &\quad - \operatorname{Re} \int_0^L (wu)_s \bar{\delta}^2 + \gamma \int_0^L |\delta|^4 \\ &\quad - \mu \operatorname{Im} \int_0^L \eta_s \bar{\delta} - 2\gamma\mu \operatorname{Im} \int_0^L \eta \bar{\delta}. \end{aligned}$$

We estimate as before to obtain

$$\|\delta_x\|^2 \leq \frac{(c(\mu + \gamma)\mathcal{R}_\infty + c\gamma^{-\frac{1}{3}}\mathcal{R}_\infty^{\frac{2}{3}}\mathcal{R}'^{\frac{4}{3}})}{\gamma} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2 + (\mu + 3\gamma\mu)|\eta|_X \sup_{s \in \mathbb{R}} \|\delta(s)\|.$$

Multiply (5.15) with $\bar{\delta}$, integrate, and take the imaginary parts to obtain

$$\frac{d}{ds} \|\delta\|^2 + 2\gamma \|\delta\|^2 + 2\mu \|P_m \delta\|^2 = 2\mu \operatorname{Re} \int_0^L \eta \bar{\delta} - 2 \operatorname{Im} \int_0^L w\bar{w} \bar{\delta}^2.$$

We make similar estimates again, and take advantage of the condition (5.13) to get

$$\begin{aligned} \frac{d}{ds} \|\delta\|^2 + 2\gamma \|\delta\|^2 &\leq \frac{cL^2 K_{10}}{\gamma(m+1)^2} \sup_{s \in \mathbb{R}} \|\delta(s)\|^2 \\ &\quad + \frac{c\mathcal{R}_\infty(\mu + 3\gamma\mu)}{(\frac{2\pi}{L})^2(m+1)^2} |\eta|_X \sup_{s \in \mathbb{R}} \|\delta(s)\|, \end{aligned}$$

where

$$K_{10} = \mathcal{R}_\infty [(\mu + \gamma)\mathcal{R}_\infty + \gamma^{-\frac{1}{3}} \mathcal{R}_\infty^{\frac{2}{3}} \mathcal{R}'^{\frac{4}{3}}]. \quad (5.18)$$

By (5.14) we have

$$\sup_{s \in \mathbb{R}} \|\delta(s)\| \leq \frac{c\mathcal{R}_\infty(\mu + 3\gamma\mu)}{(\frac{2\pi}{L})^2(m+1)^2\gamma} |\eta|_X. \quad (5.19)$$

Note that (5.19) implies that the W -map is well-defined. Now,

$$\|P_m \delta_{xx}\| \leq m^2 \|P_m \delta\| \leq m^2 \|\delta\| \leq \frac{c\mathcal{R}_\infty(\mu + 3\gamma\mu)}{(\frac{2\pi}{L})^2\gamma} |\eta|_X. \quad (5.20)$$

From (5.15) and (5.20),

$$\begin{aligned} \|P_m \delta_s\| &\leq \|P_m \delta_{xx}\| + \|P_m (|w|^2 \delta + w \tilde{w} \bar{\delta} + |\tilde{w}|^2 \delta)\| + (\gamma + \mu) \|P_m \delta\| + \mu \|\eta\| \\ &\leq m^2 \|\delta\| + c\mathcal{R}_\infty \|\delta\| + (\gamma + \mu) \|\delta\| + \mu \|\eta\| \\ &\leq \left((m^2 + c\mathcal{R}_\infty + \gamma + \mu) \left(\frac{c\mathcal{R}_\infty(\mu + 3\gamma\mu)}{(\frac{2\pi}{L})^2(m+1)^2\gamma} \right) + \mu \right) |\eta|_X, \end{aligned}$$

so that

$$\|P_m \delta\| + \|P_m \delta_s\| \leq L_W |\eta|_X,$$

where

$$L_W := (m^2 + c\mathcal{R}_\infty + \gamma + \mu + 1) \left(\frac{c\mathcal{R}_\infty(\mu + 3\gamma\mu)}{(\frac{2\pi}{L})^2(m+1)^2\gamma} \right) + \mu. \quad (5.21)$$

Thus,

$$|P_m \delta|_X \leq L_W |\eta|_X,$$

i.e.,

$$|P_m W(v) - P_m W(\tilde{v})|_X \leq L_W |v - \tilde{v}|_X. \quad \square$$

6. The determining form

For every trajectory u in the global attractor, \mathcal{A} , we have

$$|u|_X \leq R,$$

where

$$R := \mathcal{R}_0^0 + \mathcal{R}'^0, \quad (6.1)$$

with $\mathcal{R}_0^0 = \mathcal{R}_0|_{\mu=0}$ and $\mathcal{R}'^0 = \mathcal{R}'|_{\mu=0}$. Let u^* be a steady state of the damped, driven NLS (2.1). Adapting the suggestion given in [9], we propose the following *determining form* for the damped-driven NLS:

$$\frac{dv}{dt} = -|v - P_m W(v)|_{X,0}^2 (v - P_m u^*), \quad (6.2)$$

where $|\cdot|_{X,0}$ is defined in (3.1). The specific conditions on m and its dependence on R to guarantee the existence of a Lipschitz map $P_m W(v)$ are stated in Theorem 6.1 below.

Theorem 6.1. *Suppose that the conditions of Theorem 5.3 hold for $\rho = 4R$, where R is defined in (6.1).*

- (1) *The vector field in the determining form (6.2) is a Lipschitz map from the ball $\mathcal{B}_X^\rho(0) = \{v \in X: |v|_X < \rho\}$ into X . Thus, (6.2) is actually an ODE, in the Banach space X , which has a short time existence and uniqueness for the initial data in $\mathcal{B}_X^\rho(0) = \{v \in X: |v|_X < \rho\}$.*
- (2) *The ball $\mathcal{B}_X^{3R}(P_m u^*) = \{v \in X: |v - P_m u^*|_X < 3R\} \subset \mathcal{B}_X^\rho(0)$ is forward invariant in time, under the dynamics of the determining form (6.2). Consequently, (6.2) has global existence and uniqueness for all initial data in $\mathcal{B}_X^{3R}(P_m u^*)$.*
- (3) *Every solution of the determining form (6.2), with initial data in $\mathcal{B}_X^{3R}(P_m u^*)$, converges to a steady state of the determining form (6.2).*
- (4) *All the steady states of the determining form, (6.2), that are contained in the ball $\mathcal{B}_X^\rho(0)$, are given by the form $v(s) = P_m u(s)$, for all $s \in \mathbb{R}$, where $u(s)$ is a trajectory that lies on the global attractor, \mathcal{A} , of (2.1).*

Proof. We use the fact that $P_m W$ is a locally Lipschitz map to prove item (1) above. For items (2) and (3), we use dissipative property of (6.2). To prove item (4), we realize that the right-hand side of (6.2) is zero when either $v = P_m u^*$ or $v = P_m w$. In either case, we show that $v(s) = P_m u(s)$, for all $s \in \mathbb{R}$, where $u(s)$ is a trajectory that lies on the global attractor, \mathcal{A} , of (2.1). For details see [9]. \square

7. A new proof of the determining modes property

Here we give the proof of Theorem 3.1:

Proof of Theorem 3.1. We assume $u(s)$ and $\tilde{u}(s)$ are trajectories on the global attractor, \mathcal{A} , of (2.1), and $P_m(u(s)) = P_m(v(s))$ for all time $s \in \mathbb{R}$, and for some $m \in \mathbb{N}$ to be chosen later. Then,

$$\begin{aligned} iu_s + u_{xx} + |u|^2u + i\gamma u &= f, \\ i\tilde{u}_s + \tilde{u}_{xx} + |\tilde{u}|^2\tilde{u} + i\gamma\tilde{u} &= f. \end{aligned}$$

Subtract, denoting $\delta := u - \tilde{u}$, to obtain

$$i\delta_s + \delta_{xx} + |\delta|^2\delta + 2\operatorname{Re}(u\tilde{u})\delta + u\tilde{u}\bar{\delta} + i\gamma\delta = 0,$$

which is precisely (5.4), but with $\mu = 0$, w replaced by u , and \bar{u} replaced by $\bar{\tilde{u}}$. Following the proof of Proposition 5.1, we obtain the analog of (5.11) with $\mu = 0$:

$$\|\delta_x(s)\| \leq K_{11} \sup_{s \in \mathbb{R}} \|\delta(s)\|,$$

where

$$K_{11} = \sqrt{\frac{(c\gamma\mathcal{R}_\infty^0 + c\gamma^{-\frac{1}{3}}(\mathcal{R}_\infty^0)^{\frac{2}{3}}(\mathcal{R}'^0)^{\frac{4}{3}})}{\gamma}}. \quad (7.1)$$

Then, since $P_m\delta = 0$, we have

$$\begin{aligned} \|Q_m\delta\| &\leq \frac{L}{2\pi(m+1)} \|\delta_x\| \leq \frac{L}{2\pi(m+1)} K_{11} \sup_{s \in \mathbb{R}} \|\delta\| \\ &= \frac{L}{2\pi(m+1)} K_{11} \sup_{s \in \mathbb{R}} \|Q_m\delta\|. \end{aligned}$$

Thus, if we choose

$$m \geq \frac{L}{2\pi} K_{11} - 1,$$

we obtain that $Q_m\delta = 0$. As a result, $u(s) = \tilde{u}(s)$. \square

Remark 7.1.

- (1) By tracking the $\|f\|$ and γ dependence of the bounds throughout the paper, we have that $\mathcal{R}_0 = O(\|f\|, \gamma^{-1})$, $\mathcal{R}_0^0 = O(\|f\|, \gamma^{-1})$, $\mathcal{R}_1 = O(\|f\|^3, \gamma^{-3.5})$, $\mathcal{R}_1^0 = O(\|f\|^3, \gamma^{-3})$, $\mathcal{R}_2 = O(\|f\|^{13}, \gamma^{-17})$, $\mathcal{R}_2^0 = O(\|f\|^{13}, \gamma^{-15})$, $\mathcal{R}' = O(\|f\|^{13}, \gamma^{-17})$, $\mathcal{R}'^0 = O(\|f\|^{13}, \gamma^{-15})$, $\mathcal{R}_\infty = O(\|f\|^4, \gamma^{-4.5})$, $\mathcal{R}_\infty^0 = O(\|f\|^4, \gamma^{-4})$, $K_8 = O(\|f\|^{20}, \gamma^{-\frac{77}{3}})$, $K_8^0 = O(\|f\|^{20}, \gamma^{-23})$, $K_9 = O(\|f\|^{24}, \gamma^{-\frac{359}{12}})$, $K_9^0 = O(\|f\|^{24}, \gamma^{-27})$, $K_{10} = O(\|f\|^{24}, \gamma^{-\frac{61}{2}})$, $K_{10}^0 = O(\|f\|^{24}, \gamma^{-27})$, as $\|f\| \rightarrow \infty$ and $\gamma \rightarrow 0$.
- (2) Since $\mathcal{R}'^0 = O(\|f\|^{13})$ and $\mathcal{R}_\infty^0 = O(\|f\|^4)$, from (7.1) we have $K_{11} = O(\|f\|^{10})$ as $\|f\| \rightarrow \infty$. Thus, from (3.5), a sufficient number of determining modes is of order $m = O(\|f\|^{10})$.
- (3) Similarly, since $\mathcal{R}'^0 = O(\gamma^{-15})$ and $\mathcal{R}_\infty^0 = O(\gamma^{-4})$, we have $K_{11} = O(\gamma^{-12})$ as $\gamma \rightarrow 0$. Thus a sufficient number of determining modes is of order $m = O(\gamma^{-12})$.

- (4) Following the analysis in the earlier work of Goubet [13], one can show that a sufficient number of the determining modes is of order $O(\gamma^{-12.5})$ as $\gamma \rightarrow 0$ and $O(\|f\|^{12})$ as $\|f\| \rightarrow \infty$.

Acknowledgments

E.S.T. would like to acknowledge the kind hospitality of the Instituto Nacional de Matemática Pura e Aplicada (IMPA), Brazil, where part of this work was completed. The work of M.S.J. is supported in part by National Science Foundation (NSF) Grant Nos. DMS-1008661 and DMS-1109638, and that of E.S.T. is supported in part by the NSF grants DMS-1009950, DMS-1109640 and DMS-1109645, as well as by the CNPq-CsF grant # 401615/2012-0, through the program Ciência sem Fronteiras.

References

- [1] A. Azouani, E. Olson, E.S. Titi, Continuous data assimilation using general interpolant observables, *J. Nonlinear Sci.* 24 (2) (2014) 277–304.
- [2] A. Azouani, E.S. Titi, Feedback control of nonlinear dissipative systems by finite determining parameters—a reaction–diffusion paradigm, *arXiv:1301.6992v1*.
- [3] W. Bao, D. Jaksch, An explicit unconditionally stable numerical method for solving damped nonlinear Schrödinger equations with a focusing nonlinearity, *SIAM J. Numer. Anal.* 4 (4) (2003) 1406–1426.
- [4] K.J. Blow, N.J. Doran, Global and local chaos in the pumped nonlinear Schrödinger equation, *Phys. Rev. Lett.* 52 (7) (1984) 526–539.
- [5] J. Bourgain, Fourier transformation restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, Part I: Schrödinger equations, *Geom. Funct. Anal.* 3 (1993) 107–156.
- [6] P. Constantin, C. Foias, *Navier–Stokes Equations*, University of Chicago Press, Chicago, IL, 1988.
- [7] G. Fibich, Self-focusing in the damped nonlinear Schrödinger equation, *SIAM J. Appl. Math.* 61 (5) (2001) 1680.
- [8] C. Foias, M.S. Jolly, R. Kravchenko, E.S. Titi, A determining form for the 2D Navier–Stokes equations — the Fourier modes case, *J. Math. Phys.* 53 (2012) 115623.
- [9] C. Foias, M.S. Jolly, R. Kravchenko, E.S. Titi, A unified approach to determining forms for the 2D Navier–Stokes equations — the general interpolant case, *Uspekhi Mat. Nauk* 69 (2(416)) (2014) 177–200, *arXiv:1309.0247*.
- [10] C. Foias, O. Manley, R. Rosa, R. Temam, *Navier–Stokes Equations and Turbulence*, Cambridge University Press, 2001.
- [11] C. Foias, G. Prodi, Sur le comportement global des solutions non-stationnaires des equations de Navier–Stokes en dimension 2, *Rend. Semin. Mat. Univ. Padova* 39 (1967) 1–34.
- [12] J.-M. Ghidaglia, Finite dimensional behavior for weakly damped driven Schrödinger equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 5 (4) (1988) 365–405.
- [13] O. Goubet, Regularity of the attractor for a weakly damped nonlinear Schrödinger equation, *Appl. Anal.* 60 (1996) 99–119.
- [14] J.K. Hale, G. Raugel, Regularity, determining modes and Galerkin methods, *J. Math. Pures Appl.* 82 (2003) 1075–1136.
- [15] D.A. Jones, E.S. Titi, Upper bounds on the number of determining modes, nodes and volume elements for the Navier–Stokes equations, *Indiana Univ. Math. J.* 42 (3) (1993).
- [16] M. Oliver, E.S. Titi, Analyticity of the attractor and the number of determining nodes for a weakly damped driven nonlinear Schrödinger equation, *Indiana Univ. Math. J.* 47 (1998) 49–73.
- [17] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, second ed., *Appl. Math. Sci.*, vol. 68, Springer-Verlag, New York, 1997.
- [18] X. Wang, An energy equation for the weakly damped driven nonlinear Schrödinger equations and its application to their attractors, *Phys. D* 88 (1995) 167–175.