

The Fredholm alternative for parabolic evolution equations with inhomogeneous boundary conditions [☆]

Lahcen Maniar ^{a,1}, Roland Schnaubelt ^{b,*}

^a *Cadi Ayyad University, Faculty of Sciences Semlalia, B.P. 2390, Marrakesh, Morocco*

^b *Institut für Analysis, Fakultät für Mathematik, Universität Karlsruhe, 76128 Karlsruhe, Germany*

Received 30 June 2006

Available online 8 January 2007

Abstract

We study the Fredholm properties of parabolic evolution equations on \mathbb{R} with inhomogeneous boundary values. These problems are transformed into evolution equations with inhomogeneities taking values in certain extrapolation spaces. Assuming that the underlying homogeneous problem is asymptotically hyperbolic, we show the Fredholm alternative for these equations. The results are applied to parabolic partial differential equations.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Fredholm operator; Index; Exponential dichotomy; Inhomogeneous evolution equation; Evolution family; Parabolic initial-boundary value problem; Inter- and extrapolation

1. Introduction

In recent years the Fredholm properties of evolution equations

$$u'(t) = A(t)u(t) + f(t), \quad t \in \mathbb{R}, \quad (1.1)$$

on a Banach space X have attracted considerable interest. In this work we establish a Fredholm alternative for a large class of parabolic inhomogeneous boundary value problems, see (1.4),

[☆] This work is part of a cooperation project supported by DFG (Germany) and CNRST (Morocco).

^{*} Corresponding author. Fax: +49 721 6087650.

E-mail addresses: maniar@ucam.ac.ma, lama@fa.uni-tuebingen.de (L. Maniar), schnaubelt@math.uni-karlsruhe.de (R. Schnaubelt).

¹ L.M. was supported by a Georg Forster grant of Alexander von Humboldt Foundation (AvH).

which can be transformed into a problem similar to (1.1) with inhomogeneities f taking values in spaces $X_{\alpha-1}^t$ larger than X . Before discussing the contents of our paper, we first want to recall related results concerning (1.1) with $f: \mathbb{R} \rightarrow X$.

A main line of research concentrates on parabolic problems, where the operators $A(t)$ generate an evolution family $U(t, s)$, $t \geq s$, having regularity properties similar to those of analytic semigroups. Moreover, it is assumed that (1.1) possesses *maximal regularity* on a space F of functions $f: \mathbb{R} \rightarrow X$ (cf. [7]). Roughly speaking, this notion means that the operator $G^0 u = -u' + A(\cdot)u$ is closed in F on the ‘minimal’ domain $D(G^0) = D(d/dt) \cap D(A(\cdot)) = \{u \in F: u(t) \in D(A(t)), u', A(\cdot)u \in F\}$. This property typically requires function spaces such as $F = L^p(\mathbb{R}, X)$ or $C^\alpha(\mathbb{R}, X)$ with $p \in (1, \infty)$ or $\alpha \in (0, 1)$ (the choice $F = L^p$ leads to additional restrictions on X and $A(t)$). Finally, one supposes that the operators $A(t)$ converge to operators $A_{\pm\infty}$ as $t \rightarrow \pm\infty$ in a suitable sense and that $i\mathbb{R}$ belongs to the resolvent sets of $A_{\pm\infty}$, i.e., the problem is ‘asymptotically hyperbolic.’ It is then known that $U(\cdot, \cdot)$ has an exponential dichotomy on intervals $[T, +\infty)$ and $(-\infty, -T]$ for possibly large $T \geq 0$, see [8,32,34]. In this setting, the (semi-)Fredholmity of G^0 was characterized in terms of properties of the stable and unstable subspaces of $U(t, s)$ at $t = T$, see [1,15–17,26,30,31], and the references therein. (Compare also Theorem 3.6 below and e.g. [23,25,27] for similar results on functional-differential equations.) This characterization implies that G^0 is Fredholm if the unstable subspaces of $A_{\pm\infty}$ have finite dimensions d_\pm (e.g., if $D(A_{\pm\infty})$ is compactly embedded in X), and then G^0 has the index $d_- - d_+$. As a result, one can describe the subspace of those $f \in F$ such that (1.1) has a solution $u \in D(G^0)$, and one obtains formulas for these solutions, cf. Theorem 3.10.

The above setting occurs if one linearizes a nonlinear parabolic problem on a bounded domain along a heteroclinic orbit connecting two hyperbolic equilibria. In this case the Fredholm property of G^0 is crucial to study the bifurcation behavior of the heteroclinic orbit by means of the Lyapunov–Schmidt reduction, see e.g. [15,30,31] and also [23,27]. We add that the property of maximal regularity makes it possible to show the persistence of Fredholm properties under large classes of perturbations, see [17].

If one discards the strong assumption of maximal regularity (i.e., G^0 is not required to be closed), then it seems to be most appropriate to define G via the ‘mild equation’

$$u(t) = U(t, s)u(s) + \int_s^t U(t, \tau)f(\tau) d\tau, \quad t \geq s, \quad (1.2)$$

for a given exponentially bounded ‘evolution family’ $U(t, s)$, $t \geq s$, with time interval \mathbb{R} (i.e., (2.3) below holds and $(t, s) \mapsto U(t, s)$ is strongly continuous for $t \geq s$). We say that a function $u \in F$ belongs to the domain $D(G)$ and $Gu = f$ if there is a function $f \in F$ such that (1.2) holds for all $t \geq s$ in \mathbb{R} . If the Cauchy problem

$$u'(t) = A(t)u(t), \quad t \geq s, \quad u(s) = x, \quad (1.3)$$

is well-posed, then G is the closure of G^0 as defined above, where $F = C_0(\mathbb{R}, X)$ or $F = L^p(\mathbb{R}, X)$ with $1 \leq p < \infty$, cf. [11,33]. In the recent paper [21] it is shown that G is Fredholm on F if and only if $U(\cdot, \cdot)$ has exponential dichotomies on intervals $(-\infty, a]$ and $[b, +\infty)$ and a certain ‘node operator’ connecting the dichotomies is Fredholm in X . We refer to [22] for somewhat stronger results under stronger assumptions and also to [9]. In fact, the ‘if’ implication of the results from [9,21,22] coincides with the corresponding assertions in [15–17], see

[15, §5.3]. We further mention that the invertibility of G on F is equivalent to the exponential dichotomy of $U(\cdot, \cdot)$ on \mathbb{R} , see [11].

In the present paper we study the (semi-)Fredholm properties of the parabolic inhomogeneous boundary value problem

$$\begin{aligned} u'(t) &= A_m(t)u(t) + g(t), \quad t \in \mathbb{R}, \\ B(t)u(t) &= h(t), \quad t \in \mathbb{R}. \end{aligned} \quad (1.4)$$

Here the linear operators $A_m(t)$ and $B(t)$ are defined on a subspace Z_t of X (e.g., $Z_t = W_p^2(\Omega)$ if $X = L^p(\Omega)$), $A_m(t)$ maps Z_t into the state space X , and $B(t)$ maps Z_t into a ‘boundary space’ Y such as $W_p^{1-1/p}(\partial\Omega)$. The inhomogeneities g and h are continuous with values in X and Y , respectively. Typically, $A_m(t)$ is an elliptic differential operator on Ω and $B(t)$ is a differential boundary operator of lower order, see Example 4.5. It is assumed that the restrictions $A(t)$ of $A_m(t)$ to the kernel of $B(t)$ satisfy the so-called Acquistapace–Terreni conditions stated in (2.1) and (2.2). These conditions are quite flexible in so far they only require a Hölder condition in t and they allow for nondense and time varying domains $D(A(t))$. Under these conditions the family $A(\cdot)$ generates an evolution family $U(\cdot, \cdot)$ on X having parabolic regularity due to [3] and [4], as described in the following section.

For a fixed operator $A(t)$ and $\alpha \in (0, 1)$, we further define the real interpolation spaces X_α^t of order (α, ∞) between $D(A(t))$ and X . In Section 2 we also introduce the corresponding extrapolation spaces $X_{\alpha-1}^t$ which are larger than X . In general, both X_α^t and $X_{\alpha-1}^t$ depend on t . The operator $A(t)$ possesses an extension $A_{\alpha-1}(t): X_{\alpha-1}^t \rightarrow X_{\alpha-1}^t$. We further suppose that the abstract boundary value problem

$$(\omega - A_m(t))v = 0, \quad B(t)v = \varphi,$$

has a unique solution $v = D(t)\varphi$ for $\varphi \in Y$ and that $Z_t \hookrightarrow X_\alpha^t$ for some $\alpha \in (0, 1)$. (Here ω is a fixed large real number.) As we see in Section 4, one can rewrite (1.4) as the evolution equation

$$u'(t) = A_{\alpha-1}(t)u(t) + f(t), \quad t \in \mathbb{R}, \quad (1.5)$$

where $f := g + (\omega - A_{\alpha-1}(\cdot))D(\cdot)h$. This reformulation of a boundary value problem seems to go back to work in boundary control theory, see e.g. [12, 29]. We also refer to [2, 7, 10, 13, 19] and [24, §5.1] for related results and techniques. We then show that f belongs to the space $E_{\alpha-1}$ for some $\alpha \in (0, 1)$ which is the extrapolation space for the multiplication operator $A(\cdot)$ defined on $E := C_0(\mathbb{R}, X)$. It is crucial for our approach that the operators $U(t, s)$ have locally uniformly bounded extensions $U_{\alpha-1}(t, s): X_{\alpha-1}^s \rightarrow X_{\alpha-1}^t$ which map $X_{\alpha-1}^s$ into X with norm less than $c(t-s)^{\alpha-1}$ for $0 < t-s \leq 1$, see Proposition 2.1 and Lemma A.1.

Thus we can define an operator $G_{\alpha-1}$ as in (1.2): a function $u \in E$ belongs to $D(G_{\alpha-1})$ and $G_{\alpha-1}u = f$ if there is an $f \in E_{\alpha-1}$ such that

$$u(t) = U(t, s)u(s) + \int_s^t U_{\alpha-1}(t, \tau)f(\tau)d\tau \quad \forall t \geq s \text{ in } \mathbb{R}. \quad (1.6)$$

A function $u \in C(\mathbb{R}, X)$ satisfying (1.6) is called a ‘mild solution’ of (1.5). In Proposition 2.6 we show that a function u satisfying (1.6) indeed solves (1.5) pointwise in the space $X_{\beta-1}^t$ for

every $\beta \in (0, \alpha)$. In so far the ‘mild definition’ of $G_{\alpha-1}$ is justified. However, in this work we will concentrate on the asymptotic behavior of (1.5), and we will not study the local regularity of the solutions to (1.5) in further details. These matters are treated in depth in [7, §V.2] assuming that for some $\alpha \in (0, 1)$ the spaces X_α^t and $X_{\alpha-1}^t$ do not depend on t , see also [2] and [14].

We further suppose that $U(\cdot, \cdot)$ has exponential dichotomies on half lines $(-\infty, -T]$ and $[T, +\infty)$ for some $T \geq 0$. (This property holds in the asymptotically hyperbolic case where the resolvents $R(\omega, A(t))$ converge in norm as $t \rightarrow \pm\infty$ to the resolvents of operators $A_{\pm\infty}$ with $i\mathbb{R} \subset \rho(A_{\pm\infty})$, see [34] and also [8,32].) We prove in Proposition 2.2 that $U_{\alpha-1}(\cdot, \cdot)$ inherits the exponential dichotomies of $U(\cdot, \cdot)$.

We characterize the (semi-)Fredholm properties of $G_{\alpha-1}$ in terms of the stable and unstable subspaces of $U(t, s)$ at T in Theorem 3.6. In the asymptotically hyperbolic case, $G_{\alpha-1}$ is Fredholm with index $d_- - d_+$ if the unstable subspaces of $A_{\pm\infty}$ have finite dimensions d_\pm . We further describe the kernel and range of $G_{\alpha-1}$ in Propositions 3.5 and 3.8. We point out that our conditions do not involve the extrapolated spaces $X_{\alpha-1}^t$. These results lead to a Fredholm alternative for the mild solutions $u \in C_0(\mathbb{R}, X)$ of (1.5) in Theorem 3.10. This theorem in turn implies a Fredholm alternative for the mild solutions of (1.4) stated in Theorem 4.4. In Example 4.5 we study a variant of this result, namely a diffusion equation formulated in the space $X = C(\overline{\Omega})$. In this case, the spaces X_α^t and $X_{\alpha-1}^t$ will depend on t , in general.

Our arguments are based on the properties of the extrapolated evolution family $U_{\alpha-1}(\cdot, \cdot)$, and they are inspired by the techniques of [16] and [17]. The main difference arises from the fact that we work with an ‘integral’ definition of $G_{\alpha-1}$ instead of the more explicit definition $G^0 = -d/dt + A(\cdot)$. The approach via G^0 would run into severe difficulties here. First, even if we consider homogeneous boundary conditions $h = 0$ in (1.4) (i.e., (1.5) on $E = C_0(\mathbb{R}, X)$ with $\alpha = 1$), we cannot expect that (1.5) has maximal regularity since we work with sup norm in time. This means that G^0 is not closed, and typically its closure G has a rather complicated domain. Second and more importantly, we want to allow for f taking values in time depending extrapolation spaces $X_{\alpha-1}^t$ so that a direct treatment of the differential equation (1.5) is quite inconvenient, cf. Appendix A. Fortunately, the mild description (1.6) of $G_{\alpha-1}$ suffices for the questions studied in this paper. On the other hand, the results from [21] or [22] do not apply since we work in extrapolation spaces and $(t, s) \mapsto U(t, s)$ need not to be strongly continuous at $t = s$.

In the next section we collect the background material for our investigations. We further show several auxiliary facts concerning the extrapolated evolution family $U_{\alpha-1}(t, s)$, its exponential dichotomies, and the bounded solvability of Cauchy problems on halflines. The third section contains our main results on the operator $G_{\alpha-1}$ which are based on a careful analysis of the behavior of its restrictions to the intervals $[T, +\infty)$ and $(-\infty, T]$. Here the main difficulty comes from the fact that in general $U(t, s)$ only has dichotomies on disjoint intervals $(-\infty, -T]$ and $[T, +\infty)$, see [16,21], and [34, §4.2] for a discussion of this phenomenon. In Section 4 we translate the results of Section 3 to the boundary value problem (1.4). The last section contains a proof of the regularity result Proposition 2.6. In a forthcoming paper we will treat perturbation results for the Fredholm index.

2. Notations, assumptions, and preliminaries

We denote by $D(A)$, $N(A)$, $R(A)$, $\sigma(A)$, $\rho(A)$ the domain, kernel, range, spectrum and resolvent set of a linear operator A . Moreover, $R(\lambda, A) := (\lambda I - A)^{-1} = (\lambda - A)^{-1}$ for $\lambda \in \rho(A)$ and

$\mathcal{L}(X)$ is the space of bounded linear operators on a Banach space X . By $c(\alpha, \dots)$ we designate a generic constant depending on quantities α, \dots .

We investigate linear operators $A(t)$, $t \in \mathbb{R}$, on a Banach space X subject to the following hypotheses introduced by P. Acquistapace and B. Terreni in [3] and [4]. There are constants $\omega \in \mathbb{R}$, $\theta \in (\pi/2, \pi)$, $K > 0$ and $\mu, \nu \in (0, 1]$ such that $\mu + \nu > 1$ and

$$\lambda \in \rho(A(t) - \omega), \quad \|R(\lambda, A(t) - \omega)\| \leq \frac{K}{1 + |\lambda|}, \quad (2.1)$$

$$\|(A(t) - \omega)R(\lambda, A(t) - \omega)[R(\omega, A(t)) - R(\omega, A(s))]\| \leq K \frac{|t - s|^\mu}{|\lambda|^\nu} \quad (2.2)$$

for all $t \in \mathbb{R}$ and $\lambda \in \mathbb{C} \setminus \{0\}$ with $|\arg(\lambda)| \leq \theta$. Observe that the domains $D(A(t))$ are not required to be dense. These conditions imply that the operators $A(\cdot)$ generate an evolution family $U(t, s)$, $t \geq s$, $t, s \in \mathbb{R}$. More precisely, for $t > s$ the map $(t, s) \mapsto U(t, s) \in \mathcal{L}(X)$ is continuous and continuously differentiable in t , $U(t, s)X \subseteq D(A(t))$, and $\partial_t U(t, s) = A(t)U(t, s)$. We further have

$$U(t, s)U(s, r) = U(t, r) \quad \text{and} \quad U(t, t) = I \quad \text{for } t \geq s \geq r. \quad (2.3)$$

Moreover, for $s \in \mathbb{R}$ and $x \in \overline{D(A(s))}$, the function $t \mapsto u(t) = U(t, s)x$ is continuous at $t = s$ and u is the unique solution in $C([s, \infty), X) \cap C^1((s, \infty), X)$ of the Cauchy problem

$$u'(t) = A(t)u(t), \quad t > s, \quad u(s) = x.$$

These facts have been established in [3] and [4], see also [2,7,24,35,36].

Before stating additional regularity properties of $U(t, s)$, we have to introduce the inter- and extrapolation spaces for $A(t)$. We refer to [7,18], and [24] for proofs and further information. Let A be a sectorial operator on X (i.e., (2.1) holds with $A(t)$ replaced by A) and $\alpha \in (0, 1)$. We make use of the real interpolation space

$$X_\alpha^A := \left\{ x \in X : \|x\|_\alpha^A := \sup_{r>0} \|r^\alpha (A - \omega)R(r, A - \omega)x\| < \infty \right\},$$

which is a Banach space endowed with the norm $\|\cdot\|_\alpha^A$. For convenience we further write $X_0^A := X$, $\|x\|_0^A := \|x\|$, $X_1^A := D(A)$ and $\|x\|_1^A := \|(\omega - A)x\|$. We also need the closed subspace $\hat{X}^A := \overline{D(A)}$ of X . Moreover, we define the extrapolation space X_{-1}^A as the completion of \hat{X}^A with respect to the norm $\|x\|_{-1}^A := \|R(\omega, A)x\|$. Then A has a unique continuous extension $A_{-1} : \hat{X}^A \rightarrow X_{-1}^A$. The operator A_{-1} satisfies (2.1) in X_{-1}^A , it is densely defined, it has the same spectrum as A , and it generates the semigroup $e^{tA_{-1}}$ on X_{-1}^A being the extension of e^{tA} . As above, we can then define the space

$$X_{\alpha-1}^A := (X_{-1})_\alpha^{A_{-1}} \quad \text{with the norm } \|x\|_{\alpha-1}^A := \|x\|_\alpha^{A_{-1}} = \sup_{r>0} \|r^\alpha R(r, A_{-1} - \omega)x\|.$$

The restriction $A_{\alpha-1} : X_\alpha^A \rightarrow X_{\alpha-1}^A$ of A_{-1} is sectorial in $X_{\alpha-1}^A$ with the same type as A , it has the same spectrum as A , and the semigroup $e^{tA_{\alpha-1}}$ on $X_{\alpha-1}^A$ is the extension of e^{tA} . Observe

that $\omega - A_{\alpha-1} : X_{\alpha}^A \rightarrow X_{\alpha-1}^A$ is an isometric isomorphism. We will frequently use the continuous embeddings

$$\begin{aligned} D(A) &\hookrightarrow X_{\beta}^A \hookrightarrow D((\omega - A)^{\alpha}) \hookrightarrow X_{\alpha}^A \hookrightarrow \hat{X}^A \subset X, \\ X &\hookrightarrow X_{\beta-1}^A \hookrightarrow D((\omega - A_{-1})^{\alpha}) \hookrightarrow X_{\alpha-1}^A \hookrightarrow X_{-1}^A \end{aligned} \quad (2.4)$$

for all $0 < \alpha < \beta < 1$, where the fractional powers are defined as usually. In general, $D(A)$ is not dense in the spaces X_{α}^A and X , and X is not dense in $X_{\alpha-1}^A$, but we have the inclusions

$$X_{\beta}^A \hookrightarrow \overline{D(A)}^{\|\cdot\|_{\alpha}^A} \quad \text{and} \quad X_{\beta-1}^A \hookrightarrow \overline{(\hat{X}^A)}^{\|\cdot\|_{\alpha-1}^A} \quad (2.5)$$

for $0 < \alpha < \beta < 1$. More precisely, one has the following fact: for $x \in X_{\beta-1}^A$, the vectors $x_n = nR(n, A_{-1})x$, $n > \omega$, belong to \hat{X}^A , $\|x_n\|_{\beta-1}^A \leq c\|x\|_{\beta-1}^A$ and $x_n \rightarrow x$ in $X_{\alpha-1}^A$. Moreover, \hat{X}^A is dense in $D((\omega - A_{-1})^{\alpha})$ and X_{-1}^A .

Given operators $A(t)$, $t \in \mathbb{R}$, satisfying (2.1), we set

$$X_{\alpha}^t := X_{\alpha}^{A(t)}, \quad X_{\alpha-1}^t := X_{\alpha-1}^{A(t)}, \quad \hat{X}^t := \hat{X}^{A(t)}$$

for $0 \leq \alpha \leq 1$ and $t \in \mathbb{R}$, with the corresponding norms. Then the embeddings in (2.4) hold with constants independent of $t \in \mathbb{R}$. Let $J \subset \mathbb{R}$ be a closed interval. We further define on $E = E(J) := C_0(J, X)$ (the space of continuous functions, vanishing at infinity if J is unbounded) the multiplication operator $A(\cdot)$ by

$$(A(\cdot)f)(t) := A(t)f(t) \quad \text{for } t \in J, \quad D(A(\cdot)) := \{f \in E : f(t) \in D(A(t)), A(\cdot)f \in E\}.$$

It is clear that the operator $A(\cdot)$ is also sectorial. We can thus introduce the spaces

$$E_{\alpha} := E_{\alpha}^{A(\cdot)}, \quad E_{\alpha-1} := E_{\alpha-1}^{A(\cdot)}, \quad \text{and} \quad \hat{E} := \overline{D(A(\cdot))}$$

for $\alpha \in [0, 1]$, where $E_0 := E$ and $E_1 := D(A(\cdot))$. We observe that $E_{-1} \subseteq \prod_{t \in J} X_{-1}^t$ and that the extrapolated operator $A(\cdot)_{-1} : \hat{E} \rightarrow E_{-1}$ is given by $(A(\cdot)_{-1}f)(t) := A_{-1}(t)f(t)$ for $t \in J$ and $f \in E$. Further, $E_{\alpha-1}$ has the norm

$$\|f\|_{\alpha-1} := \sup_{r>0} \sup_{s \in J} \|r^{\alpha} R(r, A_{-1}(s) - \omega)f(s)\|.$$

Let (2.1) and (2.2) hold. Then there exists a constant $C = C(t_0) > 0$ such that

$$\|(\omega - A(t))^{\alpha} e^{\tau A(t)}\| \leq C\tau^{-\alpha}, \quad (2.6)$$

$$\|U(t, s)x\|_{\alpha}^t \leq C(t-s)^{\beta-\alpha} \|x\|_{\beta}^s, \quad (2.7)$$

$$\|U(t, s)(\omega - A(s))^{\theta} y\| \leq C(\mu - \theta)^{-1} (t-s)^{-\theta} \|y\|, \quad (2.8)$$

$$\|(\omega - A(s))^{\gamma} (R(\omega, A(s)) - R(\omega, A(t)))\| \leq C(t-s)^{\mu}, \quad (2.9)$$

$$\|(\omega - A(t))^{\gamma-1} - (\omega - A(s))^{\gamma-1}\| \leq C(t-s)^{\mu} \quad (2.10)$$

for all $t, s \in \mathbb{R}$ and $t_0 > 0$ with $0 < t - s \leq t_0$ and all $0 < \tau \leq t_0$, $0 \leq \beta \leq \alpha \leq 1$, $0 \leq \theta < \mu$, $0 \leq \gamma < \nu$, $x \in X_\beta^s$, and $y \in D((\omega - A(s))^\theta)$. Here, (2.6) is well known, (2.7) follows from [4, Theorem 2.3] by interpolation, and (2.8) was proved in [36, Theorem 2.1] in a slightly different setting, but the proof also works under the present assumptions. Finally, (2.9) and (2.10) are straightforward consequences of (2.1) and (2.2), cf. [34] and [35]. We state an easy consequence of (2.8) which is crucial for our work, see also Lemma A.1.

Proposition 2.1. *Assume that (2.1) and (2.2) hold and let $1 - \mu < \alpha < 1$ and $0 \leq \beta \leq 1$. Then the following assertions hold for $s < t \leq s + t_0$ and $t_0 > 0$ with constants possibly depending on t_0 .*

(i) *The operators $U(t, s)$ have continuous extensions $U_{\alpha-1}(t, s) : X_{\alpha-1}^s \rightarrow X$ satisfying*

$$\|U_{\alpha-1}(t, s)\|_{\mathcal{L}(X_{\alpha-1}^s, X)} \leq c(\alpha)(t-s)^{\alpha-1}, \quad (2.11)$$

and $U_{\alpha-1}(t, s)x = U_{\gamma-1}(t, s)x$ for $1 - \mu < \gamma < \alpha < 1$ and $x \in X_{\alpha-1}^s$.

(ii) *The map $\{(t, s) : t > s\} \ni (t, s) \mapsto U_{\alpha-1}(t, s)f(s) \in X$ is continuous for $f \in E_{\alpha-1}$.*

(iii) *For $x \in X_{\alpha-1}^s$ we have*

$$\|U_{\alpha-1}(t, s)x\|_\beta^t \leq c(\alpha)(t-s)^{\alpha-\beta-1}\|x\|_{\alpha-1}^s. \quad (2.12)$$

Proof. Let $s < t \leq s + t_0$. Due to (2.8), we can uniquely extend $U(t, s)$ to operators from $D((\omega - A_{-1})^{\alpha \pm \varepsilon})$ to X , with norms bounded by $c(t-s)^{\alpha-1 \pm \varepsilon}$, where $1 - \mu < \alpha \pm \varepsilon < 1$. Assertion (i) now follows by reiteration employing (2.4) and e.g. Theorem 1.2.15 and Proposition 2.2.15 in [24]. The map $\Phi : (t, s) \mapsto U_{\alpha-1}(t, s)f(s) \in X$ is continuous for $t > s$ if $f \in E$. For $f \in E_{\alpha-1}$, the continuity of Φ is shown by approximation using (2.11) and (2.5). Finally, (2.7) and (2.11) yield

$$\begin{aligned} \|U_{\alpha-1}(t, s)x\|_\beta^t &= \left\| U\left(t, \frac{1}{2}(t+s)\right) U_{\alpha-1}\left(\frac{1}{2}(t+s), s\right)x \right\|_\beta^t \\ &\leq 2^\beta C(t-s)^{-\beta} \left\| U_{\alpha-1}\left(\frac{1}{2}(t+s), s\right)x \right\| \leq c(\alpha)(t-s)^{\alpha-\beta-1}\|x\|_{\alpha-1}^s \end{aligned}$$

for $x \in X_{\alpha-1}^s$. \square

Exponential dichotomies are another important tool in our study, cf. [11, 24, 33, 34]. We recall that an evolution family $U(\cdot, \cdot)$ is said to have an *exponential dichotomy* in an interval $J \subset \mathbb{R}$ if there exists a family of projections $P(t) \in \mathcal{L}(X)$, $t \in J$, being strongly continuous with respect to t , and numbers $\delta, N > 0$ such that

$$\begin{aligned} (a) \quad & U(t, s)P(s) = P(t)U(t, s), \\ (b) \quad & U(t, s) : \mathcal{Q}(s)(X) \rightarrow \mathcal{Q}(t)(X) \text{ is invertible with inverse } \tilde{U}(s, t), \\ (c) \quad & \|U(t, s)P(s)\| \leq Ne^{-\delta(t-s)}, \\ (d) \quad & \|\tilde{U}(s, t)\mathcal{Q}(t)\| \leq Ne^{-\delta(t-s)}, \end{aligned} \quad (2.13)$$

for all $s, t \in J$ with $s \leq t$, where $Q(t) := I - P(t)$ is the ‘unstable projection.’ In the parabolic case one easily obtains regularity properties of the exponential dichotomy, see e.g. [34, Proposition 3.18]. For instance, $\|A(t)Q(t)\| \leq c(\eta)$ for $t \in J$, $t - \eta > \inf J$ and each $\eta > 0$ since $A(t)Q(t) = A(t)U(t, t - \eta)\tilde{U}(t - \eta, t)Q(t)$. In the next proposition we state some results concerning extrapolation spaces. We use the convention $\pm\infty + r = \pm\infty$ for $r \in \mathbb{R}$, and we set $J' = J \setminus \{\sup J\}$, i.e., $J = J'$ if J is unbounded from above. Moreover, we write $U_0(t, s) := U(t, s)$, $P_0(t) := P(t)$, and $Q_0(t) := Q(t)$, where $X_0^t = X$ by definition.

Proposition 2.2. *Assume that (2.1) and (2.2) hold and that $U(t, s)$ has an exponential dichotomy on an interval J . Let $\eta > 0$ and $1 - \mu < \alpha \leq 1$. Then the operators $P(t)$ and $Q(t)$ admit continuous extensions $P_{\alpha-1}(t): X_{\alpha-1}^t \rightarrow X_{\alpha-1}^t$ and $Q_{\alpha-1}(t): X_{\alpha-1}^t \rightarrow X$, respectively, for $t \in J'$; which are uniformly bounded for $t < \sup J - \eta$. Moreover, the following assertions hold for $t, s \in J'$ with $t \geq s$.*

- (a) $Q_{\alpha-1}(t)X_{\alpha-1}^t = Q(t)X$;
- (b) $U_{\alpha-1}(t, s)P_{\alpha-1}(s) = P_{\alpha-1}(t)U_{\alpha-1}(t, s)$;
- (c) $U_{\alpha-1}(t, s): Q_{\alpha-1}(s)(X_{\alpha-1}^s) \rightarrow Q_{\alpha-1}(t)(X_{\alpha-1}^t)$ is invertible with inverse $\tilde{U}_{\alpha-1}(s, t)$;
- (d) $\|U_{\alpha-1}(t, s)P_{\alpha-1}(s)x\| \leq N(\alpha, \eta) \max\{(t-s)^{\alpha-1}, 1\}e^{-\delta(t-s)}\|x\|_{\alpha-1}^s$ for $x \in X_{\alpha-1}^s$ and $s < t < \sup J - \eta$;
- (e) $\|\tilde{U}_{\alpha-1}(s, t)Q_{\alpha-1}(t)x\| \leq N(\alpha, \eta)e^{-\delta(t-s)}\|x\|_{\alpha-1}^t$ for $x \in X_{\alpha-1}^t$ and $s \leq t < \sup J - \eta$;
- (f) Let $J_0 \subset J'$ be a closed interval and $f \in E_{\alpha-1}(J_0)$. Then $P(\cdot)f \in E_{\alpha-1}(J_0)$ and $Q(\cdot)f \in C_0(J_0, X)$.

Proof. Let $t \in J$ such that $t + \eta < \sup J$, $1 - \beta < \theta < \mu$, and $x \in D((\omega - A(t))^\theta)$. The estimates (2.8) and (2.13)(d) imply that

$$\|Q(t)(\omega - A(t))^\theta x\| = \|\tilde{U}(t, t + \eta)Q(t + \eta)U(t + \eta, t)(\omega - A(t))^\theta x\| \leq c(\eta)\|x\|.$$

The embeddings (2.4) thus yield

$$\|Q(t)y\| \leq c(\eta)\|(\omega - A_{-1}(t))^{-\theta}y\| = c(\eta)\|(\omega - A_{-1}(t))^{1-\theta}y\|_{-1}^t \leq c(\eta)\|y\|_{\beta-1}^t \quad (2.14)$$

for all $y \in X$. Observe that (2.14) is true for $\alpha = \beta$, in particular. Taking $\beta < \alpha$ and using the remarks after (2.5) (with reversed roles of α and β), we see that $Q(t)$ has a uniformly bounded extension $Q_{\alpha-1}(t): X_{\alpha-1}^t \rightarrow X$ for $t < \sup J - \eta$. Then the operator $I - Q_{\alpha-1}(t) \in \mathcal{L}(X_{\alpha-1}^t)$ is a uniformly bounded extension of $P(t)$.

Assertion (a) is a consequence of the fact that $Q_{\alpha-1}(t)$ has values in X and that it is a projection. Assertion (b) follows from (2.13)(a) by approximation using (2.5) and (2.11). To show (c), let $y \in Q_{\alpha-1}(t)X_{\alpha-1}^t = Q(t)X$. Due to (2.13)(b), there is a unique vector $x \in Q(s)X = Q_{\alpha-1}(s)X_{\alpha-1}^s$ such that $y = U(t, s)x = U_{\alpha-1}(t, s)x$.

Let $t \geq s + 1$ and $x \in X_{\alpha-1}^s$. Using the exponential dichotomy of U and the estimate (2.11), we obtain

$$\|U_{\alpha-1}(t, s)P_{\alpha-1}(s)x\| = \|U(t, s+1)P(s+1)U_{\alpha-1}(s+1, s)x\| \leq ce^{-\delta(t-s)}\|x\|_{\alpha-1}^s.$$

If $0 \leq t - s \leq 1$, assertion (d) follows from (b) and (2.11). Assertion (e) is a consequence of (a), (2.13), and (2.14).

Let $f \in E_{\alpha-1}(J_0)$. Then there are $f_n \in C_0(J_0, X)$ converging to f in $E_{\beta-1}(J_0)$ for $\beta \in (1 - \mu, \alpha)$. Then $Q(\cdot)f_n$ converges in $C_0(J_0, X)$ to $Q_{\alpha-1}(\cdot)f$, whence (f) follows. \square

We further use the operator family

$$\Gamma_{\alpha-1}(t, s) = \begin{cases} U_{\alpha-1}(t, s)P_{\alpha-1}(s), & t \geq s, t, s \in J', \\ -\tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s), & t < s, t, s \in J'. \end{cases} \quad (2.15)$$

In some results we shall assume that $A(\cdot)$ is *asymptotically hyperbolic*, i.e., there are two operators $A_{-\infty} : D(A_{-\infty}) \rightarrow X$ and $A_{+\infty} : D(A_{+\infty}) \rightarrow X$ satisfying (2.1) and

$$\lim_{t \rightarrow -\infty} R(\omega, A(t)) = R(\omega, A_{-\infty}), \quad \lim_{t \rightarrow +\infty} R(\omega, A(t)) = R(\omega, A_{+\infty}) \quad (\text{in } \mathcal{L}(X)); \quad (2.16)$$

$$\sigma(A_{+\infty}) \cap i\mathbb{R} = \sigma(A_{-\infty}) \cap i\mathbb{R} = \emptyset. \quad (2.17)$$

Under assumptions (2.1), (2.2), (2.16), (2.17), there exists $T \geq 0$ such that $U(t, s)$ has exponential dichotomies in $(-\infty, -T]$ and in $[T, +\infty)$. For the interval $[T, +\infty)$, this has been shown in Theorem 4.3 of [34]. The proofs given there extend in a straightforward way to the interval $(-\infty, -T]$. The case of dense domains was treated before in [8] and, for a slightly stronger version of (2.16), in [32]. Moreover, we have

$$\dim Q(t)X = \dim Q_{+\infty}X, \quad t \geq T, \quad \dim Q(t)X = \dim Q_{-\infty}X, \quad t \leq -T, \quad (2.18)$$

by [34, Theorem 3.2], where $Q_{\pm\infty}$ are the projections for $A_{\pm\infty}$. Due to Proposition 2.2, our extrapolated evolution family $U_{\alpha-1}(t, s)$ has then exponential dichotomies in $(-\infty, -T)$ and in $[T, +\infty)$. From (2.18) and Proposition 2.2(a), we conclude that

$$\dim Q_{\alpha-1}(t)X_{\alpha-1}^t = \dim Q_{+\infty}X, \quad t \geq T, \quad \dim Q_{\alpha-1}(t)X_{\alpha-1}^t = \dim Q_{-\infty}X, \quad t < -T,$$

if (2.1), (2.2), (2.16), and (2.17) hold.

Remark 2.3. In the proof of Theorem 4.3 of [34], the projections $P(t)$ (for $t \geq T$ and $t \leq -T$, respectively) are obtained as the restriction of projections for a parabolic evolution family having an exponential dichotomy on $J = \mathbb{R}$. Hence, assumptions (2.1), (2.2), (2.16), and (2.17) imply that

$$U(\cdot, \cdot) \text{ has exponential dichotomies on } [T, +\infty) \text{ and } (-\infty, -T] \text{ for some } T \geq 0 \\ \text{and the assertions of Proposition 2.2 are true with } \eta = 0. \quad (2.19)$$

Definition 2.4. We assume that (2.1) and (2.2) hold, take $1 - \mu < \alpha \leq 1$, and let $J \subset \mathbb{R}$ be a closed interval. Let $f(t) \in X_{\alpha-1}^t$, $t \in J$, such that $f|_{[a, b]} \in E_{\alpha-1}([a, b])$ for all subintervals $[a, b] \subseteq J$. We say that $u \in C(J, X)$ is a *mild solution* of

$$u'(t) = A_{-1}(t)u(t) + f(t), \quad t \in J, \quad (2.20)$$

if the equation

$$u(t) = U(t, s)u(s) + \int_s^t U_{\alpha-1}(t, \sigma)f(\sigma) d\sigma \quad (2.21)$$

holds for all $t \geq s$ in J . If in addition $u \in E(J)$ and $f \in E_{\alpha-1}(J)$, then we write $u \in D(G_{\alpha-1})$ and $G_{\alpha-1}u = f$, where $G_0 =: G$. If u is a mild solution of (2.20) for $J = [t_0, +\infty)$, respectively $J = (-\infty, t_0]$, with $u(t_0) = x$, then we call u a *mild solution* of the initial, respectively final, value problems

$$u'(t) = A_{-1}(t)u(t) + f(t), \quad t \geq t_0, \quad u(t_0) = x; \quad \text{respectively,} \quad (2.22)$$

$$u'(t) = A_{-1}(t)u(t) + f(t), \quad t \leq t_0, \quad u(t_0) = x. \quad (2.23)$$

Remark 2.5. We make the assumptions stated in Definition 2.4. Then there always exists a unique mild solution of (2.22) with $u(t_0) = x \in \hat{X}^{t_0}$. Moreover, a function $u \in C(J, X)$ can be the mild solution of (2.20) for at most one f , so that $G_{\alpha-1}$ is single-valued. Finally, $G_{\alpha-1} : D(G_{\alpha-1}) \subset E(J) \rightarrow E_{\alpha-1}(J)$ is a closed linear operator.

Proof. The first assertion follows easily from Proposition 2.1. For the second assertion, take f and g such that $f(t), g(t) \in X_{\alpha-1}^t$ for $t \in J$, $f|_{[a, b]}, g|_{[a, b]} \in E_{\alpha-1}([a, b])$ for all subintervals $[a, b] \subseteq J$, and (2.21) holds for some $u \in C(J, X)$ and both f and g . Setting $h = f - g$, we thus obtain

$$\int_s^t U_{\alpha-1}(t, \sigma)h(\sigma) d\sigma = 0 \quad \forall t, s \in J \text{ with } t \geq s,$$

and hence $U_{\alpha-1}(t, s)h(s) = 0$ for all $t > s$ due to Proposition 2.1(ii). Take $\theta \in (1 - \nu, \mu)$ such that $\theta > 1 - \alpha$. Then $(\omega - A_{-1}(\cdot))^{-\theta}h \in \hat{E}([a, b])$ by (2.4), and thus Lemma A.1 yields $h = 0$, i.e., $f = g$. (We can take any $\bar{\alpha} \in (1 - \mu, 1 - \theta)$ when applying Lemma A.1. We point out that in the proof of this lemma we use no results established after Proposition 2.1.) The last assertion is a straightforward consequence of (2.11). \square

The next proposition shows that a mild solution of (2.20) is in fact a differentiable solution of (2.20) in a slightly weaker topology, see Appendix A for the proof. However, it is more convenient for us to work with the integral equation (2.21).

Proposition 2.6. Assume that (2.1) and (2.2) hold and that $f \in E_{\alpha-1}(J)$ for $1 - \mu < \alpha \leq 1$ and some closed interval $J \subset \mathbb{R}$. Let $u \in C(J, X)$ be a mild solution of (2.20) and let $0 \leq \beta < \min\{\alpha, \nu\}$. Then $u(t) \in X_{\beta}^t$, the map $s \mapsto u(s)$ is differentiable at $s = t$ in the norm of $X_{\beta-1}^t$, and (2.20) holds pointwise in $X_{\beta-1}^t$, for each $t \in J \setminus \inf J$.

Employing exponential dichotomies on halflines, we can derive existence results for forward and backward Cauchy problems with inhomogeneities in extrapolation spaces.

Proposition 2.7. Assume that (2.1) and (2.2) hold, $1 - \mu < \alpha \leq 1$, and that $U(t, s)$ has an exponential dichotomy on an interval $[T, +\infty)$. Let $t_0 \geq T$, $f \in E_{\alpha-1}([T, +\infty))$, and $x \in \bar{D}(A(t_0))$. Then the mild solution $u \in C([t_0, \infty), X)$ of (2.22) is bounded on $[t_0, \infty)$ if and only if

$$Q(t_0)x = - \int_{t_0}^{+\infty} \tilde{U}_{\alpha-1}(t_0, s) Q_{\alpha-1}(s) f(s) ds, \quad (2.24)$$

in which case u is given by

$$u(t) = U(t, t_0)P(t_0)x + \int_{t_0}^t U_{\alpha-1}(t, s)P_{\alpha-1}(s)f(s)ds - \int_t^{+\infty} \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)f(s)ds. \quad (2.25)$$

Proof. Let $t_0 \geq T$. The mild solution u of (2.22) satisfies

$$u(t) = U_{\alpha-1}(t, t_0)u(t_0) + \int_{t_0}^t U_{\alpha-1}(t, s)f(s)ds, \quad t \geq t_0.$$

Using Proposition 2.2 and (2.15), we can write this equality as

$$\begin{aligned} u(t) &= U(t, t_0)u(t_0) + \int_{t_0}^t U_{\alpha-1}(t, s)P_{\alpha-1}(s)f(s)ds - \int_t^{+\infty} \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)f(s)ds \\ &\quad + \int_{t_0}^{+\infty} \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)f(s)ds \\ &= U(t, t_0) \left[u(t_0) + \int_{t_0}^{+\infty} \tilde{U}_{\alpha-1}(t_0, s)Q_{\alpha-1}(s)f(s)ds \right] + \int_t^{+\infty} \Gamma_{\alpha-1}(t, s)f(s)ds \end{aligned} \quad (2.26)$$

for $t \geq t_0$. Proposition 2.2 and the boundedness of f on $[t_0, +\infty)$ show that u is bounded if and only if the term in brackets $[\dots]$ belongs to $P(t_0)X$ which is equivalent to (2.24). In this case, (2.25) follows directly from (2.26). \square

In the next proposition we may also take $t_0 = -T$ in the situation of Remark 2.3.

Proposition 2.8. Assume that (2.1) and (2.2) hold, $1 - \mu < \alpha \leq 1$, and that $U(t, s)$ has an exponential dichotomy on an interval $(-\infty, -T]$. Let $t_0 < -T$, $f \in E_{\alpha-1}((-\infty, t_0])$ and $x \in X$. Then there is a bounded mild solution $u \in C((-\infty, t_0], X)$ of (2.23) on $(-\infty, t_0]$ if and only if

$$P(t_0)x = \int_{-\infty}^{t_0} U_{\alpha-1}(t_0, s)P_{\alpha-1}(s)f(s)ds, \quad (2.27)$$

in which case u is given by

$$u(t) = \tilde{U}(t, t_0)Q(t_0)x - \int_t^{t_0} \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)f(s)ds + \int_{-\infty}^t U_{\alpha-1}(t, s)P_{\alpha-1}(s)f(s)ds. \quad (2.28)$$

Proof. It is straightforward to check that (2.28) gives a bounded mild solution of (2.23) satisfying (2.27). Let $t_0 < -T$, $s \leq t \leq t_0$, and let u be a bounded mild solution of (2.23). As in Proposition 2.7, we can write

$$\begin{aligned} u(t) = U(t, s) \left[P(s)u(s) - \int_{-\infty}^s U_{\alpha-1}(s, \tau)P_{\alpha-1}(\tau)f(\tau)d\tau \right] &+ \int_{-\infty}^t U_{\alpha-1}(t, \tau)P_{\alpha-1}(\tau)f(\tau)d\tau \\ &+ U(t, s)Q(s)u(s) + \int_s^t U_{\alpha-1}(t, \tau)Q_{\alpha-1}(\tau)f(\tau)d\tau. \end{aligned}$$

Since $U(t, s)Q(s)u(s) + \int_s^t U_{\alpha-1}(t, \tau)Q_{\alpha-1}(\tau)f(\tau)d\tau = Q(t)u(t)$, we have

$$\begin{aligned} P(t)u(t) = U(t, s)P(s) \left[P(s)u(s) - \int_{-\infty}^s U_{\alpha-1}(s, \tau)P_{\alpha-1}(\tau)f(\tau)d\tau \right] \\ + \int_{-\infty}^t U_{\alpha-1}(t, \tau)P_{\alpha-1}(\tau)f(\tau)d\tau. \end{aligned} \quad (2.29)$$

Due to Proposition 2.2, the boundedness of u and f implies that the term in $[\dots]$ is bounded for $s \leq t_0$. Therefore, letting $s \rightarrow -\infty$ in (2.29), we deduce from (2.13) that

$$P(t)u(t) = \int_{-\infty}^t U_{\alpha-1}(t, \tau)P_{\alpha-1}(\tau)f(\tau)d\tau, \quad (2.30)$$

and in particular the condition (2.27) for $t = t_0$. Moreover, it holds

$$\begin{aligned} Q(t_0)u(t_0) &= U(t_0, t)Q(t)u(t) + \int_t^{t_0} U_{\alpha-1}(t_0, \tau)Q_{\alpha-1}(\tau)f(\tau)d\tau, \\ Q(t)u(t) &= \tilde{U}(t, t_0)Q(t_0)u(t_0) - \int_t^{t_0} \tilde{U}_{\alpha-1}(t, \tau)Q_{\alpha-1}(\tau)f(\tau)d\tau. \end{aligned}$$

The last equation together with (2.30) yield the formula (2.28). \square

3. Properties of the operator $G_{\alpha-1}$

In this section we assume that the operators $A(t)$, $t \in \mathbb{R}$, on X satisfy the hypotheses (2.1), (2.2), and (2.19) (where the latter condition follows from (2.16) and (2.17)). Again, $U(t, s)$ is the evolution family on X generated by $A(\cdot)$ and $U_{\alpha-1}(t, s)$ is its extrapolated evolution family on $X_{\alpha-1}^s$. Both families have exponential dichotomies on $(-\infty, -T]$ and $[T, +\infty)$ for some $T \geq 0$ with projections $P(\cdot)$ and $P_{\alpha-1}(\cdot)$, respectively. To study the operator $G_{\alpha-1}$ on $J = \mathbb{R}$, we introduce the stable and unstable subspaces of $U_{\alpha-1}(\cdot, \cdot)$.

Definition 3.1. Let $t_0 \in \mathbb{R}$. We define the *stable space* at t_0 by

$$X_s(t_0) := \left\{ x \in X_{\alpha-1}^{t_0} : \lim_{t \rightarrow +\infty} \|U_{\alpha-1}(t, t_0)x\| = 0 \right\},$$

and the *unstable space* at t_0 by

$$X_u(t_0) := \{x \in X : \exists \text{ a mild solution } u \in C_0((-\infty, t_0], X) \text{ of (2.23) with } f = 0\}.$$

Observe that the function u in the definition of $X_u(t_0)$ satisfies $u(t) = U(t, s)u(s)$ for $s \leq t \leq t_0$ and $u(t_0) = x$.

Lemma 3.2. Assume that the assumptions (2.1), (2.2), and (2.19) are satisfied and that $1 - \mu < \alpha \leq 1$. Then the following assertions hold.

- (a) $X_s(t_0) = P_{\alpha-1}(t_0)X_{\alpha-1}^{t_0}$ for $t_0 \geq T$;
- (b) $X_u(t_0) = Q(t_0)X$ for $t_0 \leq -T$;
- (c) $U_{\alpha-1}(t, t_0)X_s(t_0) \subseteq X_s(t)$ for $t \geq t_0$ in \mathbb{R} ;
- (d) $U(t, t_0)X_u(t_0) = X_u(t)$ for $t \geq t_0$ in \mathbb{R} ;
- (e) $X_s(t_0)$ is closed in $X_{\alpha-1}^{t_0}$ for $t_0 \in \mathbb{R}$.

Proof. The inclusions ‘ \supseteq ’ in (a) and (b) are clear. Let $t \geq t_0 + 1 > t_0 \geq T$ and $x \in X_s(t_0)$. Due to Proposition 2.2, we obtain

$$\begin{aligned} c &\geq \|U_{\alpha-1}(t, t_0)x\| \geq \|U_{\alpha-1}(t, t_0)Q_{\alpha-1}(t_0)x\| - \|U_{\alpha-1}(t, t_0)P_{\alpha-1}(t_0)x\| \\ &\geq N^{-1}e^{\delta(t-t_0)}\|Q_{\alpha-1}(t_0)x\| - Ne^{-\delta(t-t_0)}\|P_{\alpha-1}(t_0)x\|_{\alpha-1}^{t_0}. \end{aligned}$$

Letting $t \rightarrow \infty$, this estimate implies that $Q_{\alpha-1}(t_0)x = 0$; i.e., (a) is verified. Let $t \leq t_0 - 1 < t_0 \leq -T$ and $x \in X_u(t_0)$. Let u be as in Definition 3.1. We then have $P(t_0)u(t_0) = U(t_0, t)P(t)u(t)$, and thus

$$\|P(t_0)x\| \leq Ne^{-\delta(t_0-t)}\|u(t)\| \leq ce^{\delta t}.$$

Letting $t \rightarrow -\infty$, we deduce $P(t_0)x = 0$ so that (b) holds. The assertions (c) and (d) are easy consequences of Definition 3.1. To show (e), let $t_0 \in \mathbb{R}$. If $t_0 \geq T$, the closeness of $X_s(t_0)$ in

$X_{\alpha-1}^{t_0}$ follows from (a). If $t_0 < T$, take $x_n \in X_s(t_0)$ such that $x_n \rightarrow x$ in $X_{\alpha-1}^{t_0}$. Then assertions (a) and (c) and estimate (2.11) imply that

$$\|U_{\alpha-1}(t, t_0)x\| = \lim_{n \rightarrow \infty} \|U(t, T)P(T)U_{\alpha-1}(T, t_0)x_n\| \leq cNe^{-\delta(t-T)}$$

for $t \geq T$. Thus $x \in X_s(t_0)$. \square

Let $1 - \mu < \alpha \leq 1$. The restrictions $G_{\alpha-1}^+$ and $G_{\alpha-1}^-$ of $G_{\alpha-1}$ to the halflines $[T, +\infty)$ and $(-\infty, T]$ are defined as in Definition 2.4: a function $u \in C_0([T, +\infty), X)$ (respectively $u \in C_0((-\infty, T], X)$) belongs to $D(G_{\alpha-1}^+)$ (respectively $D(G_{\alpha-1}^-)$) if there is a function $f \in E_{\alpha-1}([T, +\infty))$ (respectively $f \in E_{\alpha-1}((-\infty, T])$) such that

$$u(t) = U(t, s)u(s) + \int_s^t U_{\alpha-1}(t, \sigma)f(\sigma)d\sigma$$

holds for all $t \geq s \geq T$ (respectively for all $s \leq t \leq T$).

As in [16] and [17], we introduce on $E_{\alpha-1}([T, +\infty))$ and on $E_{\alpha-1}((-\infty, T])$ the right inverses $R_{\alpha-1}^+$ and $R_{\alpha-1}^-$ for $G_{\alpha-1}^+$ and $G_{\alpha-1}^-$, respectively, by setting

$$\begin{aligned} (R_{\alpha-1}^+h)(t) &= -\int_t^\infty \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)h(s)ds + \int_T^t U_{\alpha-1}(t, s)P_{\alpha-1}(s)h(s)ds, \quad t \geq T, \\ (R_{\alpha-1}^-h)(t) &= \begin{cases} \int_{-\infty}^t U_{\alpha-1}(t, s)P_{\alpha-1}(s)h(s)ds - \int_t^{-T} \tilde{U}_{\alpha-1}(t, s)Q_{\alpha-1}(s)h(s)ds, & t \leq -T, \\ \int_{-\infty}^{-T} U_{\alpha-1}(t, s)P_{\alpha-1}(s)h(s)ds + \int_{-T}^t U_{\alpha-1}(t, s)h(s)ds, & -T \leq t \leq T. \end{cases} \end{aligned}$$

Proposition 3.3. Assume that the assumptions (2.1), (2.2), and (2.19) are satisfied and that $1 - \mu < \alpha \leq 1$. Then the following assertions hold.

- (a) $R_{\alpha-1}^+ : E_{\alpha-1}([T, +\infty)) \rightarrow C_0([T, +\infty), X)$ is bounded and $G_{\alpha-1}^+ R_{\alpha-1}^+ h = h$ for each $h \in E_{\alpha-1}([T, +\infty))$.
- (b) $R_{\alpha-1}^- : E_{\alpha-1}((-\infty, T]) \rightarrow C_0((-\infty, T], X)$ is bounded and $G_{\alpha-1}^- R_{\alpha-1}^- h = h$ for each $h \in E_{\alpha-1}((-\infty, T])$.

Proof. (a) Proposition 2.2 shows that $\sup_{t \geq T} \|R_{\alpha-1}^+ h(t)\|_\infty \leq c\|h\|_{\alpha-1}$ for a constant $c > 0$ and $h \in E_{\alpha-1}$. Moreover, $R_{\alpha-1}^+ h \in C_0([T, +\infty), X)$ if $h \in C_0([T, +\infty), X)$. For $h \in E_{\alpha-1}([T, +\infty))$ and $1 - \mu < \beta < \alpha$, there are $h_n \in C_0([T, +\infty), X)$ converging to h in $E_{\beta-1}([T, +\infty))$ due to (2.5). Therefore $R_{\beta-1}^+ h_n \rightarrow R_{\beta-1}^+ h = R_{\alpha-1}^+ h$ in $C_b([T, +\infty), X)$ (the space of bounded continuous functions), and the first part of (a) is shown. For $t \geq s \geq T$, we further compute

$$\begin{aligned} &U(t, s)R_{\alpha-1}^+ h(s) + \int_s^t U_{\alpha-1}(t, \tau)h(\tau)d\tau \\ &= \int_s^t U_{\alpha-1}(t, \tau)h(\tau)ds - \int_s^{+\infty} \tilde{U}_{\alpha-1}(t, \tau)Q_{\alpha-1}(\tau)h(\tau)d\tau + \int_T^s U_{\alpha-1}(t, \tau)P_{\alpha-1}(\tau)h(\tau)d\tau \end{aligned}$$

$$\begin{aligned}
&= \int_T^t U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau - \int_t^{+\infty} \tilde{U}_{\alpha-1}(t, \tau) Q_{\alpha-1}(\tau) h(\tau) d\tau \\
&= R_{\alpha-1}^+ h(t).
\end{aligned}$$

Hence, $R_{\alpha-1}^+ h \in D(G_{\alpha-1}^+)$ and $G_{\alpha-1}^+ R_{\alpha-1}^+ h = h$.

(b) The first part of (b) follows similarly as in (a). For $h \in E_{\alpha-1}((-\infty, T])$ and $s \leq t \leq -T$, we calculate

$$\begin{aligned}
&\int_s^t U_{\alpha-1}(t, \tau) h(\tau) d\tau + U(t, s) R_{\alpha-1}^- h(s) \\
&= \int_s^t U_{\alpha-1}(t, \tau) h(\tau) ds + \int_{-\infty}^s U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau - \int_s^{-T} \tilde{U}_{\alpha-1}(t, \tau) Q_{\alpha-1}(\tau) h(\tau) d\tau \\
&= \int_{-\infty}^t U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau - \int_t^{-T} \tilde{U}_{\alpha-1}(t, \tau) Q_{\alpha-1}(\tau) h(\tau) d\tau = R_{\alpha-1}^- h(t).
\end{aligned}$$

For $s \leq -T \leq t$, it holds

$$\begin{aligned}
&\int_s^t U_{\alpha-1}(t, \tau) h(\tau) d\tau + U(t, s) R_{\alpha-1}^- h(s) \\
&= \int_s^t U_{\alpha-1}(t, \tau) h(\tau) ds + \int_{-\infty}^s U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau - \int_s^{-T} \tilde{U}_{\alpha-1}(t, \tau) Q_{\alpha-1}(\tau) h(\tau) d\tau \\
&= \int_{-\infty}^{-T} U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau + \int_{-T}^t U_{\alpha-1}(t, \tau) h(\tau) d\tau \\
&= R_{\alpha-1}^- h(t).
\end{aligned}$$

Finally, for $-T \leq s \leq t \leq T$, we compute

$$\begin{aligned}
&\int_s^t U_{\alpha-1}(t, \tau) h(\tau) d\tau + U(t, s) R_{\alpha-1}^- h(s) \\
&= \int_s^t U_{\alpha-1}(t, \tau) h(\tau) ds + \int_{-\infty}^{-T} U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau + \int_{-T}^s U_{\alpha-1}(t, \tau) h(\tau) d\tau
\end{aligned}$$

$$\begin{aligned}
&= \int_{-\infty}^{-T} U_{\alpha-1}(t, \tau) P_{\alpha-1}(\tau) h(\tau) d\tau + \int_{-T}^t U_{\alpha-1}(t, \tau) h(\tau) d\tau \\
&= R_{\alpha-1}^- h(t).
\end{aligned}$$

As a result, $R_{\alpha-1}^- h \in D(G_{\alpha-1}^-)$ and $G_{\alpha-1}^- R_{\alpha-1}^- h = h$. \square

Lemma 3.4. Assume that (2.1), (2.2), and (2.19) hold and that $1 - \mu < \alpha \leq 1$. Let $x \in Q(T)(X)$. Then there exists $u \in D(G_{\alpha-1})$ such that $R_{\alpha-1}^+ u(T) = x$, $R_{\alpha-1}^- u(T) = 0$, and $\|u\|_E + \|G_{\alpha-1} u\|_E \leq K \|x\|$, where $K \geq 0$ is a constant independent of x .

Proof. We fix a test function φ with $\varphi(t) = 0$ for $t \leq T$ and $\int_T^\infty \varphi(s) ds = -1$, and define the functions

$$\begin{aligned}
u(t) &:= \varphi(t) U(t, T)x, & t \geq T, & & u(t) &:= 0, & t \leq T, \\
f(t) &:= \varphi'(t) U(t, T)x, & t \geq T, & & f(t) &:= 0, & t < T.
\end{aligned}$$

It is easy to check that $R_{\alpha-1}^+ u(T) = x$ and $R_{\alpha-1}^- u(T) = 0$. We further obtain

$$\begin{aligned}
u(t) - U(t, s)u(s) &= (\varphi(t) - \varphi(s))U(t, T)x \\
&= \int_s^t U_{\alpha-1}(t, \tau) \varphi'(\tau) U(\tau, T)x d\tau = \int_s^t U_{\alpha-1}(t, \tau) f(\tau) d\tau
\end{aligned}$$

for $t \geq s \geq T$. The case $s < T$ is treated similarly. As a result, $u \in D(G_{\alpha-1})$ and $G_{\alpha-1} u = f$, so that the asserted estimate follows. \square

We can now describe the range and the kernel of $G_{\alpha-1}$.

Proposition 3.5. Assume that (2.1), (2.2), and (2.19) are satisfied and that $1 - \mu < \alpha \leq 1$. Then the following assertions hold for $G_{\alpha-1}$ defined on $E_{\alpha-1} = E_{\alpha-1}(\mathbb{R})$.

- (a) $N(G_{\alpha-1}^+) = \{u \in C_0([T, +\infty), X): u(t) = U(t, T)x \ \forall t \geq T, \ x \in P(T)\hat{X}^T\}$;
- (b) $N(G_{\alpha-1}^-) = \{u \in C_0((-\infty, T]): u(t) = U(t, s)u(s) \ \forall s \leq t \leq T, \ u(T) \in X_u(T)\}$;
- (c) $N(G_{\alpha-1}) = \{u \in C_0(\mathbb{R}, X): u(t) = U(t, s)u(s) \ \forall t \geq s, \ u(T) \in P(T)X \cap X_u(T)\}$;
- (d) $R(G_{\alpha-1}) = \{f \in E_{\alpha-1}: R_{\alpha-1}^+ f(T) - R_{\alpha-1}^- f(T) \in P(T)X + X_u(T)\}$, where for $f \in R(G_{\alpha-1})$ a function $u \in D(G_{\alpha-1})$ with $G_{\alpha-1} u = f$ is given by (3.1) below;
- (e) $\overline{R(G_{\alpha-1})} = \{f \in E_{\alpha-1}: R_{\alpha-1}^+ f(T) - R_{\alpha-1}^- f(T) \in \overline{P(T)X + X_u(T)}\}$, where the closure on the left- (right-) hand side is taken in $E_{\alpha-1}$ (in X).

Proof. Assertions (a), (b) and (c) follow from Lemma 3.2 and $P(T)X \cap X_u(T) = P(T)\hat{X}^T \cap X_u(T)$. To show (d), let $G_{\alpha-1} u = f \in E_{\alpha-1}$ for some $u \in D(G_{\alpha-1})$. Then the restrictions of

f to $[T, +\infty)$ and $(-\infty, T]$ belong to $R(G_{\alpha-1}^+)$ and to $R(G_{\alpha-1}^-)$, respectively. Proposition 3.3 shows that the functions

$$v_+ = (u|[T, +\infty)) - R_{\alpha-1}^+ f \quad \text{and} \quad v_- = (u|(-\infty, T]) - R_{\alpha-1}^- f$$

belong to the kernel of $G_{\alpha-1}^+$ and $G_{\alpha-1}^-$, respectively. Thus

$$(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T) = v_-(T) - v_+(T) \in X_u(T) + P(T)\hat{X}^T$$

by (a) and (b). Conversely, let $f \in E_{\alpha-1}$ with $(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T) = y_s + y_u \in P(T)X + X_u(T)$. Set $x_0 := (R_{\alpha-1}^+ f)(T) - y_s = y_u + (R_{\alpha-1}^- f)(T)$ and

$$u(t) := \begin{cases} u_+(t) := -U(t, T)y_s + (R_{\alpha-1}^+ f)(t), & t \geq T, \\ u_-(t) := \tilde{v}(t) + (R_{\alpha-1}^- f)(t), & t \leq T, \end{cases} \quad (3.1)$$

where $\tilde{v} \in N(G_{\alpha-1}^-)$ such that $\tilde{v}(T) = y_u$. Using Propositions 2.1 and 2.2, one checks that $R_{\alpha-1}^\pm f(T) \in X_\varepsilon^T$ for $0 < \varepsilon < \alpha$, so that $y_s \in X_\varepsilon^t \subseteq \hat{X}^T$. Hence, $u \in C_0(\mathbb{R}, X)$. Proposition 3.3 further yields

$$u_\pm(t) = U(t, s)u_\pm(s) + \int_s^t U_{\alpha-1}(t, \tau)f(\tau) d\tau$$

for all $t \geq s \geq T$ and $s \leq t \leq T$, respectively. Let now $s \leq T \leq t$. Since $u_+(T) = u_-(T) = x_0$, we have

$$\begin{aligned} u(t) &= u_+(t) = U(t, T)u_-(T) + \int_T^t U_{\alpha-1}(t, \tau)f(\tau) d\tau \\ &= U(t, T) \left[U(T, s)u_-(s) + \int_s^T U_{\alpha-1}(T, \tau)f(\tau) d\tau \right] + \int_T^t U_{\alpha-1}(t, \tau)f(\tau) d\tau \\ &= U(t, s)u(s) + \int_s^t U_{\alpha-1}(t, \tau)f(\tau) d\tau. \end{aligned}$$

Therefore $G_{\alpha-1}u = f$, and (d) is established.

The inclusion ‘ \subseteq ’ in assertion (e) follows from (d) and Proposition 3.3. Take $f \in E_{\alpha-1}$ and $z := (R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T)$ such that there is a sequence $z_n \in P(T)X + X_u(T)$ converging to z in X as $n \rightarrow \infty$. Set $y_n := z - z_n$ and $x_n := Q(T)y_n$. Lemma 3.4 yields a function $f_n \in D(G_{\alpha-1})$ such that $(R_{\alpha-1}^+ f_n)(T) = x_n$, $(R_{\alpha-1}^- f_n)(T) = 0$, and $\|f_n\|_E \leq K\|x_n\|$ for a constant K independent of n . Then the vector

$$(R_{\alpha-1}^+(f - f_n))(T) - (R_{\alpha-1}^-(f - f_n))(T) = z - x_n = z_n + P(T)y_n$$

belongs to $P(T)X + X_u(T)$, so that $f - f_n \in R(G_{\alpha-1})$ by (d). Since $E \hookrightarrow E_{\alpha-1}$, we can estimate

$$\|f - (f - f_n)\|_{E_{\alpha-1}} \leq c \|f_n\|_E \leq cK \|x_n\| \leq c \|z - z_n\|,$$

and thus assertion (e) is shown. \square

Using the above results, we are able to describe other properties of the operator $G_{\alpha-1}$, in particular its Fredholmity, in terms of properties of the subspaces $X_s(T)$ and $X_u(T)$, using similar arguments as in [16], see also [17] for L^p spaces. For the convenience of the readers, we give the complete proof. Recall that subspaces V and W of a Banach space E are called a *semi-Fredholm couple* if $V + W$ is closed and if at least one of the dimensions $\dim(V \cap W)$ and $\operatorname{codim}(V + W)$ is finite. The *index* of (V, W) is defined by $\operatorname{ind}(V, W) := \dim(V \cap W) - \operatorname{codim}(V + W)$. If the index is finite, then (V, W) is a *Fredholm couple*. Observe that in the next theorem the operator $U(T, -T)|_{Q(-T)(X)}$ is trivially injective if $T = 0$.

Theorem 3.6. Assume that (2.1), (2.2), and (2.19) are satisfied and that $1 - \mu < \alpha \leq 1$. Then the following assertions hold for $G_{\alpha-1}$ defined on $E_{\alpha-1} = E_{\alpha-1}(\mathbb{R})$.

- (a) $R(G_{\alpha-1})$ is closed in $E_{\alpha-1}$ if and only if $P(T)X + X_u(T)$ is closed in X .
- (b) $G_{\alpha-1}$ is surjective if and only if $P(T)X + X_u(T) = X$.
- (c) If $G_{\alpha-1}$ is injective, then $P(T)X \cap X_u(T) = \{0\}$. The converse is true if $U(T, -T)|_{Q(-T)(X)}$ is injective, in addition.
- (d) If $G_{\alpha-1}$ is invertible, then $P(T)X \oplus X_u(T) = X$. The converse is true if $U(T, -T)|_{Q(-T)(X)}$ is injective, in addition.
- (e) $\dim N(G_{\alpha-1}) = \dim(P(T)X \cap X_u(T)) + \dim N(U(T, -T)|_{Q(-T)(X)})$.
If $R(G_{\alpha-1})$ is closed in $E_{\alpha-1}$, then $\operatorname{codim}(P(T)X + X_u(T)) = \operatorname{codim} R(G_{\alpha-1})$.
- (f) If $G_{\alpha-1}$ is a semi-Fredholm operator, then $(P(T)X, X_u(T))$ is a semi-Fredholm couple, and $\operatorname{ind}(P(T)X, X_u(T)) \leq \operatorname{ind} G_{\alpha-1}$. If in addition the kernel of $U(T, -T)|_{Q(-T)(X)}$ is finite-dimensional, then

$$\operatorname{ind}(P(T)X, X_u(T)) = \operatorname{ind} G_{\alpha-1} - \dim N(U(T, -T)|_{Q(-T)(X)}). \quad (3.2)$$

Conversely, if $(P(T)X, X_u(T))$ is a semi-Fredholm couple and the kernel of $U(T, -T)|_{Q(-T)(X)}$ is finite-dimensional, then $G_{\alpha-1}$ is a semi-Fredholm operator and (3.2) holds.

Proof. The ‘if’ part of assertion (a) is a direct consequence of Proposition 3.5(d) and (e). Assume that $R(G_{\alpha-1})$ is closed in $E_{\alpha-1}$. Take $y_n^s \in P(T)X$ and $y_n^u \in X_u(T)$ with $y_n^s + y_n^u \rightarrow y$ in X as $n \rightarrow \infty$. Set $z = Q(T)y$. By Lemma 3.4, there is a function $h \in D(G_{\alpha-1})$ such that $R_{\alpha-1}^+ h(T) = z$ and $R_{\alpha-1}^- h(T) = 0$. Since

$$z = \lim_{n \rightarrow \infty} Q(T)(y_n^s + y_n^u) = \lim_{n \rightarrow \infty} Q(T)y_n^u = \lim_{n \rightarrow \infty} (y_n^u - P(T)y_n^u),$$

we obtain $R_{\alpha-1}^+ h(T) - R_{\alpha-1}^- h(T) = z \in \overline{P(T)X + X_u(T)}$. Proposition 3.5 implies that $h \in \overline{R(G_{\alpha-1})} = R(G_{\alpha-1})$, and thus $z \in P(T)X + X_u(T)$ by Proposition 3.5(d). As a result, $y =$

$P(T)y + z \in P(T)X + X_u(T)$, and so (a) holds. The ‘if’ part of assertion (b) follows from Proposition 3.5(d), and the converse can be shown as in statement (a).

Proposition 3.5(c) yields the first part of (c). For the converse, assume that $U(T, -T)|_{Q(-T)(X)}$ is injective and $P(T)X \cap X_u(T) = \{0\}$, and let $u \in N(G_{\alpha-1})$. Then $u(t) = U(t, s)u(s)$ for all $t \geq s$, and so $u(T) = 0$ by Proposition 3.5(c). From Lemma 3.2(b) we further deduce $u(-T) \in Q(-T)(X)$. Since $0 = u(T) = U(T, -T)|_{Q(-T)(X)}u(-T)$, our assumption yields $u(-T) = 0$ and thus $u(t) = 0$ for $t \geq -T$. Finally, $u(t) = \tilde{U}(t, -T)u(-T)$ for all $t \leq -T$ by Proposition 2.8, so that $u = 0$. We have thus shown (c). Assertion (d) is an easy consequence of (b) and (c).

To show the first equality in (e), we define $\Gamma := \{u \in N(G_{\alpha-1}): u(t) = 0, t \geq T\}$ and the linear mapping

$$K : N(G_{\alpha-1})/\Gamma \rightarrow P(T)X \cap X_u(T), \quad [u] \mapsto u(T).$$

Proposition 3.5(c) implies that K is well defined and bijective. Since also $\dim \Gamma = \dim N(U(T, -T)|_{Q(-T)(X)})$, the first identity holds. We next assume that $R(G_{\alpha-1})$ is closed in $E_{\alpha-1}$. Hence, $P(T)X + X_u(T)$ is closed X by (a). Define the linear map

$$J : E_{\alpha-1}/R(G_{\alpha-1}) \rightarrow X/(P(T)X + X_u(T)), \quad [f] \mapsto [(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T)].$$

Due to Proposition 3.5(d), J is well defined and injective. Take $x \in X$. By Lemma 3.4 there is a function $f \in E_{\alpha-1}$ such that $(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T) = Q(T)x = x - P(T)x$. Hence, $J[f] = [(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T)] = [x]$. Consequently, J is also surjective and thus the second equality in (e) follows. Assertion (f) is a consequence of (a) and (e). \square

Using (2.18) and the same arguments as in [16], we obtain the following sufficient condition for the Fredholmity of $G_{\alpha-1}$.

Corollary 3.7. *Assume that (2.1), (2.2), (2.16), and (2.17) are satisfied and that $1 - \mu < \alpha \leq 1$. Further suppose that $\dim Q_{\pm\infty}X < \infty$ (which holds if $D(A_{\pm\infty})$ are compactly embedded in X). Then $G_{\alpha-1}$ is Fredholm and $\text{ind } G_{\alpha-1} = \dim Q_{-\infty}X - \dim Q_{+\infty}X$.*

We next characterize the range of $G_{\alpha-1}$ in terms of the dual problem, see Remark 3.9 below. Related results have been shown in [17] and [21] for other settings by different methods. We start with a simple observation. Let $0 \leq \theta < \alpha \leq 1$. Then $X_{\alpha-1}^t$ is densely embedded in $D((\omega - A_{-1}(t))^\theta)$ by (2.4). Since $D((\omega - A(t)^*)^{1-\theta}) \hookrightarrow [D((\omega - A_{-1}(t))^\theta)]^*$, we deduce that $D((\omega - A(t)^*)^{1-\theta}) \hookrightarrow (X_{\alpha-1}^t)^*$ for $t \in \mathbb{R}$ with a uniform embedding constant. We denote by \mathcal{V} the space of $v \in C(\mathbb{R}, X^*)$ such that $v(s) = U_{\alpha-1}(t, s)^*v(t)$, $v(t) \in D((\omega - A(t)^*)^{1-\theta})$, and $(\omega - A(\cdot)^*)^{1-\theta}v \in L^1(\mathbb{R}, X^*)$ for all $t \geq s$ in \mathbb{R} .

Proposition 3.8. *Assume that (2.1), (2.2), and (2.19) hold and that $1 - \mu < \theta < \alpha \leq 1$. Then the closure of $R(G_{\alpha-1})$ is equal to the space*

$$\mathcal{E} := \left\{ f \in E_{\alpha-1}: \int_{\mathbb{R}} \langle f(s), v(s) \rangle_{X_{\alpha-1}^s} ds = 0 \text{ for all } v \in \mathcal{V} \right\}.$$

Proof. We first show that under our assumptions it holds

$$\mathcal{V} = \{v \in L^1(\mathbb{R}, X^*): v(s) = U_{\alpha-1}(t, s)^* v(t) \forall t \geq s\} =: \mathcal{V}'. \quad (3.3)$$

Clearly, $\mathcal{V} \subset \mathcal{V}'$. Take $v \in \mathcal{V}'$. Then $v \in C(\mathbb{R}, X^*)$ since $U(t, s)$ is norm continuous for $t > s$. We denote by $V(t, s)$ the extension of $U(t, s)(\omega - A(s))^{1-\theta}$ to $\mathcal{L}(X)$. For $x \in D((\omega - A(s))^{1-\theta})$, we then obtain

$$\begin{aligned} \langle (\omega - A(s))^{1-\theta} x, v(s) \rangle &= \langle (\omega - A(s))^{1-\theta} x, U_{\alpha-1}(s+1, s)^* v(s+1) \rangle \\ &= \langle V(s+1, s)x, v(s+1) \rangle, \\ |\langle (\omega - A(s))^{1-\theta} x, v(s) \rangle| &\leq c \|v(s+1)\|_{X^*} \|x\| \end{aligned} \quad (3.4)$$

due to (2.8). The estimate (3.4) yields

$$v(s) \in D((\omega - A(s)^*)^{1-\theta}) \quad \text{and} \quad \|(\omega - A(s)^*)^{1-\theta} v(s)\|_{X^*} \leq c \|v(s+1)\|_{X^*}.$$

Thus $v \in \mathcal{V}$ and (3.3) is true. We now come to the main part of the proof. Proposition 3.5 shows that

$$f \in \overline{R(G_{\alpha-1})} \iff z := R_{\alpha-1}^+ f(T) - R_{\alpha-1}^- f(T) \in \overline{P(T)X + X_u(T)}.$$

Employing also [28, Theorem 4.7] and [20, (IV.4.11)], we deduce

$$\begin{aligned} f \in \overline{R(G_{\alpha-1})} &\iff z \in {}^\perp((P(T)X + X_u(T))^\perp) \\ &\iff z \in {}^\perp((P(T)X)^\perp \cap (X_u(T))^\perp) \end{aligned}$$

where $M^\perp := \{x^* \in X^*: \langle x, x^* \rangle = 0 \forall x \in M\}$ for $M \subseteq X$ and ${}^\perp N := \{x \in X: \langle x, x^* \rangle = 0 \forall x^* \in N\}$ for $N \subseteq X^*$. Straightforward duality arguments imply that $U(t, s)^*$ has an exponential dichotomy on $[T, +\infty)$ and $(-\infty, -T]$ with projections $P(t)^*$ and that

$$(P(T)X)^\perp = Q(T)^* X^*, \quad (X_u(T))^\perp = N(Q(-T)^* U(T, -T)^*), \quad (3.5)$$

using also $X_u(T) = U(T, -T)Q(-T)X$, see Lemma 3.2. We further compute

$$\begin{aligned} \langle z, y^* \rangle &= - \int_T^{+\infty} \langle \tilde{U}_{\alpha-1}(T, s) Q_{\alpha-1}(s) f(s), y^* \rangle_X ds - \int_{-T}^T \langle U_{\alpha-1}(T, s) f(s), y^* \rangle_X ds \\ &\quad - \int_{-\infty}^{-T} \langle U_{\alpha-1}(T, s) P_{\alpha-1}(s) f(s), y^* \rangle_X ds \\ &= - \int_{\mathbb{R}} \langle f(s), v(s) \rangle_{X_{\alpha-1}^s} ds \end{aligned}$$

for all $y^* \in (P(T)X)^\perp \cap (X_u(T))^\perp$, where v is given by

$$v(s) := \begin{cases} \tilde{U}(T, s)^* Q(T)^* y^* = \tilde{U}(T, s)^* y^*, & s \geq T, \\ U(T, s)^* y^*, & -T \leq s \leq T, \\ U(-T, s)^* P(-T)^* U(T, -T)^* y^* = U(T, s)^* y^*, & s \leq -T. \end{cases} \quad (3.6)$$

(Here we have used (3.5).) Summing up, we have shown that $f \in \overline{R(G_{\alpha-1})}$ if and only if

$$\int_{\mathbb{R}} \langle f(s), v(s) \rangle_{X_{\alpha-1}^s} ds = 0$$

for all v as in (3.6) with $y^* \in Q(T)^* X^* \cap N(Q(-T)^* U(T, -T)^*)$. It remains to show that \mathcal{V} consists precisely of the functions defined in (3.6).

First, one verifies by a duality argument that each function v in (3.6) belongs to $\mathcal{V}' = \mathcal{V}$, recall (3.3). Conversely, let $v \in \mathcal{V}$. Then we have

$$P(T)^* v(T) = U(t, T)^* P(t)^* v(t), \quad \|P(T)^* v(T)\| \leq N e^{-\delta(t-T)} \|v(t)\|$$

for $t \geq T$. There is a sequence $t_n \rightarrow \infty$ such that $\|v(t_n)\|$ is bounded since $v \in L^1(\mathbb{R}, X^*)$. Therefore, $P(T)^* v(T) = 0$. For $s \leq -T$, one obtains

$$Q(s)^* v(s) = U(-T, s)^* Q(-T)^* v(-T) = U(-T, s)^* Q(-T)^* U(T, -T)^* v(T), \quad (3.7)$$

$$\|Q(-T)^* U(T, -T)^* v(T)\| = \|\tilde{U}(s, -T)^* Q(s)^* v(s)\| \leq N e^{-\delta(-T-s)} \|v(s)\|. \quad (3.8)$$

As above, it follows that $Q(-T)^* U(T, -T)^* v(T) = 0$. Consequently, v is of the form (3.6) with $y^* = v(T) \in Q(T)^* X^* \cap N(Q(-T)^* U(T, -T)^*)$. \square

Remark 3.9. One can see that the functions $v \in \mathcal{V}$, see (3.3), solve the dual evolution equation

$$-v'(s) = A(s)^* v(s), \quad s \in \mathbb{R}, \quad (3.9)$$

in a weak sense. The function v is a classical solution of (3.9) if also the adjoint operators $A(t)^*$ satisfy the Acquistapace–Terreni conditions (2.1) and (2.2), see [2, Proposition 2.9].

Theorem 3.6, Corollary 3.7, and Propositions 3.5 and 3.8 now yield the following Fredholm alternative, where we focus on a simplified setting.

Theorem 3.10. Assume that (2.1), (2.2), (2.16) and (2.17) are true, that $\dim Q_{\pm\infty} X < \infty$, and that $1 - \mu < \alpha \leq 1$. Let $f \in E_{\alpha-1} = E_{\alpha-1}(\mathbb{R})$. Then there is a mild solution $u \in C_0(\mathbb{R}, X)$ of (2.20) if and only if

$$\int_{\mathbb{R}} \langle f(s), w(s) \rangle_{X_{\alpha-1}^s} ds = 0$$

for each $w \in L^1(\mathbb{R}, X^*)$ with $w(s) = U_{\alpha-1}(t, s)^* w(t)$ for all $t \geq s$. The mild solutions u are given by

$$\begin{aligned} u(t) &= v(t) - U(t, T)y_s + (R_{\alpha-1}^+ f)(t), \quad t \geq T, \\ u(t) &= v(t) + \tilde{v}(t) + (R_{\alpha-1}^- f)(t), \quad t \leq T, \end{aligned}$$

where $R_{\alpha-1}^\pm$ were defined before Proposition 3.3, $(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T) = y_s + y_u \in P(T)X + X_u(T)$, $\tilde{v} \in C_0((-\infty, T], X)$ with $\tilde{v}(T) = y_u$ and $\tilde{v}(t) = U(t, s)\tilde{v}(s)$ for all $T \geq t \geq s$, and $v \in C_0(\mathbb{R}, X)$ with $v(t) = U(t, s)v(s)$ for all $t \geq s$.

Note that in the above result we obtain mild solutions which are unique modulo the finite-dimensional subspace $N(G_{\alpha-1})$. We further remark that if $U(\cdot, \cdot)$ has an exponential dichotomy on \mathbb{R} with projections $P(t)$, $t \in \mathbb{R}$, then we can take $T = 0$ and we have $X_u(0) = (I - P(0))X$. Hence, $G_{\alpha-1}$ is invertible by Theorem 3.6(d). As a result, for each $f \in E_{\alpha-1}$ we obtain a unique mild solution of $u \in C_0(\mathbb{R}, X)$ of (2.20) which is given by

$$u(t) = \int_{\mathbb{R}} \Gamma_{\alpha-1}(t, \tau) f(\tau) d\tau, \quad t \in \mathbb{R},$$

due to (3.1); cf. [11] for this formula in the case $\alpha = 1$. We conclude this section with two remarks indicating straightforward variants of the results established so far. The details are left to the reader.

Remark 3.11. Note that we allow for the case $\alpha = 1$, i.e., $G_0 = G$ on $E_0 = E = C_0(\mathbb{R}, X)$, in this section. In fact, in this case the results shown in this section remain valid for each exponentially bounded evolution family $U(t, s)$, $t \geq s$ (i.e., (2.3) holds) such that $(t, s) \mapsto U(t, s)$ is strongly continuous for $t \geq s$ and $U(\cdot, \cdot)$ has exponential dichotomies on halflines $(-\infty, -T]$ and $[T, +\infty)$. (Here one sets $\hat{X}^t = X$.)

Remark 3.12. All results established in this and the previous section remain valid with the slightly simplified proofs if we replace the function spaces $C_0(J, X)$ by $C_b(J, X)$ in the assertions and in Definitions 2.4 and 3.1 and set $X_s(t_0) = \{x \in X_{\alpha-1}^{t_0} : \sup_{t \geq t_0+1} \|U_{\alpha-1}(t, t_0)x\| < \infty\}$. Moreover, one can replace throughout the space $X_{\alpha-1}^t$ by the closure of X in $X_{\alpha-1}^t$.

4. Nonautonomous parabolic boundary evolution equations

In this section we study the nonautonomous parabolic boundary evolution equation

$$\begin{aligned} u'(t) &= A_m(t)u(t) + g(t), \quad t \geq t_0, \\ B(t)u(t) &= h(t), \quad t \geq t_0, \\ u(t_0) &= u_0, \end{aligned} \tag{4.1}$$

and its variant on the line

$$\begin{aligned} u'(t) &= A_m(t)u(t) + g(t), \quad t \in \mathbb{R}, \\ B(t)u(t) &= h(t), \quad t \in \mathbb{R}. \end{aligned} \tag{4.2}$$

Here $t_0 \in \mathbb{R}$, $u_0 \in X$, and the operators $A_m(t)$ and $B(t)$, $t \in \mathbb{R}$, are defined on a Banach space $Z_t \hookrightarrow X$ and map into the state space X and the ‘boundary space’ Y , respectively. The inhomogeneities g and h take values in X and Y , respectively. In the typical applications $A_m(t)$ is a differential operator with ‘maximal’ domain not containing boundary conditions and $B(t)$ are boundary operators. We further introduce the operators

$$A(t)u := A_m(t)u, \quad u \in D(A(t)) := \{u \in Z_t : B(t)u = 0\}.$$

More precisely, we make the following assumptions.

- (A1) For every $t \in \mathbb{R}$ there is a Banach space $Z_t \hookrightarrow X$ such that $A_m(t) \in \mathcal{L}(Z_t, X)$. The operators $B(t) \in \mathcal{L}(Z_t, Y)$ are surjective for $t \in \mathbb{R}$.
 (A2) The operators $A(t) = A_m(t)|_{N(B(t))}$, $t \in \mathbb{R}$, satisfy (2.1) and (2.2).

Under these hypotheses, there is an evolution family $(U(t, s))_{t \geq s}$ solving the problem with homogeneous conditions $g = h = 0$. Moreover, by [19, Lemma 1.2] there exists the *Dirichlet map* $D(t)$ for $\omega - A_m(t)$, i.e., $v = D(t)y$ is the unique solution of the abstract boundary value problem

$$(\omega - A_m(t))v = 0, \quad B(t)v = y,$$

for each $y \in Y$. Fixing $\alpha \in (1 - \mu, 1]$ (where μ is given by (2.2)), we further assume that

- (A3) $Z_t \subset X'_\alpha$, $\sup_{t \in \mathbb{R}} \|D(t)\|_{\mathcal{L}(Y, X'_\alpha)} < \infty$ and $\mathbb{R} \ni t \mapsto D(t)y$ is continuous in X for each $y \in Y$.

If (A1)–(A3) hold with \mathbb{R} replaced by a closed interval J , we may extend $A_m(t)$, $B(t)$, and Z_t constantly to $t \in \mathbb{R}$, and then (A1)–(A3) hold on \mathbb{R} for this extension. Hypotheses (A1)–(A3) describe one convenient general setting for the application of our results, in particular suited for parabolic problems formulated on L^p or C^B spaces. But our approach is more flexible. So we treat in Example 4.5 an initial boundary value problem on the state space $X = C(\bar{\Omega})$ which does not fit in the above setting. We add a simple observation.

Lemma 4.1. Assume that assumptions (A1)–(A3) hold and that $h \in C_0(J, Y)$ for a closed interval J . Then $(\omega - A_{-1}(\cdot))D(\cdot)h \in E_{\alpha-1}(J)$.

Proof. Assumption (A3) yields $D(\cdot)h \in E_\alpha(J)$ which implies the assertion. \square

In order to apply the results from the previous sections to the boundary evolution equation (4.1), we write it as the inhomogeneous Cauchy problem

$$\begin{aligned} u'(t) &= A_{-1}(t)u(t) + f(t), \quad t \geq t_0, \\ u(t_0) &= u_0, \end{aligned} \tag{4.3}$$

setting $f := g + (\omega - A_{-1}(\cdot))D(\cdot)h$. We also consider the evolution equation

$$u'(t) = A_{-1}(t)u(t) + f(t), \quad t \in \mathbb{R}. \tag{4.4}$$

If $g \in C_0(J, X)$ and $h \in C_0(J, Y)$, then $f \in E_{\alpha-1}(J)$ by Lemma 4.1. As in Definition 2.4, we call a function $u \in C(J, X)$ a *mild solution* of (4.2) and (4.4) on J if the equation

$$u(t) = U(t, s)u(s) + \int_s^t U_{\alpha-1}(t, \sigma) [g(\sigma) + (\omega - A_{-1}(\sigma))D(\sigma)h(\sigma)] d\sigma \quad (4.5)$$

holds for all $t \geq s$ in J . The function u is called a *mild solution* of (4.1) and (4.3) if in addition $u(t_0) = u_0$ and $J = [t_0, \infty)$. Mild solutions for the corresponding final value problems are defined in the same way. We note that a function $u \in C^1(J, X)$ with $u(t) \in Z_t$ satisfies (4.1), respectively (4.2), if and only if it satisfies (4.3), respectively (4.4), and then it is given by (4.5). These facts can be shown as in Proposition 4.2 of [13].

Propositions 2.7 and 2.8 immediately imply two results on the existence of bounded mild solutions for forward and backward boundary evolution equations.

Proposition 4.2. *Assume that assumptions (A1)–(A3) hold with $1 - \mu < \alpha \leq 1$ and that $U(t, s)$ has an exponential dichotomy on an interval $[T, \infty)$. Let $t_0 \geq T$, $g \in C_0([T, \infty), X)$, $h \in C_0([T, +\infty), Y)$, and $u_0 \in \overline{D(A(t_0))}$. Then the mild solution $u \in C([t_0, +\infty), X)$ of the boundary evolution equation (4.1) is bounded on $[t_0, \infty)$ if and only if*

$$Q(t_0)u_0 = - \int_{t_0}^{+\infty} \tilde{U}_{\alpha-1}(t_0, s) Q_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds.$$

In this case u is given by

$$u(t) = U(t, t_0)P(t_0)u_0 + \int_{t_0}^t U_{\alpha-1}(t, s)P_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds \\ - \int_t^{\infty} \tilde{U}_{\alpha-1}(t, s) Q_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds, \quad t \geq t_0.$$

Proposition 4.3. *Assume that assumptions (A1)–(A3) hold with $1 - \mu < \alpha \leq 1$ and that $U(t, s)$ has an exponential dichotomy on an interval $(-\infty, -T]$. Let $t_0 < -T$, $g \in C_0((-\infty, -T], X)$, $h \in C_0((-\infty, -T], Y)$, and $u_0 \in X$. Then there is a bounded mild solution $u \in C((-\infty, t_0], X)$ of the backward boundary evolution equation*

$$u'(t) = A_m(t)u(t) + g(t), \quad t \leq t_0, \\ B(t)u(t) = h(t), \quad t \leq t_0, \\ u(t_0) = u_0,$$

if and only if

$$P(t_0)u_0 = \int_{-\infty}^{t_0} U_{\alpha-1}(t_0, s) P_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds.$$

In this case u is given by

$$\begin{aligned} u(t) = & \tilde{U}(t, t_0) Q(t_0) u_0 - \int_t^{t_0} \tilde{U}_{\alpha-1}(t, s) Q_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds \\ & + \int_{-\infty}^t U_{\alpha-1}(t, s) P_{\alpha-1}(s) [g(s) + (\omega - A_{-1}(s))D(s)h(s)] ds, \quad t \leq t_0. \end{aligned}$$

Moreover, Theorem 3.10 implies the following Fredholm alternative for the mild solutions of (4.2).

Theorem 4.4. Assume that assumptions (A1)–(A3) hold with $1 - \mu < \alpha \leq 1$, that (2.16) and (2.17) are true, and that $\dim Q_{\pm\infty} X < \infty$. Let $g \in C_0(\mathbb{R}, X)$ and $h \in C_0(\mathbb{R}, Y)$. Then there is a mild solution $u \in C_0(\mathbb{R}, X)$ of (4.2) if and only if

$$\int_{\mathbb{R}} \langle f(s), w(s) \rangle_{X_{\alpha-1}^s} ds = 0$$

for $f = g + (\omega - A_{-1}(\cdot))D(\cdot)h$ and all $w \in L^1(\mathbb{R}, X^*)$ with $w(s) = U_{\alpha-1}(t, s)^* w(t)$ for all $t \geq s$. The mild solutions u are given by

$$\begin{aligned} u(t) &= v(t) - U(t, T)y_s + (R_{\alpha-1}^+ f)(t), \quad t \geq T, \\ u(t) &= v(t) + \tilde{v}(t) + (R_{\alpha-1}^- f)(t), \quad t \leq T, \end{aligned}$$

where $R_{\alpha-1}^{\pm}$ were defined before Proposition 3.3, $(R_{\alpha-1}^+ f)(T) - (R_{\alpha-1}^- f)(T) = y_s + y_u \in P(T)X + X_u(T)$, $\tilde{v} \in C_0((-\infty, T], X)$ with $\tilde{v}(T) = y_u$ and $\tilde{v}(t) = U(t, s)\tilde{v}(s)$ for all $T \geq t \geq s$, and $v \in C_0(\mathbb{R}, X)$ with $v(t) = U(t, s)v(s)$ for all $t \geq s$.

We add an example dealing with a parabolic PDE in a sup norm context. One could treat more general problems, in particular systems, cf. [17], and one could weaken the regularity assumptions.

Example 4.5. We study the boundary value problem

$$\begin{aligned} \partial_t u(t, x) &= A(t, x, D)u(t, x) + g(t, x), \quad t \in \mathbb{R}, x \in \Omega, \\ B(t, x, D)u(t, x) &= h(t, x), \quad t \in \mathbb{R}, x \in \partial\Omega, \end{aligned} \tag{4.6}$$

on a bounded domain $\Omega \subseteq \mathbb{R}^n$ with boundary $\partial\Omega$ of class C^2 and outer unit normal vector $\nu(x)$, employing the differential expressions

$$A(t, x, D) = \sum_{k,l} a_{kl}(t, x) \partial_k \partial_l + \sum_k a_k(t, x) \partial_k + a_0(t, x),$$

$$B(t, x, D) = \sum_k b_k(t, x) \partial_k + b_0(t, x).$$

We require that $a_{kl} = a_{lk}$ and b_k are real-valued, $a_{kl}, a_k, a_0 \in C_b^\mu(\mathbb{R}, C(\overline{\Omega}))$, $b_k, b_0 \in C_b^\mu(\mathbb{R}, C^1(\partial\Omega))$,

$$\sum_{k,l=1}^n a_{kl}(t, x) \xi_k \xi_l \geq \eta |\xi|^2, \quad \text{and} \quad \sum_{k=1}^n b_k(t, x) \nu_k(x) \geq \beta$$

for constants $\mu \in (1/2, 1)$, $\beta, \eta > 0$ and all $\xi \in \mathbb{R}^n$, $k, l = 1, \dots, n$, $t \in \mathbb{R}$, $x \in \overline{\Omega}$ respectively $x \in \partial\Omega$. (C_b^μ is the space of bounded, globally Hölder continuous functions.) We set $X = C(\overline{\Omega})$,

$$Z_t = \left\{ u \in \bigcap_{p>1} W_p^2(\Omega) : A(t, \cdot, D)u \in C(\overline{\Omega}) \right\},$$

$A_m(t)u = A(t, \cdot, D)u$ and $B(t)u = B(t, \cdot, D)u$ for $u \in Z_t$, and $A(t) = A_m(t)|N(B(t))$, i.e.,

$$D(A(t)) = \left\{ u \in \bigcap_{p>1} W_p^2(\Omega) : A(t, \cdot, D)u \in C(\overline{\Omega}), B(t, \cdot, D)u = 0 \text{ on } \partial\Omega \right\},$$

for $t \in \mathbb{R}$. It is known that the operators $A(t)$, $t \in \mathbb{R}$, satisfy (2.1) and (2.2), see [4,24], or [34, Example 2.9]. Thus $A(\cdot)$ generates an evolution family $U(\cdot, \cdot)$ on X . Let us fix numbers $\alpha \in (1 - \mu, 1/2)$ and $p > \frac{n}{2(1-\alpha)}$. Then $X_\alpha^t = C^{2\alpha}(\overline{\Omega})$ with uniformly equivalent constants due to Theorem 3.1.30 in [24]. So Sobolev's embedding theorem yields $W_p^2(\Omega) \hookrightarrow X_\alpha^t$ with a uniform constant. Standard elliptic theory tells us that for each $\varphi \in W_p^{1-1/p}(\partial\Omega)$ there is a unique $D(t)\varphi := u \in W_p^2(\Omega)$ such that

$$(\omega - A(t, \cdot, D))u = 0 \quad \text{on } \Omega, \quad B(t, \cdot, D)u = \varphi \quad \text{on } \partial\Omega,$$

where $D(t) : W_p^{1-1/p}(\partial\Omega) \rightarrow W_p^2(\Omega)$ is bounded uniformly in $t \in \mathbb{R}$, see [6, Theorem 15.2]. (The Slobodetskij spaces $W_p^s(\partial\Omega)$ are defined in e.g. [5, §7.51].) For $\varphi \in W_p^{1-1/p}(\partial\Omega)$, the properties of $D(t)$ yield

$$(\omega - A(t, \cdot, D))(D(t)\varphi - D(s)\varphi) = (A(t, \cdot, D) - A(s, \cdot, D))D(s)\varphi,$$

$$B(t, \cdot, D)(D(t)\varphi - D(s)\varphi) = (B(s, \cdot, D) - B(t, \cdot, D))D(s)\varphi,$$

so that [6, Theorem 15.2] implies that

$$\begin{aligned}
& \|D(t)\varphi - D(s)\varphi\|_{W_p^2(\Omega)} \\
& \leq c(\|(A(t, \cdot, D) - A(s, \cdot, D))D(s)\varphi\|_{L^p(\Omega)} + \|(B(s, \cdot, D) - B(t, \cdot, D))D(s)\varphi\|_{W_p^{1-1/p}(\partial\Omega)}) \\
& \leq c|t - s|^\mu \|D(s)\varphi\|_{W_p^2(\Omega)} \leq c|t - s|^\mu \|\varphi\|_{W_p^{1-1/p}(\partial\Omega)}
\end{aligned}$$

for constants independent of $t, s \in \mathbb{R}$ (using [5, §7.51]). So we see that $D(\cdot)h \in E_\alpha$ if $h \in C_0(\mathbb{R}, W_p^{1-1/p}(\partial\Omega))$. Further let $g \in C_0(\mathbb{R}, X)$. We define mild solutions of (4.6) again by (4.5). (Observe that a solution $u \in C^1(\mathbb{R}, C(\overline{\Omega}))$ of (4.6) with $u(t) \in Z_t$ for $t \in \mathbb{R}$ solves (4.6) formulated on $X = L^p(\Omega)$. On this state space, (A1)–(A3) hold with $Z_t = W_p^2(\Omega)$ and $Y = W_p^{1-1/p}(\partial\Omega)$ by the above mentioned results. In this setting we have already justified the concept of mild solutions given (4.5).) We further assume that

$$a_\alpha(t, \cdot) \rightarrow a_\alpha(\pm\infty, \cdot) \quad \text{in } C(\overline{\Omega}) \quad \text{and} \quad b_j(t, \cdot) \rightarrow b_j(\pm\infty, \cdot) \quad \text{in } C^1(\partial\Omega)$$

as $t \rightarrow \pm\infty$, where $\alpha = (k, l)$ or $\alpha = j$ for $k, l = 1, \dots, n$ and $j = 0, \dots, n$. We define the sectorial operators $A_{\pm\infty}$ in the same way as $A(t)$. As in [17, Example 5.1] one can check that (2.16) holds. Finally we assume that $i\mathbb{R} \subset \rho(A_{\pm\infty})$. (Observe that the operators $A_{\pm\infty}$ have compact resolvent so that the spectrum consists only of eigenvalues. The spectrum of $A_{\pm\infty}$ was studied in [17, Example 5.1].) Then the Fredholm alternative Theorem 4.4 holds for mild solutions of (4.6) on $X = C(\overline{\Omega})$ for $g \in C_0(J, X)$ and $h \in C_0(\mathbb{R}, W_p^{1-1/p}(\partial\Omega))$ due to the results from Section 3.

Appendix A. Proof of Proposition 2.6

We start with a lemma giving an additional estimate on $U_{\alpha-1}(t, s)$.

Lemma A.1. *Assume that (2.1) and (2.2) hold. Let $s < t \leq s + t_0$, $t_0 > 0$, $1 - \nu < \theta < \mu$, and $1 - \mu < \bar{\alpha} < 1 - \theta$. Then the operators $V(t, s) := (\omega - A(t))^{-\theta} U_{\bar{\alpha}-1}(t, s) (\omega - A_{-1}(s))^\theta$ defined on X belong to $\mathcal{L}(X)$ with norms bounded by a constant $c(t_0, \theta)$. We further set $V(s, s) := I$. Then the map $(t, s) \mapsto V(t, s)f(s)$ is continuous for $t \geq s$ and every $f \in \hat{E}(J)$, where $J \subset \mathbb{R}$ is a closed interval. For $1 - \mu < \alpha \leq 1$ the operators $U_{\alpha-1}(t, s) : X_{\alpha-1}^s \rightarrow X_{\alpha-1}^t$ are locally uniformly bounded for $s \leq t \leq s + t_0$.*

Proof. Let $s < t \leq s + t_0$, $t_0 > 0$, and $1 - \nu < \theta < \mu$. By rescaling, we may assume that (2.1) and (2.2) hold for some $\omega < 0$. Then the Yosida approximations $A_n(t) = nA(t)R(n, A(t))$, $t \in \mathbb{R}$, fulfill (2.1) and (2.2) with $\omega = 0$ and possibly different, but n -independent constants, for sufficiently large $n \in \mathbb{N}$. Thus $A_n(\cdot)$ generates an evolution family $U_n(\cdot, \cdot)$ with estimates independent of n . These evolution families satisfy

$$\begin{aligned}
V_n(t, s) &:= (-A_n(t))^{-\theta} U_n(t, s) (-A_n(s))^\theta \\
&= e^{(t-s)A_n(s)} + [(-A_n(t))^{-\theta} - (-A_n(s))^{-\theta}] (-A_n(s))^\theta e^{(t-s)A_n(s)}
\end{aligned}$$

$$\begin{aligned}
& + \int_s^t V_n(t, \sigma) (-A_n(\sigma))^{1-\theta} [(-A_n(\sigma))^{-1} - (-A_n(s))^{-1}] (-A_n(s))^{1+\theta} \\
& \times e^{(\sigma-s)A_n(s)} d\sigma.
\end{aligned} \tag{A.1}$$

In view of the above integral equation for $V_n(t, s)$, we introduce the operators

$$\begin{aligned}
a_n(t, s) &:= [(-A_n(t))^{-\theta} - (-A_n(s))^{-\theta}] (-A_n(s))^\theta e^{(t-s)A_n(s)}, \\
k_n(t, s) &:= (-A_n(t))^{1-\theta} [(-A_n(t))^{-1} - (-A_n(s))^{-1}] (-A_n(s))^{1+\theta} e^{(t-s)A_n(s)}.
\end{aligned}$$

The estimates (2.6), (2.9) and (2.10) yield

$$\|a_n(t, s)\| \leq c(t-s)^{\mu-\theta} \quad \text{and} \quad \|k_n(t, s)\| \leq c(t-s)^{\mu-\theta-1} \tag{A.2}$$

with constants $c = c(t_0)$ independent of n . Setting $b_n(t, s) := e^{(t-s)A_n(s)} + a_n(t, s)$, we can rewrite (A.1) as

$$V_n(t, s) = b_n(t, s) + \int_s^t V_n(t, \tau) k_n(\tau, s) d\tau =: b_n(t, s) + (V_n * k_n)(t, s).$$

Theorem II.3.2.2 and Lemma II.3.2.1 of [7] now show that

$$V_n(t, s) = b_n(t, s) + \sum_{j=1}^{\infty} (b_n * [k_n *]^j)(t, s) \quad \text{and} \quad \|V_n(t, s)\| \leq c \tag{A.3}$$

for $s \leq t \leq s + t_0$ and the j -times ‘convolution’ $[k_n *]^j = k_n * \cdots * k_n$, where

$$\|[k_n *]^j(t, s)\| \leq c_j(t-s)^{-\alpha} \quad \text{with} \quad \sum_{j=1}^{\infty} c_j < \infty, \tag{A.4}$$

and the constants $c = c(t_0)$ and $c_j = c_j(t_0)$ do not depend on n . It is straightforward to show that

$$\begin{aligned}
\lim_{n \rightarrow \infty} a_n(t, s) &= a(t, s) := [(-A(t))^{-\theta} - (-A(s))^{-\theta}] (-A(s))^\theta e^{(t-s)A(s)}, \\
\lim_{n \rightarrow \infty} b_n(t, s) &= b(t, s) := e^{(t-s)A(s)} + [(-A(t))^{-\theta} - (-A(s))^{-\theta}] (-A(s))^\theta e^{(t-s)A(s)}, \\
\lim_{n \rightarrow \infty} k_n(t, s) &= k(t, s) := (-A(t))^{1-\theta} [(-A(t))^{-1} - (-A(s))^{-1}] (-A(s))^{1+\theta} e^{(t-s)A(s)}
\end{aligned}$$

in $\mathcal{L}(X)$ locally uniformly for $t > s$, cf. [35, Proposition 2.1] and use Lemmas 4.1 and 4.2 of [3]. Moreover, the limit operators satisfy estimates analogous to (A.2) and (A.4). Therefore (A.3) implies that $V_n(t, s)$ converges in $\mathcal{L}(X)$ and locally uniformly for $t > s$ to an operator $V(t, s)$ satisfying $\|V(t, s)\| \leq c(t_0)$ and

$$V(t, s) = e^{(t-s)A(s)} + a(t, s) + \int_s^t V(t, \tau)k(\tau, s) d\tau \quad (\text{A.5})$$

for $s < t \leq s + t_0$ and $t_0 > 0$. Since $U_n(t, s) \rightarrow U(t, s)$ in $\mathcal{L}(X)$ by e.g. Proposition 2.5 of [34], $V_n(t, s)x$ converges to $(\omega - A(t))^{-\theta}U(t, s)(\omega - A(s))^{\theta}x$ for $x \in D((\omega - A(s))^{\theta})$. We then deduce the first assertion from Proposition 2.1 and embedding (2.4) by approximation. Further, the third assertion follows by the reiteration (see e.g. Theorem 1.2.15 and Proposition 2.2.15 in [24]). The second assertion was shown in Proposition 2.1 for $t > s$. Let $f \in \hat{E}(J)$ and $\varepsilon > 0$. Take $g \in D(A(\cdot))$ with $\|f - g\|_{\infty} \leq \varepsilon$. Using (A.5), we estimate

$$\begin{aligned} \|V(t, s)f(s) - f(r)\| &\leq \|(e^{(t-s)A(s)} - I)f(s)\| + \|f(s) - f(r)\| + c(t-s)^{\mu-\theta} \\ &\leq \|(e^{(t-s)A(s)} - I)g(s)\| + c\varepsilon + \|f(s) - f(r)\| + c(t-s)^{\mu-\theta}. \end{aligned}$$

This inequality shows that

$$\limsup_{(t,s) \rightarrow (r,r)} \|V(t, s)f(s) - f(r)\| \leq c\varepsilon,$$

and so the last assertion is established, too. \square

Proof of Proposition 2.6. By rescaling, we can assume that (2.1) and (2.2) hold with $\omega = 0$. Let $1 - \mu < \alpha \leq 1$, $f \in E_{\alpha-1}(J)$, $0 \leq \beta < \min\{\alpha, \nu\}$, $s < t$ in J , and let $u \in C(J, X)$ be a mild solution of (2.20). Formulas (2.21) and (2.12) yield

$$\begin{aligned} \|u(t)\|_{\beta}^t &\leq \|U(t, s)u(s)\|_{\beta}^t + \int_s^t \|U_{\alpha-1}(t, \sigma)f(\sigma)\|_{\beta}^t d\sigma \\ &\leq c(t-s)^{-\beta}\|u(s)\| + c \int_s^t (t-\sigma)^{\alpha-\beta-1}\|f(\sigma)\|_{\alpha-1}^s d\sigma \\ &\leq c(t-s)^{-\beta}\|u(s)\| + c(t-s)^{\alpha-\beta}\|f\|_{\alpha-1}, \end{aligned}$$

so that $u(t) \in X_{\beta}^t$. Moreover, we have

$$\frac{1}{h}(U(t+h, s) - U(t, s))u(s) \rightarrow A(t)U(t, s)u(s)$$

in X as $h \rightarrow 0$. So it remains to differentiate the term

$$v(t) := \int_s^t U_{\alpha-1}(t, \sigma)f(\sigma) d\sigma$$

for $t \in J \setminus \inf J$. Fix θ with $\max\{1 - \nu, 1 - \alpha\} < \theta < \mu$ and let $h > 0$. Then we can write

$$\begin{aligned}
& \frac{1}{h} (-A(t))^{-\theta} (v(t+h) - v(t)) \\
&= (-A(t))^{-\theta} \frac{1}{h} (U(t+h, t) - I)v(t) + \frac{1}{h} \int_t^{t+h} (-A(t+h))^{-\theta} U_{\bar{\alpha}-1}(t+h, \sigma) f(\sigma) d\sigma \\
&\quad + ((-A(t))^{-\theta} - (-A(t+h))^{-\theta}) \frac{1}{h} \int_t^{t+h} U_{\alpha-1}(t+h, \sigma) f(\sigma) d\sigma \\
&=: S_1 + S_2 + S_3,
\end{aligned}$$

where we take a number $\bar{\alpha} \in (1 - \mu, 1 - \theta)$ (thus $\bar{\alpha} < \alpha$). Since $(-A_{-1}(\cdot))^{-\theta} f(\cdot) \in \hat{E}$ by $1 - \theta < \alpha$ and (2.4), Lemma A.1 shows that

$$S_2 = \frac{1}{h} \int_t^{t+h} V(t+h, \sigma) (-A_{-1}(\sigma))^{-\theta} f(\sigma) d\sigma \rightarrow (-A_{-1}(t))^{-\theta} f(t)$$

in X as $h \rightarrow 0$. Using (2.10) and (2.11), we estimate

$$\|S_3\| \leq ch^{\mu-1} \int_t^{t+h} (t+h-\sigma)^{\alpha-1} d\sigma \|f\|_{\alpha-1} \leq ch^{\alpha+\mu-1} \rightarrow 0, \quad h \rightarrow 0.$$

We note that (A.5) applied to $x \in X_t^\beta$ can be shown also for $\theta = 0$ (where $V(t, s) = U(t, s)$ and $a(t, s) = 0$) using similar methods, cf. [36, p. 347]. The term S_1 can thus be transformed into

$$\begin{aligned}
S_1 &= \frac{1}{h} (e^{hA(t)} - I) (-A(t))^{-\theta} v(t) \\
&\quad + \frac{1}{h} \int_t^{t+h} V(t+h, \sigma) (-A(\sigma))^{1-\theta} [(-A(\sigma))^{-1} - (-A(t))^{-1}] (-A(t))^{1-\gamma} \\
&\quad \times e^{(\sigma-t)A(t)} (-A(t))^\gamma v(t) d\sigma \\
&\quad + [(-A(t))^{-\theta} - (-A(t+h))^{-\theta}] \frac{1}{h} \int_t^{t+h} U_{\alpha-1}(t+h, \sigma) (-A(\sigma))^\theta \\
&\quad \times (-A(\sigma))^{1-\theta} [(-A(\sigma))^{-1} - (-A(t))^{-1}] (-A(t))^{1-\gamma} e^{(\sigma-t)A(t)} (-A(t))^\gamma v(t) d\sigma \\
&=: S_{11} + S_{12} + S_{13}.
\end{aligned}$$

Here we take γ with $1 - \mu < 1 - \theta < \gamma < \min\{\alpha, \nu\}$. Since $v(t) \in X_\gamma^t$, the embedding (2.4) yields that $(-A(t))^{-\theta} v(t) \in D(A(t))$, and hence

$$\lim_{h \rightarrow 0} \frac{1}{h} (e^{hA(t)} - I) (-A(t))^{-\theta} v(t) = A(t) (-A(t))^{-\theta} v(t)$$

in X . Lemma A.1 and the inequalities (2.6) and (2.9) allow to estimate

$$\|S_{12}\| \leq \frac{c}{h} \int_t^{t+h} (\sigma - t)^\mu (\sigma - t)^{\gamma-1} d\sigma \|(-A(t))^\gamma v(t)\| = ch^{\gamma+\mu-1} \|(-A(t))^\gamma v(t)\| \rightarrow 0.$$

Finally, we deduce from (2.6), (2.8)–(2.10) that

$$\begin{aligned} \|S_{13}\| &\leq ch^{\mu-1} \int_t^{t+h} (t+h-\sigma)^{-\theta} (\sigma - t)^\mu (\sigma - t)^{\gamma-1} d\sigma \|(-A(t))^\gamma v(t)\| \\ &\leq ch^{2\mu+\gamma-\theta-1} \rightarrow 0, \quad h \rightarrow 0. \end{aligned}$$

Therefore S_1 converges to $A(t)(-A(t))^{-\theta}v(t)$ in X . Summarizing, we have established

$$\lim_{h \rightarrow 0} (-A(t))^{-\theta} \frac{1}{h} (u(t+h) - u(t)) = (-A_{-1}(t))^{-\theta} (A_{-1}(t)u(t) + f(t))$$

in X . By (2.4), this limit exists in $X_{-\theta}^t$, and so in $X_{\beta-1}^t$ for $0 \leq \beta \leq 1 - \theta < \min\{\alpha, v\}$. \square

References

- [1] A. Abbondandolo, P. Majer, Ordinary differential operators in Hilbert spaces and Fredholm pairs, *Math. Z.* 243 (2003) 525–562.
- [2] P. Acquistapace, F. Flandoli, B. Terreni, Initial boundary value problems and optimal control for nonautonomous parabolic systems, *SIAM J. Control Optim.* 29 (1991) 89–118.
- [3] P. Acquistapace, B. Terreni, A unified approach to abstract linear nonautonomous parabolic equations, *Rend. Sem. Mat. Univ. Padova* 78 (1987) 47–107.
- [4] P. Acquistapace, Evolution operators and strong solutions of abstract linear parabolic equations, *Differential Integral Equations* 1 (1988) 433–457.
- [5] R. Adams, *Sobolev Spaces*, Academic Press, New York, 1975.
- [6] S. Agmon, A. Douglis, L. Nirenberg, Estimates near the boundary for solutions of elliptic partial differential equations satisfying general boundary conditions, *Comm. Pure Appl. Math.* 12 (1959) 623–727.
- [7] H. Amann, *Linear and Quasilinear Parabolic Problems*, vol. 1: Abstract Linear Theory, Birkhäuser, Basel, 1995.
- [8] C.J.K. Batty, R. Chill, Approximation and asymptotic behavior of evolution families, *Differential Integral Equations* 15 (2002) 477–512.
- [9] A.G. Baskakov, On invertibility and the Fredholm property of parabolic differential operators, *Dokl. Akad. Nauk* 383 (2002) 583–585.
- [10] V. Casarino, K.-J. Engel, R. Nagel, G. Nickel, A semigroup approach to boundary feedback systems, *Integral Equations Operator Theory* 47 (2003) 289–306.
- [11] C. Chicone, Y. Latushkin, *Evolution Semigroups in Dynamical Systems and Differential Equations*, Amer. Math. Soc., Providence, RI, 1999.
- [12] W. Desch, I. Lasiecka, W. Schappacher, Feedback boundary control problems for linear semigroups, *Israel J. Math.* 51 (1985) 177–207.
- [13] W. Desch, J. Milota, W. Schappacher, Least square control problems in nonreflexive spaces, *Semigroup Forum* 62 (2001) 337–357.
- [14] G. Di Blasio, Maximal L^p regularity for nonautonomous parabolic equations in extrapolation spaces, *J. Evol. Equ.* 6 (2006) 249–264.
- [15] D. Di Giorgio, On Fredholm properties of $(\mathcal{L}u)(t) = u'(t) - A(t)u(t)$, $t \in \mathbb{R}$, for paths of sectorial operators, PhD thesis, Dipartimento di Matematica, Università di Pisa, 2005.
- [16] D. Di Giorgio, A. Lunardi, On Fredholm properties of $\mathcal{L}u = u' - A(t)u$ for paths of sectorial operators, *Proc. Roy. Soc. Edinburgh Sect. A* 135 (2005) 39–59.

- [17] D. Di Giorgio, A. Lunardi, R. Schnaubelt, Fredholm properties of abstract parabolic operators in L^p spaces on the real line, *Proc. London Math. Soc.* 91 (2005) 703–737.
- [18] K.-J. Engel, R. Nagel, *One-Parameter Semigroups for Linear Evolution Equations*, Springer-Verlag, New York, 2000.
- [19] G. Greiner, Perturbing the boundary conditions of a generator, *Houston J. Math.* 13 (1987) 213–229.
- [20] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, 1980.
- [21] Y. Latushkin, A. Pogan, R. Schnaubelt, Dichotomy and Fredholm properties of evolution equations, *J. Operator Theory*, in press.
- [22] Y. Latushkin, A. Tomilov, Fredholm differential operators with unbounded coefficients, *J. Differential Equations* 208 (2005) 388–429.
- [23] X.B. Lin, Exponential dichotomies and homoclinic orbits in functional-differential equations, *J. Differential Equations* 63 (1986) 227–254.
- [24] A. Lunardi, *Analytic Semigroups and Optimal Regularity in Parabolic Problems*, Birkhäuser, Basel, 1995.
- [25] J. Mallet-Paret, The Fredholm alternative for functional-differential equations of mixed type, *J. Dynam. Differential Equations* 11 (1999) 1–47.
- [26] P. Rabier, The Robbin–Salamon index theorem in Banach spaces with UMD, *Dyn. Partial Differ. Equ.* 1 (2004) 303–337.
- [27] S. Ruan, W. Zhang, Exponential dichotomies, the Fredholm alternative, and transverse homoclinic orbits in partial functional differential equations, *J. Dynam. Differential Equations* 17 (2005) 759–777.
- [28] W. Rudin, *Functional Analysis*, McGraw–Hill, New Delhi, 1989.
- [29] D. Salamon, Infinite-dimensional linear systems with unbounded control and observation: A functional analytic approach, *Trans. Amer. Math. Soc.* 300 (1987) 383–431.
- [30] B. Sandstede, Stability of travelling waves, in: B. Fiedler (Ed.), *Handbook of Dynamical Systems*, vol. 2, Elsevier, 2002, pp. 983–1055.
- [31] B. Sandstede, A. Scheel, On the structure of spectra of modulated travelling waves, *Math. Nachr.* 232 (2001) 39–93.
- [32] R. Schnaubelt, Asymptotically autonomous parabolic evolution equations, *J. Evol. Equ.* 1 (2001) 19–37.
- [33] R. Schnaubelt, Well-posedness and asymptotic behavior of non-autonomous linear evolution equations, in: A. Lorenzi, B. Ruf (Eds.), *Evolution Equations, Semigroups and Functional Analysis*, Birkhäuser, Basel, 2002, pp. 311–338.
- [34] R. Schnaubelt, Asymptotic behavior of parabolic nonautonomous evolution equations, in: M. Iannelli, R. Nagel, S. Piazzera (Eds.), *Functional Analytic Methods for Evolution Equations*, in: *Lecture Notes in Math.*, vol. 1855, Springer-Verlag, Berlin, 2004, pp. 401–472.
- [35] A. Yagi, Parabolic equations in which the coefficients are generators of infinitely differentiable semigroups II, *Funkcial. Ekvac.* 33 (1990) 139–150.
- [36] A. Yagi, Abstract quasilinear evolution equations of parabolic type in Banach spaces, *Boll. Unione Mat. Ital.* (7) 5 (1991) 341–368.