



On parametric Gevrey asymptotics for some nonlinear initial value Cauchy problems

A. Lastra ^{a,b,*}, S. Malek ^{a,b}

^a University of Alcalá, Departamento de Física y Matemáticas, Ap. de Correos 20, E-28871 Alcalá de Henares (Madrid), Spain

^b University of Lille 1, Laboratoire Paul Painlevé, 59655 Villeneuve d'Ascq cedex, France

Received 10 March 2014; revised 9 January 2015

Available online 8 July 2015

Abstract

We study a nonlinear initial value Cauchy problem depending upon a complex perturbation parameter ϵ with vanishing initial data at complex time $t = 0$ and whose coefficients depend analytically on (ϵ, t) near the origin in \mathbb{C}^2 and are bounded holomorphic on some horizontal strip in \mathbb{C} w.r.t. the space variable. This problem is assumed to be non-Kowalevskian in time t , therefore analytic solutions at $t = 0$ cannot be expected in general. Nevertheless, we are able to construct a family of actual holomorphic solutions defined on a common bounded open sector with vertex at 0 in time and on the given strip above in space, when the complex parameter ϵ belongs to a suitably chosen set of open bounded sectors whose union form a covering of some neighborhood Ω of 0 in \mathbb{C}^* . These solutions are achieved by means of Laplace and Fourier inverse transforms of some common ϵ -depending function on $\mathbb{C} \times \mathbb{R}$, analytic near the origin and with exponential growth on some unbounded sectors with appropriate bisecting directions in the first variable and exponential decay in the second, when the perturbation parameter belongs to Ω . Moreover, these solutions satisfy the remarkable property that the difference between any two of them is exponentially flat for some integer order w.r.t. ϵ . With the help of the classical Ramis–Sibuya theorem, we obtain the existence of a formal series (generally divergent) in ϵ which is the common Gevrey asymptotic expansion of the built up actual solutions considered above.

© 2015 Elsevier Inc. All rights reserved.

MSC: 35C10; 35C20

* Corresponding author.

E-mail addresses: alberto.lastra@uah.es (A. Lastra), Stephane.Malek@math.univ-lille1.fr (S. Malek).

¹ The author is partially supported by the project MTM2012-31439 of Ministerio de Ciencia e Innovación, Spain.

Keywords: Asymptotic expansion; Borel–Laplace transform; Fourier transform; Nonlinear integro-differential equation; Nonlinear partial differential equation; Singular perturbation

1. Introduction

In this paper, we consider a family of parameter depending nonlinear initial value Cauchy problems of the form

$$\begin{aligned} Q(\partial_z)(\partial_t u(t, z, \epsilon)) &= (Q_1(\partial_z)u(t, z, \epsilon))(Q_2(\partial_z)u(t, z, \epsilon)) + \sum_{l=1}^D \epsilon^{\Delta_l} t^{d_l} \partial_t^{\delta_l} R_l(\partial_z)u(t, z, \epsilon) \\ &\quad + c_0(t, z, \epsilon)R_0(\partial_z)u(t, z, \epsilon) + f(t, z, \epsilon) \end{aligned} \quad (1)$$

for given vanishing initial data $u(0, z, \epsilon) \equiv 0$, where $D \geq 2$, $\Delta_l, d_l, \delta_l, 1 \leq l \leq D$ are integers which satisfy the inequalities

$$\begin{aligned} 1 &= \delta_1, \quad \delta_l < \delta_{l+1}, \quad d_D = (\delta_D - 1)(k + 1), \quad \Delta_D = d_D - \delta_D + 1, \\ d_l &> (\delta_l - 1)(k + 1), \quad \delta_D \geq \delta_l + \frac{2}{k}, \quad \Delta_l + k(1 - \delta_D) + 1 \geq 0 \end{aligned}$$

for all $1 \leq l \leq D - 1$ and for some integer $k \geq 1$. Besides, $Q(X), Q_1(X), Q_2(X), R_l(X), 0 \leq l \leq D$ are polynomials submitted to the constraints

$$\begin{aligned} \deg(Q) &\geq \deg(R_D) \geq \deg(R_l), \quad \deg(R_D) \geq \deg(Q_1), \quad \deg(R_D) \geq \deg(Q_2), \\ Q(im) &\neq 0, \quad R_D(im) \neq 0 \end{aligned}$$

for all $m \in \mathbb{R}$, all $0 \leq l \leq D - 1$. The coefficient $c_0(t, z, \epsilon)$ and the forcing term $f(t, z, \epsilon)$ are bounded holomorphic functions on a product $D(0, r) \times H_\beta \times D(0, \epsilon_0)$, where $D(0, r)$ (resp. $D(0, \epsilon_0)$) is a disc centered at 0 with small radius $r > 0$ (resp. $\epsilon_0 > 0$) and $H_\beta = \{z \in \mathbb{C} / |\operatorname{Im}(z)| < \beta\}$ is some strip of width $\beta > 0$. In order to avoid cumbersome statements and to improve the readability of the computations, we have restricted our study to a quadratic nonlinearity and monomial coefficients in t in front of the derivatives with respect to t and z but the method described here can also be extended to higher order nonlinearities, with polynomial coefficients w.r.t. t in the linear part on the right handside of equation (1).

This work can be seen as a continuation of the study described in [22] where the second author has studied nonlinear integro-differential initial values problems with the shape

$$R(\partial_z)P(\partial_t, \partial_z)Y(t, z) = \int_0^t b(t-s, z) \partial_z^{s_0} Y(s, z) ds + \int_0^t \partial_z^{s_1} Y(t-s, z) \partial_z^{s_2} Y(s, z) ds \quad (2)$$

where $R(X) \in \mathbb{C}[X]$, $P(T, X) \in \mathbb{C}[T, X]$ and $s_0, s_1, s_2 \geq 0$ are non negative integers. The coefficient $b(t, z) = \sum_{k \in \mathbb{Z}} b_k(z) t^k$ is a polynomial in t and its coefficients $b_k(z)$ are the Fourier inverse transform of some function $b_k(m)$ belonging to a Banach space $E_{(\beta, \mu)}$ of continuous functions

$h : \mathbb{R} \rightarrow \mathbb{C}$ endowed with the norm $\|h(m)\|_{(\beta, \mu)} = \sup_{m \in \mathbb{R}} (1 + |m|)^\mu \exp(\beta|m|)|h(m)|$ and define bounded holomorphic functions on any strip $H_{\beta'}$, $0 < \beta' < \beta$. The initial conditions are defined by $Y(0, z) = Y_0(z)$, $(\partial_z^j Y)(0, z) \equiv 0$, for all $1 \leq j \leq \deg_T P(T, X) - 1$, where Y_0 is also assumed to be the Fourier inverse transform of some $\mathfrak{Y}_0(m)$ belonging to $E_{(\beta, \mu)}$. We focused on the case when the degree of $R(X)P(T, X)$ with respect to T is smaller than its degree in X . In that case the classical Cauchy–Kowalevski theorem (see [12]) cannot be applied and the unique formal power series solution $\hat{Y}(t, z) = \sum_{l \geq 0} Y_l(z)t^l$, with coefficients belonging to the Banach space of bounded holomorphic functions on $H_{\beta'}$ equipped with the sup norm, is in general divergent. Nevertheless, under suitable constraints on the roots of the polynomial $T \mapsto P(T^2, im)$ and for sufficiently small data $\|b_k\|_{(\beta, \mu)}$, $\|\mathfrak{Y}_0\|_{(\beta, \mu)}$, one can construct by means of the classical Borel–Laplace procedure and the Fourier inverse transform an actual holomorphic solution $Y(t, z)$ on $\mathbb{C}_+ \times H_{\beta'}$ of (2) for the given initial data (\mathbb{C}_+ denotes the set of complex numbers t such that $\operatorname{Re}(t) > 0$), which possess the formal series \hat{Y} as the Gevrey asymptotic expansion of order 1 as t tends to 0, meaning that for any compact subsector $\mathcal{W} \subset \mathbb{C}_+$ centered at 0, there exist constants $C, M > 0$ with

$$\sup_{z \in H_{\beta'}} |Y(t, z) - \sum_{l=0}^{n-1} Y_l(z)t^l| \leq CM^n n! |t|^n$$

for all $n \geq 1$, all $t \in \mathcal{W}$.

Compared to the work [22], the problem (1) now involves an additional complex parameter ϵ . Provided that $\delta_D + \deg(R_D) > \deg(Q) + 1$ holds, the problem (1) is singularly perturbed in the parameter ϵ and belongs to a class of so-called PDEs with irregular singularity at $t = 0$ in the sense of [24]. In the paper [21], the second author has already considered a similar problem of the form

$$\epsilon t^2 \partial_t \partial_z^S X_p(t, z, \epsilon) = F(t, z, \epsilon, \partial_t, \partial_z) X_p(t, z, \epsilon) + P(t, z, \epsilon, X_p(t, z, \epsilon)) \quad (3)$$

for given initial data

$$(\partial_z^j X_p)(t, 0, \epsilon) = \phi_{j,p}(t, \epsilon), \quad 0 \leq p \leq \varsigma - 1, 0 \leq j \leq S - 1, \quad (4)$$

where $S, \varsigma \geq 2$ are some positive integers, F is some differential operator with polynomial coefficients and P a polynomial. The initial data $\phi_{j,p}(t, \epsilon)$ were assumed to be holomorphic on products $\mathcal{T} \times \mathcal{E}_p \subset \mathbb{C}^2$ for some sector \mathcal{T} centered at 0 and where $\underline{\mathcal{E}} = \{\mathcal{E}_p\}_{0 \leq p \leq \varsigma-1}$ denotes a family of open bounded sectors with aperture larger than π which form a so-called *good covering* in \mathbb{C}^* , meaning that $\mathcal{E}_p \cap \mathcal{E}_{p+1} \neq \emptyset$ for all $0 \leq p \leq \varsigma - 1$ (with the convention that $\mathcal{E}_\varsigma = \mathcal{E}_0$) with the property that the intersection of any three different elements in $\{\mathcal{E}_p\}_{0 \leq p \leq \varsigma-1}$ is empty and that $\bigcup_{p=0}^{\varsigma-1} \mathcal{E}_p = \mathcal{U} \setminus \{0\}$, where \mathcal{U} is some neighborhood of 0 in \mathbb{C} . Under convenient assumptions on the shape of equation (3) and on the initial data (4), the existence of a formal series $\hat{X}(t, z, \epsilon) = \sum_{k \geq 0} h_k(t, z)\epsilon^k/k!$ solution of (3) is established with coefficients $h_k(t, z)$ belonging to the Banach space \mathbb{F} of bounded holomorphic functions on $\mathcal{T} \times D(0, \delta)$ (for some $\delta > 0$ small enough) equipped with the sup norm. This formal series $\hat{X}(t, z, \epsilon)$ is the Gevrey asymptotic expansion of order 1 of actual holomorphic solutions $X_p(t, z, \epsilon)$ of (3), (4) on \mathcal{E}_p as \mathbb{F} -valued functions, for all $0 \leq p \leq \varsigma - 1$, in other words for any closed subsector $\mathcal{W} \subset \mathcal{E}_p$ centered at 0, there exist constants $C, M > 0$ such that

$$\sup_{t \in \mathcal{T}, z \in \bar{D}(0, \delta)} |X_p(t, z, \epsilon) - \sum_{k=0}^{n-1} h_k(t, z) \epsilon^k / k!| \leq C M^n n! |\epsilon|^n$$

for all $n \geq 1$, all $\epsilon \in \mathcal{W}$.

In this work we address the same queries as in [21,22], namely our main purpose is the construction of actual holomorphic solutions $u_p(t, z, \epsilon)$ to the problem (1) on domains $\mathcal{T} \times H_{\beta'} \times \mathcal{E}_p$ using some Borel–Laplace procedure and the Fourier inverse transform and the analysis of their asymptotic expansions as ϵ tends to 0. More specifically, we can present our main statements as follows.

Main results. Assume the existence of an unbounded sector

$$S_{Q, R_D} = \{z \in \mathbb{C} / |z| \geq r_{Q, R_D}, \quad |\arg(z) - d_{Q, R_D}| \leq \eta_{Q, R_D}\}$$

with direction $d_{Q, R_D} \in \mathbb{R}$, aperture $\eta_{Q, R_D} > 0$ and radius $r_{Q, R_D} > 0$ such that the quotient $Q(im)/R_D(im)$ belongs to S_{Q, R_D} for all $m \in \mathbb{R}$. This sector S_{Q, R_D} is prescribed in such a way that there exists a set of adequate directions $\mathfrak{d}_p \in \mathbb{R}$, $0 \leq p \leq \varsigma - 1$, with the feature that the distinct complex roots $q_l(m)$, $0 \leq l \leq (\delta_D - 1)k - 1$, of the polynomial $P_m(\tau) = Q(im)k - R_D(im)k^{\delta_D} \tau^{(\delta_D - 1)k}$ fulfill estimates of the form: there exist constants $M_1, M_2 > 0$ such that

$$|\tau - q_l(m)| \geq M_1(1 + |\tau|), \quad |\tau - q_{l_0}(m)| \geq M_2|q_{l_0}(m)|$$

for all $0 \leq l \leq (\delta_D - 1)k - 1$, some integer $l_0 \in \{0, \dots, (\delta_D - 1)k - 1\}$, for all $m \in \mathbb{R}$, all $\tau \in S_{\mathfrak{d}_p} \cup \bar{D}(0, \rho)$, for some well chosen unbounded sectors $S_{\mathfrak{d}_p}$ centered at 0 with direction \mathfrak{d}_p and for some radius $\rho > 0$. Then, we choose a family $\underline{\mathcal{E}} = \{\mathcal{E}_p\}_{0 \leq p \leq \varsigma - 1}$ of sectors with aperture slightly larger than π/k which defines a good covering of \mathbb{C}^* and we take an open bounded sector \mathcal{T} centered at 0 such that for every $0 \leq p \leq \varsigma - 1$, the product ϵt belongs to a sector with direction \mathfrak{d}_p and aperture slightly larger than π/k , for all $\epsilon \in \mathcal{E}_p$, all $t \in \mathcal{T}$. We make the assumption that the coefficient $c_0(t, z, \epsilon)$ and the forcing term $f(t, z, \epsilon)$ can be written as convergent series of the special form

$$c_0(t, z, \epsilon) = \sum_{n \geq 0} c_{0,n}(z, \epsilon) (\epsilon t)^n, \quad f(t, z, \epsilon) = \sum_{n \geq 1} f_n(z, \epsilon) (\epsilon t)^n,$$

on a domain $D(0, r) \times H_{\beta'} \times D(0, \epsilon_0)$ (where $H_{\beta'}$ is a strip of width β') such that $\mathcal{T} \subset D(0, r)$, $\cup_{0 \leq p \leq \varsigma - 1} \mathcal{E}_p \subset D(0, \epsilon_0)$ and $0 < \beta' < \beta$ are given positive real numbers. The coefficients $c_{0,0}(z, \epsilon)$, $c_{0,n}(z, \epsilon)$ and $f_n(z, \epsilon)$, $n \geq 1$, are supposed to be inverse the Fourier transform of functions $m \mapsto C_{0,0}(m, \epsilon)$, $m \mapsto C_{0,n}(m, \epsilon)$ and $m \mapsto F_n(m, \epsilon)$ that belong to the Banach space $E_{(\beta, \mu)}$ for some $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ and that depend holomorphically on ϵ in $D(0, \epsilon_0)$.

Our first result stated in Theorem 1 claims that if the norm $\|C_{0,0}(m, \epsilon)\|_{(\beta, \mu)}$ and the radius ϵ_0 are chosen small enough and if the radius r_{Q, R_D} is taken sufficiently large then we can construct a family of holomorphic bounded functions $u_p(t, z, \epsilon)$, $0 \leq p \leq \varsigma - 1$, defined on the products $\mathcal{T} \times H_{\beta'} \times \mathcal{E}_p$, which solves the problem (1) with vanishing initial data $u_p(0, z, \epsilon) \equiv 0$ and which can be written as the Laplace–Fourier transform

$$u_p(t, z, \epsilon) = \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\gamma_p}} \omega_k^{\partial_p}(u, m, \epsilon) e^{-(\frac{u}{\epsilon})^k} e^{izm} \frac{du}{u} dm,$$

where the inner integration is made along some halfline $L_{\gamma_p} \subset S_{\partial_p}$ where $\omega_k^{\partial_p}(u, m, \epsilon)$ denotes a function with at most exponential growth of order k in u/ϵ and exponential decay in $m \in \mathbb{R}$ which satisfies more precisely estimates of the form

$$|\omega_k^{\partial_p}(u, m, \epsilon)| \leq C(1 + |m|)^{-\mu} e^{-\beta|m|} \frac{|\frac{u}{\epsilon}|}{1 + |\frac{u}{\epsilon}|^{2k}} \exp(v|\frac{u}{\epsilon}|^k)$$

for some constants $C, v > 0$, for all $m \in \mathbb{R}$, all $u \in S_{\partial_p} \cup \bar{D}(0, \rho)$, all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Our second main result, described in [Theorem 2](#), asserts that the functions u_p , $0 \leq p \leq \varsigma - 1$, turn out to be the k -sums on \mathcal{E}_p of a common formal power series

$$\hat{u}(t, z, \epsilon) = \sum_{m \geq 0} h_m(t, z) \frac{\epsilon^m}{m!} \in \mathbb{F}[[\epsilon]]$$

where \mathbb{F} is the Banach space of bounded holomorphic functions on $\mathcal{T} \times H_{\beta'}$ equipped with the sup norm. Namely, for any closed subsector $\mathcal{W} \subset \mathcal{E}_p$ centered at 0, there exist constants $C, M > 0$ such that

$$\sup_{t \in \mathcal{T}, z \in H_{\beta'}} |u_p(t, z, \epsilon) - \sum_{m=0}^{n-1} h_m(t, z) \frac{\epsilon^m}{m!}| \leq CM^n \Gamma(1 + \frac{n}{k}) |\epsilon|^n$$

for all $n \geq 1$, all $\epsilon \in \mathcal{W}$.

It is worth remarking that when $\deg(Q) + 1 > \delta_D + \deg(R_D)$, equation (1) is not singularly perturbed in ϵ and possess no irregular singularity at $t = 0$. However, the asymptotic expansion \hat{u} of u_p as ϵ tends to 0 on \mathcal{E}_p remains divergent in general. The reason for this phenomenon to appear relies on the way one constructs the actual solutions u_p as Laplace transforms of order k in the new variable ϵt and from the fact that for any fixed $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, the problem (1) is not Kowalevskian with respect to t at 0 (meaning that formal series solutions $\hat{v}(t, z, \epsilon) = \sum_{n \geq 1} v_n(z, \epsilon) t^n$, with coefficients $z \mapsto v_n(z, \epsilon)$ bounded holomorphic on $H_{\beta'}$, are in general divergent, as a consequence of [Propositions 8 and 9](#)) as it was already the case in our previous paper [22].

The Cauchy problem (1) we consider here comes within the new trend of research concerning Borel–Laplace summability procedures applied to partial differential equation going back to the seminal work of D. Lutz, M. Miyake and R. Schäfke on the linear complex heat equation, see [18]. We quote below some important results in this field not pretending to be exhaustive. This construction of Borel–Laplace k -summable or even multi-summable formal series solutions has been extended to general linear PDEs in two complex variables with constant coefficients by W. Balser in [3] and [4] provided that their initial data are analytic functions near the origin that can be analytically continued with exponential growth on some unbounded sectors. A similar result has been obtain for the so-called fractional linear PDEs with non-integer derivatives by S. Michalik, see [23]. Latter on, linear complex heat like equations with variable coefficients

have been explored by several authors, see [5,7,20]. Recently, general linear PDEs with time dependent coefficients taking for granted that their initial data are entire functions in \mathbb{C}^N , $N \geq 1$, have been investigated by H. Tahara and H. Yamazawa in [27]. In the context of nonlinear PDEs, we mention the work [19] of G. Lysik who constructed summable formal solutions of the one dimensional Burgers equations with the help of the so-called Cole–Hopf transform. We also point out that O. Costin and S. Tanveer have constructed summable formal series in time variable to the celebrated 3D Navier Stokes equations in [9]. We also refer to the work of S. Ouchi who constructed multisummable formal solutions to nonlinear PDEs which come from perturbations of ordinary differential equations, see [25]. We also mention the fact that, these last years, a lot of attention has been payed to singularly perturbed PDEs in the complex domain partly motived by a conjecture of B. Dubrovin which concerns the question of universal behavior of generic solutions near gradient catastrophe of singularly Hamiltonian perturbations of first order hyperbolic equations, see [10]. In this active direction, we refer namely to the works of B. Dubrovin and M. Elaeva who investigated the case of generalized Burgers equations in [11] and of T. Claeys and T. Grava in [6] who solved the problem for KdV equations. We indicate the recent important studies of T. Koike on Garnier systems, [15,16] and of S. Hirose on the reduction of general singularly perturbed holonomic systems in two complex variables to Pearcey systems normal forms, [13].

In the sequel, we explain our principal intermediate key results and the arguments needed in their proofs. In a first part, we depart from an auxiliary parameter depending initial value differential and convolution equation which is singular in its perturbation parameter ϵ at 0, see (72). This equation is formally constructed by making the change of variable $T = \epsilon t$ in equation (1) (as done in our previous works [21,17]) and by taking the Fourier transform with respect to the variable z . Under the constraint (70) and the assumption that $d_l \geq \delta_l$, $0 \leq l \leq D - 1$ (which follows from the hypothesis (69)) we can construct a formal power series solution $\hat{U}(T, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) T^n$ of (72) whose coefficients $m \mapsto U_n(m, \epsilon)$ depend holomorphically on $\epsilon \in \mathbb{C}^*$ near the origin and belong to a Banach space $E_{(\beta, \mu)}$ of continuous function with exponential decay on \mathbb{R} introduced in the paper [9] by O. Costin and S. Tanveer. This series turns out to be in general divergent as we will see below.

In the next step, we follow the strategy developed recently by H. Tahara and H. Yamazawa in [27], namely we multiply each hand side of (72) by the power T^k which transforms it into equation (76) which involves only differential operators in T of irregular type at $T = 0$ of the form $T^\beta \partial_T$ with $\beta \geq k + 1$ due to our assumption (69) on the shape of equation (72).

Then, we apply a formal Borel transform of order k (defined as a slightly modified version of the classical Borel transform of order k from the reference book [1]), that we call m_k -Borel transform in Definition 3, to the formal series \hat{U} with respect to T , denoted

$$\omega_k(\tau, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) \frac{\tau^k}{\Gamma(\frac{n}{k})}.$$

From the commutation rules of the m_k -Borel transform with respect to the weighted convolution product \star of formal series (introduced in Proposition 5) and the differential operators $T^\beta \partial_T$ for $\beta \geq k + 1$ described in Proposition 6, we get that $\omega_k(\tau, m, \epsilon)$ formally solves a convolution equation in both variables τ and m , see (80).

Under some size constraint on the $E_{(\beta, \mu)}$ -norm of the constant term $C_{0,0}$ of one coefficient of equation (80) and for all $\epsilon \in \mathbb{C}^*$ close enough to 0, we show that $\omega_k(\tau, m, \epsilon)$ is actually convergent for τ on some fixed neighborhood of 0 and can be extended to a holomorphic functions

$\omega_k^d(\tau, m, \epsilon)$ on unbounded sectors S_d centered at zero with bisecting direction d and tiny aperture provided that S_d stays away from the roots of some polynomial $P_m(\tau)$, for all $m \in \mathbb{R}$. Besides, the function $\omega_k^d(\tau, m, \epsilon)$ satisfies estimates of the form: there exist constants $\nu > 0$ and $\varpi_d > 0$ with

$$|\omega_k^d(\tau, m, \epsilon)| \leq \varpi_d (1 + |m|)^{-\mu} e^{-\beta|m|} \frac{|\frac{\tau}{\epsilon}|}{1 + |\frac{\tau}{\epsilon}|^{2k}} \exp(\nu |\frac{\tau}{\epsilon}|^k)$$

for all $\tau \in S_d, m \in \mathbb{R}$, all $\epsilon \in \mathbb{C}^*$ near the origin (see [Proposition 9](#)). The technical constraints (69) and (87) together with (81), (84) and (85) allow, by means of lower bound estimates (86) for the polynomial $P_m(\tau)$, the transformation of equation (80) into a fixed point equation $\mathcal{H}_\epsilon(\omega_k) = \omega_k$ where the map \mathcal{H}_ϵ is given by (89) for which we can find a solution ω_k^d in some Banach space of holomorphic functions $F_{(v, \beta, \mu, k, \epsilon)}^d$ studied in Section 2. It is worth noting that the formal series $\hat{U}(T, m, \epsilon)$ diverges since the function $\omega_k(\tau, m, \epsilon)$ cannot in general be extended everywhere on \mathbb{C} w.r.t. τ . But, as a result, we get that these series \hat{U} are m_k -summable w.r.t. T (see [Definition 3](#)) in all the directions d chosen as above. In other words, some Laplace transform of order k of ω_k^d denoted $U^d(T, m, \epsilon)$ can be constructed for all T belonging to a sector $S_{d, k, h|\epsilon|}$ with bisecting direction d , aperture slightly larger than π/k and radius $h|\epsilon|$ (for some $h > 0$). This function $T \mapsto U^d(T, m, \epsilon)$ is the unique $E_{(\beta, \mu)}$ -valued map which admits $\hat{U}(T, m, \epsilon)$ as the Gevrey asymptotic expansion of order $1/k$ on $S_{d, k, h|\epsilon|}$. Moreover, $U^d(T, m, \epsilon)$ solves the auxiliary problem (72) with vanishing initial data $U^d(0, m, \epsilon)$, see [Proposition 10](#).

In [Theorem 1](#), we construct a family of actual bounded holomorphic solutions $u_p(t, z, \epsilon)$, $0 \leq p \leq \varsigma - 1$ of our original problem (1) on domains of the form $\mathcal{T} \times H_{\beta'} \times \mathcal{E}_p$. The sectors \mathcal{E}_p , $0 \leq p \leq \varsigma - 1$ constitute a so-called good covering in \mathbb{C}^* ([Definition 4](#)). The strip $H_{\beta'}$ has width $0 < \beta' < \beta$ and \mathcal{T} is a fixed bounded sector centered at 0 which fulfills the constraint $\epsilon t \in S_{\mathfrak{d}_p, k}$ for all $\epsilon \in \mathcal{E}_p$, $t \in \mathcal{T}$, and $S_{\mathfrak{d}_p, k}$ is a sector of bisecting direction \mathfrak{d}_p and aperture slightly larger than π/k where \mathfrak{d}_p are suitable directions for which the unbounded sectors $S_{\mathfrak{d}_p}$ with small aperture and bisecting direction \mathfrak{d}_p satisfy the restrictions described above. Namely, the functions u_p are set as Fourier inverse transforms of $U^{\mathfrak{d}_p}$,

$$u_p(t, z, \epsilon) = \mathcal{F}^{-1}(m \mapsto U^{\mathfrak{d}_p}(\epsilon t, m, \epsilon))(z)$$

where the definition of \mathcal{F}^{-1} is pointed out in [Proposition 7](#). In addition to that, one can prove that the difference of any two neighboring functions $u_{p+1}(t, z, \epsilon) - u_p(t, z, \epsilon)$ tends to zero as $\epsilon \rightarrow 0$ on $\mathcal{E}_p \cap \mathcal{E}_{p+1}$ faster than a function with exponential decay of order k , uniformly w.r.t. $t \in \mathcal{T}$ and $z \in H_{\beta'}$, see (119).

The last section of the paper is devoted to deal with this latter growth information in order to show the existence of a common asymptotic expansion $\hat{u}(t, z, \epsilon) = \sum_{m \geq 0} h_m(t, z) \epsilon^m / m!$ of the Gevrey order $1/k$ for all the functions $u_p(t, z, \epsilon)$ as ϵ tends to 0 on \mathcal{E}_p , uniformly w.r.t. $t \in \mathcal{T}$ and $z \in H_{\beta'}$, see [Theorem 2](#). The key tool in proving the result is the classical Ramis–Sibuya theorem (Theorem (RS)).

The layout of this work reads as follows.

In Section 2, we define some weighted parameter depending Banach spaces of continuous functions on $\mathbb{C} \times \mathbb{R}$ with exponential growth on sectors w.r.t. the first variable and exponential decay on \mathbb{R} w.r.t. the second one. We study the continuity properties of several kind of linear and nonlinear integral operators acting on these spaces that will be useful in Section 4.

In Section 3, we give a definition of k -summability (that we call m_k -summability) which is a minor modification of the classical one given in the textbook [1] and which is appropriate for the

problem we have to deal with. We also give conditions for the set of m_k -sums of formal series to be a differential algebra. This fact will be important in the next section where we construct actual solutions of the auxiliary equation (72). We provide explicit commutation formulas for the m_k -Borel transform w.r.t. products and differential operators of irregular type.

In Section 4, we introduce an auxiliary differential and convolution problem (72) for which we construct a formal solution. We show that the m_k -Borel transform of this formal solution satisfies a convolution problem (80). Under suitable assumptions, we can solve uniquely this latter problem in the Banach spaces described in Section 2 using some fixed point theorem argument. Then, applying the Laplace transform, we can give a uniquely determined actual solution to (72) having the formal solution mentioned above as the Gevrey asymptotic expansion.

In Section 5, with the help of Section 4, we build a family of actual holomorphic solutions to our initial Cauchy problem (1) on a full neighborhood of the origin in \mathbb{C}^* w.r.t. the perturbation parameter ϵ . We show that the difference of any two neighboring solutions is exponentially flat for some integer order in ϵ (Theorem 1).

In Section 6, we show that the actual solutions constructed in Section 5 share a common formal series as the Gevrey asymptotic expansion as ϵ tends to 0 on sectors (Theorem 2). The result relies on the classical so-called Ramis–Sibuya theorem.

2. Banach spaces functions with exponential growth and decay

We denote by $D(0, r)$ the open disc centered at 0 with radius $r > 0$ in \mathbb{C} and by $\bar{D}(0, r)$ its closure. Let S_d be an open unbounded sector in direction $d \in \mathbb{R}$ and \mathcal{E} be an open sector with finite radius $r_{\mathcal{E}}$, both centered at 0 in \mathbb{C} . By convention, these sectors do not contain the origin in \mathbb{C} .

Definition 1. Let $\nu, \beta, \mu > 0$ and $\rho > 0$ be positive real numbers. Let $k \geq 1$ be an integer and let $\epsilon \in \mathcal{E}$. We denote $F_{(\nu, \beta, \mu, k, \epsilon)}^d$ the vector space of continuous functions $(\tau, m) \mapsto h(\tau, m)$ on $(\bar{D}(0, \rho) \cup S_d) \times \mathbb{R}$, which are holomorphic with respect to τ on $D(0, \rho) \cup S_d$ and such that

$$\|h(\tau, m)\|_{(\nu, \beta, \mu, k, \epsilon)} = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1 + |m|)^{\mu} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - \nu|\frac{\tau}{\epsilon}|^k) |h(\tau, m)|$$

is finite. One can check that the normed space $(F_{(\nu, \beta, \mu, k, \epsilon)}^d, \|\cdot\|_{(\nu, \beta, \mu, k, \epsilon)})$ is a Banach space.

Remark. These norms are appropriate modifications of the norms defined by O. Costin and S. Tanveer in [9] and by the second the author in [21] and [22].

Throughout the whole section, we assume $\epsilon \in \mathcal{E}$, $\mu, \beta, \nu > 0$ are fixed. In the next lemma, we check the continuity property by multiplication operation with bounded functions.

Lemma 1. Let $(\tau, m) \mapsto a(\tau, m)$ be a bounded continuous function on $(\bar{D}(0, \rho) \cup S_d) \times \mathbb{R}$, which is holomorphic with respect to τ on $D(0, \rho) \cup S_d$. Then, we have

$$\|a(\tau, m)h(\tau, m)\|_{(\nu, \beta, \mu, k, \epsilon)} \leq \left(\sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} |a(\tau, m)| \right) \|h(\tau, m)\|_{(\nu, \beta, \mu, k, \epsilon)} \quad (5)$$

for all $h(\tau, m) \in F_{(\nu, \beta, \mu, k, \epsilon)}^d$.

In the next proposition, we study the continuity property of some convolution operators acting on the latter Banach spaces.

Proposition 1. *Let $\gamma_2 > 0$ be a real number. Let $k \geq 1$ be an integer such that $1/k \leq \gamma_2 \leq 1$. Then, there exists a constant $C_1 > 0$ (depending on v, k, γ_2) with*

$$\| \int_0^{\tau^k} (\tau^k - s)^{\gamma_2} f(s^{1/k}, m) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \leq C_1 |\epsilon|^{k\gamma_2} \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (6)$$

for all $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$.

Proof. Let $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$. For any $\tau \in \bar{D}(0, \rho) \cup S_d$, the segment $[0, \tau^k]$ is such that the map $s \in [0, \tau^k] \rightarrow f(s^{1/k}, m)$ is well defined, provided that $m \in \mathbb{R}$. By definition, we have that

$$\begin{aligned} & \| \int_0^{\tau^k} (\tau^k - s)^{\gamma_2} f(s^{1/k}, m) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \\ &= \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1 + |m|)^\mu \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \\ & \quad \times \left| \int_0^{\tau^k} \left\{ (1 + |m|)^\mu e^{\beta|m|} \exp(-v|s|/|\epsilon|^k) \frac{1 + \frac{|s|^2}{|\epsilon|^{2k}}}{\frac{|s|^{1/k}}{|\epsilon|}} f(s^{1/k}, m) \right\} \right. \\ & \quad \left. \times \mathcal{A}(\tau, s, m, \epsilon) ds \right| \end{aligned} \quad (7)$$

where

$$\mathcal{A}(\tau, s, m, \epsilon) = \frac{1}{(1 + |m|)^\mu} e^{-\beta|m|} \frac{\exp(v|s|/|\epsilon|^k)}{1 + \frac{|s|^2}{|\epsilon|^{2k}}} \frac{|s|^{1/k}}{|\epsilon|} (\tau^k - s)^{\gamma_2} \frac{1}{s}$$

Therefore,

$$\| \int_0^{\tau^k} (\tau^k - s)^{\gamma_2} f(s^{1/k}, m) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \leq C_1(\epsilon) \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (8)$$

where

$$C_1(\epsilon) = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \times \int_0^{|\tau|^k} \frac{\exp(vh/|\epsilon|^k)}{1 + \frac{h^2}{|\epsilon|^{2k}}} \frac{h^{\frac{1}{k}-1}}{|\epsilon|} (|\tau|^k - h)^{\gamma_2} dh$$

Making the change of variable $h = |\epsilon|^k h'$ in the integral inside $C_1(\epsilon)$ yields

$$C_1(\epsilon) = |\epsilon|^{k\gamma_2} \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \\ \times \int_0^{|\frac{\tau}{\epsilon}|^k} \frac{\exp(vh')}{1+h'^2} (h')^{\frac{1}{k}-1} (|\frac{\tau}{\epsilon}|^k - h')^{\gamma_2} dh' \leq |\epsilon|^{k\gamma_2} \sup_{x \geq 0} A(x) \quad (9)$$

where

$$A(x) = \frac{1+x^2}{x^{1/k}} \exp(-vx) \int_0^x \frac{\exp(vh)}{1+h^2} h^{\frac{1}{k}-1} (x-h)^{\gamma_2} dh$$

For any $x > 0$, we have $A(x) \leq \tilde{A}(x)$, where

$$\tilde{A}(x) = (1+x^2)x^{\gamma_2-\frac{1}{k}} \exp(-vx) \int_0^x \frac{\exp(vh)}{1+h^2} h^{\frac{1}{k}-1} dh.$$

Using L'Hospital rule, we know that

$$\lim_{x \rightarrow +\infty} \tilde{A}(x) = \lim_{x \rightarrow +\infty} \frac{\exp(vx)x^{\frac{1}{k}-1}/(1+x^2)}{\partial_x \left(\frac{\exp(vx)}{(1+x^2)x^{\gamma_2-\frac{1}{k}}} \right)} \\ = \lim_{x \rightarrow +\infty} \frac{(1+x^2)x^{2(\gamma_2-\frac{1}{k})}x^{\frac{1}{k}-1}}{v(1+x^2)x^{\gamma_2-\frac{1}{k}} - (2x^{\gamma_2-\frac{1}{k}+1} + (\gamma_2 - \frac{1}{k})x^{\gamma_2-\frac{1}{k}-1}(1+x^2))}$$

and this latter limit is finite if $\gamma_2 \leq 1$ holds. Hence, we deduce that there exists a constant $\tilde{A} > 0$ such that

$$\sup_{x \geq 0} \tilde{A}(x) \leq \tilde{A} \quad (10)$$

Gathering the estimates (8), (9), (10), we see that (6) holds. \square

Proposition 2. Let $\gamma_1 \geq 0$ and $\chi_2 > -1$ be real numbers. Let $v_2 \geq 0$ be an integer. We consider a holomorphic function $a_{\gamma_1, k}(\tau)$ on $D(0, \rho) \cup S_d$, continuous on $\bar{D}(0, \rho) \cup S_d$, such that

$$|a_{\gamma_1, k}(\tau)| \leq \frac{1}{(1+|\tau|^k)^{\gamma_1}}$$

for all $\tau \in \bar{D}(0, \rho) \cup S_d$.

i) Assume that $\chi_2 \geq 0$.

If $v_2 + \chi_2 - \gamma_1 \leq 0$, then there exists a constant $C_{2.1} > 0$ (depending on v, v_2, χ_2, γ_1) such that

$$\begin{aligned} & \|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\chi_2} s^{v_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \\ & \leq C_{2.1} |\epsilon|^{k(1+v_2+\chi_2-\gamma_1)} \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \end{aligned} \quad (11)$$

for all $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$.

ii) Assume that $\chi_2 = \frac{\chi}{k} - 1$ for some real number $\chi \geq 1$.

If $v_2 + \frac{1}{k} - \gamma_1 \leq 0$, then there exists a constant $C_{2.2} > 0$ (depending $\chi, k, v, \gamma_1, v_2$) on such that

$$\begin{aligned} & \|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\chi_2} s^{v_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \\ & \leq C_{2.2} |\epsilon|^{k(1+v_2+\chi_2-\gamma_1)} \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \end{aligned} \quad (12)$$

for all $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$.

Proof. In the first part of the proof, let us assume that **i)** holds. Let $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$. By definition, we have

$$\begin{aligned} & \|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\chi_2} s^{v_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \\ & = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1 + |m|)^\mu \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \\ & \quad \times |a_{\gamma_1, k}(\tau) \int_0^{\tau^k} \{(1 + |m|)^\mu e^{\beta|m|} \exp(-v|s|/|\epsilon|^k) \frac{1 + \frac{|s|^2}{|\epsilon|^{2k}}}{\frac{|s|^{1/k}}{|\epsilon|}} f(s^{1/k}, m)\} \\ & \quad \times \mathcal{B}(\tau, s, m, \epsilon) ds| \end{aligned} \quad (13)$$

where

$$\mathcal{B}(\tau, s, m, \epsilon) = \frac{1}{(1 + |m|)^\mu} e^{-\beta|m|} \frac{\exp(v|s|/|\epsilon|^k)}{1 + \frac{|s|^2}{|\epsilon|^{2k}}} \frac{|s|^{1/k}}{|\epsilon|} (\tau^k - s)^{\chi_2} s^{v_2}.$$

Therefore,

$$\|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\chi_2} s^{\nu_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \leq C_2(\epsilon) \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (14)$$

where

$$C_2(\epsilon) = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v |\frac{\tau}{\epsilon}|^k) \\ \times \frac{1}{(1 + |\tau|^k)^{\gamma_1}} \int_0^{|\tau|^k} \frac{\exp(vh/|\epsilon|^k) h^{\frac{1}{k}}}{1 + \frac{h^2}{|\epsilon|^{2k}}} (|\tau|^k - h)^{\chi_2} h^{\nu_2} dh$$

Making the change of variable $h = |\epsilon|^k h'$ in the integral inside $C_2(\epsilon)$ yields

$$C_2(\epsilon) = |\epsilon|^{k(1+\nu_2+\chi_2)} \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v |\frac{\tau}{\epsilon}|^k) \\ \times \frac{1}{(1 + |\epsilon|^k |\frac{\tau}{\epsilon}|^k)^{\gamma_1}} \int_0^{|\frac{\tau}{\epsilon}|^k} \frac{\exp(vh')}{1 + h'^2} (h')^{\frac{1}{k}} (|\frac{\tau}{\epsilon}|^k - h')^{\chi_2} h'^{\nu_2} dh' \\ \leq |\epsilon|^{k(1+\nu_2+\chi_2)} \sup_{x \geq 0} B(x, \epsilon) \quad (15)$$

where

$$B(x, \epsilon) = \frac{1 + x^2}{x^{1/k}} \exp(-vx) \frac{1}{(1 + |\epsilon|^k x)^{\gamma_1}} \int_0^x \frac{\exp(vh)}{1 + h^2} h^{\frac{1}{k} + \nu_2} (x - h)^{\chi_2} dh.$$

For any $x > 0$, we get that $B(x, \epsilon) \leq \tilde{B}(x, \epsilon)$, where

$$\tilde{B}(x, \epsilon) = \frac{(1 + x^2)x^{\chi_2}}{(1 + |\epsilon|^k x)^{\gamma_1}} \exp(-vx) \int_0^x \frac{\exp(vh)}{1 + h^2} h^{\nu_2} dh$$

Let $x_0 > 0$. From the inequality $1 + |\epsilon|^k x \geq 1$, for all $x \in [0, x_0]$ and $\epsilon \in \mathcal{E}$, there exists a constant $\tilde{B} > 0$ such that

$$\sup_{x \in [0, x_0], \epsilon \in \mathcal{E}} \tilde{B}(x, \epsilon) \leq \tilde{B}. \quad (16)$$

On the other hand, since $1 + |\epsilon|^k x \geq |\epsilon|^k x$ holds for all $x \geq 0$ and $\epsilon \in \mathcal{E}$, we get that $\tilde{B}(x, \epsilon) \leq \tilde{B}_2(x)/|\epsilon|^{k\gamma_1}$ where

$$\tilde{B}_2(x) = (1+x^2)x^{\chi_2-\gamma_1} \exp(-vx) \int_0^x \frac{\exp(vh)}{1+h^2} h^{v_2} dh \quad (17)$$

for all $x \geq x_0$. By L'Hospital rule we get that

$$\lim_{x \rightarrow +\infty} \tilde{B}_2(x) = \lim_{x \rightarrow +\infty} \frac{(1+x^2)x^{2(\chi_2-\gamma_1)}x^{v_2}}{v(1+x^2)x^{\chi_2-\gamma_1} - (2x^{\chi_2-\gamma_1+1} + (\chi_2-\gamma_1)x^{\chi_2-\gamma_1-1}(1+x^2))}$$

which is finite if we assume that $1 \geq (1+v_2+\chi_2-\gamma_1)$. We deduce that there exists a constant $\tilde{B}_2 > 0$ such that

$$\sup_{x \geq x_0} \tilde{B}(x, \epsilon) \leq \frac{1}{|\epsilon|^{k\gamma_1}} \sup_{x \geq x_0} \tilde{B}_2(x) \leq \frac{\tilde{B}_2}{|\epsilon|^{k\gamma_1}} \quad (18)$$

Bearing in mind the estimates (14), (15), (16) and (18), we obtain (11).

In the second part of the proof, assume now that the condition **ii**) holds. Let $f(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$. By definition, we have

$$\begin{aligned} & \|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\frac{\chi}{k}-1} s^{v_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \\ &= \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1+|m|)^\mu \frac{1 + \frac{|\tau|}{\epsilon}^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \\ & \quad \times |a_{\gamma_1, k}(\tau) \int_0^{\tau^k} \{(1+|m|)^\mu e^{\beta|m|} \exp(-v|s|/|\epsilon|^k) \frac{1 + \frac{|s|^2}{|\epsilon|^{2k}}}{\frac{|s|^{1/k}}{|\epsilon|}} f(s^{1/k}, m)\} \\ & \quad \times \{\exp(-v \frac{|\tau^k - s|}{|\epsilon|^k}) \frac{1 + \frac{|\tau^k - s|^2}{|\epsilon|^{2k}}}{\frac{|\tau^k - s|^{1/k}}{|\epsilon|}} (\tau^k - s)^{\frac{\chi}{k}}\} \times \mathfrak{B}(\tau, s, m, \epsilon) ds| \quad (19) \end{aligned}$$

where

$$\begin{aligned} \mathfrak{B}(\tau, s, m, \epsilon) &= \frac{e^{-\beta|m|}}{(1+|m|)^\mu} \exp(v \frac{|s|}{|\epsilon|^k}) \exp(v \frac{|\tau^k - s|}{|\epsilon|^k}) \frac{|s|^{1/k}}{|\epsilon|} \frac{|\tau^k - s|^{1/k}}{|\epsilon|} \\ & \quad \times \frac{1}{1 + \frac{|s|^2}{|\epsilon|^{2k}}} \frac{1}{1 + \frac{|\tau^k - s|^2}{|\epsilon|^{2k}}} (\tau^k - s)^{-1} s^{v_2}. \end{aligned}$$

Hence,

$$\begin{aligned} & \|a_{\gamma_1, k}(\tau) \int_0^{\tau^k} (\tau^k - s)^{\frac{\chi}{k}-1} s^{v_2} f(s^{1/k}, m) ds\|_{(v, \beta, \mu, k, \epsilon)} \\ & \leq C_{2.2}(\epsilon) C_{2.3}(\epsilon) \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (20) \end{aligned}$$

where

$$C_{2.2}(\epsilon) = \sup_{x \geq 0} \exp\left(-v \frac{x}{|\epsilon|^k}\right) \frac{1 + \frac{x^2}{|\epsilon|^{2k}}}{\frac{x^{1/k}}{|\epsilon|}} x^{\frac{\chi}{k}},$$

$$C_{2.3}(\epsilon) = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \frac{1}{(1 + |\tau|^k)^{\gamma_1}} \\ \times \int_0^{|\tau|^k} \frac{h^{1/k}}{|\epsilon|} \frac{(|\tau|^k - h)^{1/k}}{|\epsilon|} \frac{1}{1 + \frac{h^2}{|\epsilon|^{2k}}} \frac{1}{1 + \frac{(|\tau|^k - h)^2}{|\epsilon|^{2k}}} (|\tau|^k - h)^{-1} h^{v_2} dh. \quad (21)$$

By using the classical estimates

$$\sup_{x \geq 0} x^{m_1} \exp(-m_2 x) = \left(\frac{m_1}{m_2}\right)^{m_1} e^{-m_1} \quad (22)$$

for any real numbers $m_1 \geq 0$ and $m_2 > 0$, we get that

$$C_{2.2}(\epsilon) \leq |\epsilon|^\chi \left(\left(\frac{\chi - 1}{kv}\right)^{\frac{\chi-1}{k}} e^{-(\frac{\chi-1}{k})} + \left(\frac{2 + \frac{\chi-1}{k}}{v}\right)^{2 + \frac{\chi-1}{k}} e^{-(2 + \frac{\chi-1}{k})} \right). \quad (23)$$

Making the change of variable $h = |\epsilon|^k h'$ in the integral involved in the definition of $C_{2.3}(\epsilon)$ yields

$$C_{2.3}(\epsilon) = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \frac{1}{(1 + |\epsilon|^k |\frac{\tau}{\epsilon}|^k)^{\gamma_1}} \\ \times \int_0^{|\frac{\tau}{\epsilon}|^k} (h')^{1/k} (|\frac{\tau}{\epsilon}|^k - h')^{1/k} \frac{1}{1 + (h')^2} \frac{1}{1 + (|\frac{\tau}{\epsilon}|^k - h')^2} (|\frac{\tau}{\epsilon}|^k - h')^{-1} |\epsilon|^{kv_2} (h')^{v_2} dh' \\ \leq |\epsilon|^{kv_2} \sup_{x \geq 0} B_{2.3}(x, \epsilon) \quad (24)$$

where

$$B_{2.3}(x, \epsilon) = \frac{1 + x^2}{x^{1/k}} \frac{1}{(1 + |\epsilon|^k x)^{\gamma_1}} \int_0^x \frac{1}{(1 + h^2)(1 + (x - h)^2)} \frac{1}{(x - h)^{1 - \frac{1}{k}}} h^{\frac{1}{k} + v_2} dh.$$

For any $x > 0$, we have that $B_{2.3}(x, \epsilon) \leq \tilde{B}_{2.3}(x, \epsilon)$, where

$$\tilde{B}_{2.3}(x, \epsilon) = \frac{1 + x^2}{(1 + |\epsilon|^k x)^{\gamma_1}} \int_0^x \frac{1}{(1 + h^2)(1 + (x - h)^2)} \frac{1}{(x - h)^{1 - \frac{1}{k}}} h^{v_2} dh.$$

Let $x_0 > 0$. From the inequality $1 + |\epsilon|^k x \geq 1$, for all $x \in [0, x_0]$, $\epsilon \in \mathcal{E}$, there exists a constant $\tilde{B}_{2,3} > 0$ such that

$$\sup_{x \in [0, x_0], \epsilon \in \mathcal{E}} \tilde{B}_{2,3}(x, \epsilon) \leq \tilde{B}_{2,3}. \quad (25)$$

On the other hand, since $1 + |\epsilon|^k x \geq |\epsilon|^k x$ holds for all $x \geq 0$ and $\epsilon \in \mathcal{E}$, we get that

$$\tilde{B}_{2,3}(x, \epsilon) \leq \frac{\tilde{B}_{2,4}(x)}{|\epsilon|^{k\gamma_1}} \quad (26)$$

where

$$\tilde{B}_{2,4}(x) = (1 + x^2)x^{-\gamma_1} \int_0^x \frac{1}{(1 + h^2)(1 + (x - h)^2)} \frac{1}{(x - h)^{1 - \frac{1}{k}}} h^{\nu_2} dh$$

for all $x \geq x_0$. Now, we make the change of variable $h = xu$ in the integral inside $\tilde{B}_{2,4}(x)$. We can write

$$\tilde{B}_{2,4}(x) = (1 + x^2)x^{\nu_2 + \frac{1}{k} - \gamma_1} F_k(x)$$

where

$$F_k(x) = \int_0^1 \frac{u^{\nu_2}}{(1 + x^2 u^2)(1 + x^2(1 - u)^2)(1 - u)^{1 - \frac{1}{k}}} du.$$

Using a partial fraction decomposition, we can split $F_k = F_{1,k}(x) + F_{2,k}(x)$, where

$$F_{1,k}(x) = \frac{1}{4 + x^2} \int_0^1 \frac{(2u + 1)u^{\nu_2}}{(1 + x^2 u^2)(1 - u)^{1 - \frac{1}{k}}} du,$$

$$F_{2,k}(x) = \frac{1}{4 + x^2} \int_0^1 \frac{(3 - 2u)u^{\nu_2}}{(1 + x^2(1 - u)^2)(1 - u)^{1 - \frac{1}{k}}} du.$$

In particular, we observe that there exist two constants $\mathfrak{F}_{1,k}, \mathfrak{F}_{2,k} > 0$ such that

$$F_{1,k}(x) \leq \frac{\mathfrak{F}_{1,k}}{4 + x^2}, \quad F_{2,k}(x) \leq \frac{\mathfrak{F}_{2,k}}{4 + x^2} \quad (27)$$

for all $x \geq x_0$. Hence, if one assumes that $\nu_2 + \frac{1}{k} - \gamma_1 \leq 0$, then we get a constant $\tilde{B}_{2,4,1} > 0$ such that

$$\sup_{x \geq x_0} \tilde{B}_{2,3}(x, \epsilon) \leq \frac{1}{|\epsilon|^{k\gamma_1}} \sup_{x \geq x_0} \tilde{B}_{2,4}(x) \leq \frac{\tilde{B}_{2,4,1}}{|\epsilon|^{k\gamma_1}} \quad (28)$$

Finally, gathering all the estimates (20), (23), (24), (25), (28), we get (12). \square

Proposition 3. Let $k \geq 1$ be an integer. Let $Q_1(X), Q_2(X), R(X) \in \mathbb{C}[X]$ such that

$$\deg(R) \geq \deg(Q_1), \quad \deg(R) \geq \deg(Q_2), \quad R(im) \neq 0 \quad (29)$$

for all $m \in \mathbb{R}$. Assume that $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$. Let $m \mapsto b(m)$ be a continuous function on \mathbb{R} such that

$$|b(m)| \leq \frac{1}{|R(im)|}$$

for all $m \in \mathbb{R}$. Then, there exists a constant $C_3 > 0$ (depending on Q_1, Q_2, R, μ, k, v) such that

$$\begin{aligned} & \|b(m) \int_0^{\tau^k} (\tau^k - s)^{\frac{1}{k}} \left(\int_0^s \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) f((s - x)^{1/k}, m - m_1) \right. \\ & \quad \times Q_2(im_1) g(x^{1/k}, m_1) \frac{1}{(s - x)x} dx dm_1 ds \|_{(v, \beta, \mu, k, \epsilon)} \\ & \leq C_3 |\epsilon| \|f(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \|g(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \end{aligned} \quad (30)$$

for all $f(\tau, m), g(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$.

Proof. Let $f(\tau, m), g(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$. For any $\tau \in \bar{D}(0, \rho) \cup S_d$, the segment $[0, \tau^k]$ is such that for any $s \in [0, \tau^k]$, any $x \in [0, s]$, the expressions $f((s - x)^{1/k}, m - m_1)$ and $g(x^{1/k}, m_1)$ are well defined, provided that $m, m_1 \in \mathbb{R}$. By definition, we can write

$$\begin{aligned} & \|b(m) \int_0^{\tau^k} (\tau^k - s)^{\frac{1}{k}} \left(\int_0^s \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) f((s - x)^{1/k}, m - m_1) \right. \\ & \quad \times Q_2(im_1) g(x^{1/k}, m_1) \frac{1}{(s - x)x} dx dm_1 ds \|_{(v, \beta, \mu, k, \epsilon)} \\ & = \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1 + |m|)^\mu \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \\ & \quad \times \left| \int_0^{\tau^k} (\tau^k - s)^{1/k} \left(\int_0^s \int_{-\infty}^{+\infty} \{(1 + |m - m_1|)^\mu e^{\beta|m - m_1|} \frac{1 + \frac{|s - x|^2}{|\epsilon|^{2k}}}{\frac{|s - x|^{1/k}}{|\epsilon|}} \exp(-v|s - x|/|\epsilon|^k) \right. \right. \\ & \quad \times f((s - x)^{1/k}, m - m_1)\} \times \{(1 + |m_1|)^\mu e^{\beta|m_1|} \frac{1 + \frac{|x|^2}{|\epsilon|^{2k}}}{\frac{|x|^{1/k}}{|\epsilon|}} \exp(-v|x|/|\epsilon|^k) g(x^{1/k}, m_1)\} \\ & \quad \times C(s, x, m, m_1, \epsilon) dx dm_1 ds \Big| \end{aligned}$$

where

$$\begin{aligned} \mathcal{C}(s, x, m, m_1, \epsilon) &= \frac{\exp(-\beta|m_1|)\exp(-\beta|m-m_1|)}{(1+|m-m_1|)^\mu(1+|m_1|)^\mu} b(m) Q_1(i(m-m_1)) Q_2(im_1) \\ &\quad \times \frac{\frac{|s-x|^{1/k}|x|^{1/k}}{|\epsilon|^2}}{(1+\frac{|s-x|^2}{|\epsilon|^{2k}})(1+\frac{|x|^2}{|\epsilon|^{2k}})} \times \exp(v|s-x|/|\epsilon|^k) \exp(v|x|/|\epsilon|^k) \frac{1}{(s-x)x} \end{aligned}$$

Now, we know that there exist $\mathfrak{Q}_1, \mathfrak{Q}_2, \mathfrak{R} > 0$ with

$$\begin{aligned} |Q_1(i(m-m_1))| &\leq \mathfrak{Q}_1(1+|m-m_1|)^{\deg(Q_1)}, \quad |Q_2(im_1)| \leq \mathfrak{Q}_2(1+|m_1|)^{\deg(Q_2)}, \\ |R(im)| &\geq \mathfrak{R}(1+|m|)^{\deg(R)} \end{aligned} \quad (31)$$

for all $m, m_1 \in \mathbb{R}$. Therefore,

$$\begin{aligned} &||b(m) \int_0^{\tau^k} (\tau^k - s)^{\frac{1}{k}} (\int_0^s \int_{-\infty}^{+\infty} Q_1(i(m-m_1)) f((s-x)^{1/k}, m-m_1) \\ &\quad \times Q_2(im_1) g(x^{1/k}, m_1) \frac{1}{(s-x)x} dx dm_1) ds||_{(v, \beta, \mu, k, \epsilon)} \\ &\leq C_3(\epsilon) ||f(\tau, m)||_{(v, \beta, \mu, k, \epsilon)} ||g(\tau, m)||_{(v, \beta, \mu, k, \epsilon)} \end{aligned} \quad (32)$$

where

$$\begin{aligned} C_3(\epsilon) &= \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1+|m|)^\mu \frac{1+|\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \frac{1}{\mathfrak{R}(1+|m|)^{\deg(R)}} \\ &\quad \times \int_0^{|\tau|^k} (|\tau|^k - h)^{1/k} (\int_0^h \int_{-\infty}^{+\infty} \frac{\exp(-\beta|m_1|)\exp(-\beta|m-m_1|)}{(1+|m-m_1|)^\mu(1+|m_1|)^\mu} \\ &\quad \times \mathfrak{Q}_1 \mathfrak{Q}_2 (1+|m-m_1|)^{\deg(Q_1)} (1+|m_1|)^{\deg(Q_2)} \frac{\frac{(h-x)^{1/k} x^{1/k}}{|\epsilon|^2}}{(1+\frac{(h-x)^2}{|\epsilon|^{2k}})(1+\frac{x^2}{|\epsilon|^{2k}})} \\ &\quad \times \exp(v(h-x)/|\epsilon|^k) \exp(vx/|\epsilon|^k) \frac{1}{(h-x)x} dx dm_1) dh \end{aligned} \quad (33)$$

Using the triangular inequality $|m| \leq |m_1| + |m-m_1|$, for all $m, m_1 \in \mathbb{R}$, we get that $C_3(\epsilon) \leq C_{3.1} C_{3.2}(\epsilon)$ where

$$C_{3.1} = \frac{\mathfrak{Q}_1 \mathfrak{Q}_2}{\mathfrak{R}} \sup_{m \in \mathbb{R}} (1+|m|)^{\mu - \deg(R)} \int_{-\infty}^{+\infty} \frac{1}{(1+|m-m_1|)^{\mu - \deg(Q_1)} (1+|m_1|)^{\mu - \deg(Q_2)}} dm_1 \quad (34)$$

which is finite whenever $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ under the assumption (29) using the same estimates as in Lemma 4 of [22] (see also Lemma 2.2 from [9]), and where

$$C_{3,2}(\epsilon) = \sup_{\tau \in \bar{D}(0,\rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \\ \times \int_0^{|\tau|^k} (|\tau|^k - h)^{1/k} \exp(vh/|\epsilon|^k) \int_0^h \frac{\frac{(h-x)^{1/k} x^{1/k}}{|\epsilon|^2}}{(1 + \frac{(h-x)^2}{|\epsilon|^{2k}})(1 + \frac{x^2}{|\epsilon|^{2k}})} \frac{1}{(h-x)x} dx dh. \quad (35)$$

Making the changes of variables $h = |\epsilon|^k h'$ and $x = |\epsilon|^k x'$, we get that

$$\int_0^{|\tau|^k} (|\tau|^k - h)^{1/k} \exp(vh/|\epsilon|^k) \int_0^h \frac{\frac{(h-x)^{1/k} x^{1/k}}{|\epsilon|^2}}{(1 + \frac{(h-x)^2}{|\epsilon|^{2k}})(1 + \frac{x^2}{|\epsilon|^{2k}})} \frac{1}{(h-x)x} dx dh \\ = |\epsilon| \int_0^{|\frac{\tau}{\epsilon}|^k} (|\frac{\tau}{\epsilon}|^k - h')^{1/k} \exp(vh') \int_0^{h'} \frac{1}{(1 + (h' - x')^2)(1 + x'^2)} \frac{1}{(h' - x')^{1-\frac{1}{k}} x'^{1-\frac{1}{k}}} dx' dh' \quad (36)$$

From (35) and (36), we get that $C_{3,2}(\epsilon) \leq |\epsilon| C_{3,3}$, where

$$C_{3,3} = \sup_{x \geq 0} \frac{1 + x^2}{x^{1/k}} \exp(-vx) \int_0^x (x - h')^{1/k} \exp(vh') \\ \times \left(\int_0^{h'} \frac{1}{(1 + (h' - x')^2)(1 + x'^2)} \frac{1}{(h' - x')^{1-\frac{1}{k}} x'^{1-\frac{1}{k}}} dx' \right) dh' \quad (37)$$

Again by the change of variable $x' = h'u$, for $u \in [0, 1]$, we can write

$$\int_0^{h'} \frac{1}{(1 + (h' - x')^2)(1 + x'^2)} \frac{1}{(h' - x')^{1-\frac{1}{k}} x'^{1-\frac{1}{k}}} dx' \\ = \frac{1}{h'^{1-\frac{2}{k}}} \int_0^1 \frac{1}{(1 + (h')^2(1 - u)^2)(1 + h'^2 u^2)(1 - u)^{1-\frac{1}{k}} u^{1-\frac{1}{k}}} du = J_k(h') \quad (38)$$

Using a partial fraction decomposition, we can split $J_k(h') = J_{1,k}(h') + J_{2,k}(h')$, where

$$J_{1,k}(h') = \frac{1}{h'^{1-\frac{2}{k}}(h'^2 + 4)} \int_0^1 \frac{3 - 2u}{(1 + h'^2(1 - u)^2)(1 - u)^{1-\frac{1}{k}} u^{1-\frac{1}{k}}} du \\ J_{2,k}(h') = \frac{1}{h'^{1-\frac{2}{k}}(h'^2 + 4)} \int_0^1 \frac{2u + 1}{(1 + h'^2 u^2)(1 - u)^{1-\frac{1}{k}} u^{1-\frac{1}{k}}} du \quad (39)$$

From now on, we assume that $k \geq 2$. By construction of $J_{1,k}(h')$ and $J_{2,k}(h')$, we see that there exists a constant $j_k > 0$ such that

$$J_k(h') \leq \frac{j_k}{h'^{1-\frac{2}{k}}(h'^2 + 4)} \quad (40)$$

for all $h' \geq 0$. From (37) and (40), we deduce that $C_{3,3} \leq \sup_{x \geq 0} \tilde{C}_{3,3}(x)$, where

$$\tilde{C}_{3,3}(x) = (1+x^2) \exp(-\nu x) \int_0^x \frac{j_k \exp(\nu h')}{h'^{1-\frac{2}{k}}(h'^2 + 4)} dh'. \quad (41)$$

From L'Hospital rule, we know that

$$\lim_{x \rightarrow +\infty} \tilde{C}_{3,3}(x) = \lim_{x \rightarrow +\infty} \frac{j_k}{x^{1-\frac{2}{k}}} \frac{\frac{(1+x^2)^2}{x^2+4}}{\nu(1+x^2) - 2x}$$

is finite when $k \geq 2$. Therefore, we get a constant $\tilde{C}_{3,3} > 0$ such that

$$\sup_{x \geq 0} \tilde{C}_{3,3}(x) \leq \tilde{C}_{3,3}. \quad (42)$$

Taking into account the estimates for (33), (34), (35), (37), (41) and (42), we obtain the result (30) when $k \geq 2$.

In the remaining case $k = 1$, from Corollary 4.9 of [8] one can check the existence of a constant $j_1 > 0$ such that

$$J_1(h') \leq \frac{j_1}{h'^2 + 1} \quad (43)$$

for all $h' \geq 0$. From (37) and (43), we deduce that $C_{3,3} \leq \sup_{x \geq 0} \tilde{C}_{3,3,1}(x)$, where

$$\tilde{C}_{3,3,1}(x) = (1+x^2) \exp(-\nu x) \int_0^x \frac{j_1 \exp(\nu h')}{h'^2 + 1} dh'. \quad (44)$$

From L'Hospital rule, we know that

$$\lim_{x \rightarrow +\infty} \tilde{C}_{3,3,1}(x) = \lim_{x \rightarrow +\infty} \frac{j_1(1+x^2)}{\nu(1+x^2) - 2x}$$

is finite. Therefore, we get a constant $\tilde{C}_{3,3,1} > 0$ such that

$$\sup_{x \geq 0} \tilde{C}_{3,3,1}(x) \leq \tilde{C}_{3,3,1}. \quad (45)$$

Taking into account the estimates for (33), (34), (35), (37), (44) and (45), we obtain the result (30) for $k = 1$. \square

Definition 2. Let $\beta, \mu \in \mathbb{R}$. We denote by $E_{(\beta, \mu)}$ the vector space of continuous functions $h : \mathbb{R} \rightarrow \mathbb{C}$ such that

$$\|h(m)\|_{(\beta, \mu)} = \sup_{m \in \mathbb{R}} (1 + |m|)^\mu \exp(\beta|m|)|h(m)|$$

is finite. The space $E_{(\beta, \mu)}$ equipped with the norm $\|\cdot\|_{(\beta, \mu)}$ is a Banach space.

Proposition 4. Let $k \geq 1$ be an integer. Let $Q(X), R(X) \in \mathbb{C}[X]$ be polynomials such that

$$\deg(R) \geq \deg(Q), \quad R(im) \neq 0 \quad (46)$$

for all $m \in \mathbb{R}$. Assume that $\mu > \deg(Q) + 1$. Let $m \mapsto b(m)$ be a continuous function such that

$$|b(m)| \leq \frac{1}{|R(im)|}$$

for all $m \in \mathbb{R}$. Then, there exists a constant $C_4 > 0$ (depending on Q, R, μ, k, v) such that

$$\begin{aligned} & \|b(m) \int_0^{\tau^k} (\tau^k - s)^{\frac{1}{k}} \int_{-\infty}^{+\infty} f(m - m_1) Q(im_1) g(s^{1/k}, m_1) dm_1 \frac{ds}{s}\|_{(v, \beta, \mu, k, \epsilon)} \\ & \leq C_4 |\epsilon| \|f(m)\|_{(\beta, \mu)} \|g(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \end{aligned} \quad (47)$$

for all $f(m) \in E_{(\beta, \mu)}$, all $g(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$.

Proof. The proof follows the same lines of arguments as those of [Propositions 1 and 3](#). Let $f(m) \in E_{(\beta, \mu)}$, $g(\tau, m) \in F_{(v, \beta, \mu, k, \epsilon)}^d$. We can write

$$\begin{aligned} N_2 &:= \|b(m) \int_0^{\tau^k} (\tau^k - s)^{\frac{1}{k}} \int_{-\infty}^{+\infty} f(m - m_1) Q(im_1) g(s^{1/k}, m_1) dm_1 \frac{ds}{s}\|_{(v, \beta, \mu, k, \epsilon)} \\ &= \sup_{\tau \in \bar{D}(0, \rho) \cup S_d, m \in \mathbb{R}} (1 + |m|)^\mu \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(\beta|m| - v|\frac{\tau}{\epsilon}|^k) \\ &\quad \times |b(m) \int_0^{\tau^k} \int_{-\infty}^{+\infty} \{(1 + |m - m_1|)^\mu \exp(\beta|m - m_1|) f(m - m_1)\} \\ &\quad \times \{(1 + |m_1|)^\mu \exp(\beta|m_1|) \exp(-\frac{v|s|}{|\epsilon|^k}) \frac{1 + \frac{|s|^2}{|\epsilon|^{2k}}}{\frac{|s|^{1/k}}{|\epsilon|}} g(s^{1/k}, m_1)\} \\ &\quad \times \mathcal{D}(\tau, s, m, m_1, \epsilon) dm_1 ds| \end{aligned} \quad (48)$$

where

$$\mathcal{D}(\tau, s, m, m_1, \epsilon) = \frac{Q(im_1)e^{-\beta|m_1|}e^{-\beta|m-m_1|}}{(1+|m-m_1|)^\mu(1+|m_1|)^\mu} \times \frac{\exp(\frac{\nu|s|}{|\epsilon|^k})}{1+\frac{|s|^2}{|\epsilon|^{2k}}} \frac{|s|^{1/k}}{|\epsilon|} (\tau^k - s)^{1/k} \frac{1}{s}$$

Again, we know that there exist constants $\mathfrak{Q}, \mathfrak{R} > 0$ such that

$$|Q(im_1)| \leq \mathfrak{Q}(1+|m_1|)^{\deg(Q)}, \quad |R(im)| \geq \mathfrak{R}(1+|m|)^{\deg(R)}$$

for all $m, m_1 \in \mathbb{R}$. By means of the triangular inequality $|m| \leq |m_1| + |m - m_1|$, we get that

$$N_2 \leq C_{4.1}(\epsilon) C_{4.2} \|f(m)\|_{(\beta, \mu)} \|g(\tau, m)\|_{(\nu, \beta, \mu, k, \epsilon)} \quad (49)$$

where

$$C_{4.1}(\epsilon) = \sup_{\tau \in \tilde{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-\nu |\frac{\tau}{\epsilon}|^k) \int_0^{|\tau|^k} \frac{\exp(\nu h/|\epsilon|^k)}{1 + \frac{h^2}{|\epsilon|^{2k}}} \frac{h^{\frac{1}{k}-1}}{|\epsilon|} (|\tau|^k - h)^{1/k} dh$$

and

$$C_{4.2} = \frac{\mathfrak{Q}}{\mathfrak{R}} \sup_{m \in \mathbb{R}} (1+|m|)^{\mu - \deg(R)} \int_{-\infty}^{+\infty} \frac{1}{(1+|m-m_1|)^\mu (1+|m_1|)^{\mu - \deg(Q)}} dm_1.$$

From the estimates (9) and (10), we know that there exists a constant $C_{4.1} > 0$ such that

$$C_{4.1}(\epsilon) \leq C_{4.1} |\epsilon| \quad (50)$$

and from the estimates for (34), we know that $C_{4.2}$ is finite under the assumption (46) provided that $\mu > \deg(Q) + 1$. Finally, gathering this latter bound estimates together with (49) and (50) yields the result (47). \square

In the next proposition, we show that $(E_{(\beta, \mu)}, \|\cdot\|_{(\beta, \mu)})$ is a Banach algebra for some non-commutative product \star introduced below.

Proposition 5. *Let $Q_1(X), Q_2(X), R(X) \in \mathbb{C}[X]$ be polynomials such that*

$$\deg(R) \geq \deg(Q_1), \quad \deg(R) \geq \deg(Q_2), \quad R(im) \neq 0, \quad (51)$$

for all $m \in \mathbb{R}$. Assume that $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$. Then, there exists a constant $C_5 > 0$ (depending on Q_1, Q_2, R, μ) such that

$$\begin{aligned} & \left\| \frac{1}{R(im)} \int_{-\infty}^{+\infty} Q_1(i(m-m_1)) f(m-m_1) Q_2(im_1) g(m_1) dm_1 \right\|_{(\beta, \mu)} \\ & \leq C_5 \|f(m)\|_{(\beta, \mu)} \|g(m)\|_{(\beta, \mu)} \end{aligned} \quad (52)$$

for all $f(m), g(m) \in E_{(\beta, \mu)}$. Therefore, $(E_{(\beta, \mu)}, \|\cdot\|_{(\beta, \mu)})$ becomes a Banach algebra for the product \star defined by

$$f \star g(m) = \frac{1}{R(im)} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) f(m - m_1) Q_2(im_1) g(m_1) dm_1.$$

As a particular case, when $f, g \in E_{(\beta, \mu)}$ with $\beta > 0$ and $\mu > 1$, the classical convolution product

$$f * g(m) = \int_{-\infty}^{+\infty} f(m - m_1) g(m_1) dm_1$$

belongs to $E_{(\beta, \mu)}$.

Proof. The proof is similar to the one of [Proposition 3](#). Let $f(m), g(m) \in E_{(\beta, \mu)}$. We write

$$\begin{aligned} & \left\| \frac{1}{R(im)} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) f(m - m_1) Q_2(im_1) g(m_1) dm_1 \right\|_{(\beta, \mu)} \\ &= \sup_{m \in \mathbb{R}} (1 + |m|)^\mu e^{\beta|m|} \left| \frac{1}{R(im)} \int_{-\infty}^{+\infty} \{(1 + |m - m_1|)^\mu e^{\beta|m - m_1|} f(m - m_1)\} \right. \\ & \quad \times \left. \{(1 + |m_1|)^\mu e^{\beta|m_1|} g(m_1)\} \times \mathcal{E}(m, m_1) dm_1 \right| \end{aligned} \quad (53)$$

where

$$\mathcal{E}(m, m_1) = \frac{e^{-\beta|m - m_1|} e^{-\beta|m_1|}}{(1 + |m - m_1|)^\mu (1 + |m_1|)^\mu} Q_1(i(m - m_1)) Q_2(im_1).$$

Using the triangular inequality $|m| \leq |m_1| + |m - m_1|$ and the estimates in [\(31\)](#), we get that

$$\begin{aligned} & \left\| \frac{1}{R(im)} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) f(m - m_1) Q_2(im_1) g(m_1) dm_1 \right\|_{(\beta, \mu)} \\ & \leq C_5 \|f(m)\|_{(\beta, \mu)} \|g(m)\|_{(\beta, \mu)} \end{aligned} \quad (54)$$

where

$$C_5 = \frac{\Omega_1 \Omega_2}{\Re} \sup_{m \in \mathbb{R}} (1 + |m|)^{\mu - \deg(R)} \int_{-\infty}^{+\infty} \frac{1}{(1 + |m - m_1|)^{\mu - \deg(Q_1)} (1 + |m_1|)^{\mu - \deg(Q_2)}} dm_1$$

which is finite whenever $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ provided that [\(51\)](#) holds as explained in [Proposition 3](#) (see [\(34\)](#)). \square

3. The Laplace transform, asymptotic expansions and the Fourier transform

We give a definition of k -Borel summability of formal series with coefficients in a Banach space which is a slightly modified version of the one given in [1], Section 3.2, in order to fit our necessities.

Definition 3. Let $k \geq 1$ be an integer. Let $m_k(n)$ be the sequence defined by

$$m_k(n) = \Gamma\left(\frac{n}{k}\right) = \int_0^{+\infty} t^{\frac{n}{k}-1} e^{-t} dt$$

for all $n \geq 1$. A formal series

$$\hat{X}(T) = \sum_{n=1}^{\infty} a_n T^n \in T\mathbb{E}[[T]]$$

with coefficients in a Banach space $(\mathbb{E}, \|\cdot\|_{\mathbb{E}})$ is said to be m_k -summable with respect to t in the direction $d \in [0, 2\pi)$ if

i) there exists $\rho \in \mathbb{R}_+$ such that the following formal series, called a formal m_k -Borel transform of \hat{X}

$$\mathcal{B}_{m_k}(\hat{X})(\tau) = \sum_{n=1}^{\infty} \frac{a_n}{\Gamma(\frac{n}{k})} \tau^n \in \tau\mathbb{E}[[\tau]],$$

is absolutely convergent for $|\tau| < \rho$.

ii) there exists $\delta > 0$ such that the series $\mathcal{B}_{m_k}(\hat{X})(\tau)$ can be analytically continued with respect to τ in a sector $S_{d,\delta} = \{\tau \in \mathbb{C}^* : |d - \arg(\tau)| < \delta\}$. Moreover, there exist $C > 0$ and $K > 0$ such that

$$\|\mathcal{B}_{m_k}(\hat{X})(\tau)\|_{\mathbb{E}} \leq C e^{K|\tau|^k}$$

for all $\tau \in S_{d,\delta}$.

If this is so, the vector valued Laplace transform of $\mathcal{B}_{m_k}(\hat{X})(\tau)$ in the direction d is defined by

$$\mathcal{L}_{m_k}^d(\mathcal{B}(\hat{X}))(T) = k \int_{L_\gamma} \mathcal{B}_{m_k}(\hat{X})(u) e^{-(u/T)^k} \frac{du}{u},$$

along a half-line $L_\gamma = \mathbb{R}_+ e^{i\gamma} \subset S_{d,\delta} \cup \{0\}$, where γ depends on T and is chosen in such a way that $\cos(k(\gamma - \arg(T))) \geq \delta_1 > 0$, for some fixed δ_1 . The function $\mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}))(T)$ is well defined, holomorphic and bounded in any sector

$$S_{d,\theta,R^{1/k}} = \{T \in \mathbb{C}^* : |T| < R^{1/k}, \quad |d - \arg(T)| < \theta/2\},$$

where $\frac{\pi}{k} < \theta < \frac{\pi}{k} + 2\delta$ and $0 < R < \delta_1/K$. This function is called the m_k -sum of the formal series $\hat{X}(T)$ in the direction d .

We now state some elementary properties concerning the m_k -sums of formal power series.

1) The function $\mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}))(T)$ has the formal series $\hat{X}(T)$ as Gevrey asymptotic expansion of order $1/k$ with respect to t on $S_{d,\theta,R^{1/k}}$. This means that for all $\frac{\pi}{k} < \theta_1 < \theta$, there exist $C, M > 0$ such that

$$\|\mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}))(T) - \sum_{p=1}^{n-1} a_p T^p\|_{\mathbb{E}} \leq CM^n \Gamma(1 + \frac{n}{k}) |T|^n \quad (55)$$

for all $n \geq 2$, all $T \in S_{d,\theta_1,R^{1/k}}$. Moreover, from Watson's lemma (see Proposition 11 p. 75 in [1]), we get that $\mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}))(T)$ is the unique holomorphic function that satisfies the estimates (55) on the sectors $S_{d,\theta_1,R^{1/k}}$ with large aperture $\theta_1 > \frac{\pi}{k}$.

2) Let us assume that $(\mathbb{E}, \|\cdot\|_{\mathbb{E}})$ also has the structure of a Banach algebra for a product \star . Let $\hat{X}_1(T), \hat{X}_2(T) \in T\mathbb{E}[[T]]$ be m_k -summable formal power series in direction d . Let $q_1 \geq q_2 \geq 1$ be integers. We assume that $\hat{X}_1(T) + \hat{X}_2(T)$, $\hat{X}_1(T) \star \hat{X}_2(T)$ and $T^{q_1} \partial_T^{q_2} \hat{X}_1(T)$, which are elements of $T\mathbb{E}[[T]]$, are m_k -summable in direction d . Then, the following equalities

$$\begin{aligned} \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_1))(T) + \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_2))(T) &= \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_1 + \hat{X}_2))(T), \\ \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_1))(T) \star \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_2))(T) &= \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_1 \star \hat{X}_2))(T) \\ T^{q_1} \partial_T^{q_2} \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(\hat{X}_1))(T) &= \mathcal{L}_{m_k}^d(\mathcal{B}_{m_k}(T^{q_1} \partial_T^{q_2} \hat{X}_1))(T) \end{aligned} \quad (56)$$

hold for all $T \in S_{d,\theta,R^{1/k}}$. These equalities are consequence of the unicity of the function having a given Gevrey expansion of order $1/k$ in large sectors as stated above in 1) and from the fact that the set of holomorphic functions having the Gevrey asymptotic expansion of order $1/k$ on a sector with values in the Banach algebra \mathbb{E} form a differential algebra (meaning that this set is stable with respect to the sum and product of functions and derivation in the variable T) (see Theorems 18, 19 and 20 in [1]).

In the next proposition, we give some identities for the m_k -Borel transform that will be useful in the sequel.

Proposition 6. Let $\hat{f}(t) = \sum_{n \geq 1} f_n t^n$, $\hat{g}(t) = \sum_{n \geq 1} g_n t^n$ be formal series whose coefficients f_n, g_n belong to some Banach space $(\mathbb{E}, \|\cdot\|_{\mathbb{E}})$. We assume that $(\mathbb{E}, \|\cdot\|_{\mathbb{E}})$ is a Banach algebra for some product \star . Let $k, m \geq 1$ be integers. The following formal identities hold.

$$\mathcal{B}_{m_k}(t^{k+1} \partial_t \hat{f}(t))(\tau) = k \tau^k \mathcal{B}_{m_k}(\hat{f}(t))(\tau) \quad (57)$$

$$\mathcal{B}_{m_k}(t^m \hat{f}(t))(\tau) = \frac{\tau^k}{\Gamma(\frac{m}{k})} \int_0^{\tau^k} (\tau^k - s)^{\frac{m}{k}-1} \mathcal{B}_{m_k}(\hat{f}(t))(s^{1/k}) \frac{ds}{s} \quad (58)$$

and

$$\mathcal{B}_{m_k}(\hat{f}(t) \star \hat{g}(t))(\tau) = \tau^k \int_0^{\tau^k} \mathcal{B}_{m_k}(\hat{f}(t))((\tau^k - s)^{1/k}) \star \mathcal{B}_{m_k}(\hat{g}(t))(s^{1/k}) \frac{1}{(\tau^k - s)s} ds \quad (59)$$

Proof. First, we show (57). By definition, we have that

$$\mathcal{B}_{m_k}(\tau^{\frac{k+1}{k}} \partial_t \hat{f}(t))(\tau) = \sum_{n \geq 1} \frac{\frac{n}{k} f_n}{\Gamma(\frac{n}{k} + 1)} \tau^{n+k} \quad (60)$$

By application of the addition formula for the Gamma function which yields $\Gamma(\frac{n}{k} + 1) = \frac{n}{k} \Gamma(\frac{n}{k})$ for any $n \geq 1$, we deduce (57) from (60).

Now, we prove (58). By definition, we can write

$$\mathcal{B}_{m_k}(\tau^m \hat{f}(t))(\tau) = \frac{1}{\Gamma(\frac{m}{k})} \sum_{n \geq 1} \frac{f_n}{\Gamma(\frac{n}{k})} \frac{\Gamma(\frac{m}{k}) \Gamma(\frac{n}{k})}{\Gamma(\frac{m+n}{k})} \tau^{m+n}. \quad (61)$$

Using the Beta integral formula (see Appendix B in [2]), we can write

$$\frac{\Gamma(\frac{m}{k}) \Gamma(\frac{n}{k})}{\Gamma(\frac{m+n}{k})} = \frac{\tau^k}{\tau^{m+n}} \int_0^{\tau^k} (\tau^k - s)^{\frac{m}{k}-1} s^{\frac{n}{k}-1} ds \quad (62)$$

for any $m, n \geq 1$. Plugging (62) into (61) yields (58).

Finally, we show (59). By definition, we have

$$\mathcal{B}_{m_k}(\hat{f}(t) \star \hat{g}(t))(\tau) = \sum_{n \geq 2} \left(\sum_{p+q=n} \frac{f_p}{\Gamma(\frac{p}{k})} \star \frac{g_q}{\Gamma(\frac{q}{k})} \times \frac{\Gamma(\frac{p}{k}) \Gamma(\frac{q}{k})}{\Gamma(\frac{n}{k})} \right) \tau^n \quad (63)$$

Using again the Beta integral formula, we can write

$$\frac{\Gamma(\frac{p}{k}) \Gamma(\frac{q}{k})}{\Gamma(\frac{n}{k})} = \frac{\tau^k}{\tau^n} \int_0^{\tau^k} (\tau^k - s)^{\frac{p}{k}-1} s^{\frac{q}{k}-1} ds \quad (64)$$

when $p + q = n$ and $p, q \geq 1$. By the substitution of (64) into (63), we deduce (59). \square

In the following proposition, we recall some properties of the inverse the Fourier transform

Proposition 7. Let $f \in E_{(\beta, \mu)}$ with $\beta > 0$, $\mu > 1$. The inverse Fourier transform of f is defined by

$$\mathcal{F}^{-1}(f)(x) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} f(m) \exp(ixm) dm$$

for all $x \in \mathbb{R}$. The function $\mathcal{F}^{-1}(f)$ extends to an analytic function on the strip

$$H_\beta = \{z \in \mathbb{C} / |\operatorname{Im}(z)| < \beta\}. \quad (65)$$

Let $\phi(m) = \text{im}f(m) \in E_{(\beta, \mu-1)}$. Then, we have

$$\partial_z \mathcal{F}^{-1}(f)(z) = \mathcal{F}^{-1}(\phi)(z) \quad (66)$$

for all $z \in H_\beta$.

Let $g \in E_{(\beta, \mu)}$ and let $\psi(m) = \frac{1}{(2\pi)^{1/2}} f * g(m)$, the convolution product of f and g , for all $m \in \mathbb{R}$. From Proposition 5, we know that $\psi \in E_{(\beta, \mu)}$. Moreover, we have

$$\mathcal{F}^{-1}(f)(z)\mathcal{F}^{-1}(g)(z) = \mathcal{F}^{-1}(\psi)(z) \quad (67)$$

for all $z \in H_\beta$.

Proof. Let $f \in E_{(\beta, \mu)}$. It is straight to check that $\mathcal{F}^{-1}(f)$ is well defined on the real line. The fact that $\mathcal{F}^{-1}(f)$ extends to an analytic function on the strip H_β follows from the next inequality. There exists $C > 0$ such that

$$|f(m)| |\exp(izm)| \leq \frac{C}{(1 + |m|)^\mu} \exp((\beta' - \beta)|m|)$$

for all $m \in \mathbb{R}$, $z \in H_{\beta'}$, with $\beta' < \beta$. The relations (66), (67) are classical and can be found for instance in [26]. \square

4. Formal and analytic solutions of convolution initial value problems with complex parameters

Let $k \geq 1$ and $D \geq 2$ be integers. For $1 \leq l \leq D$, let $d_l, \delta_l, \Delta_l \geq 0$ be nonnegative integers. We assume that

$$1 = \delta_1, \quad \delta_l < \delta_{l+1}, \quad (68)$$

for all $1 \leq l \leq D - 1$. We make also the assumption that

$$d_D = (\delta_D - 1)(k + 1), \quad d_l > (\delta_l - 1)(k + 1), \quad \Delta_D = d_D - \delta_D + 1 \quad (69)$$

for all $1 \leq l \leq D - 1$. Let $Q(X), Q_1(X), Q_2(X), R_l(X) \in \mathbb{C}[X]$, $0 \leq l \leq D$, be polynomials such that

$$\begin{aligned} \deg(Q) &\geq \deg(R_D) \geq \deg(R_l), \quad \deg(R_D) \geq \deg(Q_1), \quad \deg(R_D) \geq \deg(Q_2), \\ Q(im) &\neq 0, \quad R_D(im) \neq 0 \end{aligned} \quad (70)$$

for all $m \in \mathbb{R}$, all $0 \leq l \leq D - 1$. We consider sequences of functions $m \mapsto C_{0,n}(m, \epsilon)$, for all $n \geq 0$ and $m \mapsto F_n(m, \epsilon)$, for all $n \geq 1$, that belong to the Banach space $E_{(\beta, \mu)}$ for some $\beta > 0$ and $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ and which depend holomorphically on $\epsilon \in D(0, \epsilon_0)$. We assume that there exist constants $K_0, T_0 > 0$ such that

$$||C_{0,n}(m, \epsilon)||_{(\beta, \mu)} \leq K_0 \left(\frac{1}{T_0}\right)^n, \quad ||F_n(m, \epsilon)||_{(\beta, \mu)} \leq K_0 \left(\frac{1}{T_0}\right)^n \quad (71)$$

for all $n \geq 1$, for all $\epsilon \in D(0, \epsilon_0)$. We define

$$C_0(T, m, \epsilon) = \sum_{n \geq 1} C_{0,n}(m, \epsilon) T^n, \quad F(T, m, \epsilon) = \sum_{n \geq 1} F_n(m, \epsilon) T^n$$

which are convergent series on $D(0, T_0/2)$ with values in $E_{(\beta, \mu)}$. We consider the following singular initial value problem

$$\begin{aligned} & Q(im)(\partial_T U(T, m, \epsilon)) \\ &= \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) U(T, m - m_1, \epsilon) \\ &\quad \times Q_2(im_1) U(T, m_1, \epsilon) dm_1 + \sum_{l=1}^D R_l(im) \epsilon^{\Delta_l - d_l + \delta_l - 1} T^{d_l} \partial_T^{\delta_l} U(T, m, \epsilon) \\ &\quad + \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_0(T, m - m_1, \epsilon) R_0(im_1) U(T, m_1, \epsilon) dm_1 \\ &\quad + \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) R_0(im_1) U(T, m_1, \epsilon) dm_1 + \epsilon^{-1} F(T, m, \epsilon) \end{aligned} \quad (72)$$

for given initial data $U(0, m, \epsilon) = 0$.

Proposition 8. *There exists a unique formal series*

$$\hat{U}(T, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) T^n$$

solution of (72) with initial data $U(0, m, \epsilon) \equiv 0$, where the coefficients $m \mapsto U_n(m, \epsilon)$ belong to $E_{(\beta, \mu)}$ for $\beta > 0$ and $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ given above and depend holomorphically on ϵ in $D(0, \epsilon_0) \setminus \{0\}$.

Proof. From Proposition 5 and the conditions in the statement above, we get that the coefficients $U_n(m, \epsilon)$ of $\hat{U}(T, m, \epsilon)$ are well defined, belong to $E_{(\beta, \mu)}$ for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, all $n \geq 1$ and satisfy the following recursion relation

$$\begin{aligned} & (n+1)U_{n+1}(m, \epsilon) \\ &= \frac{\epsilon^{-1}}{Q(im)} \sum_{n_1+n_2=n, n_1 \geq 1, n_2 \geq 1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) U_{n_1}(m - m_1, \epsilon) Q_2(im_1) U_{n_2}(m_1, \epsilon) dm_1 \\ &\quad + \sum_{l=1}^D \frac{R_l(im)}{Q(im)} \left(\epsilon^{\Delta_l - d_l + \delta_l - 1} \Pi_{j=0}^{\delta_l - 1} (n + \delta_l - d_l - j) \right) U_{n+\delta_l-d_l}(m, \epsilon) \end{aligned}$$

$$\begin{aligned}
& + \frac{\epsilon^{-1}}{Q(im)} \sum_{n_1+n_2=n, n_1 \geq 1, n_2 \geq 1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_{0,n_1}(m-m_1, \epsilon) R_0(im_1) U_{n_2}(m_1, \epsilon) dm_1 \\
& + \frac{\epsilon^{-1}}{(2\pi)^{1/2} Q(im)} \int_{-\infty}^{+\infty} C_{0,0}(m-m_1, \epsilon) R_0(im_1) U_n(m_1, \epsilon) dm_1 + \frac{\epsilon^{-1}}{Q(im)} F_n(m, \epsilon) \quad (73)
\end{aligned}$$

for all $n \geq \max_{1 \leq l \leq D} d_l$. \square

Using the formula from [27], p. 40, we can expand the operators $T^{\delta_l(k+1)} \partial_T^{\delta_l}$ in the form

$$T^{\delta_l(k+1)} \partial_T^{\delta_l} = (T^{k+1} \partial_T)^{\delta_l} + \sum_{1 \leq p \leq \delta_l - 1} A_{\delta_l, p} T^{k(\delta_l - p)} (T^{k+1} \partial_T)^p \quad (74)$$

where $A_{\delta_l, p}$, $p = 1, \dots, \delta_l - 1$ are real numbers. We define integers $d_{l,k} \geq 0$ to satisfy

$$d_l + k + 1 = \delta_l(k + 1) + d_{l,k} \quad (75)$$

for all $1 \leq l \leq D$. Multiplying equation (72) by T^{k+1} and using (74), we can rewrite equation (72) in the form

$$\begin{aligned}
& Q(im)(T^{k+1} \partial_T U(T, m, \epsilon)) \\
& = \epsilon^{-1} T^{k+1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} Q_1(i(m-m_1)) U(T, m-m_1, \epsilon) Q_2(im_1) U(T, m_1, \epsilon) dm_1 \\
& + \sum_{l=1}^D R_l(im) \left(\epsilon^{\Delta_l - d_l + \delta_l - 1} T^{d_{l,k}} (T^{k+1} \partial_T)^{\delta_l} U(T, m, \epsilon) \right. \\
& + \sum_{1 \leq p \leq \delta_l - 1} A_{\delta_l, p} \epsilon^{\Delta_l - d_l + \delta_l - 1} T^{k(\delta_l - p) + d_{l,k}} (T^{k+1} \partial_T)^p U(T, m, \epsilon) \Big) \\
& + \epsilon^{-1} T^{k+1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_0(T, m-m_1, \epsilon) R_0(im_1) U(T, m_1, \epsilon) dm_1 \\
& + \epsilon^{-1} T^{k+1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_{0,0}(m-m_1, \epsilon) R_0(im_1) U(T, m_1, \epsilon) dm_1 + \epsilon^{-1} T^{k+1} F(T, m, \epsilon) \quad (76)
\end{aligned}$$

We denote $\omega_k(\tau, m, \epsilon)$ the formal m_k -Borel transform of $\hat{U}(T, m, \epsilon)$ with respect to T , $\varphi_k(\tau, m, \epsilon)$ the formal m_k -Borel transform of $C_0(T, m, \epsilon)$ with respect to T and $\psi_k(\tau, m, \epsilon)$ the formal m_k -Borel transform of $F(T, m, \epsilon)$ with respect to T ,

$$\omega_k(\tau, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) \frac{\tau^n}{\Gamma(\frac{n}{k})}, \quad \varphi_k(\tau, m, \epsilon) = \sum_{n \geq 1} C_{0,n}(m, \epsilon) \frac{\tau^n}{\Gamma(\frac{n}{k})}$$

$$\psi_k(\tau, m, \epsilon) = \sum_{n \geq 1} F_n(m, \epsilon) \frac{\tau^n}{\Gamma(\frac{n}{k})}$$

Using (71) we get that $\varphi_k(\tau, m, \epsilon) \in F_{(v, \beta, \mu, k, \epsilon)}^d$ and $\psi_k(\tau, m, \epsilon) \in F_{(v, \beta, \mu, k, \epsilon)}^d$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, any unbounded sector S_d centered at 0 and bisecting direction $d \in \mathbb{R}$, for some $v > 0$. Indeed, we have that

$$\|\varphi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq \sum_{n \geq 1} \|C_{0,n}(m, \epsilon)\|_{(\beta, \mu)} \left(\sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \frac{|\tau|^n}{\Gamma(\frac{n}{k})} \right),$$

$$\|\psi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq \sum_{n \geq 1} \|F_n(m, \epsilon)\|_{(\beta, \mu)} \left(\sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \frac{|\tau|^n}{\Gamma(\frac{n}{k})} \right) \quad (77)$$

By using the classical estimates (22) and the Stirling formula $\Gamma(n/k) \sim (2\pi)^{1/2} (n/k)^{\frac{n}{k} - \frac{1}{2}} e^{-n/k}$ as n tends to $+\infty$, we get two constants $A_1, A_2 > 0$ depending on v, k such that

$$\begin{aligned} & \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} \frac{1 + |\frac{\tau}{\epsilon}|^{2k}}{|\frac{\tau}{\epsilon}|} \exp(-v|\frac{\tau}{\epsilon}|^k) \frac{|\tau|^n}{\Gamma(\frac{n}{k})} \\ &= \sup_{\tau \in \bar{D}(0, \rho) \cup S_d} |\epsilon|^n (1 + |\frac{\tau}{\epsilon}|^{2k}) |\frac{\tau}{\epsilon}|^{n-1} \frac{\exp(-v|\frac{\tau}{\epsilon}|^k)}{\Gamma(\frac{n}{k})} \\ &\leq \epsilon_0^n \sup_{x \geq 0} (1 + x^2) x^{\frac{n-1}{k}} \frac{e^{-vx}}{\Gamma(\frac{n}{k})} \\ &\leq \epsilon_0^n \left(\left(\frac{n-1}{vk} \right)^{\frac{n-1}{k}} e^{-\frac{n-1}{k}} + \left(\frac{n-1}{vk} + \frac{2}{v} \right)^{\frac{n-1}{k} + 2} e^{-(\frac{n-1}{k} + 2)} \right) / \Gamma(n/k) \\ &\leq A_1 \epsilon_0^n (A_2)^n \end{aligned} \quad (78)$$

for all $n \geq 0$, all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. Therefore, if ϵ_0 fulfills $\epsilon_0 A_2 < T_0$, we get the estimates

$$\|\varphi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq A_1 \sum_{n \geq 1} \|C_{0,n}(m, \epsilon)\|_{(\beta, \mu)} (\epsilon_0 A_2)^n \leq \frac{A_1 A_2 K_0}{T_0} \frac{\epsilon_0}{1 - \frac{A_2}{T_0} \epsilon_0},$$

$$\|\psi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq A_1 \sum_{n \geq 1} \|F_n(m, \epsilon)\|_{(\beta, \mu)} (\epsilon_0 A_2)^n \leq \frac{A_1 A_2 K_0}{T_0} \frac{\epsilon_0}{1 - \frac{A_2}{T_0} \epsilon_0} \quad (79)$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Using the computation rules for the formal m_k -Borel transform in Proposition 6, we deduce the following equation satisfied by $\omega_k(\tau, m, \epsilon)$,

$$\begin{aligned}
& Q(im)(k\tau^k\omega_k(\tau, m, \epsilon)) \\
&= \epsilon^{-1} \frac{\tau^k}{\Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \\
&\quad \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} Q_1(i(m - m_1))\omega_k((s - x)^{1/k}, m - m_1, \epsilon) \right. \\
&\quad \times Q_2(im_1)\omega_k(x^{1/k}, m_1, \epsilon) \frac{1}{(s - x)x} dx dm_1 \Big) \frac{ds}{s} \\
&\quad + R_D(im) \left(k^{\delta_D} \tau^{\delta_D k} \omega_k(\tau, m, \epsilon) \right. \\
&\quad + \sum_{1 \leq p \leq \delta_D - 1} A_{\delta_D, p} \frac{\tau^k}{\Gamma(\delta_D - p)} \int_0^{\tau^k} (\tau^k - s)^{\delta_D - p - 1} (k^p s^p \omega_k(s^{1/k}, m, \epsilon)) \frac{ds}{s} \Big) \\
&\quad + \sum_{l=1}^{D-1} R_l(im) \left(\epsilon^{\Delta_l - d_l + \delta_l - 1} \frac{\tau^k}{\Gamma(\frac{d_{l,k}}{k})} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} - 1} (k^{\delta_l} s^{\delta_l} \omega_k(s^{1/k}, m, \epsilon)) \frac{ds}{s} \right. \\
&\quad + \sum_{1 \leq p \leq \delta_l - 1} A_{\delta_l, p} \epsilon^{\Delta_l - d_l + \delta_l - 1} \frac{\tau^k}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p)} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} + \delta_l - p - 1} (k^p s^p \omega_k(s^{1/k}, m, \epsilon)) \frac{ds}{s} \Big) \\
&\quad + \epsilon^{-1} \frac{\tau^k}{\Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \\
&\quad \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} \varphi_k((s - x)^{1/k}, m - m_1, \epsilon) R_0(im_1)\omega_k(x^{1/k}, m_1, \epsilon) \frac{1}{(s - x)x} dx dm_1 \right) \frac{ds}{s} \\
&\quad + \epsilon^{-1} \frac{\tau^k}{\Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \frac{1}{(2\pi)^{1/2}} \left(\int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) R_0(im_1)\omega_k(s^{1/k}, m_1, \epsilon) dm_1 \right) \frac{ds}{s} \\
&\quad + \epsilon^{-1} \frac{\tau^k}{\Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \psi_k(s^{1/k}, m, \epsilon) \frac{ds}{s} \tag{80}
\end{aligned}$$

We make the additional assumption that there exists an unbounded sector

$$S_{Q, R_D} = \{z \in \mathbb{C} / |z| \geq r_{Q, R_D}, \quad |\arg(z) - d_{Q, R_D}| \leq \eta_{Q, R_D}\}$$

with direction $d_{Q, R_D} \in \mathbb{R}$, aperture $\eta_{Q, R_D} > 0$ for some radius $r_{Q, R_D} > 0$ such that

$$\frac{Q(im)}{R_D(im)} \in S_{Q, R_D} \quad (81)$$

for all $m \in \mathbb{R}$. We factorize the polynomial $P_m(\tau) = Q(im)k - R_D(im)k^{\delta_D} \tau^{(\delta_D-1)k}$ in the form

$$P_m(\tau) = -R_D(im)k^{\delta_D} \prod_{l=0}^{(\delta_D-1)k-1} (\tau - q_l(m)) \quad (82)$$

where

$$q_l(m) = \left(\frac{|Q(im)|}{|R_D(im)|k^{\delta_D-1}} \right)^{\frac{1}{(\delta_D-1)k}} \exp(\sqrt{-1}(\arg(\frac{Q(im)}{R_D(im)k^{\delta_D-1}}) \frac{1}{(\delta_D-1)k} + \frac{2\pi l}{(\delta_D-1)k})) \quad (83)$$

for all $0 \leq l \leq (\delta_D - 1)k - 1$, all $m \in \mathbb{R}$.

We choose an unbounded sector S_d centered at 0, a small closed disc $\bar{D}(0, \rho)$ and we prescribe the sector S_{Q, R_D} in such a way that the following conditions hold.

1) There exists a constant $M_1 > 0$ such that

$$|\tau - q_l(m)| \geq M_1(1 + |\tau|) \quad (84)$$

for all $0 \leq l \leq (\delta_D - 1)k - 1$, all $m \in \mathbb{R}$, all $\tau \in S_d \cup \bar{D}(0, \rho)$. Indeed, from (81) and the explicit expression (83) of $q_l(m)$, we first observe that $|q_l(m)| > 2\rho$ for every $m \in \mathbb{R}$, all $0 \leq l \leq (\delta_D - 1)k - 1$ for an appropriate choice of r_{Q, R_D} and of $\rho > 0$. We also see that for all $m \in \mathbb{R}$, all $0 \leq l \leq (\delta_D - 1)k - 1$, the roots $q_l(m)$ remain in a union \mathcal{U} of unbounded sectors centered at 0 that do not cover a full neighborhood of the origin in \mathbb{C}^* provided that η_{Q, R_D} is small enough. Therefore, one can choose an adequate sector S_d such that $S_d \cap \mathcal{U} = \emptyset$ with the property that for all $0 \leq l \leq (\delta_D - 1)k - 1$ the quotients $q_l(m)/\tau$ lay outside some small disc centered at 1 in \mathbb{C} for all $\tau \in S_d$, all $m \in \mathbb{R}$. This yields (84) for some small constant $M_1 > 0$.

2) There exists a constant $M_2 > 0$ such that

$$|\tau - q_{l_0}(m)| \geq M_2 |q_{l_0}(m)| \quad (85)$$

for some $l_0 \in \{0, \dots, (\delta_D - 1)k - 1\}$, all $m \in \mathbb{R}$, all $\tau \in S_d \cup \bar{D}(0, \rho)$. Indeed, for the sector S_d and the disc $\bar{D}(0, \rho)$ chosen as above in 1), we notice that for any fixed $0 \leq l_0 \leq (\delta_D - 1)k - 1$, the quotient $\tau/q_{l_0}(m)$ stays outside a small disc centered at 1 in \mathbb{C} for all $\tau \in S_d \cup \bar{D}(0, \rho)$, all $m \in \mathbb{R}$. Hence (85) must hold for some small constant $M_2 > 0$.

By construction of the roots (83) in the factorization (82) and using the lower bound estimates (84), (85), we get a constant $C_P > 0$ such that

$$\begin{aligned} |P_m(\tau)| &\geq M_1^{(\delta_D-1)k-1} M_2 |R_D(im)k^{\delta_D}| \left(\frac{|Q(im)|}{|R_D(im)|k^{\delta_D-1}} \right)^{\frac{1}{(\delta_D-1)k}} (1 + |\tau|)^{(\delta_D-1)k-1} \\ &\geq M_1^{(\delta_D-1)k-1} M_2 \frac{k^{\delta_D}}{(k^{\delta_D-1})^{\frac{1}{(\delta_D-1)k}}} (r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}} |R_D(im)| \end{aligned}$$

$$\begin{aligned}
& \times \left(\min_{x \geq 0} \frac{(1+x)^{(\delta_D-1)k-1}}{(1+x^k)^{(\delta_D-1)-\frac{1}{k}}} \right) (1+|\tau|^k)^{(\delta_D-1)-\frac{1}{k}} \\
& = C_P (r_{Q,R_D})^{\frac{1}{(\delta_D-1)k}} |R_D(im)| (1+|\tau|^k)^{(\delta_D-1)-\frac{1}{k}}
\end{aligned} \tag{86}$$

for all $\tau \in S_d \cup \bar{D}(0, \rho)$, all $m \in \mathbb{R}$.

In the next proposition, we give sufficient conditions under which equation (80) has a solution $\omega_k(\tau, m, \epsilon)$ in the Banach space $F_{(v,\beta,\mu,k,\epsilon)}^d$ where β, μ are defined above.

Proposition 9. *Under the assumption that*

$$\delta_D \geq \delta_l + \frac{2}{k}, \quad \Delta_l + k(1 - \delta_D) + 1 \geq 0, \tag{87}$$

for all $1 \leq l \leq D-1$, there exists a radius $r_{Q,R_D} > 0$, a constant $\varpi > 0$ and constants $\zeta_0, \zeta_1, \zeta_2 > 0$ (depending on $Q_1, Q_2, k, C_P, \mu, v, \epsilon_0, R_l, \Delta_l, \delta_l, d_l$ for $0 \leq l \leq D$) such that if

$$\|C_{0,0}(m, \epsilon)\|_{(\beta,\mu)} \leq \zeta_0, \quad \|\varphi_k(\tau, m, \epsilon)\|_{(v,\beta,\mu,k,\epsilon)} \leq \zeta_1, \quad \|\psi_k(\tau, m, \epsilon)\|_{(v,\beta,\mu,k,\epsilon)} \leq \zeta_2 \tag{88}$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, equation (80) has a unique solution $\omega_k^d(\tau, m, \epsilon)$ in the space $F_{(v,\beta,\mu,k,\epsilon)}^d$ where $\beta, \mu > 0$ are defined in Proposition 8 which verifies $\|\omega_k^d(\tau, m, \epsilon)\|_{(v,\beta,\mu,k,\epsilon)} \leq \varpi$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Proof. We start the proof with a lemma which provides appropriate conditions in order to apply a fixed point theorem.

Lemma 2. *One can choose the constant $r_{Q,R_D} > 0$, a constant ϖ small enough and three constants $\zeta_0, \zeta_1, \zeta_2 > 0$ (depending on $Q_1, Q_2, k, C_P, \mu, v, \epsilon_0, R_l, \Delta_l, \delta_l, d_l$ for $0 \leq l \leq D$) such that if*

$$\|C_{0,0}(m, \epsilon)\|_{(\beta,\mu)} \leq \zeta_0, \quad \|\varphi_k(\tau, m, \epsilon)\|_{(v,\beta,\mu,k,\epsilon)} \leq \zeta_1, \quad \|\psi_k(\tau, m, \epsilon)\|_{(v,\beta,\mu,k,\epsilon)} \leq \zeta_2$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$ the map \mathcal{H}_ϵ defined by

$$\begin{aligned}
& \mathcal{H}_\epsilon(w(\tau, m)) \\
& := \frac{\epsilon^{-1}}{P_m(\tau)\Gamma(1+\frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \\
& \quad \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) w((s-x)^{1/k}, m - m_1) \right. \\
& \quad \times \left. Q_2(im_1) w(x^{1/k}, m_1) \frac{1}{(s-x)x} dx dm_1 \right) \frac{ds}{s}
\end{aligned}$$

$$\begin{aligned}
& + \frac{R_D(im)}{P_m(\tau)} \left\{ \sum_{1 \leq p \leq \delta_D - 1} \frac{A_{\delta_D, p}}{\Gamma(\delta_D - p)} \int_0^{\tau^k} (\tau^k - s)^{\delta_D - p - 1} (k^p s^p w(s^{1/k}, m)) \frac{ds}{s} \right\} \\
& + \sum_{l=1}^{D-1} \frac{R_l(im)}{P_m(\tau)} \left\{ \frac{\epsilon^{\Delta_l - d_l + \delta_l - 1}}{\Gamma(\frac{d_{l,k}}{k})} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} - 1} (k^{\delta_l} s^{\delta_l} w(s^{1/k}, m)) \frac{ds}{s} \right. \\
& + \left. \sum_{1 \leq p \leq \delta_l - 1} \frac{A_{\delta_l, p} \epsilon^{\Delta_l - d_l + \delta_l - 1}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p)} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} + \delta_l - p - 1} (k^p s^p w(s^{1/k}, m)) \frac{ds}{s} \right\} \\
& + \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \\
& \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} \varphi_k((s-x)^{1/k}, m - m_1, \epsilon) R_0(im_1) w(x^{1/k}, m_1) \frac{1}{(s-x)x} dx dm_1 \right) \frac{ds}{s} \\
& + \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \frac{1}{(2\pi)^{1/2}} \left(\int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) R_0(im_1) w(s^{1/k}, m_1) dm_1 \right) \frac{ds}{s} \\
& + \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \psi_k(s^{1/k}, m, \epsilon) \frac{ds}{s} \tag{89}
\end{aligned}$$

satisfy the next properties.

i) The following inclusion holds

$$\mathcal{H}_\epsilon(\bar{B}(0, \varpi)) \subset \bar{B}(0, \varpi) \tag{90}$$

where $\bar{B}(0, \varpi)$ is the closed ball of radius $\varpi > 0$ centered at 0 in $F_{(v, \beta, \mu, k, \epsilon)}^d$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

ii) We have

$$\|\mathcal{H}_\epsilon(w_1) - \mathcal{H}_\epsilon(w_2)\|_{(v, \beta, \mu, k, \epsilon)} \leq \frac{1}{2} \|w_1 - w_2\|_{(v, \beta, \mu, k, \epsilon)} \tag{91}$$

for all $w_1, w_2 \in \bar{B}(0, \varpi)$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Proof. We first check the property (90). Let $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$ and $w(\tau, m)$ be in $F_{v, \beta, \mu, k, \epsilon}^d$. We take $\zeta_0, \zeta_1, \zeta_2, \varpi > 0$ such that

$$\begin{aligned}
\|w(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} & \leq \varpi, \quad \|C_{0,0}(m, \epsilon)\|_{(\beta, \mu)} \leq \zeta_0, \quad \|\varphi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq \zeta_1, \\
\|\psi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} & \leq \zeta_2,
\end{aligned}$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Using [Lemma 1](#) and [Proposition 3](#) with the lower bound estimates [\(86\)](#) we get that

$$\begin{aligned}
 & \left\| \frac{\epsilon^{-1}}{P_m(\tau)\Gamma(1+\frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \right. \\
 & \quad \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} Q_1(i(m-m_1)) w((s-x)^{1/k}, m-m_1) \right. \\
 & \quad \times \left. Q_2(im_1) w(x^{1/k}, m_1) \frac{1}{(s-x)x} dx dm_1 \right) \frac{ds}{s} \Big\|_{(v,\beta,\mu,k,\epsilon)} \\
 & \leq \frac{1}{\Gamma(1+\frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \|w(\tau, m)\|_{(v,\beta,\mu,k,\epsilon)}^2}{C_P(r_{Q,R_D})^{\frac{1}{(\delta_D-1)k}}} \\
 & \leq \frac{1}{\Gamma(1+\frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \varpi^2}{C_P(r_{Q,R_D})^{\frac{1}{(\delta_D-1)k}}} \quad (92)
 \end{aligned}$$

Moreover, for $0 \leq p \leq \delta_D - 1$ and by means of [Proposition 2 i\)](#), we deduce

$$\begin{aligned}
 & \left\| \frac{R_D(im)}{P_m(\tau)} \frac{A_{\delta_D,p}}{\Gamma(\delta_D-p)} \int_0^{\tau^k} (\tau^k - s)^{\delta_D-p-1} (k^p s^p w(s^{1/k}, m)) \frac{ds}{s} \right\|_{(v,\beta,\mu,k,\epsilon)} \\
 & \leq \frac{A_{\delta_D,p} k^p C_{2,1} |\epsilon|}{\Gamma(\delta_D-p) C_P(r_{Q,R})^{\frac{1}{(\delta_D-1)k}}} \|w(\tau, m)\|_{(v,\beta,\mu,k,\epsilon)} \\
 & \leq \frac{A_{\delta_D,p} k^p C_{2,1} \epsilon_0}{\Gamma(\delta_D-p) C_P(r_{Q,R})^{\frac{1}{(\delta_D-1)k}}} \varpi. \quad (93)
 \end{aligned}$$

With the help of [Proposition 2 ii\)](#) and due to the assumptions of [\(87\)](#) we also get that

$$\begin{aligned}
 & \left\| \frac{R_l(im)}{P_m(\tau)} \frac{\epsilon^{\Delta_l-d_l+\delta_l-1}}{\Gamma(\frac{d_{l,k}}{k})} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k}-1} (k^{\delta_l} s^{\delta_l} w(s^{1/k}, m)) \frac{ds}{s} \right\|_{(v,\beta,\mu,k,\epsilon)} \\
 & \leq \frac{k^{\delta_l} C_{2,2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D-1)k}}} |\epsilon|^{\Delta_l-d_l+\delta_l+k(\delta_l-\delta_D)+d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w(\tau, m)\|_{(v,\beta,\mu,k,\epsilon)} \\
 & \leq \frac{k^{\delta_l} C_{2,2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D-1)k}}} \epsilon_0^{\Delta_l-d_l+\delta_l+k(\delta_l-\delta_D)+d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \varpi, \quad (94)
 \end{aligned}$$

and that

$$\begin{aligned}
& \left\| \frac{R_l(im)}{P_m(\tau)} \frac{A_{\delta_l,p} \epsilon^{\Delta_l - d_l + \delta_l - 1}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p)} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} + \delta_l - p - 1} (k^p s^p w(s^{1/k}, m)) \frac{ds}{s} \right\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_l,p}| k^p C_{2,2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}}} |\epsilon|^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_l,p}| k^p C_{2,2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \varpi. \tag{95}
\end{aligned}$$

Using [Lemma 1](#) and [Proposition 3](#) again with the lower bound estimates [\(86\)](#) we get that

$$\begin{aligned}
& \left\| \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \right. \\
& \quad \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} \varphi_k((s-x)^{1/k}, m - m_1, \epsilon) \right. \\
& \quad \times \left. R_0(im_1) w(x^{1/k}, m_1) \frac{1}{(s-x)x} dx dm_1 \right) \frac{ds}{s} \left. \right\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k}) (2\pi)^{1/2}} \frac{C_3 \|\varphi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \|w(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)}}{C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}}} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k}) (2\pi)^{1/2}} \frac{C_3 \zeta_1 \varpi}{C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}}} \tag{96}
\end{aligned}$$

Moreover, using [Proposition 4](#), we also get

$$\begin{aligned}
& \left\| \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \frac{1}{(2\pi)^{1/2}} \left(\int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) \right. \right. \\
& \quad \times \left. \left. R_0(im_1) w(s^{1/k}, m_1) dm_1 \right) \frac{ds}{s} \right\|_{(v, \beta, \mu, \epsilon)} \leq \frac{1}{\Gamma(1 + \frac{1}{k}) (2\pi)^{1/2}} \frac{C_4 \zeta_0 \varpi}{C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}}} \tag{97}
\end{aligned}$$

Finally, from [Lemma 1](#) and [Proposition 1](#), one gets

$$\begin{aligned}
& \left\| \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \psi_k(s^{1/k}, m, \epsilon) \frac{ds}{s} \right\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{C_1}{\Gamma(1 + \frac{1}{k}) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}} \min_{m \in \mathbb{R}} |R_D(im)|} \|\psi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{C_1}{\Gamma(1 + \frac{1}{k}) C_P(r_{Q,R_D})^{\frac{1}{(\delta_D - 1)k}} \min_{m \in \mathbb{R}} |R_D(im)|} \zeta_2 \tag{98}
\end{aligned}$$

Now, we choose $\varpi, \zeta_0, \zeta_1, \zeta_2 > 0$ and $r_{Q, R_D} > 0$ such that

$$\begin{aligned} & \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \varpi^2}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} + \sum_{p=1}^{\delta_D - 1} \frac{|A_{\delta_D, p}| k^p C_{2,1} \epsilon_0}{\Gamma(\delta_D - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \varpi \\ & + \sum_{l=1}^{D-1} \frac{k^{\delta_l} C_{2,2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \varpi \\ & + \sum_{p=1}^{\delta_l - 1} \frac{|A_{\delta_l, p}| k^p C_{2,2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \varpi \\ & + \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{(C_3 \zeta_1 + C_4 \zeta_0) \varpi}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \\ & + \frac{C_1}{\Gamma(1 + \frac{1}{k}) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}} \min_{m \in \mathbb{R}} |R_D(im)|} \zeta_2 \leq \varpi \end{aligned} \quad (99)$$

Gathering all the norm estimates (92), (93), (94), (95), (96), (97), (98) with the constraint (99), one gets (90).

Now, we check the second property (91). Let $w_1(\tau, m), w_2(\tau, m)$ be in $F_{(v, \beta, \mu, k, \epsilon)}^d$. We take $\varpi > 0$ such that

$$||w_l(\tau, m)||_{(v, \beta, \mu, k, \epsilon)} \leq \varpi,$$

for $l = 1, 2$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. One can write

$$\begin{aligned} & Q_1(i(m - m_1)) w_1((s - x)^{1/k}, m - m_1) Q_2(im_1) w_1(x^{1/k}, m_1) \\ & - Q_1(i(m - m_1)) w_2((s - x)^{1/k}, m - m_1) Q_2(im_1) w_2(x^{1/k}, m_1) \\ & = Q_1(i(m - m_1)) \left(w_1((s - x)^{1/k}, m - m_1) - w_2((s - x)^{1/k}, m - m_1) \right) Q_2(im_1) w_1(x^{1/k}, m_1) \\ & + Q_1(i(m - m_1)) w_2((s - x)^{1/k}, m - m_1) Q_2(im_1) \left(w_1(x^{1/k}, m_1) - w_2(x^{1/k}, m_1) \right) \end{aligned} \quad (100)$$

and using Lemma 1 and Proposition 3 with the lower bound estimates (86) we get that

$$\begin{aligned} & || \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \\ & \times \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} (Q_1(i(m - m_1)) w_1((s - x)^{1/k}, m - m_1) Q_2(im_1) w_1(x^{1/k}, m_1) \right. \\ & \left. - Q_1(i(m - m_1)) w_2((s - x)^{1/k}, m - m_1) \right) \end{aligned}$$

$$\begin{aligned}
& \times Q_2(im_1)w_2(x^{1/k}, m_1)) \frac{1}{(s-x)x} dx dm_1 \Big) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} \\
& \quad \times \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} (\|w_1(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} + \|w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)}) \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 2\varpi}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (101)
\end{aligned}$$

From the estimates (93), (94), (95), (96), (97), (98) and under the constraints (87), we deduce that

$$\begin{aligned}
& \| \frac{R_D(im)}{P_m(\tau)} \frac{A_{\delta_D, p}}{\Gamma(\delta_D - p)} \int_0^{\tau^k} (\tau^k - s)^{\delta_D - p - 1} (k^p s^p (w_1(s^{1/k}, m) - w_2(s^{1/k}, m))) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_D, p}| k^p C_{2.1} |\epsilon|}{\Gamma(\delta_D - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_D, p}| k^p C_{2.1} \epsilon_0}{\Gamma(\delta_D - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (102)
\end{aligned}$$

and

$$\begin{aligned}
& \| \frac{R_l(im)}{P_m(\tau)} \frac{\epsilon^{\Delta_l - d_l + \delta_l - 1}}{\Gamma(\frac{d_{l,k}}{k})} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} - 1} (k^{\delta_l} s^{\delta_l} (w_1(s^{1/k}, m) - w_2(s^{1/k}, m))) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{k^{\delta_l} C_{2.2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} |\epsilon|^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{k^{\delta_l} C_{2.2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (103)
\end{aligned}$$

and that

$$\begin{aligned}
& \| \frac{R_l(im)}{P_m(\tau)} \frac{A_{\delta_l, p} \epsilon^{\Delta_l - d_l + \delta_l - 1}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p)} \int_0^{\tau^k} (\tau^k - s)^{\frac{d_{l,k}}{k} + \delta_l - p - 1} \\
& \quad \times (k^p s^p (w_1(s^{1/k}, m) - w_2(s^{1/k}, m))) \frac{ds}{s} \|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_l, p}| k^p C_{2.2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D-1)k}}} |\epsilon|^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}}
\end{aligned}$$

$$\begin{aligned}
& \times \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{|A_{\delta_l, p}| k^p C_{2,2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \\
& \times \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)} \quad (104)
\end{aligned}$$

and that

$$\begin{aligned}
& \left\| \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \left(\frac{1}{(2\pi)^{1/2}} s \int_0^s \int_{-\infty}^{+\infty} \varphi_k((s-x)^{1/k}, m - m_1, \epsilon) \right. \right. \\
& \quad \left. \left. R_0(im_1)(w_1(x^{1/k}, m_1) - w_2(x^{1/k}, m_1)) \frac{1}{(s-x)x} dx dm_1 \right) \frac{ds}{s} \right\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \|\varphi_k(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)}}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \zeta_1 \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)}}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \quad (105)
\end{aligned}$$

together with

$$\begin{aligned}
& \left\| \frac{\epsilon^{-1}}{P_m(\tau) \Gamma(1 + \frac{1}{k})} \int_0^{\tau^k} (\tau^k - s)^{1/k} \frac{1}{(2\pi)^{1/2}} \left(\int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) R_0(im_1) \right. \right. \\
& \quad \left. \left. \times (w_1(s^{1/k}, m_1) - w_2(s^{1/k}, m_1)) dm_1 \right) \frac{ds}{s} \right\|_{(v, \beta, \mu, k, \epsilon)} \\
& \leq \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_4 \zeta_0 \|w_1(\tau, m) - w_2(\tau, m)\|_{(v, \beta, \mu, k, \epsilon)}}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \quad (106)
\end{aligned}$$

Now, we take ϖ and r_{Q, R_D} such that

$$\begin{aligned}
& \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 2\varpi}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \\
& + \sum_{1 \leq p \leq \delta_D - 1} \frac{|A_{\delta_D, p}| k^p C_{2,1} \epsilon_0}{\Gamma(\delta_D - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \\
& + \sum_{1 \leq l \leq D-1} \frac{k^{\delta_l} C_{2,2}}{\Gamma(\frac{d_{l,k}}{k}) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right|
\end{aligned}$$

$$\begin{aligned}
& + \sum_{1 \leq p \leq \delta_l - 1} \frac{|A_{\delta_l, p}| k^p C_{2,2}}{\Gamma(\frac{d_{l,k}}{k} + \delta_l - p) C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \epsilon_0^{\Delta_l - d_l + \delta_l + k(\delta_l - \delta_D) + d_{l,k}} \sup_{m \in \mathbb{R}} \left| \frac{R_l(im)}{R_D(im)} \right| \\
& + \frac{1}{\Gamma(1 + \frac{1}{k})(2\pi)^{1/2}} \frac{C_3 \zeta_1 + C_4 \zeta_0}{C_P(r_{Q, R_D})^{\frac{1}{(\delta_D - 1)k}}} \leq \frac{1}{2}
\end{aligned} \tag{107}$$

Bearing in mind the estimates (101), (102), (103), (104), (105), (106) with the constraint (107), one gets (91).

Finally, we choose ϖ and r_{Q, R_D} such that both (99) and (107) are satisfied. This yields our lemma. \square

We consider the ball $\bar{B}(0, \varpi) \subset F_{(v, \beta, \mu, k, \epsilon)}^d$ constructed in Lemma 2 which is a complete metric space for the norm $\|\cdot\|_{(v, \beta, \mu, k, \epsilon)}$. From the lemma above, we get that \mathcal{H}_ϵ is a contractive map from $\bar{B}(0, \varpi)$ into itself. Due to the classical contractive mapping theorem, we deduce that the map \mathcal{H}_ϵ has a unique fixed point denoted by $\omega_k(\tau, m, \epsilon)$ (i.e. $\mathcal{H}_\epsilon(\omega_k(\tau, m, \epsilon)) = \omega_k(\tau, m, \epsilon)$) in $\bar{B}(0, \varpi)$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. Moreover, the function $\omega_k(\tau, m, \epsilon)$ depends holomorphically on ϵ in $D(0, \epsilon_0) \setminus \{0\}$. By construction, $\omega_k(\tau, m, \epsilon)$ defines a solution of equation (80). This yields the proposition. \square

In the next proposition, we construct analytic solutions of equation (72).

Proposition 10. *Let the assumption (87) hold. We also choose the sectors S_d and S_{Q, R_D} in such a way that (84) and (85) hold. We take the radius r_{Q, R_D} as prescribed in Proposition 9. We also assume that the inequalities (88) hold for $\zeta_0, \zeta_1, \zeta_2$ constructed in Proposition 9. Notice that the inequalities for ζ_1, ζ_2 can be satisfied if ϵ_0 is small enough due to the estimates (79).*

Let $S_{d, \theta, h'|\epsilon|}$ be a bounded sector with aperture $\pi/k < \theta < \pi/k + 2\delta$ (where 2δ is the small aperture of the unbounded sector S_d), with direction d and radius $h'|\epsilon|$ for some $h' > 0$ independent of ϵ . We choose $0 < \beta' < \beta$.

Then, equation (72) with initial condition $U(0, m, \epsilon) \equiv 0$ has a solution $(T, m) \mapsto U(T, m, \epsilon)$ defined on $S_{d, \theta, h'|\epsilon|} \times \mathbb{R}$ for some real number $h' > 0$ for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. Let $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, then for each $T \in S_{d, \theta, h'|\epsilon|}$, the function $m \mapsto U(T, m, \epsilon)$ belongs to the space $E_{(\beta', \mu)}$ and for each $m \in \mathbb{R}$, the function $T \mapsto U(T, m, \epsilon)$ is bounded and holomorphic on $S_{d, \theta, h'|\epsilon|}$. Moreover, the function $U(T, m, \epsilon)$ can be written as a Laplace transform of order k in the direction d ,

$$U(T, m, \epsilon) = k \int_{L_\gamma} \omega_k^d(u, m, \epsilon) e^{-(\frac{u}{T})^k} \frac{du}{u} \tag{108}$$

along a halfline $L_\gamma = \mathbb{R}_+ e^{i\gamma} \in S_d \cup \{0\}$ (the direction γ may depend on T), where $\omega_k^d(\tau, m, \epsilon)$ defines a continuous function on $(\bar{D}(0, \rho) \cup S_d) \times \mathbb{R} \times D(0, \epsilon_0) \setminus \{0\}$ which is holomorphic with respect to (τ, ϵ) on $(\bar{D}(0, \rho) \cup S_d) \times D(0, \epsilon_0) \setminus \{0\}$ and satisfies the estimates: there exists a constant ϖ_d (independent of ϵ) such that

$$|\omega_k^d(\tau, m, \epsilon)| \leq \varpi_d (1 + |m|)^{-\mu} e^{-\beta|m|} \frac{|\frac{\tau}{\epsilon}|}{1 + |\frac{\tau}{\epsilon}|^{2k}} \exp(v|\frac{\tau}{\epsilon}|^k) \tag{109}$$

for all $\tau \in D(0, \rho) \cup S_d$, all $m \in \mathbb{R}$, all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$.

Proof. Taking into account the requirements stated above in Proposition 10, we get that all the assumptions of Proposition 9 are fulfilled. Therefore, the formal m_k -Borel transform $\omega_k(\tau, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) \tau^n / \Gamma(n/k)$ of the formal series $\hat{U}(T, m, \epsilon)$ constructed in Proposition 9 is convergent with respect to τ on $D(0, \rho)$ as series with coefficients in the Banach space $E_{(\beta, \mu)}$. Moreover, this function $\omega_k(\tau, m, \epsilon)$ can be extended as an analytic function with respect to τ on the sector S_d , denoted $\omega_k^d(\tau, m, \epsilon)$, that belongs to the Banach space $F_{(v, \beta, \mu, k, \epsilon)}^d$ and satisfies the bounds $\|\omega_k^d(\tau, m, \epsilon)\|_{(v, \beta, \mu, k, \epsilon)} \leq \varpi_d$ where ϖ_d is a constant independent of ϵ in $D(0, \epsilon_0) \setminus \{0\}$. This means that (109) must hold. As a result, we get that the formal series $\hat{U}(T, m, \epsilon) \in TE_{(\beta, \mu)}[[T]]$ is m_k -summable in the direction d (see Definition 3). By construction, its m_k -sum $U(T, m, \epsilon)$ in direction d defines a holomorphic function on the sector $S_{d, \theta, h'|\epsilon|}$ described above in Proposition 10 with values in $E_{(\beta, \mu)}$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. On the other hand, the series $C_0(T, m, \epsilon), F(T, m, \epsilon) \in TE_{(\beta, \mu)}[[T]]$ are convergent. Therefore, these series are m_k -summable in any direction d and their m_k -sums satisfy

$$\mathcal{L}_{m_k}^d(\varphi_k(\tau, m, \epsilon))(T) = C_0(T, m, \epsilon), \quad \mathcal{L}_{m_k}^d(\psi_k(\tau, m, \epsilon))(T) = F(T, m, \epsilon)$$

for all $T \in D(0, T_0/2)$. Finally, using the properties for the sum, product and derivative of m_k -sums described in (56), we deduce that the m_k -sum $U(T, m, \epsilon)$ in direction d satisfies equation (76) as a function of (T, m) on $S_{d, \theta, h'|\epsilon|} \times \mathbb{R}$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, since the formal series $\hat{U}(T, m, \epsilon)$ satisfies equation (76). As a result, the function $U(T, m, \epsilon)$ also satisfies equation (72) as a function of (T, m) on $S_{d, \theta, h'|\epsilon|} \times \mathbb{R}$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. \square

5. Analytic solutions of a nonlinear initial value Cauchy problem with complex parameter

Let $k \geq 1$ and $D \geq 2$ be integers. For $1 \leq l \leq D$, let $d_l, \delta_l, \Delta_l \geq 0$ be nonnegative integers. We assume that

$$1 = \delta_1, \quad \delta_l < \delta_{l+1}, \quad (110)$$

for all $1 \leq l \leq D-1$. We make also the assumption that

$$d_D = (\delta_D - 1)(k + 1), \quad d_l > (\delta_l - 1)(k + 1), \quad \Delta_D = d_D - \delta_D + 1 \quad (111)$$

for all $1 \leq l \leq D-1$. Let $Q(X), Q_1(X), Q_2(X), R_l(X) \in \mathbb{C}[X], 0 \leq l \leq D$, be polynomials such that

$$\begin{aligned} \deg(Q) &\geq \deg(R_D) \geq \deg(R_l), \quad \deg(R_D) \geq \deg(Q_1), \quad \deg(R_D) \geq \deg(Q_2), \\ Q(im) &\neq 0, \quad R_D(im) \neq 0 \end{aligned} \quad (112)$$

for all $m \in \mathbb{R}$, all $0 \leq l \leq D-1$.

We consider the following nonlinear initial value problem

$$\begin{aligned} Q(\partial_z)(\partial_t u(t, z, \epsilon)) &= (Q_1(\partial_z)u(t, z, \epsilon))(Q_2(\partial_z)u(t, z, \epsilon)) + \sum_{l=1}^D \epsilon^{\Delta_l} t^{d_l} \partial_t^{\delta_l} R_l(\partial_z)u(t, z, \epsilon) \\ &\quad + c_0(t, z, \epsilon)R_0(\partial_z)u(t, z, \epsilon) + f(t, z, \epsilon) \end{aligned} \quad (113)$$

for given initial data $u(0, z, \epsilon) \equiv 0$.

The coefficient $c_0(t, z, \epsilon)$ and the forcing term $f(t, z, \epsilon)$ are constructed as follows. We consider sequences of functions $m \mapsto C_{0,n}(m, \epsilon)$, for $n \geq 0$ and $m \mapsto F_n(m, \epsilon)$, for $n \geq 1$, that belong to the Banach space $E_{(\beta, \mu)}$ for some $\beta > 0$, $\mu > \max(\deg(Q_1) + 1, \deg(Q_2) + 1)$ and which depend holomorphically on $\epsilon \in D(0, \epsilon_0)$. We assume that there exist constants $K_0, T_0 > 0$ such that (71) hold for all $n \geq 1$, for all $\epsilon \in D(0, \epsilon_0)$. We deduce that the functions

$$\mathbf{C}_0(T, z, \epsilon) = \sum_{n \geq 0} \mathcal{F}^{-1}(m \mapsto C_{0,n}(m, \epsilon))(z)T^n, \quad \mathbf{F}(T, z, \epsilon) = \sum_{n \geq 1} \mathcal{F}^{-1}(m \mapsto F_n(m, \epsilon))(z)T^n$$

represent bounded holomorphic functions on $D(0, T_0/2) \times H_{\beta'} \times D(0, \epsilon_0)$ for any $0 < \beta' < \beta$ (where \mathcal{F}^{-1} denotes the inverse Fourier transform defined in Proposition 7). We define the coefficient $c_0(t, z, \epsilon)$ and the forcing term $f(t, z, \epsilon)$ as

$$c_0(t, z, \epsilon) = \mathbf{C}_0(\epsilon t, z, \epsilon), \quad f(t, z, \epsilon) = \mathbf{F}(\epsilon t, z, \epsilon). \quad (114)$$

The functions c_0 and f are holomorphic and bounded on $D(0, r) \times H_{\beta'} \times D(0, \epsilon_0)$ where $r\epsilon_0 < T_0/2$.

We make the additional assumption that there exists an unbounded sector

$$S_{Q, R_D} = \{z \in \mathbb{C} / |z| \geq r_{Q, R_D}, \quad |\arg(z) - d_{Q, R_D}| \leq \eta_{Q, R_D}\}$$

with direction $d_{Q, R_D} \in \mathbb{R}$, aperture $\eta_{Q, R_D} > 0$ for some radius $r_{Q, R_D} > 0$ such that

$$\frac{Q(im)}{R_D(im)} \in S_{Q, R_D} \quad (115)$$

for all $m \in \mathbb{R}$.

Definition 4. Let $\varsigma \geq 2$ be an integer. For all $0 \leq p \leq \varsigma - 1$, we consider open sectors \mathcal{E}_p centered at 0, with radius ϵ_0 and opening $\frac{\pi}{k} + \kappa_p$, with $\kappa_p > 0$ small enough such that $\mathcal{E}_p \cap \mathcal{E}_{p+1} \neq \emptyset$, for all $0 \leq p \leq \varsigma - 1$ (with the convention that $\mathcal{E}_\varsigma = \mathcal{E}_0$). Moreover, we assume that the intersection of any three different elements in $(\mathcal{E}_p)_{0 \leq p \leq \varsigma}$ is empty and that $\bigcup_{p=0}^{\varsigma-1} \mathcal{E}_p = \mathcal{U} \setminus \{0\}$, where \mathcal{U} is some neighborhood of 0 in \mathbb{C} . Such a set of sectors $\{\mathcal{E}_p\}_{0 \leq p \leq \varsigma-1}$ is called a good covering in \mathbb{C}^* .

Definition 5. Let $\{\mathcal{E}_p\}_{0 \leq p \leq \varsigma-1}$ be a good covering in \mathbb{C}^* . Let \mathcal{T} be an open bounded sector centered at 0 with radius $r_{\mathcal{T}}$ and consider a family of open sectors

$$S_{\mathfrak{d}_p, \theta, \epsilon_0 r_{\mathcal{T}}} = \{T \in \mathbb{C}^* / |T| < \epsilon_0 r_{\mathcal{T}}, \quad |\mathfrak{d}_p - \arg(T)| < \theta/2\}$$

with aperture $\theta > \pi/k$ and where $\mathfrak{d}_p \in \mathbb{R}$, for all $0 \leq p \leq \varsigma - 1$, are directions which satisfy the following constraints: Let $q_l(m)$ be the roots of the polynomials (82) defined by (83) and $S_{\mathfrak{d}_p}$, $0 \leq p \leq \varsigma - 1$ be unbounded sectors centered at 0 with directions \mathfrak{d}_p and with small aperture. We assume that

1) There exists a constant $M_1 > 0$ such that

$$|\tau - q_l(m)| \geq M_1(1 + |\tau|) \quad (116)$$

for all $0 \leq l \leq (\delta_D - 1)k - 1$, all $m \in \mathbb{R}$, all $\tau \in S_{\mathfrak{d}_p} \cup \bar{D}(0, \rho)$, for all $0 \leq p \leq \varsigma - 1$.

2) There exists a constant $M_2 > 0$ such that

$$|\tau - q_{l_0}(m)| \geq M_2 |q_{l_0}(m)| \quad (117)$$

for some $l_0 \in \{0, \dots, (\delta_D - 1)k - 1\}$, all $m \in \mathbb{R}$, all $\tau \in S_{\delta_p} \cup \bar{D}(0, \rho)$, for all $0 \leq p \leq \zeta - 1$.

3) For all $0 \leq p \leq \zeta - 1$, for all $t \in \mathcal{T}$, all $\epsilon \in \mathcal{E}_p$, we have that $\epsilon t \in S_{\delta_p, \theta, \epsilon_0 r_{\mathcal{T}}}$.

We say that the family $\{(S_{\delta_p, \theta, \epsilon_0 r_{\mathcal{T}}})_{0 \leq p \leq \zeta - 1}, \mathcal{T}\}$ is associated to the good covering $\{\mathcal{E}_p\}_{0 \leq p \leq \zeta - 1}$.

In the next first main result, we construct a family of actual holomorphic solutions to equation (113) for given initial data at $t = 0$ being identically equal to zero, defined on the sectors \mathcal{E}_p with respect to the complex parameter ϵ . We can also control the difference between any two neighboring solutions on the intersection of sectors $\mathcal{E}_p \cap \mathcal{E}_{p+1}$ and show that it is exponentially flat of order at most k .

Theorem 1. *We consider equation (113) and we assume that the constraints (110), (111), (112) and (115) hold. We also make the additional assumption that*

$$\delta_D \geq \delta_l + \frac{2}{k}, \quad \Delta_l + k(1 - \delta_D) + 1 \geq 0, \quad (118)$$

hold for all $1 \leq l \leq D - 1$. Let the coefficient $c_0(t, z, \epsilon)$ and forcing term $f(t, z, \epsilon)$ be constructed as in (114). Let a good covering $\{\mathcal{E}_p\}_{0 \leq p \leq \zeta - 1}$ in \mathbb{C}^* be given, for which a family of sectors $\{(S_{\delta_p, \theta, \epsilon_0 r_{\mathcal{T}}})_{0 \leq p \leq \zeta - 1}, \mathcal{T}\}$ associated to this good covering can be considered.

Then, there exists a radius $r_{Q, R_D} > 0$ large enough, $\epsilon_0 > 0$ small enough and a constant $\zeta_0 > 0$ small enough such that if

$$\|C_{0,0}(m, \epsilon)\|_{(\beta, \mu)} < \zeta_0$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$, then for every $0 \leq p \leq \zeta - 1$, one can construct a solution $u_p(t, z, \epsilon)$ of equation (113) with $u_p(0, z, \epsilon) \equiv 0$ which defines a bounded holomorphic function on the domain $(\mathcal{T} \cap D(0, h')) \times H_{\beta'} \times \mathcal{E}_i$ for any given $0 < \beta' < \beta$ and for some $h' > 0$. Moreover, there exist constants $0 < h'' \leq h'$, $K_p, M_p > 0$ (independent of ϵ) such that

$$\sup_{t \in \mathcal{T} \cap D(0, h''), z \in H_{\beta'}} |u_{p+1}(t, z, \epsilon) - u_p(t, z, \epsilon)| \leq K_p e^{-\frac{M_p}{|\epsilon|^k}} \quad (119)$$

for all $\epsilon \in \mathcal{E}_{p+1} \cap \mathcal{E}_p$, for all $0 \leq p \leq \zeta - 1$ (where by convention $u_{\zeta} = u_0$).

Proof. Using Proposition 10, one can choose $r_{Q, R_D} > 0$ large enough, $\epsilon_0 > 0$ small enough and $\zeta_0 > 0$ small enough such that

$$\|C_{0,0}(m, \epsilon)\|_{(\beta, \mu)} \leq \zeta_0$$

for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$ such that for each direction \mathfrak{d}_p with $0 \leq p \leq \zeta - 1$, one can construct a function $U^{\mathfrak{d}_p}(T, m, \epsilon)$ which satisfies $U^{\mathfrak{d}_p}(0, m, \epsilon) \equiv 0$ and solves the equation

$$\begin{aligned}
& Q(im)(\partial_T U(T, m, \epsilon)) \\
&= \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} Q_1(i(m - m_1)) U(T, m - m_1, \epsilon) \\
&\quad \times Q_2(im_1) U(T, m_1, \epsilon) dm_1 \\
&\quad + \sum_{l=1}^D R_l(im) \epsilon^{\Delta_l - d_l + \delta_l - 1} T^{d_l} \partial_T^{\delta_l} U(T, m, \epsilon) \\
&\quad + \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_0(T, m - m_1, \epsilon) U(T, m_1, \epsilon) dm_1 \\
&\quad + \epsilon^{-1} \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} C_{0,0}(m - m_1, \epsilon) U(T, m_1, \epsilon) dm_1 + \epsilon^{-1} F(T, m, \epsilon) \quad (120)
\end{aligned}$$

where

$$C_0(T, m, \epsilon) = \sum_{n \geq 1} C_{0,n}(m, \epsilon) T^n, \quad F(T, m, \epsilon) = \sum_{n \geq 1} F_n(m, \epsilon) T^n$$

are convergent series in $D(0, T_0/2)$ with values in $E_{(\beta, \mu)}$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. The function $(T, m) \mapsto U^{\mathfrak{d}_p}(T, m, \epsilon)$ is well defined on the domain $S_{\mathfrak{d}_p, \theta, h'|\epsilon|} \times \mathbb{R}$ where $h' > 0$ is some real number, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. Moreover, $U^{\mathfrak{d}_p}(T, m, \epsilon)$ can be written as a Laplace transform of order k in the direction \mathfrak{d}_p ,

$$U^{\mathfrak{d}_p}(T, m, \epsilon) = k \int_{L_{\gamma_p}} \omega_k^{\mathfrak{d}_p}(u, m, \epsilon) e^{-(\frac{u}{T})^k} \frac{du}{u} \quad (121)$$

along a halfline $L_{\gamma_p} = \mathbb{R}_+ e^{i\gamma_p} \in S_{\mathfrak{d}_p} \cup \{0\}$ (the direction γ_p may depend on T), where $\omega_k^{\mathfrak{d}_p}(\tau, m, \epsilon)$ defines a continuous function on $(\bar{D}(0, \rho) \cup S_{d_p}) \times \mathbb{R} \times D(0, \epsilon_0) \setminus \{0\}$ which is holomorphic with respect to (τ, ϵ) on $(\bar{D}(0, \rho) \cup S_{\mathfrak{d}_p}) \times D(0, \epsilon_0) \setminus \{0\}$ for any $m \in \mathbb{R}$ and satisfies the estimates: there exists a constant $\varpi_{\mathfrak{d}_p}$ (independent of ϵ) such that

$$|\omega_k^{\mathfrak{d}_p}(\tau, m, \epsilon)| \leq \varpi_{\mathfrak{d}_p} (1 + |m|)^{-\mu} e^{-\beta|m|} \frac{|\frac{\tau}{\epsilon}|}{1 + |\frac{\tau}{\epsilon}|^{2k}} \exp(v|\frac{\tau}{\epsilon}|^k) \quad (122)$$

for all $\tau \in D(0, \rho) \cup S_{\mathfrak{d}_p}$, all $m \in \mathbb{R}$, all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. It is worth noticing that all the functions $\tau \mapsto \omega_k^{\mathfrak{d}_p}(\tau, m, \epsilon)$ are analytic continuation on the sectors $S_{\mathfrak{d}_p}$ of a common function denoted by

$$\omega_k(\tau, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) \frac{\tau^n}{\Gamma(\frac{n}{k})}$$

which is a convergent series on $D(0, \rho)$ with coefficients in $E_{(\beta, \mu)}$ and where $U_n(m, \epsilon) \in E_{(\beta, \mu)}$ are the coefficients of the formal series $\hat{U}(T, m, \epsilon) = \sum_{n \geq 1} U_n(m, \epsilon) T^n$ solution of equation (120), for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$. Using the estimates (122), we get that the function

$$(T, z) \mapsto \mathbf{U}^{\mathfrak{d}_p}(T, z, \epsilon) = \mathcal{F}^{-1}(m \mapsto U^{\mathfrak{d}_p}(T, m, \epsilon))(z)$$

defines a bounded holomorphic function on $S_{\mathfrak{d}_p, \theta, h'|\epsilon|} \times H_{\beta'}$, for all $\epsilon \in D(0, \epsilon_0) \setminus \{0\}$ and any $0 < \beta' < \beta$. For all $0 \leq p \leq \zeta - 1$, we define

$$u_p(t, z, \epsilon) = \mathbf{U}^{\mathfrak{d}_p}(\epsilon t, z, \epsilon) = \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\gamma_p}} \omega_k^{\mathfrak{d}_p}(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm. \quad (123)$$

By construction (see 3) in Definition 5), the function $u_p(t, z, \epsilon)$ defines a bounded holomorphic function on the domain $(\mathcal{T} \cap D(0, h')) \times H_{\beta'} \times \mathcal{E}_p$. Moreover, we have $u_p(0, z, \epsilon) \equiv 0$ and using the properties of the Fourier inverse transform from Proposition 7, we deduce that $u_p(t, z, \epsilon)$ solves the main equation (113) on $(\mathcal{T} \cap D(0, h')) \times H_{\beta'} \times \mathcal{E}_p$.

Now, we proceed to the proof of the estimates (119). Let $p \in \{0, \dots, \zeta - 1\}$. Using the fact that the function $u \mapsto \omega_k(u, m, \epsilon) \exp(-(\frac{u}{\epsilon t})^k)/u$ is holomorphic on $D(0, \rho)$ for all $(m, \epsilon) \in \mathbb{R} \times (D(0, \epsilon_0) \setminus \{0\})$, its integral along the union of a segment starting from 0 to $(\rho/2)e^{i\gamma_{p+1}}$, an arc of circle with radius $\rho/2$ which connects $(\rho/2)e^{i\gamma_{p+1}}$ and $(\rho/2)e^{i\gamma_p}$ and a segment starting from $(\rho/2)e^{i\gamma_p}$ to 0, is equal to zero. Therefore, we can write the difference $u_{p+1} - u_p$ as a sum of three integrals,

$$\begin{aligned} u_{p+1}(t, z, \epsilon) - u_p(t, z, \epsilon) &= \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\rho/2, \gamma_{p+1}}} \omega_k^{\mathfrak{d}_{p+1}}(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm \\ &\quad - \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\rho/2, \gamma_p}} \omega_k^{\mathfrak{d}_p}(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm \\ &\quad + \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{C_{\rho/2, \gamma_p, \gamma_{p+1}}} \omega_k(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm \end{aligned} \quad (124)$$

where $L_{\rho/2, \gamma_{p+1}} = [\rho/2, +\infty)e^{i\gamma_{p+1}}$, $L_{\rho/2, \gamma_p} = [\rho/2, +\infty)e^{i\gamma_p}$ and $C_{\rho/2, \gamma_p, \gamma_{p+1}}$ is an arc of circle with radius connecting $(\rho/2)e^{i\gamma_p}$ and $(\rho/2)e^{i\gamma_{p+1}}$ with a well chosen orientation.

We give estimates for the quantity

$$I_1 = \left| \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\rho/2, \gamma_{p+1}}} \omega_k^{\mathfrak{d}_{p+1}}(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm \right|.$$

By construction, the direction γ_{p+1} (which depends on ϵt) is chosen in such a way that $\cos(k(\gamma_{p+1} - \arg(\epsilon t))) \geq \delta_1$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$, all $t \in \mathcal{T} \cap D(0, h')$, for some fixed $\delta_1 > 0$. From the estimates (122), we get that

$$\begin{aligned}
 I_1 &\leq \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{\rho/2}^{+\infty} \varpi_{\mathfrak{D}_{p+1}} (1 + |m|)^{-\mu} e^{-\beta|m|} \frac{\frac{r}{|\epsilon|}}{1 + (\frac{r}{|\epsilon|})^{2k}} \\
 &\quad \times \exp(v(\frac{r}{|\epsilon|})^k) \exp(-\frac{\cos(k(\gamma_{p+1} - \arg(\epsilon t)))}{|\epsilon t|^k} r^k) e^{-m \operatorname{Im}(z)} \frac{dr}{r} dm \\
 &\leq \frac{k \varpi_{\mathfrak{D}_{p+1}}}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} e^{-(\beta - \beta')|m|} dm \int_{\rho/2}^{+\infty} \frac{1}{|\epsilon|} \exp(-(\frac{\delta_1}{|t|^k} - v)(\frac{r}{|\epsilon|})^k) dr \\
 &\leq \frac{2k \varpi_{\mathfrak{D}_{p+1}}}{(2\pi)^{1/2}} \int_0^{+\infty} e^{-(\beta - \beta')m} dm \int_{\rho/2}^{+\infty} \frac{|\epsilon|^{k-1}}{(\frac{\delta_1}{|t|^k} - v)k(\frac{\rho}{2})^{k-1}} \\
 &\quad \times \frac{(\frac{\delta_1}{|t|^k} - v)k r^{k-1}}{|\epsilon|^k} \exp(-(\frac{\delta_1}{|t|^k} - v)(\frac{r}{|\epsilon|})^k) dr \\
 &\leq \frac{2k \varpi_{\mathfrak{D}_{p+1}}}{(2\pi)^{1/2}} \frac{|\epsilon|^{k-1}}{(\beta - \beta')(\frac{\delta_1}{|t|^k} - v)k(\frac{\rho}{2})^{k-1}} \exp(-(\frac{\delta_1}{|t|^k} - v)\frac{(\rho/2)^k}{|\epsilon|^k}) \\
 &\leq \frac{2k \varpi_{\mathfrak{D}_{p+1}}}{(2\pi)^{1/2}} \frac{|\epsilon|^{k-1}}{(\beta - \beta')\delta_2 k(\frac{\rho}{2})^{k-1}} \exp(-\delta_2 \frac{(\rho/2)^k}{|\epsilon|^k}) \tag{125}
 \end{aligned}$$

for all $t \in \mathcal{T} \cap D(0, h')$ and $|\operatorname{Im}(z)| \leq \beta'$ with $|t| < (\frac{\delta_1}{\delta_2 + v})^{1/k}$, for some $\delta_2 > 0$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$.

In the same way, we also give estimates for the integral

$$I_2 = \left| \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{L_{\rho/2, \gamma_p}} \omega_k^{\mathfrak{D}_p}(u, m, \epsilon) e^{-(\frac{u}{\epsilon t})^k} e^{izm} \frac{du}{u} dm \right|.$$

Namely, the direction γ_p (which depends on ϵt) is chosen in such a way that $\cos(k(\gamma_p - \arg(\epsilon t))) \geq \delta_1$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$, all $t \in \mathcal{T} \cap D(0, h')$, for some fixed $\delta_1 > 0$. Again from the estimates (122) and following the same steps as in (125), we get that

$$I_2 \leq \frac{2k \varpi_{\mathfrak{D}_p}}{(2\pi)^{1/2}} \frac{|\epsilon|^{k-1}}{(\beta - \beta')\delta_2 k(\frac{\rho}{2})^{k-1}} \exp(-\delta_2 \frac{(\rho/2)^k}{|\epsilon|^k}) \tag{126}$$

for all $t \in \mathcal{T} \cap D(0, h')$ and $|\operatorname{Im}(z)| \leq \beta'$ with $|t| < (\frac{\delta_1}{\delta_2 + v})^{1/k}$, for some $\delta_2 > 0$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$.

Finally, we give upper bound estimates for the integral

$$I_3 = \left| \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \int_{C_{\rho/2, \gamma_p, \gamma_{p+1}}} \omega_k(u, m, \epsilon) e^{-\left(\frac{u}{\epsilon t}\right)^k} e^{izm} \frac{du}{u} dm \right|.$$

By construction, the arc of circle $C_{\rho/2, \gamma_p, \gamma_{p+1}}$ is chosen in such a way that $\cos(k(\theta - \arg(\epsilon t))) \geq \delta_1$, for all $\theta \in [\gamma_p, \gamma_{p+1}]$ (if $\gamma_p < \gamma_{p+1}$), $\theta \in [\gamma_{p+1}, \gamma_p]$ (if $\gamma_{p+1} < \gamma_p$), for all $t \in \mathcal{T}$, all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$, for some fixed $\delta_1 > 0$. Bearing in mind (122) and (22), we get that

$$\begin{aligned} I_3 &\leq \frac{k}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} \left| \int_{\gamma_p}^{\gamma_{p+1}} \max_{0 \leq p \leq \varsigma-1} \varpi_{\mathfrak{D}_p} (1 + |m|)^{-\mu} e^{-\beta|m|} \frac{\frac{\rho/2}{|\epsilon|}}{1 + (\frac{\rho/2}{|\epsilon|})^{2k}} \right. \\ &\quad \times \exp(v(\frac{\rho/2}{|\epsilon|})^k) \exp(-\frac{\cos(k(\theta - \arg(\epsilon t)))}{|\epsilon t|^k} (\frac{\rho}{2})^k) e^{-m \operatorname{Im}(z)} d\theta \Big| dm \\ &\leq \frac{k(\max_{0 \leq p \leq \varsigma-1} \varpi_{\mathfrak{D}_p})}{(2\pi)^{1/2}} \int_{-\infty}^{+\infty} e^{-(\beta-\beta')|m|} dm \times |\gamma_p - \gamma_{p+1}| \frac{\rho/2}{|\epsilon|} \exp(-\frac{(\frac{\delta_1}{|t|^k} - \nu)}{2} (\frac{\rho/2}{|\epsilon|})^k) \\ &\quad \times \exp(-\frac{(\frac{\delta_1}{|t|^k} - \nu)}{2} (\frac{\rho/2}{|\epsilon|})^k) \\ &\leq \frac{2k(\max_{0 \leq p \leq \varsigma-1} \varpi_{\mathfrak{D}_p})|\gamma_p - \gamma_{p+1}|}{(2\pi)^{1/2}(\beta - \beta')} \sup_{x \geq 0} x^{1/k} e^{-(\frac{\delta_1}{|t|^k} - \nu)x} \times \exp(-\frac{(\frac{\delta_1}{|t|^k} - \nu)}{2} (\frac{\rho/2}{|\epsilon|})^k) \\ &\leq \frac{2k(\max_{0 \leq p \leq \varsigma-1} \varpi_{\mathfrak{D}_p})|\gamma_p - \gamma_{p+1}|}{(2\pi)^{1/2}(\beta - \beta')} (\frac{1/k}{\delta_2})^{1/k} e^{-1/k} \exp(-\frac{\delta_2}{2} (\frac{\rho/2}{|\epsilon|})^k) \end{aligned} \quad (127)$$

for all $t \in \mathcal{T} \cap D(0, h')$ and $|\operatorname{Im}(z)| \leq \beta'$ with $|t| < (\frac{\delta_1}{\delta_2 + \nu})^{1/k}$, for some $\delta_2 > 0$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$.

Finally, gathering the three above inequalities (125), (126) and (127), we deduce from the decomposition (124) that

$$\begin{aligned} &|u_{p+1}(t, z, \epsilon) - u_p(t, z, \epsilon)| \\ &\leq \frac{2k(\varpi_{\mathfrak{D}_p} + \varpi_{\mathfrak{D}_{p+1}})}{(2\pi)^{1/2}} \frac{|\epsilon|^{k-1}}{(\beta - \beta')\delta_2 k (\frac{\rho}{2})^{k-1}} \exp(-\delta_2 (\frac{\rho/2}{|\epsilon|})^k) \\ &\quad + \frac{2k(\max_{0 \leq p \leq \varsigma-1} \varpi_{\mathfrak{D}_p})|\gamma_p - \gamma_{p+1}|}{(2\pi)^{1/2}(\beta - \beta')} (\frac{1/k}{\delta_2})^{1/k} e^{-1/k} \exp(-\frac{\delta_2}{2} (\frac{\rho/2}{|\epsilon|})^k) \end{aligned}$$

for all $t \in \mathcal{T} \cap D(0, h')$ and $|\operatorname{Im}(z)| \leq \beta'$ with $|t| < (\frac{\delta_1}{\delta_2 + \nu})^{1/k}$, for some $\delta_2 > 0$, for all $\epsilon \in \mathcal{E}_p \cap \mathcal{E}_{p+1}$. Therefore, the inequality (119) holds. \square

6. Existence of k -summable formal series in the complex parameter of the initial value problem

6.1. k -Summable formal series and the Ramis–Sibuya theorem

We recall the definition of k -Borel summability of formal series with coefficients in a Banach space, see [1].

Definition 6. Let $k \geq 1$ be an integer. A formal series

$$\hat{X}(\epsilon) = \sum_{j=0}^{\infty} \frac{a_j}{j!} \epsilon^j \in \mathbb{F}[[\epsilon]]$$

with coefficients in a Banach space $(\mathbb{F}, ||\cdot||_{\mathbb{F}})$ is said to be k -summable with respect to ϵ in the direction $d \in \mathbb{R}$ if

i) there exists $\rho \in \mathbb{R}_+$ such that the following formal series, called formal Borel transform of \hat{X} of order k

$$\mathcal{B}_k(\hat{X})(\tau) = \sum_{j=0}^{\infty} \frac{a_j \tau^j}{j! \Gamma(1 + \frac{j}{k})} \in \mathbb{F}[[\tau]],$$

is absolutely convergent for $|\tau| < \rho$,

ii) there exists $\delta > 0$ such that the series $\mathcal{B}_k(\hat{X})(\tau)$ can be analytically continued with respect to τ in a sector $S_{d,\delta} = \{\tau \in \mathbb{C}^* : |d - \arg(\tau)| < \delta\}$. Moreover, there exist $C > 0$, and $K > 0$ such that

$$||\mathcal{B}_k(\hat{X})(\tau)||_{\mathbb{F}} \leq C e^{K|\tau|^k}$$

for all $\tau \in S_{d,\delta}$.

If this is so, the vector valued Laplace transform of order k of $\mathcal{B}_k(\hat{X})(\tau)$ in the direction d is defined by

$$\mathcal{L}_k^d(\mathcal{B}_k(\hat{X}))(\epsilon) = \epsilon^{-k} \int_{L_\gamma} \mathcal{B}_k(\hat{X})(u) e^{-(u/\epsilon)^k} k u^{k-1} du,$$

along a half-line $L_\gamma = \mathbb{R}_+ e^{i\gamma} \subset S_{d,\delta} \cup \{0\}$, where γ depends on ϵ and is chosen in such a way that $\cos(k(\gamma - \arg(\epsilon))) \geq \delta_1 > 0$, for some fixed δ_1 , for all ϵ in a sector

$$S_{d,\theta,R^{1/k}} = \{\epsilon \in \mathbb{C}^* : |\epsilon| < R^{1/k}, \quad |d - \arg(\epsilon)| < \theta/2\},$$

where $\frac{\pi}{k} < \theta < \frac{\pi}{k} + 2\delta$ and $0 < R < \delta_1/K$. The function $\mathcal{L}_k^d(\mathcal{B}_k(\hat{X}))(\epsilon)$ is called the k -sum of the formal series $\hat{X}(t)$ in the direction d . It is bounded and holomorphic on the sector $S_{d,\theta,R^{1/k}}$ and has the formal series $\hat{X}(\epsilon)$ as the Gevrey asymptotic expansion of order $1/k$ with respect to ϵ on $S_{d,\theta,R^{1/k}}$. This means that for all $\frac{\pi}{k} < \theta_1 < \theta$, there exist $C, M > 0$ such that

$$\|\mathcal{L}_k^d(\mathcal{B}_k(\hat{X}))(\epsilon) - \sum_{p=0}^{n-1} \frac{a_p}{p!} \epsilon^p\|_{\mathbb{F}} \leq CM^n \Gamma(1 + \frac{n}{k}) |\epsilon|^n$$

for all $n \geq 1$, all $\epsilon \in S_{d, \theta_1, R^{1/k}}$.

Now, we state a cohomological criterion for k -summability of formal series with coefficients in Banach spaces (see [2], p. 121 or [14], Lemma XI-2-6) which is known as the Ramis–Sibuya theorem in the literature. This result is a crucial tool in the proof of our main result (Theorem 2).

Theorem (RS). *Let $(\mathbb{F}, \|\cdot\|_{\mathbb{F}})$ be a Banach space over \mathbb{C} and $\{\mathcal{E}_p\}_{0 \leq p \leq \varsigma-1}$ be a good covering in \mathbb{C}^* . For all $0 \leq p \leq \varsigma-1$, let G_p be a holomorphic function from \mathcal{E}_p into the Banach space $(\mathbb{F}, \|\cdot\|_{\mathbb{F}})$ and let the cocycle $\Theta_p(\epsilon) = G_{p+1}(\epsilon) - G_p(\epsilon)$ be a holomorphic function from the sector $Z_p = \mathcal{E}_{p+1} \cap \mathcal{E}_p$ into \mathbb{F} (with the convention that $\mathcal{E}_{\varsigma} = \mathcal{E}_0$ and $G_{\varsigma} = G_0$). We make the following assumptions.*

- 1) *The functions $G_p(\epsilon)$ are bounded as $\epsilon \in \mathcal{E}_p$ tends to the origin in \mathbb{C} , for all $0 \leq p \leq \varsigma-1$.*
- 2) *The functions $\Theta_p(\epsilon)$ are exponentially flat of order $1/k$ on Z_p , for all $0 \leq p \leq \varsigma-1$. This means that there exist constants $C_p, A_p > 0$ such that*

$$\|\Theta_p(\epsilon)\|_{\mathbb{F}} \leq C_p e^{-A_p/|\epsilon|^k}$$

for all $\epsilon \in Z_p$, all $0 \leq p \leq \varsigma-1$.

Then, for all $0 \leq p \leq \nu-1$, the functions $G_p(\epsilon)$ are the k -sums on \mathcal{E}_p of a common k -summable formal series $\hat{G}(\epsilon) \in \mathbb{F}[[\epsilon]]$.

6.2. Construction of k -summable formal series in the complex parameter of the initial value problem

In this subsection, we establish the second main result of our work, namely the existence of a formal power series in the parameter ϵ whose coefficients are bounded holomorphic functions on the product of a sector with small radius centered at 0 and a strip in \mathbb{C}^2 , that is a solution of equation (113) and which is the common Gevrey asymptotic expansion of order $1/k$ of the actual solutions $u_p(t, z, \epsilon)$ of (113) constructed in Theorem 1.

The second main result of this work can be stated as follows.

Theorem 2. *Let us assume that the hypotheses of Theorem 1 hold. Then, there exists a formal power series*

$$\hat{u}(t, z, \epsilon) = \sum_{m \geq 0} h_m(t, z) \epsilon^m / m!$$

solution of equation (113), whose coefficients $h_m(t, z)$ belong to the Banach space \mathbb{F} of bounded holomorphic functions on $(\mathcal{T} \cap D(0, h'')) \times H_{\beta'}$ equipped with supremum norm, where $h'' > 0$ is constructed in Theorem 1, and such that the functions $u_p(t, z, \epsilon)$ defined in Theorem 1, seen as holomorphic functions from \mathcal{E}_p into \mathbb{F} , are its k -sums on the sectors \mathcal{E}_p , for all $0 \leq p \leq \varsigma-1$. In other words, for all $0 \leq p \leq \varsigma-1$, there exist two constants $C_p, M_p > 0$ such that

$$\sup_{t \in \mathcal{T} \cap D(0, h''), z \in H_{\beta'}} |u_p(t, z, \epsilon) - \sum_{m=0}^{n-1} h_m(t, z) \frac{\epsilon^m}{m!}| \leq C_p M_p^n \Gamma(1 + \frac{n}{k}) |\epsilon|^n \quad (128)$$

for all $n \geq 1$, all $\epsilon \in \mathcal{E}_p$.

Proof. We consider the family of functions $u_p(t, z, \epsilon)$, $0 \leq p \leq \varsigma - 1$ constructed in [Theorem 1](#). For all $0 \leq p \leq \varsigma - 1$, we define $G_p(\epsilon) := (t, z) \mapsto u_p(t, z, \epsilon)$, which is by construction a holomorphic and bounded function from \mathcal{E}_p into the Banach space \mathbb{F} of bounded holomorphic functions on $(\mathcal{T} \cap D(0, h'')) \times H_{\beta'}$ equipped with the supremum norm, where \mathcal{T} is introduced in [Definition 5](#), $h'' > 0$ is set in [Theorem 1](#) and $\beta' > 0$ is the width of the strip $H_{\beta'}$ on which the coefficients c_0 and f are defined with respect to z (see (114)). Bearing in mind the estimates (119), we see that the cocycle $\Theta_p(\epsilon) = G_{p+1}(\epsilon) - G_p(\epsilon)$ is exponentially flat of order k on $Z_p = \mathcal{E}_p \cap \mathcal{E}_{p+1}$, for any $0 \leq p \leq \varsigma - 1$.

From Theorem (RS) stated above, there exists a formal power series $\hat{G}(\epsilon) \in \mathbb{F}[[\epsilon]]$ such that the functions $G_p(\epsilon)$ are the k -sums on \mathcal{E}_p of $\hat{G}(\epsilon)$ as \mathbb{F} -valued functions, for all $0 \leq p \leq \varsigma - 1$. We set

$$\hat{G}(\epsilon) = \sum_{m \geq 0} h_m(t, z) \epsilon^m / m! =: \hat{u}(t, z, \epsilon).$$

It remains to show that the formal series $\hat{u}(t, z, \epsilon)$ satisfies the main equation (113). Since the functions $G_p(\epsilon)$ are the k -sums of $\hat{G}(\epsilon)$, we have in particular that

$$\lim_{\epsilon \rightarrow 0, \epsilon \in \mathcal{E}_p} \sup_{t \in \mathcal{T} \cap D(0, h''), z \in H_{\beta'}} |\partial_\epsilon^m u_p(t, z, \epsilon) - h_m(t, z)| = 0 \quad (129)$$

for all $0 \leq p \leq \varsigma - 1$, all $m \geq 0$. Now, we choose some $p \in \{0, \dots, \varsigma - 1\}$. By construction, the function $u_p(t, z, \epsilon)$ is a solution of (113). We take the derivative of order $m \geq 0$ with respect to ϵ on the left and right handside of equation (113). From the Leibniz rule, we deduce that $\partial_\epsilon^m u_p(t, z, \epsilon)$ verifies the following equation

$$\begin{aligned} Q(\partial_z)(\partial_t \partial_\epsilon^m u_p(t, z, \epsilon)) &= \sum_{m_1+m_2=m} \frac{m!}{m_1!m_2!} (Q_1(\partial_z) \partial_\epsilon^{m_1} u_p(t, z, \epsilon)) (Q_2(\partial_z) \partial_\epsilon^{m_2} u_p(t, z, \epsilon)) \\ &+ \sum_{l=1}^D \left(\sum_{m_1+m_2=m} \frac{m!}{m_1!m_2!} \partial_\epsilon^{m_1} (\epsilon^{\Delta_l}) t^{d_l} \partial_t^{\delta_l} R_l(\partial_z) \partial_\epsilon^{m_2} u_p(t, z, \epsilon) \right) \\ &+ \sum_{m_1+m_2=m} \frac{m!}{m_1!m_2!} \partial_\epsilon^{m_1} c_0(t, z, \epsilon) R_0(\partial_z) \partial_\epsilon^{m_2} u_p(t, z, \epsilon) + \partial_\epsilon^m f(t, z, \epsilon) \end{aligned} \quad (130)$$

for all $m \geq 0$, all $(t, z, \epsilon) \in (\mathcal{T} \cap D(0, h'')) \times H_{\beta'} \times \mathcal{E}_p$. If we let ϵ tend to zero in (130) and if we use (129), we get the recursion

$$\begin{aligned}
Q(\partial_z)(\partial_t h_m(t, z)) &= \sum_{m_1+m_2=m} \frac{m!}{m_1!m_2!} (Q_1(\partial_z)h_{m_1}(t, z))(Q_2(\partial_z)h_{m_2}(t, z)) \\
&+ \sum_{l=1}^D \frac{m!}{(m-\Delta_l)!} t^{d_l} \partial_t^{\delta_l} R_l(\partial_z)h_{m-\Delta_l}(t, z) \\
&+ \sum_{m_1+m_2=m} \frac{m!}{m_1!m_2!} (\partial_\epsilon^{m_1} c_0)(t, z, 0) R_0(\partial_z)h_{m_2}(t, z) + (\partial_\epsilon^m f)(t, z, 0)
\end{aligned} \tag{131}$$

for all $m \geq \max_{1 \leq l \leq D} \Delta_l$, all $(t, z) \in (\mathcal{T} \cap D(0, h'')) \times H_{\beta'}$. Since the functions $c_0(t, z, \epsilon)$ and $f(t, z, \epsilon)$ are analytic with respect to ϵ at 0, we know that

$$c_0(t, z, \epsilon) = \sum_{m \geq 0} \frac{(\partial_\epsilon^m c_0)(t, z, 0)}{m!} \epsilon^m, \quad f(t, z, \epsilon) = \sum_{m \geq 0} \frac{(\partial_\epsilon^m f)(t, z, 0)}{m!} \epsilon^m \tag{132}$$

for all $\epsilon \in D(0, \epsilon_0)$, all $z \in H_{\beta'}$. On other hand, one can check by direct inspection from the recursion (131) and the expansions (132) that the formal series $\hat{u}(t, z, \epsilon) = \sum_{m \geq 0} h_m(t, z) \epsilon^m / m!$ solves equation (113). \square

References

- [1] W. Balser, From Divergent Power Series to Analytic Functions: Theory and Application of Multisummable Power Series, Lecture Notes in Mathematics, vol. 1582, Springer-Verlag, Berlin, 1994, x+108 pp.
- [2] W. Balser, Formal Power Series and Linear Systems of Meromorphic Ordinary Differential Equations, Universitext, Springer-Verlag, New York, 2000, xviii+299 pp.
- [3] W. Balser, Summability of power series that are formal solutions of partial differential equations with constant coefficients, J. Math. Sci. (N. Y.) 124 (4) (2004) 5085–5097.
- [4] W. Balser, Multisummability of formal power series solutions of partial differential equations with constant coefficients, J. Differential Equations 201 (1) (2004) 63–74.
- [5] W. Balser, M. Loday-Richaud, Summability of solutions of the heat equation with inhomogeneous thermal conductivity in two variables, Adv. Dyn. Syst. Appl. 4 (2) (2009) 159–177.
- [6] T. Claeys, T. Grava, Universality of the break-up profile for the KdV equation in the small dispersion limit using the Riemann–Hilbert approach, Comm. Math. Phys. 286 (3) (2009) 979–1009.
- [7] O. Costin, H. Park, Y. Takei, Borel summability of the heat equation with variable coefficients, J. Differential Equations 252 (4) (2012) 3076–3092.
- [8] O. Costin, S. Tanveer, Existence and uniqueness for a class of nonlinear higher-order partial differential equations in the complex plane, Comm. Pure Appl. Math. 53 (9) (2000) 1092–1117.
- [9] O. Costin, S. Tanveer, Short time existence and Borel summability in the Navier–Stokes equation in \mathbb{R}^3 , Comm. Partial Differential Equations 34 (7–9) (2009) 785–817.
- [10] B. Dubrovin, On Hamiltonian perturbations of hyperbolic systems of conservation laws. II. Universality of critical behaviour, Comm. Math. Phys. 267 (1) (2006) 117–139.
- [11] B. Dubrovin, M. Elaeva, On the critical behavior in nonlinear evolutionary PDEs with small viscosity, Russ. J. Math. Phys. 19 (4) (2012) 449–460.
- [12] G. Folland, Introduction to Partial Differential Equations, second edition, Princeton University Press, Princeton, NJ, 1995, xii+324 pp.
- [13] S. Hirose, On a WKB theoretic transformation for a completely integrable system near a degenerate point where two turning points coalesce, 2013, preprint RIMS.
- [14] P. Hsieh, Y. Sibuya, Basic Theory of Ordinary Differential Equations, Universitext, Springer-Verlag, New York, 1999.

- [15] T. Koike, On the Hamiltonian structures of the second and the fourth Painlevé hierarchies, and the degenerate Garnier systems, in: Algebraic, Analytic and Geometric Aspects of Complex Differential Equations and Their Deformations. Painlevé Hierarchies, in: RIMS Kôkyûroku Bessatsu, vol. B2, Res. Inst. Math. Sci. (RIMS), Kyoto, 2007, pp. 99–127.
- [16] T. Koike, On new expressions of the Painlevé hierarchies, in: Algebraic Analysis and the Exact WKB Analysis for Systems of Differential Equations, in: RIMS Kôkyûroku Bessatsu, vol. B5, Res. Inst. Math. Sci. (RIMS), Kyoto, 2008, pp. 153–198.
- [17] A. Lastra, S. Malek, J. Sanz, On Gevrey solutions of threefold singular nonlinear partial differential equations, J. Differential Equations 255 (10) (2013) 3205–3232.
- [18] D. Lutz, M. Miyake, R. Schäfke, On the Borel summability of divergent solutions of the heat equation, Nagoya Math. J. 154 (1999) 1–29.
- [19] G. Lysik, Borel summable solutions of the Burgers equation, Ann. Polon. Math. 95 (2) (2009) 187–197.
- [20] G. Lysik, The Borel summable solutions of heat operators on a real analytic manifold, J. Math. Anal. Appl. 410 (1) (2014) 117–123.
- [21] S. Malek, On the summability of formal solutions for doubly singular nonlinear partial differential equations, J. Dyn. Control Syst. 18 (1) (2012) 45–82.
- [22] S. Malek, On Gevrey asymptotics for some nonlinear integro-differential equations, J. Dyn. Control Syst. 16 (3) (2010) 377–406.
- [23] S. Michalik, Summability and fractional linear partial differential equations, J. Dyn. Control Syst. 16 (4) (2010) 557–584.
- [24] T. Mandai, Existence and nonexistence of null-solutions for some non-Fuchsian partial differential operators with T -dependent coefficients, Nagoya Math. J. 122 (1991) 115–137.
- [25] S. Ouchi, Multisummability of formal power series solutions of nonlinear partial differential equations in complex domains, Asymptot. Anal. 47 (3–4) (2006) 187–225.
- [26] W. Rudin, Functional Analysis, second edition, International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991.
- [27] H. Tahara, H. Yamazawa, Multisummability of formal solutions to the Cauchy problem for some linear partial differential equations, J. Differential Equations 255 (10) (15 November 2013) 3592–3637.