



Homogenization of generalized second-order elliptic difference operators [☆]

Alexandre B. Simas ^{a,*}, Fábio J. Valentim ^{b,*}

^a Departamento de Matemática, Universidade Federal da Paraíba, Cidade Universitária – Campus I, 58051-970, João Pessoa, PB, Brazil

^b Departamento de Matemática, Universidade Federal do Espírito Santo, Av. Fernando Ferrari, 514, Goiabeiras, 29075-910, Vitória, ES, Brazil

Received 7 February 2017; revised 14 May 2018

Available online 22 May 2018

Abstract

Consider a function $W(x_1, \dots, x_d) = \sum_{k=1}^d W_k(x_k)$, where each $W_k : \mathbb{R} \rightarrow \mathbb{R}$ is a strictly increasing right continuous function with left limits. Given a matrix function $A = \text{diag}\{a_1, \dots, a_d\}$, let $\nabla A \nabla_W = \sum_{k=1}^d \partial_{x_k} (a_k \partial_{W_k})$ be a generalized second-order differential operator. Our chief goal is to study the homogenization of generalized second-order difference operators, that is, we are interested in the convergence of the sequence of solutions of

$$\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f^N$$

to the solution of

$$\lambda u - \nabla A \nabla_W u = f,$$

where the superscript N stands for some sort of discretization. In the continuous case we study the problem in the context of W -Sobolev spaces, whereas in the discrete case we develop the theoretical context in the present paper. The main result is a homogenization result. Under minor assumptions regarding weak convergence and ellipticity of these matrices A^N , we show that every such sequence admits a homogenization. We provide two examples of matrix functions verifying these assumptions: the first one consists of fixing a matrix function A under minor regularity assumptions, and taking a convenient discretization A^N ;

[☆] Research supported by CNPq (grant No. 309947/2015-4) and FAPES (grant No. 53551150/2011).

^{*} Corresponding authors.

E-mail addresses: alexandre@mat.ufpb.br (A.B. Simas), fabio.valentim@ufes.br (F.J. Valentim).

the second one consists on the case where A^N represents a random environment associated to an ergodic group, a case in which we then show that the homogenized matrix A does not depend on the realization ω of the environment. Finally, we provide an application geared towards the hydrodynamical limit of certain gradient processes.

© 2018 Elsevier Inc. All rights reserved.

MSC: 46E35; 35J15

Keywords: Homogenization; Difference operators; Differential operators

1. Introduction

In the '50s William Feller introduced a generalized notion of differential operator of the type $(d/dW)(d/dV)$ where, typically, W and V are strictly increasing functions and V (but not necessarily W) is continuous. In this paper we are interested in the formal adjoint of $(d/dW)(d/dV)$, which is given by $(d/dV)(d/dW)$, in the case $V(x) = x$ is the identity function. For further details on these, so called, Feller operators, we refer the reader to [3,4,10].

Recently, the formal adjoint operator $(d/dx)(d/dW)$ and some non-linear versions of it were obtained as scaling limits of interacting particle systems in inhomogeneous media. They may model diffusions with permeable membranes at the points of discontinuities of W , see [1,2,5,7,14] for further details. In [14], for instance, the author introduced an extension of the formal adjoint operator to higher dimensions and provided some results regarding this extension.

We will now briefly describe the operator on which we are interested. Fix a strictly increasing right continuous functions with left limits and periodic increments, $W_k : \mathbb{R} \rightarrow \mathbb{R}$, $k = 1, \dots, d$, and let $W(x) = \sum_{k=1}^d W_k(x_k)$ for $x \in \mathbb{R}^d$. We say that the operator $\mathcal{L}_W = \sum_{k=1}^d \partial_{x_k} \partial_{W_k}$ is a *generalized Laplacian operator*. Note that if $W_k(x) = x$, $k = 1, \dots, d$, then we recover the standard Laplacian operator. Additionally, in [12], the space of functions f having *weak generalized gradients* $\nabla_W f = (\partial_{W_1} f, \dots, \partial_{W_d} f)$, referred to as W -Sobolev spaces, have been studied and several properties, analogous to the classical Sobolev spaces, were proved. Moreover, in [13], the analysis has been further extended to W -Sobolev spaces of higher order where elliptic regularity results were obtained.

Our goal in this paper is to approximate the unique solution of a certain partial differential equation driven by the generalization of the Laplacian operator by a sequence of discrete functions arising as the unique solutions of discrete versions of the partial differential equation.

A first contribution of the present paper, presented in subsection 4.1, is to build a theory in the discrete setup analogous to the theory introduced in [12]. In particular, we introduce the notion of W -interpolation and present results relating discrete and continuous Sobolev spaces.

Our main contribution rests in Section 5 and it pertains the homogenization discussed early in this introduction. In a nutshell: motivated by discrete problems of the type $\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f^N$, where A^N is a sequence of diagonal matrix functions; we will define the notion of homogenization of the sequence A^N and prove that, under minor assumptions regarding weak convergence and ellipticity every such sequence admits homogenization – see Theorem 5.9.

Amongst several possible applications, we will readily apply Theorem 5.9 in two distinct contexts. Firstly, in Proposition 5.10, we are able to associate, certain matrices A to a convenient discretization A^N and correspondent solutions u_N , then we show that u_N converges to a func-

tion u solving $\lambda u - \nabla A \nabla_W u = f$. Secondly, we look at an example of stochastic homogenization, where a sequence of random matrices A^N satisfying an ergodicity condition is shown, in Theorem 5.11, to admit homogenization A (independent of realization) of the random environment. The focus of this approach is to study the asymptotic behavior of the coefficients for a family of random difference schemes, whose coefficients can be obtained by the discretization of random high-contrast lattice structures. In this sense, we want to extend the theory of homogenization of random difference operators developed in [11], which was also tackled in seminal paper by Kozlov [8]. We also want to generalize its main Theorem (Theorem 2.16) to the context in which we have weak generalized derivatives in the sense of W -Sobolev spaces.

Finally, as an application of this theory, we prove a hydrodynamic limit result, which is, per se, an interesting result, since it extends the main result obtained by [14]. More precisely, one must observe that this is a non-trivial extension, since, in the case of a non-constant matrix A , large difficulties arise in the application of the method presented there. In fact, the method presented in [14] consisted in studying the spectral decomposition and the resolvent operator of $\mathcal{L}_W = \sum_{k=1}^d \partial_{x_k} \partial_{W_k}$. Thus if one tries to apply such method to the case of a non-constant matrix function A , one would need to study the spectral decomposition and the resolvent operator of $\sum_{k=1}^d \partial_{x_k} (a_k \partial_{W_k})$.

The remaining of the article is organized as follows: in Section 2 we provide a brief review on W -Sobolev spaces; in Section 3, we define a new space of test functions that is needed in subsequent sections; in Section 4 we introduce a discrete analogous to the continuous W -Sobolev spaces, and then we provide interpolation and projection results connecting the discrete and continuous W -Sobolev spaces; in Section 5 we provide the main results of this article; in Section 6 we apply the results of this article to prove a hydrodynamic limit for gradient processes with conductances. Finally, we provide an Appendix with a proof of an auxiliary result.

2. W -Sobolev space

In this Section we recall some notation and results from [5,12,14]. Denote by $\mathbb{T}^d = (\mathbb{R}/\mathbb{Z})^d = [0, 1)^d$ the d -dimensional torus and fix a function $W : \mathbb{R}^d \rightarrow \mathbb{R}$ such that

$$W(x_1, \dots, x_d) = \sum_{k=1}^d W_k(x_k), \quad (1)$$

where each $W_k : \mathbb{R} \rightarrow \mathbb{R}$ is a *strictly increasing* right continuous function with left limits (càdlàg), with periodic increments, in the sense that $W_k(u+1) - W_k(u) = W_k(1) - W_k(0)$ for all $u \in \mathbb{R}$.

Define the generalized partial derivative ∂_{W_k} of a function $f : \mathbb{T}^d \rightarrow \mathbb{R}$ by

$$\partial_{W_k} f(x_1, \dots, x_k, \dots, x_d) = \lim_{\epsilon \rightarrow 0} \frac{f(x_1, \dots, x_k + \epsilon, \dots, x_d) - f(x_1, \dots, x_k, \dots, x_d)}{W_k(x_k + \epsilon) - W_k(x_k)}, \quad (2)$$

if the above limit exists. Denote the generalized gradient of f by $\nabla_W f = (\partial_{W_1} f, \dots, \partial_{W_d} f)$, if the generalized partial derivatives ∂_{W_k} exist for all $k = 1, \dots, d$.

Let us remember the definition of the space $H_{1,W}(\mathbb{T}^d)$, called W -Sobolev space. We denote by $\langle \cdot, \cdot \rangle$ the inner product of the Hilbert space $L^2(\mathbb{T}^d)$ and by $\| \cdot \|$ its norm. Let $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ be the Hilbert space of measurable functions $f : \mathbb{T}^d \rightarrow \mathbb{R}$ such that

$$\int_{\mathbb{T}^d} f^2 d(x^k \otimes W_k) < \infty,$$

where $d(x^k \otimes W_k)$ represents the product measure on \mathbb{T}^d obtained from Lebesgue's measure in \mathbb{T}^{d-1} and the measure induced by W_k on \mathbb{T} :

$$d(x^k \otimes W_k) = dx_1 \cdots dx_{k-1} dW_k dx_{k+1} \cdots dx_d.$$

Denote by $\langle \cdot, \cdot \rangle_{x^k \otimes W_k}$ the inner product of $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$:

$$\langle f, g \rangle_{x^k \otimes W_k} = \int_{\mathbb{T}^d} f g d(x^k \otimes W_k),$$

and by $\| \cdot \|_{x^k \otimes W_k}$ the norm induced by this inner product.

The set \mathcal{A}_{W_k} of the eigenvectors of $\frac{d}{dx} \frac{d}{dW_k}$ forms a complete orthonormal system in $L^2(\mathbb{T})$ (the reader is referred to [5] for further details). Let $\mathbb{D}_W := \text{span}(\mathcal{A}_W)$, where

$$\mathcal{A}_W = \{ f; f = f_1 \otimes \cdots \otimes f_d, \quad f_k \in \mathcal{A}_{W_k}, k = 1, \dots, d \},$$

and \otimes denotes the tensor product. That is, for functions $f_1 \in \mathcal{A}_{W_1}, \dots, f_d \in \mathcal{A}_{W_d}$, their tensor product is the function $f = f_1 \otimes \cdots \otimes f_d$ that satisfies

$$f(x_1, \dots, x_d) = \prod_{k=1}^d f_k(x_k).$$

Also, $\text{span}(A)$ stands for the linear space generated by the set A .

Define the operator $\mathcal{L}_W : \mathbb{D}_W \rightarrow L^2(\mathbb{T}^d)$ as follows: for $f = \prod_{k=1}^d f_k \in \mathcal{A}_W$,

$$\mathcal{L}_W(f)(x_1, \dots, x_d) = \sum_{k=1}^d \prod_{j=1, j \neq k}^d f_j(x_j) \mathcal{L}_{W_k} f_k(x_k), \quad (3)$$

and extend to \mathbb{D}_W by linearity. In [14] it is proved that the set \mathcal{A}_W forms a complete, orthonormal, countable system of eigenvectors for the operator \mathcal{L}_W . In particular, \mathbb{D}_W is dense in $L^2(\mathbb{T}^d)$.

Let $L^2_{x^k \otimes W_k, 0}(\mathbb{T}^d)$ be the closed subspace of $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ consisting of the functions f that have zero mean with respect to the measure $d(x^k \otimes W_k)$:

$$\int_{\mathbb{T}^d} f d(x^k \otimes W_k) = 0.$$

A function $g \in L^2(\mathbb{T}^d)$ has *W-generalized weak derivative* if for each $k = 1, \dots, d$ there exists a function $G_k \in L^2_{x^k \otimes W_k, 0}(\mathbb{T}^d)$ satisfying the following integral by parts identity:

$$\int_{\mathbb{T}^d} (\partial_{x_k} \partial_{W_k} f) g \, dx = - \int_{\mathbb{T}^d} (\partial_{W_k} f) G_k d(x^k \otimes W_k), \quad (4)$$

for every function $f \in \mathbb{D}_W$.

The W -Sobolev space $H_{1,W}(\mathbb{T}^d)$ is the set of functions in $L^2(\mathbb{T}^d)$ having W -generalized weak derivative. Note that $\mathbb{D}_W \subset H_{1,W}(\mathbb{T}^d)$ and if $g \in \mathbb{D}_W$ then $G_k = \partial_{W_k} g$. For this reason, for each function $g \in H_{1,W}(\mathbb{T}^d)$ we denote G_k simply by $\partial_{W_k} g$, and we call it the k -th *generalized weak derivative* of the function g with respect to W . In [12] it is shown that G_k is unique almost everywhere and the set $H_{1,W}(\mathbb{T}^d)$ is a Hilbert space with respect to the inner product

$$\langle f, g \rangle_{1,W} = \langle f, g \rangle + \sum_{k=1}^d \int_{\mathbb{T}^d} (\partial_{W_k} f)(\partial_{W_k} g) d(x^k \otimes W_k),$$

and denote by $\|\cdot\|_{1,W}$ its induced norm. Furthermore, $H_{1,W}^{-1}(\mathbb{T}^d)$, the dual space to $H_{1,W}(\mathbb{T}^d)$ (that is, the set of bounded linear functionals on $H_{1,W}(\mathbb{T}^d)$), has the following characterization:

Lemma 2.1. $f \in H_{1,W}^{-1}(\mathbb{T}^d)$ if and only if there exist functions $f_0 \in L^2(\mathbb{T}^d)$ and $f_k \in L^2_{x^k \otimes W_k, 0}(\mathbb{T}^d)$, such that

$$f = f_0 - \sum_{k=1}^d \partial_{x_k} f_k, \quad (5)$$

in the sense that for $v \in H_{1,W}(\mathbb{T}^d)$

$$f(v) := \int_{\mathbb{T}^d} f_0 v \, dx + \sum_{k=1}^d \int_{\mathbb{T}^d} f_k (\partial_{W_k} v) d(x^k \otimes W_k).$$

Furthermore,

$$\|f\|_{-1,W} = \inf \left\{ \left(\int_{\mathbb{T}^d} |f_0|^2 \, dx + \sum_{k=1}^d \int_{\mathbb{T}^d} |f_k|^2 d(x^k \otimes W_k) \right)^{1/2} ; \quad f \text{ satisfies (5)} \right\}.$$

3. A new space of test functions

In this paper we deal with discrete approximations of functions defined on the torus \mathbb{T}^d . In many applications, one will eventually need to apply these discretizations to test functions of the W -Sobolev space. The problem is that these test functions are defined as functions in $L^2(\mathbb{T}^d)$, and thus, the discretization procedure obtained by restricting the test function to \mathbb{T}^d_N is not well-defined. Therefore, we will define a new space of test functions suitable for discretizations, which we will denote by $\mathcal{D}_W(\mathbb{T}^d)$. A first application of this result will appear in the proof of the Compensated Compactness Theorem (Lemma 5.4). We will also use this space, as the space of test functions in the proof of the hydrodynamic limit given in Section 6. Another motivation for such

definition is of independent interest: to develop a classical theory of functions that admit generalized derivatives. In fact, these test functions will be such that one may apply the operator $\partial_{x_k} \partial_{W_k}$ in the classical sense.

Note that in [12] we considered another space for test functions, namely \mathbb{D}_W , whose definition was given in the previous Section.

To this end, recall from (1) the definition of W . For $f : \mathbb{T} \rightarrow \mathbb{R}$, let $D(f)$ be the set of its discontinuity points. For $k = 1, \dots, d$, let $C_{W_k}(\mathbb{T})$ be the set of càdlàg functions $f : \mathbb{T} \rightarrow \mathbb{R}$ such that $D(f) \subset D(W_k)$. We endow $C_{W_k}(\mathbb{T})$ with the sup norm $\|\cdot\|_\infty$. Note that equation (2) in the one-dimensional case becomes

$$\frac{df}{dW_k}(x) = \lim_{\epsilon \rightarrow 0} \frac{f(x + \epsilon) - f(x)}{W_k(x + \epsilon) - W_k(x)},$$

if the above limit exists. Let \mathfrak{D}_{W_k} be the set of functions f in $C_{W_k}(\mathbb{T})$ such that $\frac{df}{dW_k}(x)$ is well-defined and differentiable, and that $\frac{d}{dx}(\frac{df}{dW_k})$ belongs to $C_{W_k}(\mathbb{T})$.

In [5], it is proved that \mathfrak{D}_{W_k} is the set of functions f in $C_{W_k}(\mathbb{T})$ such that

$$f(x) = a + bW_k(x) + \int_{(0,x]} dW_k(y) \int_0^y g(z) dz \quad (6)$$

for some function g in $C_{W_k}(\mathbb{T})$, where a, b are real numbers satisfying

$$bW_k(1) + \int_{\mathbb{T}} dW_k(y) \int_0^y g(z) dz = 0, \quad \int_{\mathbb{T}} g(z) dz = 0. \quad (7)$$

The first requirement corresponds to the boundary condition $f(1) = f(0)$ and the second one to the boundary condition $df/dW_k(1) = df/dW_k(0)$.

Let us now define a d -dimensional counterpart to the sets $C_{W_k}(\mathbb{T})$ and \mathfrak{D}_{W_k} , respectively:

$$C_W(\mathbb{T}^d) = \text{span} \{f; f = f_1 \otimes \dots \otimes f_d, \quad f_k \in C_{W_k}(\mathbb{T}), k = 1, \dots, d\} \quad \text{and} \quad (8)$$

$$\mathfrak{D}_W(\mathbb{T}^d) = \text{span} \{f; f = f_1 \otimes \dots \otimes f_d, \quad f_k \in \mathfrak{D}_{W_k}, k = 1, \dots, d\}. \quad (9)$$

In other words, $C_W(\mathbb{T}^d)$ and $\mathfrak{D}_W(\mathbb{T}^d)$ are the spaces generated by functions of the form $f_1 \otimes \dots \otimes f_d$.

Note that the closure of $\mathfrak{D}_W(\mathbb{T}^d)$ is the tensor product of the spaces $\mathfrak{D}_{W_1}, \dots, \mathfrak{D}_{W_d}$.

Remark 3.1. Note that if $f \in \mathfrak{D}_W(\mathbb{T}^d)$, then $\partial_{W_k} f$ admits a partial derivative in the k th direction, therefore, if $a_k \in C_W(\mathbb{T}^d)$ is such that its partial derivative in the k th direction exists, then, the function $a_k \partial_{W_k} f$ also admits a partial derivative in the k th direction and we have

$$\partial_{x_k} (a_k \partial_{W_k} f) = (\partial_{W_k} f)(\partial_{x_k} a_k) + a_k \partial_{x_k} \partial_{W_k} f$$

in the strong sense.

The above remark motivates the following definition. Let $\mathbb{M}_W \subset (C_W(\mathbb{T}^d))^d$ be the space of functions $a = (a_1, \dots, a_d)$ such that for every $k = 1, \dots, d$, $\partial_{x_k} a_k$ exists. By convenience, we will also say that a diagonal matrix $A = \text{diag}\{a_1, \dots, a_d\}$ belongs to \mathbb{M}_W to mean that the function $a = (a_1, \dots, a_d)$ belongs to \mathbb{M}_W .

Remark 3.2. It is easy to see that if $f \in \mathfrak{D}_W(\mathbb{T}^d)$, one may apply the operator $\partial_{x_k} \partial_{W_k}$ in the classical sense. Actually, we have that for $f \in \mathfrak{D}_W(\mathbb{T}^d)$, and $a \in \mathbb{M}_W$, $\nabla A \nabla_W f \in C_W(\mathbb{T}^d)$, where $\nabla A \nabla_W f = \sum_{k=1}^d \partial_{x_k} (a_k \partial_{W_k} f)$.

The following result shows that $\mathfrak{D}_W(\mathbb{T}^d)$ can be used as a space of test functions:

Proposition 3.3. *The space $\mathfrak{D}_W(\mathbb{T}^d)$ satisfies:*

- i) $\mathfrak{D}_W(\mathbb{T}^d)$ is dense in $L^2(\mathbb{T}^d)$ in the L^2 -norm;
- ii) For $k = 1, \dots, d$, $\mathfrak{D}_W(\mathbb{T}^d)$ is dense in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ in the $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ -norm;
- iii) Functions in $C(\mathbb{T}^d)$, the space of continuous functions in \mathbb{T}^d , can be approximated by functions in $\mathfrak{D}_W(\mathbb{T}^d)$ in the sup norm $\|\cdot\|_\infty$.

Proof. We begin by noting that the first and second statements follow directly from the third statement. We will, thus, prove the third statement.

Note that from Proposition A.1 in Appendix, \mathfrak{D}_{W_k} is dense in $C(\mathbb{T})$ in the sup norm. Since \mathbb{T} is a compact Hausdorff space we may use the Stone–Weierstrass Theorem to extend this result to the d -dimensional case. Indeed, the Stone–Weierstrass Theorem implies that

$$C(\mathbb{T}^d) = \bigotimes_{k=1}^d C(\mathbb{T}),$$

where $\bigotimes_{k=1}^d C(\mathbb{T})$ is the tensor product of the spaces $C(\mathbb{T})$, that is, $\bigotimes_{k=1}^d C(\mathbb{T})$ is the closure of the space generated by functions of the form $f_1 \otimes \dots \otimes f_d$, with $f_k \in C(\mathbb{T})$, $k = 1, \dots, d$.

Since \mathfrak{D}_{W_k} is dense in $C(\mathbb{T})$, it is easy to see that $\mathfrak{D}_W(\mathbb{T}^d)$ is dense in $C(\mathbb{T}^d)$. \square

We say that a function $f : \mathbb{T}^d \rightarrow \mathbb{R}$ is right-continuous if for all $x \in \mathbb{T}^d$, we have $\lim_{y \downarrow x} f(y) = f(x)$, where $y \downarrow x$ means that for each $k = 1, \dots, d$, $y_k \downarrow x_k$. Observe that for $f_k \in C_{W_k}$ we have $\lim_{y_k \downarrow x_k} f_k(y_k) = f_k(x_k)$. Thus, it is clear that for $f = f_1 \otimes \dots \otimes f_d$, we have $\lim_{y \downarrow x} f(y) = f(x)$. We then have the following remark:

Remark 3.4. Every function $f \in C_W(\mathbb{T}^d)$ is right-continuous. In particular, since $\mathfrak{D}_W(\mathbb{T}^d) \subset C_W(\mathbb{T}^d)$, every function $f \in \mathfrak{D}_W(\mathbb{T}^d)$ is right-continuous.

Lemma 3.5. *Let $u \in C_W(\mathbb{T}^d)$, then u is bounded. In particular, if $u \in \mathfrak{D}_W(\mathbb{T}^d)$, then u is bounded.*

Proof. From the definition of $C_W(\mathbb{T}^d)$, there exist $b_j \in \mathbb{R}$, and functions $u_k^j \in C_{W_k}(\mathbb{T})$, $k = 1, \dots, d$, $j = 1, \dots, m$, for some $m \in \mathbb{N}$, such that

$$u(x) = \sum_{j=1}^m b_j u_1^j \otimes \cdots \otimes u_d^j(x).$$

Therefore, it is enough to show that each function u_k^j is bounded in \mathbb{T} . Suppose that u_k^j is not bounded, then, there exists a sequence (t_n) in \mathbb{T} such that

$$|u_k^j(t_n)| \xrightarrow{n \rightarrow \infty} \infty.$$

By compactness of \mathbb{T} and using the identification $\mathbb{T} = [0, 1)$, we can find a monotone subsequence t_{n_i} , such that $\lim_i t_{n_i} \in \mathbb{T}$. If t_{n_i} is increasing, then the existence of left limits of u_k^j shows that $|u_k^j(t_{n_i})|$ is bounded, whereas if t_{n_i} is decreasing, the right-continuity shows that $|u_k^j(t_{n_i})|$ is bounded. This contradiction concludes the proof. \square

4. The discrete W -Sobolev space

In this Section, we introduce the notion of discrete W -Sobolev spaces. This can be seen as a counterpart to the continuous case considered in [12]. Furthermore, we introduce the W -interpolation, and we also deal with other types of interpolations. Finally, we present some results in order to get connections between the discrete and continuous Sobolev spaces.

4.1. The space $H_{1,W}(\mathbb{T}_N^d)$

Denote by $\mathbb{T}_N^d = (\mathbb{Z}/N\mathbb{Z})^d = \{0, \dots, N-1\}^d$ the d -dimensional discrete torus with N^d points. We will now define and obtain some results on a discrete version of the W -Sobolev space.

For $f : \frac{1}{N}\mathbb{T}_N^d \rightarrow \mathbb{R}$, let the operators $\partial_{x_k}^N$, the standard difference operator in the k th canonical direction, and $\partial_{W_k}^N$, the W_k -difference operator, be given, respectively, by:

$$\partial_{x_k}^N f\left(\frac{x}{N}\right) = N \left[f\left(\frac{x+e_k}{N}\right) - f\left(\frac{x}{N}\right) \right] \quad \text{and} \quad \partial_{W_k}^N f\left(\frac{x}{N}\right) = \frac{f\left(\frac{x+e_k}{N}\right) - f\left(\frac{x}{N}\right)}{W\left(\frac{x+e_k}{N}\right) - W\left(\frac{x}{N}\right)}, \quad (10)$$

for $x \in \mathbb{T}_N^d$.

Denote by $L^2(\mathbb{T}_N^d)$, $L_{W_k}^2(\mathbb{T}_N^d)$ and $H_{1,W}(\mathbb{T}_N^d)$ the Hilbert spaces of the functions defined on $\frac{1}{N}\mathbb{T}_N^d$ with respect to the following inner products and their induced norms:

$$\begin{aligned} \langle f, g \rangle_N &:= \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} f(x/N) g(x/N), \\ \langle f, g \rangle_{W_k, N} &:= \frac{1}{N^{d-1}} \sum_{x \in \mathbb{T}_N^d} f(x/N) g(x/N) (W((x+e_k)/N) - W(x/N)), \\ \langle f, g \rangle_{1, W, N} &:= \langle f, g \rangle_N + \sum_{k=1}^d \langle \partial_{W_k}^N f, \partial_{W_k}^N g \rangle_{W_k, N}, \end{aligned}$$

and

$$\|f\|_N^2 = \langle f, f \rangle_N, \quad \|f\|_{W_k, N}^2 = \langle f, f \rangle_{W_k, N} \text{ and } \|f\|_{1, W, N}^2 = \langle f, f \rangle_{1, W, N},$$

respectively.

These norms are natural discretizations of the norms considered in the continuous version. The Lemma below is a discrete version of the Poincaré inequality. The proof is omitted since it is analogous to the proof of its continuous version, given in Corollary 2.7 in [12].

Lemma 4.1 (Discrete Poincaré inequality). *There exists a finite constant C depending on d and W , but not on N , such that*

$$\left\| f - \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} f(x/N) \right\|_N \leq C \|\nabla_W^N f\|_{W, N},$$

for all $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, where

$$\|\nabla_W^N f\|_{W, N}^2 = \sum_{k=1}^d \|\partial_{W_k}^N f\|_{W_k, N}^2.$$

We will now provide results on discrete elliptic equations. We would like to remark that the proofs of these lemmas are identical to the ones in the continuous case, and the reader is then referred to [12].

Let $\lambda \geq 0$ and $A = \text{diag}\{a_1(x), \dots, a_d(x)\}$, $x \in \mathbb{T}^d$, be a diagonal matrix function satisfying the ellipticity condition:

$$\begin{aligned} &\text{there exists a constant } \theta > 0 \text{ such that } \theta^{-1} \leq a_k(x) \leq \theta, \\ &\text{for every } x \in \mathbb{T}^d \text{ and every } k = 1, \dots, d. \end{aligned} \quad (11)$$

Given a function $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, we are interested in studying the problem

$$T_\lambda^N u = f, \quad (12)$$

where $u : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ is the unknown function and T_λ^N denotes the discrete generalized elliptic operator

$$T_\lambda^N u := \lambda u - \nabla^N A \nabla_W^N u, \quad (13)$$

with

$$\nabla^N A \nabla_W^N u := \sum_{k=1}^d \partial_{x_k}^N \left(a_k(x/N) \partial_{W_k}^N u \right).$$

The bilinear form $B^N[\cdot, \cdot]$ associated with the elliptic operator T_λ^N is given by

$$B^N[u, v] = \lambda \langle u, v \rangle_N + \frac{1}{N^{d-1}} \sum_{k=1}^d \sum_{x \in \mathbb{T}_N^d} a_k(x/N) (\partial_{W_k}^N u) (\partial_{W_k}^N v) [W_k((x_k + 1)/N) - W_k(x_k/N)], \quad (14)$$

where $u, v : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$.

A function $u : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ is said to be a weak solution of (12) if

$$B^N[u, v] = \langle f, v \rangle_N \text{ for all } v : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}.$$

Denote by $H_{1,W}^\perp(\mathbb{T}_N^d)$ the subspace of $H_{1,W}(\mathbb{T}_N^d)$ formed by functions $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ which are orthogonal to the constant functions with respect to the inner product $\langle \cdot, \cdot \rangle_{1,W,N}$. Note that $H_{1,W}^\perp(\mathbb{T}_N^d)$ is the set of functions $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ such that

$$\frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} f(x/N) = 0.$$

Remark 4.2. Note that in the discrete Sobolev space $H_{1,W}(\mathbb{T}_N^d)$ we have a “Dirac measure” concentrated at a point x as a function: the function that takes value N^d in x and zero elsewhere. Therefore, we may integrate these weak solutions with respect to this function to obtain that every weak solution is, in fact, a strong solution. Hence, a function u is a weak solution of the discrete problem (12) if, and only if, it is a strong solution. We chose to present weak solutions of discrete problems because it is easy to obtain existence and uniqueness results following this approach, and the results are stated in a very similar fashion to those in the continuous case, thus making the analogy with the continuous case more evident.

Lemma 4.3. Given a function $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, the equation

$$\nabla^N A \nabla_{W^N}^N u = f,$$

has a weak solution $u : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ if and only if $f \in H_{1,W}^\perp(\mathbb{T}_N^d)$. In this case we have uniqueness of the solution disregarding addition by constants. Moreover, if $u \in H_{1,W}^\perp(\mathbb{T}_N^d)$ we have the bound

$$\|u\|_{1,W,N} \leq C \|f\|_N,$$

where $C > 0$ does not depend on f nor on N .

Motivated by Lemma 2.1 we are able to obtain a characterization of the elements of the discrete dual Sobolev space.

Lemma 4.4. Let f belong to $H_{1,W}^{-1}(\mathbb{T}_N^d)$, the discrete dual Sobolev space and $\|\cdot\|_{-1,W,N}$ its norm. Then, there exist unique functions $f_k : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, $k = 0, 1, \dots, d$ such that for all $v : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$, the action of f over v is given by

$$\begin{aligned}
 f(v) &:= \langle f_0, v \rangle_N + \sum_{k=1}^d \langle f_k, \partial_{W_k}^N v_k \rangle_{W_k, N} \\
 &= \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} f_0(x/N) v(x/N) + \frac{1}{N^{d-1}} \sum_{k=1}^d \sum_{x \in \mathbb{T}_N^d} f_k(x/N) \partial_{W_k}^N v(x/N) \left[W((x + e_k/n) - W(x)) \right].
 \end{aligned}$$

Moreover, $\|f\|_{-1, W, N}^2 = \|f_0\|_N^2 + \sum_{k=1}^d \|f_k\|_{W_k, N}^2$.

In notation of the Lemma above, we denote the functional f by

$$f = f_0 - \sum_{k=1}^d \partial_{x_k}^N f_k.$$

Proof. Let u be the unique weak solution obtained by Lemma 4.5 when $\lambda = 1$, A the identity matrix and, $f \in H_{1, W}^{-1}(\mathbb{T}_N^d)$. In particular, we have $(f, v) = \langle u, v \rangle_{1, W, N}$. Thus, it suffices to consider $f_0 = u$ and $f_k = \partial_{W_k}^N u$, $k = 1, \dots, d$. \square

Lemma 4.5. Let $\lambda > 0$ and $f : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$. There exists a unique weak solution $u : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ of the equation

$$\lambda u - \nabla^N A \nabla_{W, N}^N u = f. \quad (15)$$

Moreover,

$$\|u\|_{1, W, N} \leq C \|f\|_{-1, W, N}, \text{ and } \|u\|_N \leq \lambda^{-1} \|f\|_{-1, W, N},$$

where $C > 0$ does not depend neither on f nor on N .

4.2. Connections between the discrete and continuous Sobolev spaces

For each $x \in \frac{1}{N} \mathbb{T}_N^d$, let $Q_N(x)$ be the set

$$Q_N(x) = \{y \in \mathbb{T}^d; x_k \leq y_k < x_k + 1/N, k = 1, 2, \dots, d\}. \quad (16)$$

We will now introduce some different interpolation schemes, that is, procedures to extend a mesh function $u_N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ from $\frac{1}{N} \mathbb{T}_N^d$ to the continuous torus \mathbb{T}^d . The first scheme is the *piecewise-constant interpolation* $\tilde{u}_N : \mathbb{T}^d \rightarrow \mathbb{R}$ given by

$$\tilde{u}_N(y) = \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} u_N(x) 1_{Q_N(x)}(y),$$

where 1_A denotes the indicator function of the set A . However, it is not straightforward to extend a function u_N from $\frac{1}{N} \mathbb{T}_N^d$ to \mathbb{T}^d in such a way that it belongs to $H_{1, W}(\mathbb{T}^d)$. To do so, we need the definition of W -interpolation. Motivated by the standard construction of the d -dimensional linear interpolation, see for instance [9, chapter VI], we define the W -interpolation $u_N^* : \mathbb{T}^d \rightarrow \mathbb{R}$ by

$$\begin{aligned}
u_N^*(y) &= u_N(x) + \sum_{k=1}^d \partial_{W_k}^N u_N(x) (W_k(y_k) - W(x_k)) + \cdots \\
&+ \sum_{k=1}^d \partial_{W_1}^N \cdots \partial_{W_{k-1}}^N \partial_{W_{k+1}}^N \cdots \partial_{W_d}^N u_N(x) \prod_{\substack{i=1 \\ i \neq k}}^d (W_i(y_i) - W_i(x_i)) \\
&+ \partial_{W_1}^N \cdots \partial_{W_d}^N u_N(x) \prod_{k=1}^d (W_k(y_k) - W_k(x_k)),
\end{aligned}$$

where $y \in Q_N(x)$.

Let us consider a third type of interpolation, which are actually d (one for each $m = 1, \dots, d$) interpolations:

$$\begin{aligned}
u_N^{(m)}(y) &= u_N(x) + \sum_{\substack{k=1 \\ k \neq m}}^d \partial_{W_k}^N u_N(x) (W_k(y_k) - W(x_k)) + \cdots \\
&+ \partial_{W_1}^N \cdots \partial_{W_{m-1}}^N \partial_{W_{m+1}}^N \cdots \partial_{W_d}^N u_N(x) \prod_{\substack{i=1 \\ i \neq m}}^d (W_i(y_i) - W_i(x_i)).
\end{aligned}$$

A simple calculation shows that

$$\frac{\partial u_N^*}{\partial W_m}(y) = \left(\partial_{W_m}^N u_N \right)^{(m)}(y). \quad (17)$$

We now establish the connection between the discrete and continuous spaces by showing how a sequence of functions defined in $\frac{1}{N}\mathbb{T}_N^d$ can converge to a function defined on \mathbb{T}^d .

We will say that the sequence of mesh functions $(f_N)_{N \in \mathbb{N}}$ in $L^2(\mathbb{T}_N^d)$ or $L_{W_k}^2(\mathbb{T}_N^d)$, $k = 1, \dots, d$, converges strongly (weakly) to the function $f \in L^2(\mathbb{T}^d)$ or $L_{W_k}^2(\mathbb{T}^d)$, as $N \rightarrow \infty$, if $\tilde{f}_N \rightarrow f$ in $L^2(\mathbb{T}^d)$ or $L_{W_k}^2(\mathbb{T}^d)$ strongly (weakly), respectively. Similarly, a sequence $(u_N)_{N \in \mathbb{N}}$ in $H_{1,W}(\mathbb{T}_N^d)$ converges strongly (weakly) to $u \in H_{1,W}(\mathbb{T}^d)$ if $u_N^* \rightarrow u$ strongly (weakly) in $H_{1,W}(\mathbb{T}^d)$.

With respect to convergence of functionals, we say that a sequence of functionals $f_N \in H_W^{-1}(\mathbb{T}_N^d)$, with representation

$$f_N = f_{0,N} - \sum_{k=1}^d \partial_{x_k}^N f_{k,N}$$

converges weakly (resp. strongly) to $f \in H_W^{-1}(\mathbb{T}^d)$ if the sequence of functionals F_N given by

$$F_N = \tilde{f}_{0,N} - \sum_{k=1}^d \partial_{x_k} \tilde{f}_{k,N}$$

converges weakly (resp. strongly) to f in $H_W^{-1}(\mathbb{T}^d)$.

Lemma 4.6. Let (u_N) be a sequence of functions $u_N : \frac{1}{N}\mathbb{T}_N^d \rightarrow \mathbb{R}$:

a) If there exists a constant $C > 0$ satisfying

$$\|u_N\|_N^2 = \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N^2(x/N) \leq C, \quad \text{for all } N \geq 1, \quad (18)$$

and one of the sequences (u_N^*) , (\tilde{u}_N) or $(u_N^{(m)})$ (for some $m = 1, \dots, d$) converges weakly in $L^2(\mathbb{T}^d)$ to a function $u : \mathbb{T}^d \rightarrow \mathbb{R}$ as $N \rightarrow \infty$. Then, the remaining sequences also converge to u in the same manner.

b) If, for some $k \in \{1, 2, \dots, d\}$, there exists a constant $C_k > 0$ satisfying

$$\|u_N\|_{W_k, N}^2 = \frac{1}{N^{d-1}} \sum_{x \in \mathbb{T}_n^d} u_N^2(x/N) [W((x + e_k)/N) - W(x/N)] \leq C_k, \quad \text{for all } N \geq 1, \quad (19)$$

and one of the sequences (u_N^*) , (\tilde{u}_N) or $(u_N^{(m)})$ (for some $m = 1, \dots, d$) converges weakly in $L_{W_k}^2(\mathbb{T}^d)$ to a function $u : \mathbb{T}^d \rightarrow \mathbb{R}$ as $N \rightarrow \infty$. Then, the remaining sequences also converge to u in the same manner.

Proof. We begin by proving item a). Suppose (u_N^*) converges weakly to u in $L^2(\mathbb{T}^d)$. We will prove that (\tilde{u}_N) also converges to u . The remaining cases can be handled in a similar manner. Note that (18) implies the uniform boundedness of the norms $\|u_N^*\|$, $\|\tilde{u}_N\|$ and $\|u_N^{(m)}\|$ because the values of the functions u_N^* , \tilde{u}_N and $u_N^{(m)}$ in each cell $Q_N(x)$, $x \in \mathbb{T}_N^d$, lies between the largest and smallest values of u_N at the vertices of $Q_N(x)$. Hence, each of these sequences are weakly compact in $L^2(\mathbb{T}^d)$. From this, to prove the convergence we are interested, it is enough to prove the convergence $\int_{\mathbb{T}^d} \tilde{u}_N \Phi \, dy \rightarrow \int_{\mathbb{T}^d} u \Phi \, dy$ for all $\Phi \in C^\infty(\mathbb{T}^d)$ (notice that we have, by assumption, that $\int_{\mathbb{T}^d} u_N^* \Phi \, dy \rightarrow \int_{\mathbb{T}^d} u \Phi \, dy$).

Let $\tilde{\Phi}_N$ be the piecewise-constant function which coincides with Φ at the lattice-points in $\frac{1}{N}\mathbb{T}_N^d$. By uniform convergence of $\tilde{\Phi}_N$ to Φ , we have $\int_{\mathbb{T}^d} u_N^* \tilde{\Phi}_N \, dy \rightarrow \int_{\mathbb{T}^d} u \Phi \, dy$. Let us consider

$$R_N := \int_{\mathbb{T}^d} (u_N^* - \tilde{u}_N) \tilde{\Phi}_N \, dy = \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} \Phi(x) \int_{Q_N(x)} (u_N^* - \tilde{u}_N) \, dy.$$

By the explicit form of the interpolations, we have

$$R_N = \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} \Phi(x) \left\{ \sum_{k=1}^d \partial_{W_k}^N u_N(x) \int_{Q_N(x)} (W_k(y_k) - W(x_k)) \, dy + \dots \right.$$

$$\begin{aligned}
& + \sum_{k=1}^d \partial_{W_1}^N \cdots \partial_{W_{k-1}}^N \partial_{W_{k+1}}^N \cdots \partial_{W_d}^N u_N(x) \int_{Q_N(x)} \prod_{\substack{i=1 \\ i \neq k}}^d (W_i(y_i) - W_i(x_i)) dy \\
& + \partial_{W_1}^N \cdots \partial_{W_d}^N u_N(x) \int_{Q_N(x)} \prod_{i=1}^d (W_i(y_i) - W_i(x_i)) dy \Big\}.
\end{aligned}$$

After a summation by parts we transfer, in each term inside the brackets, one of the W -differences from u_N to Φ and thus, the resulting expression is

$$\begin{aligned}
R_N = - \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \Big\{ \sum_{k=1}^d u_N(x) \partial_{W_k}^N \Phi(x) \int_{Q_N(x)} (W_k(y_k) - W(x_k)) dy + \cdots \\
+ \partial_{W_1}^N \Phi(x) \partial_{W_2}^N \cdots \partial_{W_d}^N u_N(x) \prod_{i=1}^d \int_{x_i}^{x_i + \frac{1}{N}} (W_i(y_i) - W_i(x_i)) dy_i \Big\},
\end{aligned}$$

where

$$\partial_{W_k}^N f\left(\frac{x}{N}\right) = \frac{f\left(\frac{x}{N}\right) - f\left(\frac{x-e_k}{N}\right)}{W\left(\frac{x+e_k}{N}\right) - W\left(\frac{x}{N}\right)}, \quad \text{for } x \in \mathbb{T}_N^d,$$

and the exchange of the order of integration is due to Fubini's Theorem. Note that

$$\begin{aligned}
& \left(\partial_{W_{i_1}}^N \Phi(x) \partial_{W_{i_2}}^N \cdots \partial_{W_{i_s}}^N u_N(x) \prod_{j=1}^s \int_{x_{i_j}}^{x_{i_j} + 1/N} [W_{i_j}(y_{i_j}) - W_{i_j}(x_{i_j})] dy_{i_j} \right)^2 \\
& \leq \left(\frac{1}{N^d} \partial_{W_{i_1}}^N \Phi(x) \partial_{W_{i_2}}^N \cdots \partial_{W_{i_s}}^N u_N(x) \prod_{j=1}^s [W_{i_j}(x_{i_j} + 1/N) - W_{i_j}(x_{i_j})] \right)^2 \\
& = \left(\frac{1}{N^d} \partial_{W_{i_1}}^N \Phi(x) [W_{i_1}(x_{i_1} + 1/N) - W_{i_1}(x_{i_1})] \right)^2 \\
& \quad \times \left(\partial_{W_{i_2}}^N \cdots \partial_{W_{i_s}}^N u_N(x) \prod_{j=2}^s [W_{i_j}(x_{i_j} + 1/N) - W_{i_j}(x_{i_j})] \right)^2 \\
& = \left(\frac{1}{N^d} \partial_{x_{i_1}}^N \Phi(x) \frac{1}{N} \right)^2 \left(C \sum_{k=1}^d u_N^2(x + e_k/N) \right)^2
\end{aligned}$$

where in the last equality the constant $C > 0$ does not depend on N .

In this way, $|R_N|$ is bounded by a finite sum (that depends on d) of terms of the form

$$\begin{aligned} \frac{C}{N} \frac{1}{N^d} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} |\partial_{\tilde{x}_{i_1}}^N \Phi(x)| \left(\sum_{k=1}^d u_N^2(x + e_k/N) \right)^{1/2} &\leq \\ \frac{C}{N^{d+1}} \left(\sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \partial_{\tilde{x}_{i_1}}^N \Phi^2(x) \right)^{1/2} \left(\sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \sum_{k=1}^d u_N^2(x + e_k/N) \right)^{1/2} &\leq \\ \frac{Cd}{N} \left(\frac{1}{N^d} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \partial_{\tilde{x}_{i_1}}^N \Phi^2(x) \right)^{1/2} \left(\frac{1}{N^d} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} u_N^2(x) \right)^{1/2}, \end{aligned}$$

where the previous estimates follow from Cauchy–Schwarz inequality. By hypothesis (18) and the fact that $\Phi \in C^\infty(\mathbb{T}^d)$ the sums inside the brackets are bounded. Thus, R_N goes to zero as $N \rightarrow \infty$. Therefore, we have shown that

$$\int_{\mathbb{T}^d} \tilde{u}_N \tilde{\Phi}_N dx \rightarrow \int_{\mathbb{T}^d} u \Phi dx.$$

Hence

$$\int_{\mathbb{T}^d} \tilde{u}_N \Phi dx = \int_{\mathbb{T}^d} \tilde{u}_N \tilde{\Phi}_N dx + \int_{\mathbb{T}^d} \tilde{u}_N [\Phi - \tilde{\Phi}_N] dx \rightarrow \int_{\mathbb{T}^d} u \Phi dx.$$

This concludes the proof of item a).

Now, let us consider the item b). The proof is similar to the previous one. To keep notation simple fix, without loss of generality, $k = 1$. The beginning of the proof is exactly the same as in the previous case, one just needs to replace the Lebesgue measure dy by the product measure $d(y_1 \otimes W_1)$. The essential difference comes in the estimation of the term

$$\left(\partial_{\tilde{W}_{i_1}}^N \Phi(x) \partial_{\tilde{W}_{i_2}}^N \cdots \partial_{\tilde{W}_{i_s}}^N u_N(x) \int_{Q_N(x)} \prod_{j=1}^s [W_{i_j}(y_{i_j}) - W_{i_j}(x_{i_j})] d(y_1 \otimes W_1) \right)^2.$$

We will now provide this estimation. The previous term is bounded above by

$$\begin{aligned} &\left(\frac{1}{N^{d-1}} \partial_{\tilde{W}_{i_1}}^N \Phi(x) \partial_{\tilde{W}_{i_2}}^N \cdots \partial_{\tilde{W}_{i_s}}^N u_N(x) [W_1(x_1 + 1/N) - W_1(x_1)] \times \right. \\ &\quad \left. \times \prod_{j=1}^s [W_{i_j}(x_{i_j} + 1/N) - W_{i_j}(x_{i_j})] \right)^2 \\ &= \left(\frac{[W_1(x_1 + \frac{1}{N}) - W_1(x_1)]}{N^{d-1}} \partial_{\tilde{W}_{i_1}}^N \Phi(x) [W_{i_1}(x_{i_1} + \frac{1}{N}) - W_{i_1}(x_{i_1})] \right)^2 \times \end{aligned}$$

$$\begin{aligned} & \times \left(\partial_{W_{i_2}}^N \cdots \partial_{W_{i_s}}^N u_N(x) \prod_{j=2}^s [W_{i_j}(x_{i_j} + \frac{1}{N}) - W_{i_j}(x_{i_j})] \right)^2 \\ & \leq \left(\frac{[W_1(x_1 + \frac{1}{N}) - W_1(x_1)]}{N^{d-1}} \partial_{x_{i_1}}^N \Phi(x) \frac{1}{N} \right)^2 \left(C_1 \sum_{k=1}^d u_N^2(x + e_k/N) \right)^2. \end{aligned}$$

So, the new term $|R_N|$ is bounded by a finite sum (that depends on d) of terms of the form

$$\begin{aligned} & \frac{C_1}{N} \left(\frac{1}{N^{d-1}} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} [W_1(x_1 + \frac{1}{N}) - W_1(x_1)] \partial_{x_{i_1}}^N \Phi^2(x) \right)^{1/2} \\ & \times \left(\frac{1}{N^{d-1}} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} [W_1(x_1 + \frac{1}{N}) - W_1(x_1)] u_N^2(x) \right)^{1/2}. \end{aligned}$$

By hypothesis (19) and the fact that $\Phi \in C^\infty(\mathbb{T}^d)$ the sums inside the brackets are bounded and thus R_N goes to zero as $N \rightarrow \infty$. Therefore, we have shown that

$$\int_{\mathbb{T}^d} \tilde{u}_N \tilde{\Phi}_N d(y_1 \otimes W_1) \rightarrow \int_{\mathbb{T}^d} u \Phi d(y_1 \otimes W_1).$$

Hence

$$\begin{aligned} \int_{\mathbb{T}^d} \tilde{u}_N \Phi d(y_1 \otimes W_1) &= \int_{\mathbb{T}^d} \tilde{u}_N \tilde{\Phi}_N d(y_1 \otimes W_1) + \int_{\mathbb{T}^d} \tilde{u}_N [\Phi - \tilde{\Phi}_N] d(y_1 \otimes W_1) \\ &\rightarrow \int_{\mathbb{T}^d} u \Phi d(y_1 \otimes W_1). \end{aligned}$$

This concludes the proof. \square

We will now state and prove the strong counterpart to the above Lemma:

Lemma 4.7. Let $u_N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ be a sequence of functions and $C > 0$ be a constant such that

$$\|u_N\|_{1,W,N} \leq C, \quad \text{for all } N \geq 1. \quad (20)$$

Then, if one of the sequences (u_N^*) , (\tilde{u}_N) or $(u_N^{(m)})$ (for some $m = 1, \dots, d$) converges in $L^2(\mathbb{T}^d)$ to a function $u : \mathbb{T}^d \rightarrow \mathbb{R}$ as $N \rightarrow \infty$, then the remaining sequences will converge to u in the same manner.

Proof. Let us consider the case $u_N^* \rightarrow u$ in $L^2(\mathbb{T}^d)$ and we will prove that $u_N^{(m)} \rightarrow u$ in $L^2(\mathbb{T}^d)$. The remaining cases can be proved analogously. Recall the notation introduced at the beginning

of this subsection and note that

$$R_N = \int_{\mathbb{T}^d} (u_N^* - u_N^{(m)})^2 dy \leq \sum_{x \in \frac{1}{N} \mathbb{T}_N^d \cap \mathcal{Q}_N(x)} \int (u_N^* - u_N^{(m)})^2 dy.$$

A simple computation shows that the quantity $|u_N^*(y) - u_N^{(m)}(y)|$ assumes its largest value over the closure of $\mathcal{Q}_N(x)$ at one of its vertices, and that this value is equal to $(W_m(x_m + 1/N) - W_m(x_m)) |\partial_{W_m}^N u_N(y)|$. That is,

$$\max_{y \in \bar{\mathcal{Q}}_N(x)} |u_N^*(y) - u_N^{(m)}(y)| = (W_m(x_m + 1/N) - W_m(x_m)) \max_{y: |y-x|=1/N, y_m=x_m} |\partial_{W_m}^N u_N(y)|.$$

In fact,

$$\begin{aligned} u_N^*(y) - u_N^{(m)}(y) &= (W_m(y_m) - W_m(x_m)) \left\{ \partial_{W_m}^N u_N(x) \right. \\ &\quad + \sum_{j=1; j \neq m}^d \partial_{W_j}^N \partial_{W_m}^N u_N(x) (W_j(y_j) - W(x_j)) + \cdots \\ &\quad + \sum_{j=1; j \neq m}^d \partial_{W_1}^N \cdots \partial_{W_{j-1}}^N \partial_{W_{j+1}}^N \cdots \partial_{W_d}^N \partial_{W_m}^N u_N(x) \prod_{\substack{i=1 \\ i \neq j}}^d (W_i(y_i) - W_i(x_i)) \\ &\quad \left. + \partial_{W_1}^N \cdots \partial_{W_d}^N \partial_{W_m}^N u_N(x) \prod_{i=1; i \neq m}^d (W_i(y_i) - W_i(x_i)) \right\} = \\ &\quad (W_m(y_m) - W_m(x_m)) (\partial_{W_m}^N u_N)^*(y_1, \dots, y_{m-1}, y_{m+1}, \dots, y_d), \end{aligned}$$

where $(\partial_{W_m}^N u_N)^*$ denotes the W -interpolation of the function $\partial_{W_m}^N u_N$ in the $(d-1)$ -dimensional setup.

In this way, R_N is bounded above by

$$\begin{aligned} &\sum_{x \in \frac{1}{N} \mathbb{T}_N^d \cap \bar{\mathcal{Q}}_N(x)} \int \left([W_m(x_m + 1/N) - W_m(x_m)] \max_{z: |z-x|=1/N, z_m=x_m} |\partial_{W_m}^N u_N(z)| \right)^2 dy = \\ &\sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \frac{1}{N^d} \left([W_m(x_m + 1/N) - W_m(x_m)] \max_{z: |z-x|=1/N, z_m=x_m} |\partial_{W_m}^N u_N(z)| \right)^2 \leq \\ &C_1 \frac{W_m(1) - W_m(0)}{N} \sum_{z \in \mathbb{T}_N^d} \left(\partial_{W_m}^N u_N(z) \right)^2 [W_m(z_m + 1/N) - W_m(z)] \frac{1}{N^{d-1}} = \\ &C_1 \frac{W_m(1) - W_m(0)}{N} \|\partial_{W_m}^N u_N\|_{W_m, N}^2 \leq C_1 \frac{W_m(1) - W_m(0)}{N} \|u_N\|_{1, W, N}^2 \end{aligned}$$

where, in the previous expression, C_1 is a constant that depends on d (one may take, for instance, $C_1 = 2^d$). So, by (20), $\lim_{N \rightarrow \infty} R_N = 0$.

Thus, it follows that

$$\int_{\mathbb{T}^d} (u_N^{(m)} - u)^2 dy \leq 2 \int_{\mathbb{T}^d} (u_N^* - u_N^{(m)})^2 + (u_N^* - u)^2 dy \rightarrow 0,$$

as $N \rightarrow \infty$. This concludes the proof. \square

Finally, we have the following Lemma regarding strong and weak compactness:

Lemma 4.8. *Let $u_N : \mathbb{T}^d \rightarrow \mathbb{R}$ be a sequence of functions such that there exists some constant $C \geq 0$ such that, for every $N \geq 1$,*

$$\|u_N\|_{1,W,N} \leq C. \quad (21)$$

Then, the sequence (u_N^) forms a uniformly bounded set in $H_{1,W}(\mathbb{T}^d)$, which is therefore strongly precompact in $L^2(\mathbb{T}^d)$ and weakly precompact in $H_{1,W}(\mathbb{T}^d)$.*

Proof. To prove the lemma, it is enough to show that (21) implies the uniform boundedness of $\|u_N^*\|_{1,W,N}$, and then apply [12, Proposition 2.9].

Note that

$$\begin{aligned} \|u_N^*\|^2 &= \int_{\mathbb{T}^d} (u_N^*(y))^2 dy = \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} \int_{Q_N(x)} (u_N^*(y))^2 dy \leq \\ &\sum_{x \in \frac{1}{N}\mathbb{T}_N^d} \int_{Q_N(x)} \sum_{z \in \frac{1}{N}\mathbb{T}_N^d; |z-x|=1/N} (u_N(z))^2 dy \leq \frac{2^d}{N^d} \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} u_N^2(x) = 2^d \|u_N\|_N^2, \end{aligned}$$

where z , in the previous expression, represents the summation over all vertices of the cell $Q_N(x)$. Further, for $m = 1, 2, \dots, d$, we have, by an explicit computation,

$$\begin{aligned} \partial_{W_m} u_N^*(y) &= \partial_{W_m}^N u_N(x) + \sum_{j=1; j \neq m}^d \partial_{W_j}^N \partial_{W_m}^N u_N(x) (W_j(y_j) - W(x_j)) + \dots \\ &+ \sum_{j=1; j \neq m}^d \partial_{W_1}^N \dots \partial_{W_{j-1}}^N \partial_{W_{j+1}}^N \dots \partial_{W_d}^N \partial_{W_m}^N u_N(x) \prod_{\substack{i=1 \\ i \neq j}}^d (W_i(y_i) - W_i(x_i)) \\ &+ \partial_{W_1}^N \dots \partial_{W_d}^N \partial_{W_m}^N u_N(x) \prod_{i=1; i \neq m}^d (W_i(y_i) - W_i(x_i)) = \\ &(\partial_{W_m}^N u_N)^*(y_1, \dots, y_{m-1}, y_{m+1}, \dots, y_d), \end{aligned}$$

where $(\partial_{W_m}^N u_N)^{*'}$ denotes the W -interpolation of the function $\partial_{W_m}^N u_N$ in the $(d-1)$ -dimensional setup.

In this way,

$$\begin{aligned} \|\partial_{W_m} u_N^*\|_{x^m \otimes W_m}^2 &= \int_{\mathbb{T}^d} (\partial_{W_m} u_N^*(y))^2 d(y^m \otimes W_m) \\ &= \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \int_{Q_N(x)} (\partial_{W_m} u_N^*(y))^2 d(y^m \otimes W_m) \\ &\leq \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} \int \sum_{z \in \frac{1}{N} \mathbb{T}_N^d; |z-x|=1/N} (\partial_{W_m}^N u_N(z))^2 d(y^m \otimes W_m) \\ &\leq \frac{2^d}{N^{d-1}} \sum_{x \in \frac{1}{N} \mathbb{T}_N^d} (\partial_{W_m}^N u_N(x))^2 \left(W_m\left(\frac{x_m+1}{N}\right) - W_m\left(\frac{x_m}{N}\right) \right) \\ &= 2^d \|\partial_{W_m}^N u_N\|_{W_m, N}^2. \end{aligned}$$

Thus we have shown that

$$\|u_N^*\|_{1, W} \leq 2^d \|u_N\|_{1, W, N} < 2^d C,$$

where the last inequality follows from (21), and this concludes the proof. \square

We will now obtain the converse procedure, that is, how to use a measurable function f in $L^2(\mathbb{T}^d)$ to properly define a mesh function in \mathbb{T}_N^d . This is not straightforward, since the restriction of f to the set $\{0, 1/N, \dots, (N-1)/N\}^d$ is not well-defined (this is a set of Lebesgue measure zero).

For arbitrary functions $f \in L^2(\mathbb{T}^d)$ and $g \in L^2_{x^k \otimes W_k, 0}(\mathbb{T}^d)$, consider the mesh functions defined on the discrete torus $f_N, g_N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ given by

$$f_N(x) = N^d \int_{Q_N(x)} f(y) dy, \quad \text{and} \quad (22)$$

$$g_N(x) = \frac{N^{d-1}}{W_k(x_k + 1/N) - W_k(x_k)} \int_{Q_N(x)} g(y) d(y^k \otimes W_k). \quad (23)$$

For a generalized function $f \in H_{1, W}^{-1}(\mathbb{T}^d)$ in the form (5), $f = f_0 - \sum_{k=1}^d \partial_{x_k} f_k$, where $f_0 \in L^2(\mathbb{T}^d)$, and $f_k \in L^2_{x^k \otimes W_k, 0}(\mathbb{T}^d)$, we consider

$$f_N(x) = f_{0, N}(x) - \sum_{k=1}^d \partial_{x_k}^N f_{k, N}(x), \quad (24)$$

where $x \in \frac{1}{N}\mathbb{T}_N^d$, $f_{0,N}$ is the discretization given by (22), and $f_{k,N}$ ($k = 1, \dots, d$) is the discretization given by (23). The function f_N clearly belongs to $H_W^{-1}(\mathbb{T}_N^d)$.

The following lemma shows that this procedure yields a good approximation for f .

Lemma 4.9. *Let $f \in L^2(\mathbb{T}^d)$, and $g \in L_{x^k \otimes W_k}^2(\mathbb{T}^d)$ for some $k = 1, 2, \dots, d$. Let f_N and g_N be the mesh functions defined in equations (22) and (23). Let \tilde{f}_N and \tilde{g}_N be the piecewise-constant interpolation for f_N and g_N . Then,*

$$\|f - \tilde{f}_N\| \rightarrow 0,$$

and

$$\|g - \tilde{g}_N\|_{x^k \otimes W_k} \rightarrow 0,$$

as $N \rightarrow \infty$. In particular, if $F \in H_W^{-1}(\mathbb{T}^d)$, then $F_N \rightarrow F$ strongly in $H_W^{-1}(\mathbb{T}^d)$.

Proof. We will prove the first assertion. The proof of the second one is analogous. Recall the notation used in the previous lemmas. Since the continuous functions are dense in $L^2(\mathbb{T}^d)$ (and also in $L_{x^j \otimes W_j}^2(\mathbb{T}^d)$) it is enough to consider f continuous. Thus,

$$\begin{aligned} \|f - \tilde{f}_N\|^2 &= \int_{\mathbb{T}^d} \left(f(y) - \tilde{f}_N(y) \right)^2 dy = \sum_{k=1}^{N^d} \int_{Q_N(x^k)} \left(f(y) - N^d \int_{Q_N(x^k)} f(z) dz \right)^2 dy = \\ &= \sum_{k=1}^{N^d} \int_{Q_N(x^k)} \left(N^d \int_{Q_N(x^k)} [f(y) - f(z)] dz \right)^2 dy \leq \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} N^d \int_{Q_N(x)} \int_{Q_N(x)} [f(y) - f(z)]^2 dz dy, \end{aligned}$$

where the last inequality follows from Hölder's inequality. Thus, the previous expression is bounded above by

$$\begin{aligned} &\sum_{x \in \frac{1}{N}\mathbb{T}_N^d} N^d \int_{\|\eta\|_\infty \leq 1/N} d\eta \int_{Q_N(x)} [f(y + \eta) - f(y)]^2 dy \\ &\leq 2^d \sum_{x \in \frac{1}{N}\mathbb{T}_N^d} \sup_{\substack{\eta; |\eta_i| < 1/N \\ i=1, \dots, d}} \int_{Q_N(x)} [f(y + \eta) - f(y)]^2 dy. \end{aligned}$$

To conclude, note that by compactness of \mathbb{T}^d the continuous function f is, in fact, uniformly continuous. So, for a fixed $\epsilon > 0$, there exists N_0 such that $\|\eta\|_\infty < 1/N_0$ implies that $|f(y + \eta) - f(y)| < \epsilon^{1/2}/2^d$. Therefore, the previous expression is bounded above by ϵ , and the proof of the Lemma follows. \square

5. Homogenization

Our main goal in this Section is to prove the convergence of energies, namely, Proposition 5.7, and also the Homogenization Theorem, namely, Theorem 5.9. This last result is presented with fairly general hypotheses on the matrices A^N . In Proposition 5.10 we provide an example of a very large class of functions that admit homogenization and in subsection 5.3 we consider a scenario in which A^N represents a random environment. We begin with some definitions and auxiliary results.

5.1. Definitions and auxiliary results

We now focus on the analysis of the asymptotic behavior of the sequence (u_N) given by the solutions of the equations

$$\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f^N, \quad \lambda \geq 0, \quad (25)$$

where $f^N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ are fixed functions, $\nabla^N = (\partial_{x_1}^N, \dots, \partial_{x_d}^N)$ and $\nabla_W^N = (\partial_{W_1}^N, \dots, \partial_{W_d}^N)$ are the difference operators and, $A^N = \text{diag}\{a_1^N, \dots, a_d^N\}$ is a sequence of diagonal matrices of order d satisfying the ellipticity condition (11).

The continuous counterpart of the theory developed in subsection 4.1 can be found in [12]. More precisely, one can find results on existence, uniqueness and boundedness of weak solutions of the problem

$$\lambda u - \nabla A \nabla_W u = f, \quad \lambda \geq 0. \quad (26)$$

We say that the sequence of matrices $A^N = \text{diag}\{a_1^N, \dots, a_d^N\}$ *H-converges* to the matrix $A = \text{diag}\{a_1, \dots, a_d\}$, denoted by $A^N \xrightarrow{H} A$, if for every sequence f^N of functionals on $H_{1,W}(\mathbb{T}_N^d)$ and $f \in H_W^{-1}(\mathbb{T}^d)$, such that $f^N \rightarrow f$ as $N \rightarrow \infty$ strongly in $H_W^{-1}(\mathbb{T}^d)$, we have

- $u_N \rightarrow u$ weakly in $H_{1,W}(\mathbb{T}^d)$ as $N \rightarrow \infty$,
- $a_k^N \partial_{W_k}^N u_N \rightarrow a_k \partial_{W_k} u$ weakly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ for each $k = 1, \dots, d$,

where $u_N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ is the solution of the (25) and $u \in H_{1,W}(\mathbb{T}^d)$ is the unique weak solution of the (26). In this case, we say that the diagonal matrix A is a *homogenization* of the sequence of the random matrices A^N . We also say that the operator $\nabla A \nabla_W$ is a *homogenization* of the sequence of random operators $\nabla^N A^N \nabla_W^N$.

Next, we present an estimate that will be useful in this Section.

Lemma 5.1. *Let $f_N \rightarrow f$ as $N \rightarrow \infty$ weakly in $H_W^{-1}(\mathbb{T}^d)$. Then, if $u_N : \frac{1}{N} \mathbb{T}_N^d \rightarrow \mathbb{R}$ is the solution of (25), there exists $C \geq 0$, not depending on N , such that*

$$\|u_N\|_{1,W,N} \leq C.$$

Proof. Using Lemma 4.5, we obtain a constant C_1 , not depending on N such that

$$\|u_N\|_{1,W,N} \leq C_1 \|f_N\|_{-1,W,N}.$$

Note that $f_N \rightarrow f$ weakly in $H_W^{-1}(\mathbb{T}^d)$ means that $\tilde{f}_N \rightarrow f$ weakly in $H_W^{-1}(\mathbb{T}^d)$. Since $\|f_N\|_{-1,W,N} = \|\tilde{f}_N\|_{-1,W}$, and \tilde{f}_N is weakly convergent, it is bounded, and thus there exists a constant $C_2 \geq 0$, such that

$$\|f_N\|_{-1,W,N} = \|\tilde{f}_N\|_{-1,W} \leq C_2.$$

Thus, for all $N \geq 1$,

$$\|u_N\|_{1,W,N} \leq C_1 C_2 = C. \quad \square$$

Let us prove an auxiliary lemma that will be needed in the proof of the convergence of energies of a sequence of homogenized matrices (Proposition 5.7).

Lemma 5.2. *Let X be a Banach space and X^* its dual space. If $f_N \in X^*$ is such that $f_N \rightarrow f \in X^*$, and $u_N \in X$ is such that $u_N \rightarrow u \in X$ weakly. Then,*

$$f_N(u_N) \rightarrow f(u),$$

as $N \rightarrow \infty$.

Proof. Since u_N converges weakly in X , it forms a bounded sequence. That is, there exists $C \geq 0$ such that, for each $N \geq 1$, $\|u_N\| \leq C$. We also have

$$|f_N(u_N) - f(u_N)| \leq \|f_N - f\|_{X^*} \|u_N\|_X \leq C \|f_N - f\|_{X^*},$$

which tends to zero as $N \rightarrow \infty$. On the other hand, from weak convergence, $f(u_N) \rightarrow f(u)$ as $N \rightarrow \infty$. Therefore, $f_N(u_N) \rightarrow f(u)$ as $N \rightarrow \infty$. This concludes the proof. \square

The following lemma will be useful in proving the compensated compactness theorem (Theorem 5.4) and also in proving the main result of this paper (Theorem 5.9).

Lemma 5.3. *Fix k in $\{1, \dots, d\}$ and let $v_{k,N} \rightarrow v_k$ as $N \rightarrow \infty$ weakly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. If there exists $C > 0$ such that $|v_{k,N}(x)| < C$ (resp. $v_{k,N} > C$) for every $N \in \mathbb{N}$ and $x \in \mathbb{T}^d$, then $|v_k(x)| \leq C$ (resp. $v_k(x) \geq C$) for $d(x^k \otimes W_k)$ -almost every x in \mathbb{T}^d .*

Proof. Let us prove the statement $|v_k(x)| \leq C$, the other being analogous. If this is statement is false, there exists a measurable set A , with positive $d(x^k \otimes W_k)$ -measure, such that $v_k(x) > C$ or $v_k(x) < -C$ in A . Let $B_1, B_2 \subset A$ be such that $v_k(x) > C$ in B_1 and $v_k(x) < -C$ in B_2 . Suppose, without loss of generality, that $d(x^k \otimes W_k)(B_1) > 0$. Thus, denoting by $1_{B_1}(x)$ the indicator function of the set B_1 , we have that $1_{B_1} \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, and

$$\int_{\mathbb{T}^d} v_{k,N} 1_{B_1} d(x^k \otimes W_k) \xrightarrow{N \rightarrow \infty} \int_{\mathbb{T}^d} v_k 1_{B_1} d(x^k \otimes W_k).$$

On one hand, for every $N \in \mathbb{N}$,

$$\int_{\mathbb{T}^d} v_{k,N} 1_{B_1} d(x^k \otimes W_k) \leq C d(x^k \otimes W_k)(B_1),$$

and on the other hand

$$\int_{\mathbb{T}^d} v_k 1_{B_1} d(x^k \otimes W_k) > C d(x^k \otimes W_k)(B_1).$$

Since $d(x^k \otimes W_k)(B_1) > 0$, we have a contradiction. Therefore, the statement is true. \square

We will now prove a simple version of the compensated compactness Theorem.

Lemma 5.4 (Compensated compactness in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$). For $k = 1, \dots, d$, let $(g_{k,N})$ and $(w_{k,N})$ be sequences of functions in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ such that one of these sequences is uniformly bounded in x and N . That is, there exists $C > 0$ such that

$$\forall N \in \mathbb{N}, \forall x \in \mathbb{T}^d, \quad |g_{k,N}(x)| < C$$

or

$$\forall N \in \mathbb{N}, \forall x \in \mathbb{T}^d, \quad |w_{k,N}(x)| < C.$$

Let, also,

$$g_{k,N} \rightarrow g_k, \quad \text{strongly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d) \quad \text{and} \quad w_{k,N} \rightarrow w_k, \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d),$$

where $g_k, w_k \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. Then, $g_k w_k \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, and

$$g_{k,N} w_{k,N} \rightarrow g_k w_k \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d).$$

Proof. Since $g_{k,N}$ converges strongly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, it also converges weakly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. Thus, from Lemma 5.3, we have that g_k or w_k is uniformly bounded in x , which implies that $g_k w_k \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$.

Let $\phi \in \mathcal{D}_W(\mathbb{T}^d)$, then

$$\int_{\mathbb{T}^d} g_{k,N} w_{k,N} \phi d(x^k \otimes W_k) = \int_{\mathbb{T}^d} (g_{k,N} - g_k) w_{k,N} \phi d(x^k \otimes W_k) + \int_{\mathbb{T}^d} g_k w_{k,N} \phi d(x^k \otimes W_k).$$

Let us deal with each term in the right-hand side of the previous equation. By Cauchy–Schwarz Inequality

$$\left| \int_{\mathbb{T}^d} (g_{k,N} - g_k) w_{k,N} \phi d(x^k \otimes W_k) \right| \leq \|g_{k,N} - g_k\|_{x^k \otimes W_k} \|w_{k,N} \phi\|_{x^k \otimes W_k}.$$

From Lemma 3.5, ϕ is bounded, and since $(w_{k,N})$ is a weakly convergent sequence, its norm is uniformly bounded. Therefore, the previous equation tends to zero as $N \rightarrow \infty$.

We now deal with the other term. Begin by recalling that ϕ is bounded. Thus, $q_k \phi \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, since \mathbb{T}^d is a set of finite $d(x^k \otimes W_k)$ -measure. Therefore, the weak convergence of $w_{k,N}$ to w_k implies that

$$\int_{\mathbb{T}^d} g_k w_{k,N} \phi d(x^k \otimes W_k) = \int_{\mathbb{T}^d} w_{k,N} (g_k \phi) d(x^k \otimes W_k) \rightarrow \int_{\mathbb{T}^d} w_k g_k \phi d(x^k \otimes W_k).$$

Now, notice that since $(g_{k,N})$ is strongly convergent in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ and $(w_{k,N})$ is weakly convergent in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, both sequences are uniformly bounded in the $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ -norm. Since one of these sequences is also uniformly bounded in x and in N , we have that there exists $M > 0$ such that for every $N \in \mathbb{N}$

$$\|g_{k,N} w_{k,N}\|_{x^k \otimes W_k} \leq M.$$

Finally, let $\varphi \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. Note that, by Proposition 3.3, $\mathfrak{D}_W(\mathbb{T}^d)$ is dense in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ we may find a sequence $\phi_m \in \mathfrak{D}_W(\mathbb{T}^d)$ such that $\phi_m \rightarrow \varphi$ in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$.

Therefore, fix $\varepsilon > 0$ and choose $m \in \mathbb{N}$ such that

$$\|\phi_m - \varphi\|_{x^k \otimes W_k} \leq \varepsilon \cdot \min \left\{ \frac{1}{3(\|g_k w_k\|_{x^k \otimes W_k} + 1)}, \frac{1}{3M} \right\}.$$

Now, take $N_0 \in \mathbb{N}$ such that $N \geq N_0$ implies

$$\left| \int_{\mathbb{T}^d} g_{k,N} w_{k,N} \phi_m d(x^k \otimes W_k) - \int_{\mathbb{T}^d} g_k w_k \phi_m d(x^k \otimes W_k) \right| < \frac{\varepsilon}{3}.$$

Thus, if $N \geq N_0$,

$$\begin{aligned} & \left| \int_{\mathbb{T}^d} g_{k,N} w_{k,N} \varphi d(x^k \otimes W_k) - \int_{\mathbb{T}^d} g_k w_k \varphi d(x^k \otimes W_k) \right| \\ & \leq \left| \int_{\mathbb{T}^d} g_{k,N} w_{k,N} (\varphi - \phi_m) d(x^k \otimes W_k) \right| + \left| \int_{\mathbb{T}^d} (g_{k,N} w_{k,N} - g_k w_k) \phi_m d(x^k \otimes W_k) \right| \\ & \quad + \left| \int_{\mathbb{T}^d} g_k w_k (\phi_m - \varphi) d(x^k \otimes W_k) \right| \\ & \leq \|g_{k,N} w_{k,N}\|_{x^k \otimes W_k} \|\varphi - \phi_m\|_{x^k \otimes W_k} + \frac{\varepsilon}{3} + \|g_k w_k\|_{x^k \otimes W_k} \|\phi_m - \varphi\|_{x^k \otimes W_k} < \varepsilon. \end{aligned}$$

This concludes the proof. \square

We conclude this subsection with a version of Compensated Compactness Theorem for discrete approximations.

Corollary 5.5 (*Compensated compactness for discrete approximations*). *Let $k \in \{1, \dots, d\}$, and let $(q_{k,N})$ and $(v_{k,N})$ be sequences of functions on $L^2_{x^k \otimes W_k}(\mathbb{T}^d_N)$ such that one of these sequences is uniformly bounded in x and N , and also*

$$q_{k,N} \rightarrow q_k, \quad \text{strongly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d) \quad \text{and} \quad v_{k,N} \rightarrow v_k, \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d),$$

where $q_k, v_k \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. Then, $q_k v_k \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ and

$$q_{k,N} v_{k,N} \rightarrow q_k v_k \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d).$$

Proof. Let $w_{k,N} = \tilde{v}_{k,N}$ and $g_{k,N} = \tilde{q}_{k,N}$. Thus, by assumption, $w_{k,N}$ converges weakly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ to v_k , and $g_{k,N}$ converges strongly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$ to q_k . Therefore, we may apply Lemma 5.4 to conclude that

$$\widetilde{v_{k,N} q_{k,N}} = \tilde{v}_{k,N} \tilde{q}_{k,N} = w_{k,N} g_{k,N} \rightarrow v_k q_k \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d). \quad \square$$

Remark 5.6. One should notice that Lemma 5.4 is, indeed, a version of the Compensated Compactness Theorem. In fact, the classical assumptions would be $g_{k,N} \rightarrow g_k$ and $w_{k,N} \rightarrow w_k$ weakly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, and $\partial_{x_k} g_{k,N} \rightarrow h$ strongly in $H^{-1}_W(\mathbb{T}^d)$, where $\partial_{x_k} g_{k,N}$ should be understood as in Lemma 2.1. However, in our setup, the functional induced by $g_{k,N}$ coincides with $-\partial_{x_k} g_{k,N}$, and thus, since $\|\partial_x g_{k,N}\|_{-1,W} = \|g_{k,N}\|_{x^k \otimes W_k}$, we recover strong convergence for $g_{k,N}$. The reason why the functional induced by $g_{k,N}$ is $-\partial_{x_k} g_{k,N}$ instead of the standard functional is because the standard functional given by

$$\phi \mapsto \int_{\mathbb{T}^d} g_{k,N} \phi dx$$

is not well-defined for $g_{k,N} \in L^2_{x^k \otimes W_k}(\mathbb{T}^d)$. In fact, $g_{k,N}$ may not belong to $L^2(\mathbb{T}^d)$.

5.2. Main results

In this subsection we will state and prove the homogenization of the difference operators introduced in the previous Section.

We begin by showing, in the following proposition, that even though the H -convergence only requires weak convergence in its definition, it yields a convergence in a strong sense (convergence in the L^2 -norm for the piecewise-constant interpolation).

Proposition 5.7. *Let $A^N \xrightarrow{H} A$, as $N \rightarrow \infty$, with u_N being the solution of (25), where $f \in H^{-1}_W(\mathbb{T}^d)$ is fixed, $f^N \rightarrow f$ strongly in $H^{-1}_W(\mathbb{T}^d)$ and, u is the weak solution of (26). Then, the following limit relations hold true:*

$$u_N \rightarrow u \text{ in } L^2(\mathbb{T}^d),$$

$$\begin{aligned} & \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N^2\left(\frac{x}{N}\right) \xrightarrow{N \rightarrow \infty} \int_{\mathbb{T}^d} u^2(x) dx, \quad \text{and} \\ & \frac{1}{N^{d-1}} \sum_{k=1}^d \sum_{x \in \mathbb{T}_N^d} a_{kk}^N\left(\frac{x}{N}\right) (\partial_{W_k}^N u_N\left(\frac{x}{N}\right))^2 \left[W_k\left(\frac{x_k+1}{N}\right) - W_k\left(\frac{x_k}{N}\right) \right] \\ & \xrightarrow{N \rightarrow \infty} \sum_{k=1}^d \int_{\mathbb{T}^d} a_{kk}(x) (\partial_{W_k} u(x))^2 d(x^k \otimes W_k). \end{aligned}$$

Proof. We begin by proving that

$$f^N(u_N) \rightarrow f(u), \quad \text{as} \quad N \rightarrow \infty. \quad (27)$$

In order to apply Lemma 4.6, we need to obtain the required bound. By Lemma 5.1, the sequence u_N is uniformly bounded with respect to the norm $\|\cdot\|_{1,W}$. In particular, there exist constants $C_1, C_2^k > 0, k = 1, \dots, d$, such that, for all $N \geq 1$, we have

$$\frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N^2(x/N) \leq C_1, \quad (28)$$

and

$$\frac{1}{N^{d-1}} \sum_{x \in \mathbb{T}_N^d} (\partial_{W_k}^N u_N(x/N))^2 (W_k((x_k+1)/N) - W_k(x_k/N)) \leq C_2^k. \quad (29)$$

Now, observe that from Lemma 4.4, there exist functions $f_{0,N}, \dots, f_{d,N}$ such that

$$f_N(u_N) = \langle f_{0,N}, u_N \rangle_N + \sum_{k=1}^d \langle f_{k,N}, \partial_{W_k}^N u_N \rangle_{W_k,N}.$$

This motivates us to define the functionals $g_{i,N} \in H_W^{-1}(\mathbb{T}_N^d)$, $i = 0, 1, \dots, d$, by $g_{0,N}(v) = \langle f_{0,N}, v \rangle_N$, and $g_{k,N}(v) = \langle f_{k,N}, \partial_{W_k}^N v \rangle_{W_k,N}$, for $k = 1, \dots, d$. Note that

$$g_{0,N}(v) = \langle \tilde{f}_{0,N}, \tilde{v} \rangle, \quad \text{and} \quad g_{k,N}(v) = \langle \tilde{f}_{k,N}, \widehat{\partial_{W_k}^N v} \rangle_{x^k \otimes W_k}.$$

We have, by hypothesis, that $f_N \rightarrow f$ strongly, which means

$$\|\tilde{f}_N - f\|_{-1,W}^2 = \|\tilde{f}_{N,0} - f_0\|^2 + \sum_{k=1}^d \|\tilde{f}_{k,N} - f_k\|_{x^k \otimes W_k}^2 \rightarrow 0,$$

as $N \rightarrow \infty$. Thus, $g_{0,N} \rightarrow f_0$ strongly in $L^2(\mathbb{T}^d)$, $g_{k,N} \rightarrow f_k$ strongly in $L_{x^k \otimes W_k}^2(\mathbb{T}^d)$.

Observe that $u_N \rightarrow u$ weakly in $H_W^1(\mathbb{T}^d)$ means that $u_N^* \rightarrow u$ weakly in $H_W^1(\mathbb{T}^d)$, which in turn implies that

$$u_N^* \rightarrow u, \quad \text{weakly in } L^2(\mathbb{T}^d), \quad (30)$$

and

$$\partial_{W_k} u_N^* \rightarrow \partial_{W_k} u, \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d).$$

Nevertheless, by equation (17), we have

$$\partial_{W_k} u_N^* = \left(\partial_{W_k}^N u_N \right)^{(k)},$$

and thus

$$\left(\partial_{W_k}^N u_N \right)^{(k)} \rightarrow \partial_{W_k} u, \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d). \quad (31)$$

Using Lemma 4.6, equation (30) implies that

$$\tilde{u}_N \rightarrow u, \quad \text{weakly in } L^2(\mathbb{T}^d), \quad (32)$$

and equation (31) implies that

$$\widetilde{\partial_{W_k}^N u_N} \rightarrow \partial_{W_k} u, \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d). \quad (33)$$

Now, since $\tilde{g}_{0,N} \rightarrow f_0$ strongly in $L^2(\mathbb{T}^d)$ and $\tilde{g}_{k,N} \rightarrow f_k$ strongly in $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$, we may apply Lemma 5.2 to equations (32) and (33) to obtain that

$$\tilde{g}_{0,N}(\tilde{u}_N) \rightarrow f_0(u), \quad \text{as } N \rightarrow \infty \text{ and,} \quad (34)$$

$$\tilde{g}_{k,N}(\widetilde{\partial_{W_k}^N u_N}) \rightarrow f_k(\partial_{W_k} u), \quad \text{as } N \rightarrow \infty. \quad (35)$$

Note that

$$f(u) = f_0(u) + \sum_{k=1}^d f_k(\partial_{W_k}(u)) \quad \text{and}$$

$$f_N(u_N) = g_{0,N}(u_N) + \sum_{k=1}^d g_{k,N}(\partial_{W_k}^N(u_N)) = \tilde{g}_{0,N}(\tilde{u}_N) + \sum_{k=1}^d \tilde{g}_{k,N}(\widetilde{\partial_{W_k}^N u_N}).$$

Thus, from (34) and (35), $f_N(u_N)$ converges to $f(u)$. Since, by assumption, $A^N \xrightarrow{H} A$ and u is the weak solution of (26). Thus, we have,

$$f(u) = \lambda \int_{\mathbb{T}^d} u^2 dx + \sum_{k=1}^d \int_{\mathbb{T}^d} a_{kk}(\partial_{W_k} u)^2 d(x^k \otimes W_k).$$

Note also that

$$\begin{aligned}
 f^N(u_N) &= \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} (\lambda u_N(x/N) - \nabla^N A^N \nabla_W^N u_N(x/N)) u_N(x/N) \\
 &= \frac{\lambda}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N^2(x/N) - \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N(x/N) \nabla^N A^N \nabla_W^N u_N(x/N),
 \end{aligned}$$

which, after a summation by parts in the above expressions, and using that $f_N(u_N) \rightarrow f(u)$, we obtain

$$\begin{aligned}
 \frac{\lambda}{N^d} \sum_{x \in \mathbb{T}_N^d} u_N^2(x/N) + \frac{1}{N^{d-1}} \sum_{k=1}^d \sum_{x \in \mathbb{T}_N^d} a_{kk}^N (\partial_{W_k}^N u_N(x/N))^2 [W_k((x_k+1)/N) - W_k(x_k/N)] \\
 \xrightarrow{N \rightarrow \infty} \lambda \int_{\mathbb{T}^d} u^2 dx + \sum_{k=1}^d \int_{\mathbb{T}^d} a_{kk} (\partial_{W_k} u)^2 d(x^k \otimes W_k). \quad (36)
 \end{aligned}$$

Suppose that u_N does not converge to u in $L^2(\mathbb{T}^d)$. That is, there exist $\epsilon > 0$ and a subsequence (u_{N_j}) such that

$$\|\tilde{u}_{N_j} - u\| > \epsilon,$$

for all j . By Lemma 4.8, we have that there exist $v \in L^2(\mathbb{T}^d)$ and a further subsequence (also denoted by u_{N_j}) such that

$$u_{N_j}^* \xrightarrow{j \rightarrow \infty} v, \quad \text{in } L^2(\mathbb{T}^d).$$

Using Lemma 4.7, we further obtain that

$$\tilde{u}_{N_j} \xrightarrow{j \rightarrow \infty} v, \quad \text{in } L^2(\mathbb{T}^d).$$

This implies that

$$\tilde{u}_{N_j} \rightarrow v, \quad \text{weakly in } L^2(\mathbb{T}^d),$$

but this is a contradiction. Indeed, from H convergence of A_N to A , we have that $u_N \rightarrow u$ weakly in $H_{1,W}(\mathbb{T}^d)$, which means that $u_N^* \rightarrow u$ in $H_{1,W}(\mathbb{T}^d)$. Thus $u_N^* \rightarrow u$ weakly in $L^2(\mathbb{T}^d)$. Using Lemma 4.6, this implies that $\tilde{u}_N \rightarrow u$ weakly in $L^2(\mathbb{T}^d)$. In particular,

$$u_{N_j} \rightarrow u, \quad \text{weakly in } L^2(\mathbb{T}^d),$$

and at the same time $\|v - u\| \geq \epsilon$. Therefore, $u_N \rightarrow u$ in $L^2(\mathbb{T}^d)$. The proof thus follows from expression (36). \square

Corollary 5.8. Let $u \in \mathfrak{D}_W(\mathbb{T}^d)$ and $u_N : \frac{1}{N}\mathbb{T}_N^d \rightarrow \mathbb{R}$ such that $\lim u_N = u$ in $L^2(\mathbb{T}^d)$. Then,

$$\|u - u_N\| \longrightarrow 0.$$

Proof. We have that $u_N \rightarrow u$ in $L^2(\mathbb{T}^d)$ implies that

$$\|\widetilde{u}_N - u\| \longrightarrow 0, \quad (37)$$

as $N \rightarrow \infty$. On the other hand, from the definition of the set $\mathcal{Q}_N(x)$ in subsection 4.2 and the fact that the functions in $\mathfrak{D}_W(\mathbb{T}^d)$ are right-continuous (see Remark 3.4), we have that

$$\widetilde{u|_{\mathbb{T}_N^d}} \longrightarrow u$$

pointwise as $N \rightarrow \infty$. From Lemma 3.5, u is bounded, and thus $\widetilde{u|_{\mathbb{T}_N^d}}$ is bounded, and therefore integrable (\mathbb{T}^d has finite Lebesgue measure). Thus, we can use the Dominated Convergence Theorem, to conclude that

$$\|\widetilde{u|_{\mathbb{T}_N^d}} - u\| \longrightarrow 0, \quad (38)$$

as $N \rightarrow \infty$. Therefore, using (37) and (38), we obtain

$$\begin{aligned} \|u_N - u\| &= \|\widetilde{u}_N - \widetilde{u|_{\mathbb{T}_N^d}}\| \\ &\leq \|\widetilde{u}_N - u\| + \|\widetilde{u|_{\mathbb{T}_N^d}} - u\| \longrightarrow 0, \end{aligned}$$

as N goes to ∞ . This concludes the proof. \square

We will now state and prove the main result of this paper.

Theorem 5.9. Let $A^N = \text{diag}\{a_1^N, \dots, a_d^N\}$ be a sequence of diagonal matrices and $\theta > 0$, such that $\theta^{-1} \leq a_k^N \leq \theta$. If

$$1/a_k^N \rightarrow b_k \text{ weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d),$$

then, A^N admits a homogenization, with homogenized matrix $A = \text{diag}\{1/b_1, \dots, 1/b_d\}$.

Proof of Theorem 5.9. Fix $f \in H_W^{-1}(\mathbb{T}^d)$, and consider the problem

$$\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f_N, \quad (39)$$

where f_N is the discretization of f obtained by using equation (24). From Lemma 4.9, $f_N \rightarrow f$ strongly in $H_W^{-1}(\mathbb{T}^d)$. Using Lemma 5.1, we obtain that there exists a unique weak solution u_N of the problem (39) such that its $H_{1,W}(\mathbb{T}_N^d)$ -norm is uniformly bounded in N . That is, there exists a constant $C > 0$ such that

$$\|u_N\|_{1,W,N} \leq C.$$

From Lemma 4.8, there exists a convergent subsequence of u_N (which we will also denote by u_N) such that

$$u_N \rightarrow u, \quad \text{weakly in } H_{1,W}(\mathbb{T}^d).$$

In particular,

$$\partial_{W_k}^N u_N \xrightarrow{N \rightarrow \infty} \partial_{W_k} u \quad \text{weakly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d). \quad (40)$$

Applying (39) to u_N , we obtain

$$\lambda \|u_N\|_N^2 + \sum_{k=1}^d \|a_{kk}^N \partial_{W_k}^N u_N\|_{W_k,N}^2 = f_N(u_N) \leq \|f_N\|_{-1,W,N} \|u_N\|_{1,W,N}.$$

Thus using Lemma 5.1 we have

$$\|a_k^N \partial_{W_k}^N u_N\|_{W_k,N}^2 \leq \|f_N\|_{-1,W,N} \cdot C, \quad \text{for each } k = 1, \dots, d.$$

By using the same argument (from weak convergence of the functionals) we can find a constant $C_1 \geq 0$ such that $\|f_N\|_{-1,W,N} \leq C_1$. Therefore,

$$\|a_k^N \partial_{W_k}^N u_N\|_{W_k,N} \leq C_1 C.$$

This, in turn, implies that

$$\|a_k^N \partial_{W_k}^N u_N\|_{x^k \otimes W_k} = \|a_k^N \partial_{W_k}^N u_N\|_{W_k,N}$$

is uniformly bounded in N . Thus, since $L_{x^k \otimes W_k}^2(\mathbb{T}^d)$ is a separable Hilbert space, we can find a further subsequence (also denoted by u_N), such that

$$a_k^N \partial_{W_k}^N u_N \rightarrow v_{0,k} \quad \text{weakly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d), \quad (41)$$

as $N \rightarrow \infty$, $v_{0,k}$ being some function in $L_{x^k \otimes W_k}^2(\mathbb{T}^d)$.

Since (u_N) is uniformly bounded in the Sobolev-norm and $L^2(\mathbb{T}^d)$ is precompact in this space we have that

$$u_N \rightarrow u \quad \text{strongly in } L^2(\mathbb{T}^d).$$

In particular,

$$u_N \rightarrow u \quad \text{strongly in } H_W^{-1}(\mathbb{T}^d).$$

On the other hand, $(\lambda u_N - \nabla^N A^N \nabla_W^N u_N)$ converges strongly to f in $H_W^{-1}(\mathbb{T}^d)$. Therefore,

$$\nabla^N A^N \nabla_{W_k}^N u_N \rightarrow v_0 \quad \text{strongly in } H_W^{-1}(\mathbb{T}^d).$$

From the very definition of the functional $\nabla^N A^N \nabla_{W_k}^N u_N$, the previous convergence means that for each k ,

$$a_k^N \partial_{W_k}^N u_N \rightarrow v_{0,k} \quad \text{strongly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d).$$

Since $\theta^{-1} < a_k^N(x) < \theta$, we have that $1/a_k^N$ is uniformly bounded, and $1/a_k^N > \theta$. Therefore, from Lemma 5.3,

$$b_k \geq \theta > 0.$$

From the Compensated Compactness Theorem (take $q_k^N = a_k^N \partial_{W_k}^N u_N$ and $v_k^N = 1/a_k^N$ in Corollary 5.5, and notice that (v_k^N) is uniformly bounded in x and N):

$$\frac{1}{a_k^N} a_k^N \partial_{W_k}^N u_N \rightarrow b_k v_{0,k}, \quad \text{weakly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d).$$

On the other hand, by (40),

$$\frac{1}{a_k^N} a_k^N \partial_{W_k}^N u_N = \partial_{W_k}^N u_N \rightarrow \partial_{W_k} u, \quad \text{weakly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d).$$

From uniqueness of the weak limit, we have that $\partial_{W_k} u = b_k v_{0,k}$. Since $b_k \neq 0$, we have that

$$v_{0,k} = \frac{1}{b_k} \partial_{W_k} u.$$

Thus, we can summarize our findings:

$$\begin{aligned} u_N &\rightarrow u \quad \text{strongly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d), \\ \partial_{W_k} u_N &\rightarrow \partial_{W_k} u \quad \text{weakly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d), \quad \text{and} \\ a_k^N \partial_{W_k} u_N &\rightarrow \frac{1}{b_k} \partial_{W_k} u \quad \text{strongly in } L_{x^k \otimes W_k}^2(\mathbb{T}^d). \end{aligned}$$

Therefore, u solves the problem

$$\lambda u - \nabla A \nabla_W u = f,$$

where $A = \text{diag}\{1/b_1, \dots, 1/b_d\}$.

To conclude the proof it remains to be shown that we can pass from the subsequence to the sequence. This follows from uniqueness of weak solutions of the problem (26), see [12, Proposition 3.4]. The fact that any converging subsequence is a solution of the same problem, and the fact that u_N is uniformly bounded in the Sobolev norm, thus we can find a convergent subsequence (thus a sequence that do not converge to the solution, must converge to somewhere else, since the limit point is also a solution, uniqueness shows the result). \square

We will now provide an example of a very large class of matrix functions that admits homogenization. Recall the definition of the space \mathbb{M}_W given below Remark 3.1: $\mathbb{M}_W \subset (C_W(\mathbb{T}^d))^d$ is the space of functions $a = (a_1, \dots, a_d)$ such that for every $k = 1, \dots, d$, $\partial_{x_k} a_k$ exists.

Proposition 5.10. *Let $A = \text{diag}\{a_1, \dots, a_d\} \in \mathbb{M}_W$ be a diagonal matrix such that $a_k \geq \theta^{-1}$, for some $\theta > 0$, and the discretization $A^N = \text{diag}\{a_1^N, \dots, a_d^N\}$ be the sequence of diagonal matrices obtained from applying (23) to the entries of A . Then, the sequence (A^N) admits a homogenization, with homogenized matrix A .*

Proof. It is clear that $0 < \theta^{-1} \leq a_k^N$. Further, it is clear that the right-continuity implies the pointwise convergence of \tilde{a}_k^N to a_k . Finally, from Lemma 3.5, the functions a_k are bounded, and thus the sequences a_k^N are bounded. From the Dominated Convergence Theorem

$$a_k^N \rightarrow a_k \text{ strongly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d).$$

On the other hand, we also have the pointwise convergence of $\widetilde{1/a_k^N}$ to $1/a_k$, and the bound $1/a_k^N \leq \theta$ implies that

$$1/a_k^N \rightarrow 1/a_k \text{ strongly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d).$$

Therefore, the result follows from Theorem 5.9. \square

5.3. Homogenization of Random difference operators

In this subsection we consider the homogenization problem when the matrix A^N represents a random environment. More precisely, we focus on the analysis of the asymptotic behavior of the sequence (u_N) given by the solutions of the equations

$$\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f^N,$$

where f^N are fixed functions defined on $\frac{1}{N}\mathbb{T}^d$, $\nabla^N = (\partial_{x_1}^N, \dots, \partial_{x_d}^N)$ and $\nabla_W^N = (\partial_{W_1}^N, \dots, \partial_{W_d}^N)$ are the difference operators and, the random diagonal matrices $A^N = \text{diag}\{a_1, \dots, a_d\}$ of order d represent the statistically homogeneous rapidly oscillating coefficients. Therefore these equations are driven by the random difference operators $\nabla^N A^N \nabla_W^N$, and to fully understand them, we need to understand the random matrices A^N . Thus, let $(\Omega, \mathcal{F}, \mu)$ be a standard probability space and $\{T_x : \Omega \rightarrow \Omega; x \in \mathbb{Z}^d\}$ be a group of \mathcal{F} -measurable and ergodic transformations which preserve the measure μ :

- $T_x : \Omega \rightarrow \Omega$ is \mathcal{F} -measurable for all $x \in \mathbb{Z}^d$,
- $\mu(T_x \mathbf{A}) = \mu(\mathbf{A})$, for any $\mathbf{A} \in \mathcal{F}$ and $x \in \mathbb{Z}^d$,
- $T_0 = I$, $T_x \circ T_y = T_{x+y}$,
- if $f \in L^1(\Omega)$ is such that $f(T_x \omega) = f(\omega)$ for each $x \in \mathbb{Z}^d$, and almost every ω , then f is constant almost surely.

Note that the last condition implies that the group T_x is ergodic. We call the underlying probability space $(\Omega, \mathcal{F}, \mu)$ *random environment*, and a point $\omega \in \Omega$ a *realization* of the random environment.

Let us now introduce the vector-valued \mathcal{F} -measurable functions $\{b_k(\omega); k = 1, \dots, d\}$ such that there exists $\theta > 0$ with

$$\theta^{-1} \leq b_k(\omega) \leq \theta, \quad (42)$$

for all $\omega \in \Omega$ and $k = 1, \dots, d$. Define the *random* diagonal matrices B^N whose elements are given by

$$b_k^N(x) := b_k(T_{Nx}\omega), \quad x \in \frac{1}{N}\mathbb{T}_N^d, \quad k = 1, \dots, d. \quad (43)$$

Let us show some weak convergences associated to the random environment (b_{kk}^N) . First, note that by the Birkhoff's Ergodic Theorem, we have

$$b_k^N \longrightarrow E[b_k] \quad \text{and} \quad 1/b_k^N \longrightarrow E[1/b_k] \quad \text{weakly in } L^2(\mathbb{T}^d) \quad \text{almost surely,} \quad (44)$$

for $k = 1, \dots, d$. We will need a similar result for $L^2_{x^k \otimes W_k}(\mathbb{T}^d)$.

Denote by μ_{W_k} the measure induced by the function W_k . By Lebesgue decomposition, there exists a function g such that

$$\mu_{W_k} = g\lambda + \lambda^\perp,$$

where $g\lambda$ and λ^\perp are mutually singular measures and λ denotes the Lebesgue measure. Let $V_k \subset \mathbb{T}$ be the support of λ^\perp and $\mathfrak{V}^k = \mathbb{T} \times \dots \times \mathbb{T} \times V_k \times \mathbb{T} \times \dots \times \mathbb{T} \subset \mathbb{T}^d$, V_k in the k -th component.

Recall the definition of $Q_N(x)$ in (16) and define $a_k^N : \frac{1}{N}\mathbb{T}_N^d \rightarrow \mathbb{R}$ as

$$a_k^N(x) = \begin{cases} b_k^N(x) & \text{if } \mathfrak{V}^k \cap Q_N(x) = \emptyset \\ (E[1/b_k])^{-1} & \text{if } \mathfrak{V}^k \cap Q_N(x) \neq \emptyset \end{cases}. \quad (45)$$

Consider the problem

$$\lambda u_N - \nabla^N A^N \nabla_W^N u_N = f^N, \quad (46)$$

where $A^N = (a_{kk}^N)$. The main result of this subsection is the following.

Theorem 5.11. *Let A^N be a sequence random matrices, as defined previously in (46). Then, almost surely, $A^N(\omega, x)$ admits a homogenization, where the homogenized matrix A does not depend on the realization ω , and is constant in x .*

Proof. The proof follows from Lemma 5.12 below that ensures the hypothesis of the Theorem 5.9 are valid for this sequence. \square

Lemma 5.12. Let $a_k^N : \frac{1}{N}\mathbb{T}_N^d \rightarrow \mathbb{R}$ be as defined above. Then,

$$1/a_k^N \longrightarrow E[1/b_k] \quad \text{weakly in } L^2_{x^k \otimes W_k}(\mathbb{T}^d) \quad \text{almost surely.}$$

Proof. Following the notation introduced above, let

$$\mathfrak{V}_N^k = \bigcup_{\substack{x \in \frac{1}{N}\mathbb{T}_N^d \\ \mathfrak{V}^k \cap Q_N(x) \neq \emptyset}} Q_N(x).$$

We have that

$$1_{\mathfrak{V}_N^k} \rightarrow 1_{\mathfrak{V}^k} \quad \text{pointwise.}$$

Therefore,

$$1_{\mathfrak{V}_N^k} \rightarrow 1_{\mathfrak{V}^k} \quad \text{strongly in } L^2(\mathbb{T}^d). \quad (47)$$

Let $\phi \in \mathfrak{D}_W(\mathbb{T}^d)$ be fixed. By Lebesgue decomposition we have,

$$\int_{\mathbb{T}^d} \widetilde{1/a_k^N} \phi d(x^k \otimes W_k) = \int_{\mathbb{T}^d} \widetilde{1/a_k^N} \phi g dx + \int_{\mathbb{T}^d} \widetilde{1/a_k^N} \phi d(x^k \otimes \lambda^\perp).$$

Note that the support of the measure $d(x^k \otimes \lambda^\perp)$ is confined in the set \mathfrak{V}^k . Since $\mathfrak{V}^k \subset \mathfrak{V}_N^k$, we have that $\widetilde{1/a_k^N}$ is almost everywhere constant, namely $\widetilde{1/a_k^N} = E[1/b_k]$, with respect to the measure $d(x^k \otimes \lambda^\perp)$. Thus, the second integral in the right-hand side in the previous expression is equal to

$$\int_{\mathbb{T}^d} E[1/b_k] \phi d(x^k \otimes \lambda^\perp).$$

On the other hand, by equations (44) and (47), we have

$$1/a_k^N 1_{[\mathfrak{V}_N^k]^c} = 1/b_k^N 1_{[\mathfrak{V}_N^k]^c} \longrightarrow E[1/b_k] 1_{[\mathfrak{V}^k]^c} \quad \text{weakly in } L^2(\mathbb{T}^d) \quad \text{almost surely,} \quad (48)$$

where X^c denotes the complementary set of X .

So, the first integral in the right-hand side in the above expression is equal to

$$\int_{\mathbb{T}^d} \widetilde{1/a_k^N} 1_{\mathfrak{V}_N^k} \phi g dx + \int_{\mathbb{T}^d} \widetilde{1/a_k^N} 1_{[\mathfrak{V}_N^k]^c} \phi g dx = \int_{\mathbb{T}^d} E[1/b_k] 1_{\mathfrak{V}_N^k} \phi g dx + \int_{\mathbb{T}^d} \widetilde{1/b_k^N} 1_{[\mathfrak{V}_N^k]^c} \phi g dx.$$

From the previous convergence, (47) and (48), the right hand-side in the previous expression converges to

$$\int_{\mathbb{T}^d} E[1/b_k] 1_{\overline{\mathfrak{B}}^k} \phi g dx + \int_{\mathbb{T}^d} E[1/b_k] 1_{[\overline{\mathfrak{B}}^k]^c} \phi g dx = \int_{\mathbb{T}^d} E[1/b_k] \phi g dx.$$

Then, we have shown that

$$\begin{aligned} \lim_{N \rightarrow \infty} \int_{\mathbb{T}^d} \widetilde{1/a_k^N} \phi d(x^k \otimes W_k) &= \int_{\mathbb{T}^d} E[1/b_k] \phi d(x^k \otimes \lambda^\perp) + \int_{\mathbb{T}^d} E[1/b_k] \phi g dx \\ &= \int_{\mathbb{T}^d} E[1/b_k] \phi d(x^k \otimes W_k) \end{aligned}$$

and this concludes the proof of the first statement. \square

Remark 5.13. One should notice that since we were dealing with a diagonal matrix, our operator is the tensor product of one-dimensional operators. Thus, in the proof of Theorem 5.9 we were able to split the analysis of the operator in \mathbb{T}^d into d analysis on essentially one-dimensional spaces. Finally, we then gathered all the results together to recover the result for \mathbb{T}^d . With this, the result obtained by Lemma 5.12 is remarkably similar to that of the one-dimensional homogenization, which is known to not fully generalize to the d -dimensional setup for $d > 1$. The result was similar to its one-dimensional counterpart due to the diagonality of the matrix. In a non-diagonal matrix, we would not be able to tackle the coordinates independently, and thus the result would be very different.

We had to consider diagonal matrices, because a non-diagonal matrix implies the usage of an operator of the form $\partial_{x_i} \partial_{W_j}$, with $i \neq j$. To apply such an operator in the strong sense, the function W_j cannot be discontinuous at any point. This makes us, for instance, lose the interpretation of the discontinuities as permeable membranes, which was the motivation of [5] for introducing this operator for the one-dimensional setup. Nevertheless, the situation involving $\partial_{x_i} \partial_{W_j}$, with $i \neq j$, is challenging from the mathematical point-of-view and is still interesting from the physical point-of-view.

6. Application

To conclude the paper we will provide an application of a new result on probability theory which is an improvement of the result obtained in [14] in two directions: first, it considers a more general model; second, it has a more natural and simpler proof. It is also noteworthy that the homogenization results obtained in this paper were the key results in proving the main Theorem in the article [2]. For each choice of diagonal matrix function A^N satisfying the hypotheses of Proposition 5.10 or of Theorem 5.11, we have a corresponding version of hydrodynamic limit.

6.1. The hydrodynamic limit

We will begin by recalling some definitions. Recall in (1) the definition of function W , consider a sequence of operators $\nabla^N A^N \nabla_W^N$ that satisfies the hypotheses of Proposition 5.10 or of Theorem 5.11. For each $x \in \mathbb{T}_N^d$ and $k = 1, \dots, d$, define the symmetric rate $\xi_{x, x+e_k} = \xi_{x+e_k, x}$ by

$$\xi_{x,x+e_k} = \frac{a_k^N(x/N)}{N[W((x+e_k)/N) - W(x/N)]} = \frac{a_k^N(x/N)}{N[W_k((x_k+1)/N) - W_k(x_k/N)]}, \quad (49)$$

where $A^N = \text{diag}\{a_1^N(x), \dots, a_d^N(x)\}$, and $\{e_1, \dots, e_d\}$ is the canonical basis of \mathbb{R}^d . Let $b > -1/2$ and

$$c_{x,x+e_k}(\eta) = 1 + b\{\eta(x - e_k) + \eta(x + 2e_k)\},$$

where all sums are modulo N .

Distribute particles on \mathbb{T}_N^d in such a way that each site of \mathbb{T}_N^d is occupied at most by one particle. Denote by η the configuration of the state space $\{0, 1\}^{\mathbb{T}_N^d} = \{\eta : \mathbb{T}_N^d \rightarrow \{0, 1\}\}$ so that $\eta(x) = 0$ if site x is vacant, and $\eta(x) = 1$ if site x is occupied.

The exclusion process with conductances is a continuous-time Markov process $\{\eta_t : t \geq 0\}$ with state space $\{0, 1\}^{\mathbb{T}_N^d}$, whose generator L_N acts on functions $f : \{0, 1\}^{\mathbb{T}_N^d} \rightarrow \mathbb{R}$ as

$$(L_N f)(\eta) = \sum_{k=1}^d \sum_{x \in \mathbb{T}_N^d} \xi_{x,x+e_k} c_{x,x+e_k}(\eta) \{f(\sigma^{x,x+e_k} \eta) - f(\eta)\}, \quad (50)$$

where $\sigma^{x,x+e_k} \eta$ is the configuration obtained from η by exchanging the variables $\eta(x)$ and $\eta(x + e_k)$:

$$(\sigma^{x,x+e_k} \eta)(y) = \begin{cases} \eta(x + e_k) & \text{if } y = x, \\ \eta(x) & \text{if } y = x + e_k, \\ \eta(y) & \text{otherwise.} \end{cases} \quad (51)$$

We consider the Markov process $\{\eta_t : t \geq 0\}$ on the configurations $\{0, 1\}^{\mathbb{T}_N^d}$ associated to the generator L_N in the diffusive scale, i.e., L_N is speeded up by N^2 .

We now describe the stochastic evolution of the process. Let $x = (x_1, \dots, x_d) \in \mathbb{T}_N^d$. After a time given by an exponential distribution, at rate $\xi_{x,x+e_k} c_{x,x+e_k}(\eta)$ the occupation variables $\eta(x)$ and $\eta(x + e_k)$ are exchanged. Note that only nearest neighbor jumps are allowed. If W is differentiable at $x/N \in [0, 1]^d$, the rate at which particles are exchanged is of order 1 for each direction, but if some W_k is discontinuous at x_k/N , it no longer holds. In fact, assume, to fix ideas, that W_k is discontinuous at x_k/N , and smooth on the segments $(x_k/N, x_k/N + \varepsilon e_k)$ and $(x_k/N - \varepsilon e_k, x_k/N)$. Assume, also, that W_j is differentiable in a neighborhood of x_j/N for $j \neq k$. In this case, the rate at which particles jump over the bonds $\{y - e_k, y\}$, with $y_k = x_k$, is of order $1/N$, whereas in a neighborhood of size N of these bonds, particles jump at rate 1. Thus, note that a particle at site $y - e_k$ jumps to y at rate $1/N$ and jumps at rate 1 to each one of the $2d - 1$ other options. Particles, therefore, tend to avoid the bonds $\{y - e_k, y\}$. However, since time will be scaled diffusively, and since on a time interval of length N^2 a particle spends a time of order N at each site y , particles will be able to cross the slower bond $\{y - e_k, y\}$. The scaling limits of this interacting particle systems in inhomogeneous media may, for instance, model diffusions in which permeable membranes, at the points of discontinuities of the conductances W , tend to reflect particles, creating space discontinuities in the density profiles. For more details see [14].

The effect of the factor $c_{x,x+e_k}(\eta)$ is the following: if the parameter b is positive, the presence of particles in the neighboring sites of the bond $\{x, x+e_k\}$ speeds up the exchange rate by a factor of order one, and if the parameter b is negative, the presence of particles in the neighboring sites slows down the exchange rate also by a factor of order one.

Let $A = \text{diag}\{a_1, \dots, a_d\}$ be the homogenization of the sequence of matrices A^N . Note that A is a diagonal matrix with $a_k > 0$, $k = 1, \dots, d$, and recall from subsection 3, the operator defined on $\mathfrak{D}_W(\mathbb{T}^d)$:

$$\nabla A \nabla_W := \sum_{k=1}^d \partial_{x_k} a_k \partial_{W_k}.$$

A sequence of probability measures $\{\mu_N : N \geq 1\}$ on $\{0, 1\}^{\mathbb{T}_N^d}$ is said to be associated to a profile $\rho_0 : \mathbb{T}^d \rightarrow [0, 1]$ if for every $\delta > 0$ and every function $H \in \mathfrak{D}_W(\mathbb{T}^d)$:

$$\lim_{N \rightarrow \infty} \mu_N \left\{ \eta; \left| \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} H(x/N) \eta(x) - \int H(u) \rho_0(u) du \right| > \delta \right\} = 0. \quad (52)$$

Let $\gamma : \mathbb{T}^d \rightarrow [l, r]$ be a bounded density profile and consider the parabolic differential equation

$$\begin{cases} \partial_t \rho = \nabla A \nabla_W \Phi(\rho) \\ \rho(0, \cdot) = \gamma(\cdot) \end{cases}, \quad (53)$$

where the function $\Phi : [l, r] \rightarrow \mathbb{R}$ has bounded derivative, its derivative is also away from zero, and $t \in [0, T]$, for $T > 0$ fixed.

A function $\rho : [0, T] \times \mathbb{T}^d \rightarrow [l, r]$ is said to be a weak solution of the parabolic differential equation (53) if the following conditions hold. $\Phi(\rho(\cdot, \cdot))$ and $\rho(\cdot, \cdot)$ belong to $L^2([0, T], H_{1,W}(\mathbb{T}^d))$, and we have the integral identity

$$\int_{\mathbb{T}^d} \rho(t, u) H(u) du - \int_{\mathbb{T}^d} \rho(0, u) H(u) du = \int_0^t \int_{\mathbb{T}^d} \Phi(\rho(s, u)) \nabla A \nabla_W H(u) du ds,$$

for every function $H \in \mathfrak{D}_W(\mathbb{T}^d)$ and all $t \in [0, T]$.

Existence of such weak solutions follows from the tightness of the process, which is proved in subsection 6.2, and from the energy estimate given in [14, Lemma 6.2]. Uniqueness of weak solutions was proved in [12].

The main result of this Section is the following.

Theorem 6.1. Fix a continuous initial profile $\rho_0 : \mathbb{T}^d \rightarrow [0, 1]$ and consider a sequence of probability measures μ_N on $\{0, 1\}^{\mathbb{T}_N^d}$ associated to ρ_0 , in the sense of (52). Then, for any $t \geq 0$,

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left\{ \left| \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} H(x/N) \eta_t(x) - \int H(u) \rho(t, u) du \right| > \delta \right\} = 0$$

for every $\delta > 0$ and every function $H \in \mathfrak{D}_W(\mathbb{T}^d)$. Here, ρ is the unique weak solution of the non-linear equation (53) with $l = 0$, $r = 1$, $\gamma = \rho_0$ and $\Phi(\alpha) = \alpha + \alpha\alpha^2$.

6.2. Proof of Theorem 6.1

A simple computation shows that the Bernoulli product measures $\{\nu_\alpha^N : 0 \leq \alpha \leq 1\}$ are invariant, and in fact reversible, for the dynamics. The measure ν_α^N is obtained by placing a particle at each site, independently from the other sites, with probability α . Thus, ν_α^N is a product measure over $\{0, 1\}^{\mathbb{T}_N^d}$ with marginals given by $\nu_\alpha^N\{\eta : \eta(x) = 1\} = \alpha$, for x in \mathbb{T}_N^d . For more details see [6, chapter 2].

Consider the random walk $\{X_t\}_{t \geq 0}$ of a particle in \mathbb{T}_N^d induced by the generator L_N given as follows. Let $\xi_{x, x+e_k}$ be given by (49). If the particle is on a site $x \in \mathbb{T}_N^d$, it will jump to $x + e_k$ with rate $N^2 \xi_{x, x+e_k}$. Furthermore, only nearest neighbor jumps are allowed. The generator \mathbb{L}_N of the random walk $\{X_t\}_{t \geq 0}$ acts on functions $f : \mathbb{T}_N^d \rightarrow \mathbb{R}$ as

$$\mathbb{L}_N f\left(\frac{x}{N}\right) = \sum_{k=1}^d \mathbb{L}_N^k f\left(\frac{x}{N}\right),$$

where,

$$\mathbb{L}_N^k f\left(\frac{x}{N}\right) = N^2 \left\{ \xi_{x, x+e_k} \left[f\left(\frac{x+e_k}{N}\right) - f\left(\frac{x}{N}\right) \right] + \xi_{x-e_k, x} \left[f\left(\frac{x-e_k}{N}\right) - f\left(\frac{x}{N}\right) \right] \right\}$$

It is not difficult to see that the following equality holds:

$$\mathbb{L}_N f(x/N) = \sum_{k=1}^d \partial_{x_k}^N (a_k^N \partial_{W_k}^N f)(x) := \nabla^N A^N \nabla_W^N f(x). \quad (54)$$

The counting measure m_N on $\frac{1}{N} \mathbb{T}_N^d$ is reversible for this process. This random walk plays an important role in the proof of the hydrodynamic limit of the process η_t .

Let $D(\mathbb{R}_+, \{0, 1\}^{\mathbb{T}_N^d})$ be the path space of càdlàg trajectories with values in $\{0, 1\}^{\mathbb{T}_N^d}$. For a measure μ_N on $\{0, 1\}^{\mathbb{T}_N^d}$, denote by \mathbb{P}_{μ_N} the probability measure on $D(\mathbb{R}_+, \{0, 1\}^{\mathbb{T}_N^d})$ induced by the initial state μ_N and the Markov process $\{\eta_t : t \geq 0\}$. Expectation with respect to \mathbb{P}_{μ_N} is denoted by \mathbb{E}_{μ_N} .

Let \mathcal{M} be the space of positive measures on \mathbb{T}^d with total mass bounded by one endowed with the weak topology. Recall that $\pi_t^N \in \mathcal{M}$ stands for the empirical measure at time t . This is the measure on \mathbb{T}^d obtained by rescaling space by N and by assigning mass $1/N^d$ to each particle:

$$\pi_t^N = \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} \eta_t(x) \delta_{x/N}, \quad (55)$$

where δ_u is the Dirac measure concentrated on u .

For a function $H : \mathbb{T}^d \rightarrow \mathbb{R}$, $\langle \pi_t^N, H \rangle$ stands for the integral of H with respect to π_t^N :

$$\langle \pi_t^N, H \rangle = \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} H(x/N) \eta_t(x).$$

This notation is not to be mistaken with the inner product in $L^2(\mathbb{T}^d)$ introduced earlier. Also, when π_t has a density ρ , $\pi(t, du) = \rho(t, u)du$.

Fix $T > 0$ and let $D([0, T], \mathcal{M})$ be the space of \mathcal{M} -valued càdlàg trajectories $\pi : [0, T] \rightarrow \mathcal{M}$ endowed with the *uniform* topology. Note that \mathcal{M} is endowed with the weak topology, which is metrizable, since \mathbb{T}^d is a compact metric space. Thus, this uniform topology is well-defined. For each probability measure μ_N on $\{0, 1\}^{\mathbb{T}_N^d}$, denote by $\mathbb{Q}_{\mu_N}^{W,N}$ the measure on the path space $D([0, T], \mathcal{M})$ induced by the measure μ_N and the process π_t^N introduced in (55).

Fix a continuous profile $\rho_0 : \mathbb{T}^d \rightarrow [0, 1]$ and consider a sequence $\{\mu_N : N \geq 1\}$ of measures on $\{0, 1\}^{\mathbb{T}_N^d}$ associated to ρ_0 in the sense of (52). Further, we denote by \mathbb{Q}_W the probability measure on $D([0, T], \mathcal{M})$ concentrated on the deterministic path $\pi(t, du) = \rho(t, u)du$, where ρ is the unique weak solution of (53) with $\gamma = \rho_0$, $l_k = 0$, $r_k = 1$, $k = 1, \dots, d$ and $\Phi(\alpha) = \alpha + b\alpha^2$.

In subsection 6.2 we show that the sequence $\{\mathbb{Q}_{\mu_N}^{W,N} : N \geq 1\}$ is tight, and we characterize the limit points of this sequence.

Tightness: The goal of this part is to prove tightness of sequence $\{\mathbb{Q}_{\mu_N}^{W,N} : N \geq 1\}$. Fix $\lambda > 0$ and consider, initially, the auxiliary \mathcal{M} -valued Markov process $\{\Pi_t^{\lambda,N} : t \geq 0\}$ defined by

$$\Pi_t^{\lambda,N}(H) = \langle \pi_t^N, H_\lambda^N \rangle = \frac{1}{N^d} \sum_{x \in \mathbb{Z}^d} H_\lambda^N(x/N) \eta_t(x),$$

for H in $\mathfrak{D}_W(\mathbb{T}^d)$, where H_λ^N is the unique weak solution in $H_{1,W}(\mathbb{T}_N^d)$ of

$$\lambda H_\lambda^N - \nabla^N A^N \nabla_W^N H_\lambda^N = \lambda H - \nabla A \nabla_W H,$$

with the right-hand side being understood as the restriction of the function to the lattice \mathbb{T}_N^d , which is well-defined, since $H \in \mathfrak{D}_W(\mathbb{T}^d)$ and from Remark 3.2 under the hypotheses of Proposition 5.10 or by noticing that the homogenized matrix is constant under the hypotheses of Theorem 5.11, we have that $\nabla A \nabla_W H$ belongs to $C_W(\mathbb{T}^d)$. Thus the right-hand side belongs to $C_W(\mathbb{T}^d)$ and it is right-continuous, Remark 3.4.

We first prove tightness of the process $\{\Pi_t^{\lambda,N} : 0 \leq t \leq T\}$. Then we show that $\{\Pi_t^{\lambda,N} : 0 \leq t \leq T\}$ and $\{\pi_t^N : 0 \leq t \leq T\}$ are not far apart.

It is well known [6] that to prove tightness of $\{\Pi_t^{\lambda,N} : 0 \leq t \leq T\}$ it is enough to show tightness of the real-valued processes $\{\Pi_t^{\lambda,N}(H) : 0 \leq t \leq T\}$ for a set of test functions $H : \mathbb{T}^d \rightarrow \mathbb{R}$ dense in $C(\mathbb{T}^d)$ for the uniform topology. We may use, for instance, $\mathfrak{D}_W(\mathbb{T}^d)$.

Fix a function $H : \mathbb{T}^d \rightarrow \mathbb{R}$ in $\mathfrak{D}_W(\mathbb{T}^d)$. Keep in mind that $\Pi_t^{\lambda,N}(H) = \langle \pi_t^N, H_\lambda^N \rangle$, and denote by $M_t^{N,\lambda}$ the martingale defined by

$$M_t^{N,\lambda} = \Pi_t^{\lambda,N}(H) - \Pi_0^{\lambda,N}(H) - \int_0^t ds N^2 L_N \langle \pi_s^N, H_\lambda^N \rangle. \quad (56)$$

Clearly, tightness of $\Pi_t^{\lambda,N}(H)$ follows from tightness of the martingale $M_t^{N,\lambda}$ and tightness of the additive functional $\int_0^t ds N^2 L_N \langle \pi_s^N, H_\lambda^N \rangle$.

A long computation, albeit simple, shows that the quadratic variation $\langle M^{N,\lambda} \rangle_t$ of the martingale $M_t^{N,\lambda}$ is given by:

$$\begin{aligned} & \frac{1}{N^{2d-1}} \sum_{k=1}^d \sum_{x \in \mathbb{T}^d} [\partial_{W_k}^N H_\lambda^N(x/N)]^2 [W((x+e_k)/N) - W(x/N)] \\ & \times \int_0^t c_{x,x+e_k}(\eta_s) [\eta_s(x+e_k) - \eta_s(x)]^2 ds. \end{aligned}$$

In particular, by Lemma 4.5,

$$\langle M^{N,\lambda} \rangle_t \leq \frac{C_0 t}{N^d} \sum_{k=1}^d \|H_\lambda^N\|_{W_k,N}^2 \leq \frac{C(H)t}{\lambda N^d},$$

for some finite constant $C(H)$, which depends only on H . Thus, by Doob inequality, for every $\lambda > 0$, $\delta > 0$,

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left[\sup_{0 \leq t \leq T} |M_t^{N,\lambda}| > \delta \right] = 0. \quad (57)$$

In particular, the sequence of martingales $\{M_t^{N,\lambda} : N \geq 1\}$ is tight for the uniform topology.

It remains to examine the additive functional of the decomposition (56). The generator of the exclusion process L_N can be decomposed in terms of the generator of the random walk \mathbb{L}_N . By a long but simple computation, we obtain that $N^2 L_N \langle \pi^N, H_\lambda^N \rangle$ is equal to

$$\begin{aligned} & \sum_{k=1}^d \left\{ \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} (\mathbb{L}_N^k H_\lambda^N)(x/N) \eta(x) - \frac{b}{N^d} \sum_{x \in \mathbb{T}_N^d} (\mathbb{L}_N^k H_\lambda^N)(x/N) (\tau_x h_{2,k})(\eta) \right\} \\ & + \frac{b}{N^d} \sum_{x \in \mathbb{T}_N^d} [(\mathbb{L}_N^k H_\lambda^N)((x+e_k)/N) + (\mathbb{L}_N^k H_\lambda^N)(x/N)] (\tau_x h_{1,k})(\eta), \end{aligned}$$

where $\{\tau_x : x \in \mathbb{Z}^d\}$ is the group of translations, so that $(\tau_x \eta)(y) = \eta(x+y)$ for x, y in \mathbb{Z}^d , and the sum is understood modulo N . Also, $h_{1,k}, h_{2,k}$ are the cylinder functions

$$h_{1,k}(\eta) = \eta(0)\eta(e_k), \quad h_{2,k}(\eta) = \eta(-e_k)\eta(e_k).$$

Since H_λ^N is the weak solution of the discrete equation, we have by Remark 4.2 that it is also a strong solution. Then, we may replace $\mathbb{L}_N H_\lambda^N$ by $U_\lambda^N = \lambda(H_\lambda^N - H) + \nabla A \nabla_W H$ in the previous formula. In particular, for all $0 \leq s < t \leq T$,

$$\left| \int_s^t dr N^2 L_N \langle \pi_r^N, H_\lambda^N \rangle \right| \leq \frac{(1 + 3|b|)(t - s)}{N^d} \sum_{x \in \mathbb{T}_N^d} |U_\lambda^N(x/N)|.$$

It follows from the estimate given in Lemma 4.5, see Remark 3.2, Lemma 3.5, and from Schwartz inequality, that the right hand side of the previous expression is bounded above by $C(H, b)(t - s)$ uniformly in N , where $C(H, b)$ is a finite constant depending only on b and H . This proves that the additive part of the decomposition (56) is tight for the uniform topology and, therefore, that the sequence of processes $\{\Pi_t^{\lambda, N} : N \geq 1\}$ is tight.

Lemma 6.2. *The sequence of measures $\{\mathbb{Q}_{\mu^N}^{W, N} : N \geq 1\}$ is tight for the uniform topology.*

Proof. Fix $\lambda > 0$. It is enough to show that for every function $H \in \mathfrak{D}_W(\mathbb{T}^d)$ and every $\epsilon > 0$, we have

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\mu^N} \left[\sup_{0 \leq t \leq T} |\Pi_t^{\lambda, N}(H) - \langle \pi_t^N, H \rangle| > \epsilon \right] = 0,$$

whence tightness of π_t^N follows from tightness of $\Pi_t^{\lambda, N}$. By Chebyshev's inequality, the last expression is bounded above by

$$\frac{1}{\epsilon^2} \mathbb{E}_{\mu^N} \left[\sup_{0 \leq t \leq T} |\Pi_t^{\lambda, N}(H) - \langle \pi_t^N, H \rangle|^2 \right] \leq \frac{2}{\epsilon^2} \|H_\lambda^N - H\|^2,$$

since there exists at most one particle per site. By Theorem 5.9, Proposition 5.7 and Corollary 5.8, $\|H_\lambda^N - H\|^2 \rightarrow 0$ as $N \rightarrow \infty$, and the proof follows. \square

The reader should compare the proof given in this Section to the proof given in [14], to check that the one given here follows closely the standard approach given, for instance, in [6], whereas the approach given in [14] is non-standard. This means that the theory provided in this article made the study of hydrodynamic behavior of exclusion processes with conductances tractable in the standard fashion. Thus, future work on this field will be simplified.

Uniqueness of limit points: We prove in this subsection that all limit points \mathbb{Q}^* of the sequence $\mathbb{Q}_{\mu^N}^{W, N}$ are concentrated on absolutely continuous trajectories $\pi(t, du) = \rho(t, u)du$, whose density $\rho(t, u)$ is a weak solution of the hydrodynamic equation (53) with $l = 0$, $r = 1$ and $\Phi(\alpha) = \alpha + a\alpha^2$.

We now state a result necessary to prove the uniqueness of limit points. Let, for a local function $g : \{0, 1\}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$, $\tilde{g} : [0, 1] \rightarrow \mathbb{R}$ be the expected value of g under the stationary states:

$$\tilde{g}(\alpha) = E_{v_\alpha}[g(\eta)].$$

For $\ell \geq 1$ and d -dimensional integer $x = (x_1, \dots, x_d)$, denote by $\eta^\ell(x)$ the empirical density of particles in the box $\mathbb{B}_+^\ell(x) = \{(y_1, \dots, y_d) \in \mathbb{Z}^d; 0 \leq y_i - x_i < \ell\}$:

$$\eta^\ell(x) = \frac{1}{\ell^d} \sum_{y \in \mathbb{B}_+^\ell(x)} \eta(y).$$

Let \mathbb{Q}^* be a limit point of the sequence $\mathbb{Q}_{\mu_N}^{W,N}$ and assume, without loss of generality, that $\mathbb{Q}_{\mu_N}^{W,N}$ converges to \mathbb{Q}^* .

Since there is at most one particle per site, it is clear that \mathbb{Q}^* is concentrated on trajectories $\pi_t(du)$ which are absolutely continuous with respect to the Lebesgue measure, $\pi_t(du) = \rho(t, u)du$, and whose density ρ is non-negative and bounded by 1. The reader is referred to [6, Chapter 4] for further details.

Fix a function $H \in \mathcal{D}_W(\mathbb{T}^d)$ and $\lambda > 0$. Recall the definition of the martingale $M_t^{N,\lambda}$ introduced in the previous Section. From (57) we have that for every $\delta > 0$,

$$\lim_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left[\sup_{0 \leq t \leq T} |M_t^{N,\lambda}| > \delta \right] = 0,$$

and from (56), for fixed $0 < t \leq T$ and $\delta > 0$, we have

$$\lim_{N \rightarrow \infty} \mathbb{Q}_{\mu_N}^{W,N} \left[\left| \langle \pi_t^N, H_\lambda^N \rangle - \langle \pi_0^N, H_\lambda^N \rangle - \int_0^t ds N^2 L_N \langle \pi_s^N, H_\lambda^N \rangle \right| > \delta \right] = 0.$$

Note that the expression $N^2 L_N \langle \pi_s^N, H_\lambda^N \rangle$ has been computed in the previous subsection in terms of generator \mathbb{L}_N . On the other hand, $\mathbb{L}_N H_\lambda^N = \lambda H_\lambda^N - \lambda H + \nabla A \nabla_W H$. Since there is at most one particle per site, we may apply Theorem 5.9 along with Proposition 5.7 and Corollary 5.8, to replace $\langle \pi_t^N, H_\lambda^N \rangle$ and $\langle \pi_0^N, H_\lambda^N \rangle$ by $\langle \pi_t, H \rangle$ and $\langle \pi_0, H \rangle$, respectively, and replace $\mathbb{L}_N H_\lambda^N$ by $\nabla A \nabla_W H$ plus a term that vanishes as $N \rightarrow \infty$.

Since $E_{\nu_\alpha}[h_{i,k}] = \alpha^2$, $i = 1, 2$ and $k = 1, \dots, d$, we have by the Replacement Lemma [14, Corollary 5.4] that, for every $t > 0$, $\lambda > 0$, $\delta > 0$, $i = 1, 2$,

$$\lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{P}_{\mu_N} \left[\left| \int_0^t ds \frac{1}{N^d} \sum_{x \in \mathbb{T}_N^d} \nabla A \nabla_W H(x/N) \times \left\{ \tau_x h_{i,k}(\eta_s) - [\eta_s^{\varepsilon N}(x)]^2 \right\} \right| > \delta \right] = 0.$$

Since $\eta_s^{\varepsilon N}(x) = \varepsilon^{-d} \pi_s^N(\prod_{k=1}^d [x_k/N, x_k/N + \varepsilon e_k])$, we obtain, from the previous considerations, that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \limsup_{N \rightarrow \infty} \mathbb{Q}_{\mu_N}^{W,N} \left[\left| \langle \pi_t, H \rangle - \langle \pi_0, H \rangle - \int_0^t ds \left\langle \Phi(\varepsilon^{-d} \pi_s^N(\prod_{k=1}^d [\cdot, \cdot + \varepsilon e_k])), \nabla A \nabla_W H \right\rangle \right| > \delta \right] \\ & = 0. \end{aligned}$$

Using the fact that $\mathbb{Q}_{\mu_N}^{W,N}$ converges in the uniform topology to \mathbb{Q}^* , we have that

$$\lim_{\varepsilon \rightarrow 0} \mathbb{Q}^* \left[\left| \langle \pi_t, G_\lambda H \rangle - \langle \pi_0, G_\lambda H \rangle - \int_0^t ds \left\langle \Phi(\varepsilon^{-d} \pi_s(\prod_{k=1}^d [\cdot, \cdot + \varepsilon e_k])) , U_\lambda \right\rangle \right| > \delta \right] = 0.$$

Recall that \mathbb{Q}^* is concentrated on absolutely continuous paths $\pi_t(du) = \rho(t, u)du$ with positive density bounded by 1. Therefore, $\varepsilon^{-d} \pi_s(\prod_{k=1}^d [\cdot, \cdot + \varepsilon e_k])$ converges in $L^1(\mathbb{T}^d)$ to $\rho(s, \cdot)$ as $\varepsilon \downarrow 0$. Thus,

$$\mathbb{Q}^* \left[\left| \langle \pi_t, H \rangle - \langle \pi_0, H \rangle - \int_0^t ds \langle \Phi(\rho_s), \nabla A \nabla_W H \rangle \right| > \delta \right] = 0.$$

Letting $\delta \downarrow 0$, we see that, \mathbb{Q}^* almost surely,

$$\int_{\mathbb{T}^d} \rho(t, u) H(u) du - \int_{\mathbb{T}^d} \rho(0, u) H(u) du = \int_0^t \int_{\mathbb{T}^d} \Phi(\rho(s, u)) \nabla A \nabla_W H(u) du ds.$$

This identity can be extended to a countable set of times t . Taking this set to be dense we obtain, by continuity of the trajectories π_t , that it holds for all $0 \leq t \leq T$.

From [14, Lemma 6.2], which we may easily adapt to our setup by using the uniform ellipticity condition (42) of the environment, we may conclude that all limit points have, almost surely, finite energy, and therefore, by [12, Lemma 4.1], $\Phi(\rho(\cdot, \cdot)) \in L^2([0, T], H_{1,W}(\mathbb{T}^d))$. Analogously, it is possible to show that $\rho(\cdot, \cdot)$ has finite energy and hence it belongs to $L^2([0, T], H_{1,W}(\mathbb{T}^d))$.

Proposition 6.3. *As $N \uparrow \infty$, the sequence of probability measures $\mathbb{Q}_{\mu_N}^{W,N}$ converges in the uniform topology to \mathbb{Q}_W .*

Proof. In the previous subsection, we showed that the sequence of probability measures $\mathbb{Q}_{\mu_N}^{W,N}$ is tight for the uniform topology. Moreover, we just proved that all limit points of this sequence are concentrated on weak solutions of the parabolic equation (53). The proposition now follows from the uniqueness proved in [12]. \square

Proof of Theorem 6.1. Since $\mathbb{Q}_{\mu_N}^{W,N}$ converges in the uniform topology to \mathbb{Q}_W , a measure which is concentrated on a deterministic path, for each $0 \leq t \leq T$ and each continuous function $H : \mathbb{T}^d \rightarrow \mathbb{R}$, $\langle \pi_t^N, H \rangle$ converges in probability to $\int_{\mathbb{T}^d} d\rho(t, u) H(u)$, where ρ is the unique weak solution of (53) with $l_k = 0$, $r_k = 1$, $\gamma = \rho_0$ and $\Phi(\alpha) = \alpha + \alpha \alpha^2$. \square

Appendix A

In this appendix we will present, for the reader's convenience, part of the proof of the [5, Lemma 1]. This result was used in Proposition 3.3.

Proposition A.1. *The space $\mathfrak{D}_W(\mathbb{T})$ is dense in $C(\mathbb{T})$, the space of continuous functions in \mathbb{T} , in the sup norm, $\|\cdot\|_\infty$.*

Proof. Let $f : \mathbb{T} \rightarrow \mathbb{R}$ be a continuous function, and $\epsilon > 0$. From uniform continuity, there exists $\delta > 0$ such that $|f(y) - f(x)| \leq \epsilon$ whenever $|x - y| \leq \delta$. Choose an integer $n \geq \delta^{-1}$ and consider the function $g : \mathbb{T} \rightarrow \mathbb{R}$ defined by

$$g(x) = \sum_{j=0}^{n-1} \frac{f([j+1]/n) - f(j/n)}{W_k([j+1]/n) - W_k(j/n)} 1_{(j/n, (j+1)/n]}(x).$$

Observe that g can be seen as a discrete derivative of f with respect to W_k . Thus, it is natural to guess that integrating this function with respect to W_k would yield an approximation of f . Thus, let $G : \mathbb{T} \rightarrow \mathbb{R}$ be given by $G(x) = f(0) + \int_{(0,x]} g(y) W_k(dy)$. By the very definition of g , $G(j/n) = f(j/n)$ for $0 \leq j < n$.

Since $n \geq \delta^{-1}$, we can use the definition of G to obtain, for $j/n \leq x \leq (j+1)/n$,

$$|G(x) - f(x)| \leq |G(x) - G(j/n)| + |f(x) - f(j/n)| \leq 2\epsilon,$$

whence $\|G - f\|_\infty \leq 2\epsilon$. Note that

$$\int_{(0,1]} g dW_k = 0. \quad (58)$$

It remains to be shown that the function G may be approximated in the sup norm by functions in $\mathfrak{D}_W(\mathbb{T}^d)$. Note that the only restriction we had when choosing the set $\{0, 1/n, \dots, (n-1)/n\}$ is that the distance between two consecutive points is less than δ . Therefore, we may replace these n points, by any such points satisfying this restriction. Since W_k is strictly increasing, it has a countable number of discontinuities, and then, we may assume, without loss of generality, that W_k is continuous at the points $\{0, 1/n, \dots, (n-1)/n\}$ (in the sense, that we may replace these points if it is not the case). Let $\{H_m : m \geq 1\}$ be a sequence of smooth functions $H_m : \mathbb{T} \rightarrow \mathbb{R}$, with $|H_m(x)| \leq \|g\|_\infty$, for all $x \in \mathbb{T}$, and such that $\lim_m H_m(x) = g(x)$ for $xn \notin \mathbb{Z}$. Then, the Dominated Convergence Theorem implies

$$\lim_{m \rightarrow \infty} \int_{\mathbb{T}} |H_m(y) - g(y)| dW_k(y) = 0. \quad (59)$$

Let $\{F_m : m \geq 1\}$ be the sequence of functions $F_m : \mathbb{T} \rightarrow \mathbb{R}$ defined by

$$F_m(x) = f(0) + \int_{(0,x]} H(y) dW_k(y).$$

Therefore, it is immediate that

$$\|G - F_m\|_\infty \leq \int_{\mathbb{T}} |H_m(y) - g(y)| dW_k(y) \xrightarrow{m \rightarrow \infty} 0.$$

This shows that we can approximate, in the sup norm, any function $f \in C(\mathbb{T})$ by functions of the form of F_m . To conclude that \mathfrak{D}_{W_k} is dense in $C(\mathbb{T})$, we must show that $F_m \in \mathfrak{D}_{W_k}$. To this end, note that

$$\begin{aligned} F_m(x) &= f(0) + \int_{(0,x]} \left\{ b_m + \int_0^y H'_m(z) dz \right\} dW_k(y) \\ &= f(0) + b_m W_k(x) + \int_{(0,x]} dW_k(y) \int_0^y H'_m(z) dz, \end{aligned}$$

where $b_m = H_m(0) - W_k(1)^{-1} \int_{\mathbb{T}} H_m(y) dW_k(y)$. Since H'_m is continuous, $H'_m \in C_W(\mathbb{T})$. Then, using the characterization of \mathfrak{D}_{W_k} , one may conclude that F_m belongs to \mathfrak{D}_{W_k} , for each $m \geq 1$. \square

References

- [1] A. Faggionato, Random walks and exclusion processes among random conductances on random infinite clusters: homogenization and hydrodynamic limit, *Electron. J. Probab.* 13 (73) (2008) 2217–2247.
- [2] Jonathan Farfan, Alexandre B. Simas, Fábio J. Valentim, Equilibrium fluctuations for exclusion processes with conductances in random environments, *Stochastic Process. Appl.* 120 (2010) 1535–1562.
- [3] W. Feller, On second order differential operators, *Ann. of Math.* 61 (1) (1955) 90–105.
- [4] W. Feller, Generalized second order differential operators and their lateral conditions, *Illinois J. Math.* 1 (4) (1957) 459–504.
- [5] T. Franco, C. Landim, Hydrodynamic limit of gradient exclusion processes with conductances, *Arch. Ration. Mech. Anal.* 195 (2009) 409–439 (print).
- [6] C. Kipnis, C. Landim, *Scaling Limits of Interacting Particle Systems*, Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 320, Springer-Verlag, Berlin, 1999.
- [7] M. Jara, C. Landim, Quenched nonequilibrium central limit theorem for a tagged particle in the exclusion process with bond disorder, [arXiv:math/0603653](https://arxiv.org/abs/math/0603653), *Ann. Inst. Henri Poincaré Probab. Stat.* 44 (2008) 341–361.
- [8] S.M. Kozlov, Averaging of difference schemes, *Math. USSR, Sb.* 57 (2) (1987) 351–369.
- [9] O.A. Ladyzhenskaya, *The Boundary Value Problems of Mathematical Physics*, Springer, NY, 1985.
- [10] P. Mandl, *Analytical Treatment of One-Dimensional Markov Processes*, Grundlehren der mathematischen Wissenschaften, vol. 151, Springer-Verlag, Berlin, 1968.
- [11] A. Piatnitski, E. Remy, Homogenization of elliptic difference operators, *SIAM J. Math. Anal.* 33 (2001) 53–83.
- [12] A.B. Simas, F.J. Valentim, W -Sobolev spaces, *J. Math. Anal. Appl.* 382 (1) (2011) 214–230.
- [13] A.B. Simas, F.J. Valentim, W -Sobolev spaces: higher order and regularity, *Commun. Pure Appl. Anal.* 14 (2015) 597–607.
- [14] F.J. Valentim, Hydrodynamic limit of a d -dimensional exclusion process with conductances, *Ann. Inst. Henri Poincaré Probab. Stat.* 48 (1) (2012) 188–211.