

Self-Similar Solutions to a Parabolic System Modeling Chemotaxis

Yūki Naito¹

*Department of Applied Mathematics, Faculty of Engineering, Kobe University,
Kobe 657-8501, Japan
E-mail: naito@cs.kobe-u.ac.jp.*

Takashi Suzuki

Department of Mathematics, Osaka University, Toyonaka 560-0043, Japan

and

Kiyoshi Yoshida

*Division of Mathematical and Information Sciences, Faculty of Integrated Arts and Sciences,
Hiroshima University, Higashi-Hiroshima 739-8521, Japan*

Received May 15, 2001; revised November 1, 2001

We study the forward self-similar solutions to a parabolic system modeling chemotaxis

$$u_t = \nabla \cdot (\nabla u - u \nabla v), \quad \tau v_t = \Delta v + u$$

in the whole space \mathbb{R}^2 , where τ is a positive constant. Using the Liouville-type result and the method of moving planes, it is proved that self-similar solutions (u, v) must be radially symmetric about the origin. Then the structure of the set of self-similar solutions is investigated. As a consequence, it is shown that there exists a threshold in $\int_{\mathbb{R}^2} u$ for the existence of self-similar solutions. In particular, for $0 < \tau \leq 1/2$, there exists a self-similar solution (u, v) if and only if $\int_{\mathbb{R}^2} u < 8\pi$. © 2002 Elsevier Science (USA)

Key Words: self-similar solution; parabolic system; chemotaxis; radial symmetry; blow-up analysis.

¹To whom correspondence should be addressed.



1. INTRODUCTION

We are concerned with the parabolic system of the form

$$\begin{cases} \frac{\partial u}{\partial t} = \nabla \cdot (\nabla u - u \nabla v), & x \in \mathbb{R}^N, \quad t > 0, \\ \tau \frac{\partial v}{\partial t} = \Delta v + u, & x \in \mathbb{R}^N, \quad t > 0, \end{cases} \tag{1.1}$$

where $\tau > 0$ is a constant. This is a simplified system of the one given by Keller and Segel [21] describing chemotactic feature of cellular slime molds sensitive to the gradient of a chemical substance secreted by themselves. The functions $u(x, t) \geq 0$ and $v(x, t) \geq 0$ denote the cell density of cellular slime molds and the concentration of the chemical substance at place x and time t , respectively.

Backward self-similar solutions are studied in [13] for $\tau = 0$. The present paper is devoted to the forward self-similar solutions. Namely, this system is invariant under the similarity transformation

$$u_\lambda(x, t) = \lambda^2 u(\lambda x, \lambda^2 t) \quad \text{and} \quad v_\lambda(x, t) = v(\lambda x, \lambda^2 t)$$

for $\lambda > 0$, that is, if (u, v) is a solution of (1.1) globally in time, then so is (u_λ, v_λ) . A solution (u, v) is said to be *self-similar*, when the solution is invariant under this transformation, that is, $u(x, t) = u_\lambda(x, t)$ and $v(x, t) = v_\lambda(x, t)$ for all $\lambda > 0$. Letting $\lambda = 1/\sqrt{t}$, we see that (u, v) has the form

$$u(x, t) = \frac{1}{t} \phi\left(\frac{x}{\sqrt{t}}\right) \quad \text{and} \quad v(x, t) = \psi\left(\frac{x}{\sqrt{t}}\right) \tag{1.2}$$

for $x \in \mathbb{R}^N$ and $t > 0$. It follows that

$$\int_{\mathbb{R}^N} u(x, t) \, dx = t^{(N-2)/2} \int_{\mathbb{R}^N} \phi(y) \, dy \tag{1.3}$$

for $\phi \in L^1(\mathbb{R}^N)$. Therefore, self-similar solution (u, v) preserves the mass $\|u(\cdot, t)\|_{L^1(\mathbb{R}^2)}$ if and only if $N = 2$. On the other hand, the mass conservation of $u(\cdot, t)$ follows formally in the original system (1.1) in any space dimensions. Regarding this fact, we study the case $N = 2$ in this paper.

By a direct computation it is shown that (u, v) in (1.2) satisfies (1.1) if and only if (ϕ, ψ) satisfies

$$\begin{cases} \nabla \cdot (\nabla \phi - \phi \nabla \psi) + \frac{1}{2} x \cdot \nabla \phi + \phi = 0, & x \in \mathbb{R}^2, \\ \Delta \psi + \frac{\tau}{2} x \cdot \nabla \psi + \phi = 0, & x \in \mathbb{R}^2. \end{cases} \tag{1.4}$$

We are concerned with the classical solutions $(\phi, \psi) \in C^2(\mathbb{R}^2) \times C^2(\mathbb{R}^2)$ of (1.4) satisfying

$$\phi, \psi \geq 0 \quad \text{in } \mathbb{R}^2 \quad \text{and} \quad \phi(x), \psi(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty. \quad (1.5)$$

Define the solution set \mathcal{S} of (1.4) as

$$\mathcal{S} = \{(\phi, \psi) \in C^2(\mathbb{R}^2) \times C^2(\mathbb{R}^2) : (\phi, \psi) \text{ is a solution of (1.4) with (1.5)}\}. \quad (1.6)$$

Put $\phi(x) = ce^{-|x|^2/4}e^{\psi(x)}$, where c is a positive constant. Then ϕ satisfies the first equation of (1.4), and so if we find a positive solution ψ of

$$\Delta\psi + \frac{\tau}{2}x \cdot \nabla\psi + ce^{-|x|^2/4}e^{\psi} = 0 \quad \text{in } \mathbb{R}^2$$

satisfying $\psi(x) \rightarrow 0$ as $|x| \rightarrow \infty$, we can obtain the solutions $(\phi, \psi) \in \mathcal{S}$. Therefore, the existence of solutions $(\phi, \psi) \in \mathcal{S}$ follows from [26, Proposition 2; 27, Theorem 1.1]. (We also obtain $\mathcal{S} \neq \emptyset$ by Lemma 4.2 below. See also Proposition A.1 in Appendix A.) We investigate the structure of the solution set \mathcal{S} .

THEOREM 1. *Any $(\phi, \psi) \in \mathcal{S}$ is radially symmetric about the origin, and satisfies $\phi, \psi \in L^1(\mathbb{R}^2)$.*

THEOREM 2. *The solution set \mathcal{S} is expressed as a one-parameter family:*

$$\mathcal{S} = \{(\phi(s), \psi(s)) : s \in \mathbb{R}\}.$$

If $\lambda(s) = \|\phi(s)\|_{L^1(\mathbb{R}^2)}$, then $(\phi(s), \psi(s))$ and $\lambda(s)$ satisfy the following properties:

- (i) $s \mapsto (\phi(s), \psi(s)) \in C^2(\mathbb{R}^2) \times C^2(\mathbb{R}^2)$ and $s \mapsto \lambda(s) \in \mathbb{R}$ are continuous;
- (ii) $(\phi(s), \psi(s)) \rightarrow (0, 0)$ in $C^2(\mathbb{R}^2) \times C^2(\mathbb{R}^2)$ and $\lambda(s) \rightarrow 0$ as $s \rightarrow -\infty$;
- (iii) $\|\psi(s)\|_{L^\infty(\mathbb{R}^2)} \rightarrow \infty, \lambda(s) \rightarrow 8\pi$, and $\phi(s) dx \rightarrow 8\pi\delta_0(dx)$ in the sense of measure as $s \rightarrow \infty$, where $\delta_0(dx)$ denotes Dirac's delta function with the support in origin;
- (iv) $0 < \lambda(s) < 8\pi$ for $s \in \mathbb{R}$, if $0 < \tau \leq 1/2$, and $0 < \lambda(s) \leq \max\{4\pi^3/3, 4\pi^3\tau^2/3\}$ for $s \in \mathbb{R}$, if $\tau > 1/2$.

As a consequence of Theorem 2 we obtain the following:

COROLLARY. *There exists a constant λ^* satisfying $\lambda^* = 8\pi$, if $0 < \tau \leq 1/2$, and $8\pi \leq \lambda^* \leq \max\{4\pi^3/3, 4\pi^3\tau^2/3\}$, if $\tau > 1/2$, such that*

- (i) for every $\lambda \in (0, \lambda^*)$, there exists a solution $(\phi, \psi) \in \mathcal{S}$ satisfying $\|\phi\|_{L^1(\mathbb{R}^2)} = \lambda$;
- (ii) for $\lambda > \lambda^*$, there exists no solution $(\phi, \psi) \in \mathcal{S}$ satisfying $\|\phi\|_{L^1(\mathbb{R}^2)} = \lambda$.

Remark. Biler [1] has shown that system (1.4) with $\tau = 1$ has a radial solution (ϕ, ψ) satisfying $\|\phi\|_{L^1(\mathbb{R}^2)} = \lambda$ for every $\lambda \in (0, 8\pi)$, and has no radial solutions (ϕ, ψ) satisfying $\|\phi\|_{L^1(\mathbb{R}^2)}/2\pi \geq 7.82 \dots$

Theorem 1 is a consequence of the following:

THEOREM 3. *Assume that (ϕ, ψ) is a nonnegative solution of (1.4) satisfying $\phi, \psi \in L^\infty(\mathbb{R}^2)$. Then ϕ and ψ are positive, and there exists a constant $\sigma > 0$ such that*

$$\phi(x) = \sigma e^{-|x|^2/4} e^{\psi(x)}. \tag{1.7}$$

Assume furthermore that $\psi(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then ϕ and ψ are radially symmetric about the origin, and satisfy $\partial\phi/\partial r < 0$ and $\partial\psi/\partial r < 0$ for $r = |x| > 0$, and

$$\phi(x) = O(e^{-|x|^2/4}) \quad \text{and} \quad \psi(x) = O(e^{-\min\{\tau, 1\}|x|^2/4}) \quad \text{as } |x| \rightarrow \infty.$$

The proof of Theorem 3 consists of two steps. First, we show that (1.7) holds by employing the Liouville-type result essentially due to Meyers and Serrin [24]. Then we show the radial symmetry of solutions by the method of moving planes. This device was first developed by Serrin [36] in PDE theory, and later extended and generalized by Gidas *et al.* [8, 9]. We will obtain a symmetry result for Eq. (1.8) below with a change of variables as in [31].

By Theorem 3 it follows that under the condition $\phi, \psi \in L^\infty(\mathbb{R}^2)$, system (1.4) is reduced to the equation

$$\Delta\psi + \frac{\tau}{2}x \cdot \nabla\psi + \sigma e^{-|x|^2/4} e^{\psi} = 0 \quad \text{in } \mathbb{R}^2 \tag{1.8}$$

for some positive constant σ . Moreover, $(\phi, \psi) \in \mathcal{S}$ if and only if ψ satisfies (1.8) with

$$\psi(x) \rightarrow 0 \quad \text{as } |x| \rightarrow \infty, \tag{1.9}$$

and ϕ is given by (1.7). Let $\lambda = \|\phi\|_{L^1(\mathbb{R}^2)}$. From (1.7) we see that

$$\lambda = \sigma \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi(y)} dy.$$

Then (1.8) is rewritten as the elliptic equation with the nonlocal term,

$$\Delta\psi + \frac{\tau}{2}x \cdot \nabla\psi + \lambda e^{-|x|^2/4} e^{\psi} / \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi(y)} dy = 0 \quad \text{in } \mathbb{R}^2. \tag{1.10}$$

The proof of Theorem 2 is based on the ODE arguments to Eqs. (1.8) and (1.10). Furthermore, we employ the results by Brezis and Merle [2] concerning the asymptotic behavior of sequences of solutions of

$$-\Delta u_k = V_k(x)e^{u_k} \quad \text{in } \Omega, \quad (1.11)$$

where $\Omega \subset \mathbb{R}^2$ is a bounded domain and V_k is a nonnegative continuous function. We also need Theorem 4 in order to prove Theorem 2. Here we recall Theorem 3 in [2].

THEOREM A (Brezis and Merle [2]). *Suppose that*

$$0 \leq V_k(x) \leq C_0, \quad x \in \Omega, \quad (1.12)$$

for some positive constant C_0 . Let $\{u_k\}$ be a sequence of solutions of (1.11) satisfying

$$\limsup_{k \rightarrow \infty} \int_{\Omega} e^{u_k} dx < \infty. \quad (1.13)$$

Then there exists a subsequence (still denoted by $\{u_k\}$) satisfying one of the following alternatives:

- (i) $\{u_k\}$ is bounded in $L_{\text{loc}}^{\infty}(\Omega)$;
- (ii) $u_k \rightarrow -\infty$ uniformly on compact subset of Ω ;
- (iii) *there exists a finite blowup set $\mathcal{B} = \{a_1, \dots, a_{\ell}\} \subset \Omega$ such that, for any $1 \leq i \leq \ell$, there exist $\{x_k\} \subset \Omega$, $x_k \rightarrow a_i$, $v_k(x_k) \rightarrow \infty$, and $v_k \rightarrow -\infty$ uniformly on compact subsets of $\Omega \setminus \mathcal{B}$. Moreover, $\int_{\Omega} V_k e^{u_k} dx \rightarrow \sum_{i=1}^{\ell} \alpha_i \delta_{a_i}(dx)$ in the sense of measure with $\alpha_i \geq 4\pi$, where $\delta_{a_i}(dx)$ is Dirac's delta function with the support in $x = a_i$.*

It was conjectured in [2] that each α_i can be written as $\alpha_i = 8\pi m_i$ for some positive integer m_i . This was established by Li and Shafrir [23]. Chen has shown in [3] that any positive integer m_i can occur in the case $V_k \equiv 1$ and Ω is a unit disk. On the other hand, under more restrictive assumptions that $V_k \in C^1(\Omega)$ we obtain the following theorem. It is related to Theorem 0.3 of Li [22] and is proven in the appendix of the present paper.

THEOREM 4. *Suppose that $V_k \in C^1(\Omega)$ satisfies (1.12) and*

$$\|\nabla V_k\|_{L^{\infty}(\Omega)} \leq C_1 \quad (1.14)$$

for some positive constants C_0 and C_1 . Let $\{u_k\}$ be a sequence of solutions of (1.11) satisfying (1.13) and

$$\max_{\partial\Omega} u_k - \min_{\partial\Omega} u_k \leq C_2 \tag{1.15}$$

for some positive constant C_2 . Assume that alternative (iii) in Theorem A holds. Then $\alpha_i = 8\pi$ for each $i \in \{1, 2, \dots, \ell\}$.

Recently, attentions have been paid to blowup problems for the system

$$\begin{cases} \frac{\partial u}{\partial t} = \nabla \cdot (\nabla u - u\nabla v), & x \in \Omega, \quad t > 0, \\ \tau \frac{\partial v}{\partial t} = \Delta v - \gamma v + u, & x \in \Omega, \quad t > 0, \\ \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0, & x \in \partial\Omega, \quad t > 0, \\ u(x, 0) = u_0, \quad v(x, 0) = v_0, & x \in \Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^2$ is a bounded domain with smooth boundary $\partial\Omega$, τ and γ are positive constants, and ν is the outer normal unit vector. As concerns dynamic aspects of solutions, Nanjundiah [32] has given a conjecture that blowup solution $u(x, t)$ will form a δ -function singularity as $t \nearrow T_{\max} < +\infty$, where T_{\max} denotes the maximal time for the existence of the solution. Such a phenomenon is referred to as *chemotactic collapse*. Herrero and Velázquez [14–16] constructed a family of radially symmetric solutions on $\Omega = \{x \in \mathbb{R}^2 : |x| < 1\}$, satisfying

$$u(x, t) \, dx \rightarrow 8\pi\delta_0(dx) + f(x) \, dx$$

as $t \nearrow T_{\max} < +\infty$ in $M(\bar{\Omega})$, where $f \in C(\bar{\Omega} \setminus \{0\}) \cap L^1(\Omega)$ is a nonnegative function. If $\tau = 0$, the blowup mechanism has been clarified much more by Senba and Suzuki [34].

Childress and Percus [5] and Childress [4] have studied the stationary problem and have conjectured that there exists a threshold in $\|u_0\|_{L^1(\Omega)}$ for the blow up of the solution (u, v) . Their arguments were heuristic, while recent studies are supporting their validity rigorously, see [12, 18, 28]. In particular, it is proven that $\|u_0\|_{L^1(\Omega)} < 4\pi$ implies $T_{\max} = +\infty$ by Nagai *et al.* [30], Biler [1], and Gajewski and Zacharias [7], independently. Furthermore, the optimality of the condition $\|u_0\|_{L^1(\Omega)} < 4\pi$ for $T_{\max} = +\infty$ is shown by Nagai [29] and Senba and Suzuki [35]. For related results, we refer to Horstmann and Wang [17].

On the other hand, concerning the Cauchy problem for the semilinear parabolic equation in \mathbb{R}^N , it is asserted that self-similar solutions take an important role in the asymptotic behavior of the solutions, see, e.g.

[6, 19, 20]. From corollary, we are led to the following conjectures for problem (1.1) with $N = 2$ subject to the initial condition $u(x, 0) = u_0$ and $v(x, 0) = v_0$ in \mathbb{R}^2 .

For $0 < \tau \leq 1/2$, if $\|u_0\|_{L^1(\mathbb{R}^2)} < 8\pi$ then the solution of the Cauchy problem to (1.1) exists globally in time, and if $\|u_0\|_{L^1(\mathbb{R}^2)} > 8\pi$ then the solution can blow up in a finite time.

We organize this paper as follows. In Section 2 we show that (1.7) holds by employing the Liouville-type result. In Section 3 we show the radial symmetry of solutions by the method of moving planes, and then give the proof of Theorem 3. In Section 4 we give the ODE arguments to investigate the properties of radial solutions of (1.8). We study the behavior of sequences $\{(\phi_k, \psi_k)\} \subset \mathcal{S}$ satisfying $\|\psi_k\|_{L^\infty(\mathbb{R}^2)} \rightarrow \infty$ in Section 5. In Section 6 we investigate the upper bounds of $\|\phi\|_{L^1(\mathbb{R}^2)}$. Finally, in Section 7 we prove Theorem 2 by using the results in Sections 4–6. In Appendixes, A and B, we are concerned with the existence of solutions to problem (1.8) and (1.9), and give the proof of Theorem 4.

2. REDUCTION TO THE SINGLE EQUATION

In this section we show that system (1.4) is reduced to Eq. (1.8) if $\phi, \psi \in L^\infty(\mathbb{R}^2)$. More precisely, we have the following:

PROPOSITION 2.1. *Let (ϕ, ψ) be a nonnegative solution of (1.4) with $\phi, \psi \in L^\infty(\mathbb{R}^2)$. Then relation (1.7) holds with some constant $\sigma > 0$.*

To prove this proposition we use the Liouville-type result for second-order elliptic inequalities essentially due to Meyers and Serrin [24].

LEMMA 2.1. *Let u satisfy*

$$\Delta u + \nabla b \cdot \nabla u \geq 0 \quad \text{in } \mathbb{R}^2. \quad (2.1)$$

Assume that $x \cdot \nabla b(x) \leq 0$ for large $|x|$. If $\sup_{x \in \mathbb{R}^2} u(x) < \infty$, then u must be a constant function.

Proof. Take a function μ as $\mu(r) = 1/\log(1+r)$. Then μ satisfies the Meyers–Serrin condition

$$\int_1^\infty \frac{k(t)}{t} dt = \infty, \quad \text{where } k(t) = \exp\left(-\int_1^t \frac{\mu(s)}{s} ds\right).$$

Define v as

$$v(r) = \int_1^r \frac{k(t)}{t} dt, \quad r \geq 1.$$

Then $v(r)$ is positive and increasing for $r \in (1, \infty)$, and satisfies $v(r) \rightarrow \infty$ as $r \rightarrow \infty$. Furthermore, $v = v(|x|)$ solves

$$\Delta v + \nabla b \cdot \nabla v = \frac{k(|x|)}{|x|^2}(-\mu(|x|) + x \cdot \nabla b(x)).$$

By the assumption, there exists a large $R > 0$ such that

$$\Delta v + \nabla b \cdot \nabla v < 0 \quad \text{for } |x| \geq R. \tag{2.2}$$

Now assume to the contrary that u is not a constant function. Without loss of generality, we may assume that u is not a constant function in $|x| \leq R$. Define

$$U(r) = \sup\{u(x) : |x| = r\}.$$

Then $U(r)$ is strictly increasing for $r \geq R$. To see why, suppose $R \leq r_1 < r_2$ and $U(r_1) \geq U(r_2)$. Then u attains its maximum for $|x| \leq r_2$ at an interior point and by the strong maximum principle u is constant, which contradicts the assumption. Therefore, $U(r)$ is strictly increasing, and we have $U(R + 1) > U(R)$. Choose $\delta > 0$ so small that

$$0 < \delta < \frac{U(R + 1) - U(R)}{v(R + 1) - v(R)}. \tag{2.3}$$

Put $w(x) = u(x) - \delta v(|x|)$. Then it follows from (2.1) and (2.2) that

$$\Delta w + \nabla b \cdot \nabla w > 0 \quad \text{for } |x| \geq R. \tag{2.4}$$

From (2.3) we obtain $U(R + 1) - \delta v(R + 1) > U(R) - \delta v(R)$. This implies

$$\sup_{|x|=R+1} w(x) > \sup_{|x|=R} w(x).$$

Since $w(x) \rightarrow -\infty$ as $|x| \rightarrow \infty$, w has the maximum at a point $x_0 \in \mathbb{R}^2$, $|x_0| > R$. Then we have $\Delta w + \nabla b \cdot \nabla w \leq 0$ at $x = x_0$. This contradicts (2.4). Hence, u must be a constant function. ■

LEMMA 2.2. *Let (ϕ, ψ) be a nonnegative solution of (1.4) with $\phi, \psi \in L^\infty(\mathbb{R}^2)$. Then $\nabla \psi \in L^\infty(\mathbb{R}^2)$.*

Proof. Define u and v by (1.2). Then (u, v) solves (1.1), and it holds that

$$\|u(t)\|_{L^\infty(\mathbb{R}^2)} = \frac{1}{t} \|\phi\|_{L^\infty(\mathbb{R}^2)} \quad \text{and} \quad \|v(t)\|_{L^\infty(\mathbb{R}^2)} = \|\psi\|_{L^\infty(\mathbb{R}^2)}.$$

Take $t_0 > 0$. From the second equation of (1.1) we have

$$v(t) = e^{((t-t_0)/\tau)\Delta} v(t_0) + \frac{1}{\tau} \int_{t_0}^t e^{((t-s)/\tau)\Delta} u(s) \, ds \equiv v_1(t) + v_2(t), \quad t > t_0, \quad (2.5)$$

where $\{e^{t\Delta}\}$ is the heat semigroup. We recall the L^p - L^q estimates for the linear heat equation

$$\|\nabla e^{(t/\tau)\Delta} w\|_{L^q(\mathbb{R}^2)} \leq C t^{1/q-1/p-1/2} \|w\|_{L^p(\mathbb{R}^2)} \quad (2.6)$$

for $t > 0$ with $1 \leq p \leq q \leq \infty$, where $C = C(\tau)$ is a positive constant, see e.g. [10]. In particular we have

$$\|\nabla e^{(t/\tau)\Delta} w\|_{L^\infty(\mathbb{R}^2)} \leq C t^{-1/2} \|w\|_{L^\infty(\mathbb{R}^2)} \quad \text{for } t > 0.$$

Put $t_0 = t/2$ in (2.5). Then it follows that

$$\|\nabla v_1(t)\|_{L^\infty(\mathbb{R}^2)} \leq C t^{-1/2} \|v(t/2)\|_{L^\infty(\mathbb{R}^2)} \leq C t^{-1/2} \|\psi\|_{L^\infty(\mathbb{R}^2)} \quad (2.7)$$

and

$$\|\nabla v_2(t)\|_{L^\infty(\mathbb{R}^2)} \leq C \int_{t/2}^t (t-s)^{-1/2} \|u(s)\|_{L^\infty} \, ds \leq C \|\phi\|_{L^\infty} \int_{t/2}^t t/2 (t-s)^{-1/2} s^{-1} \, ds.$$

By the change of variable $s = t\sigma$, we have

$$\|\nabla v_2(t)\|_{L^\infty(\mathbb{R}^2)} \leq C t^{-1/2} \|\phi\|_{L^\infty} \int_{1/2}^1 (1-\sigma)^{-1/2} \sigma^{-1} \, d\sigma. \quad (2.8)$$

From (2.7) and (2.8) we obtain $\sup_{t>0} t^{1/2} \|\nabla v(t)\|_{L^\infty(\mathbb{R}^2)} < \infty$. By the definition of v it follows that $t^{1/2} \|\nabla v(t)\|_{L^\infty(\mathbb{R}^2)} = \|\nabla \psi\|_{L^\infty(\mathbb{R}^2)}$. Thus we have $\nabla \psi \in L^\infty(\mathbb{R}^2)$. ■

Proof of Proposition 2.1. Put $w(x) = -\phi(x)e^{|\mathbf{x}|^2/4} e^{-\psi(x)} \leq 0$. Then $e^{-|\mathbf{x}|^2/4} e^\psi \nabla w = -\nabla \phi - x\phi/2 + \phi \nabla \psi$. From the first equation of (1.4) we have

$$\nabla \cdot (e^{-|\mathbf{x}|^2/4} e^\psi \nabla w) = 0 \quad \text{or} \quad \Delta w + \nabla b \cdot \nabla w = 0 \quad \text{in } \mathbb{R}^2,$$

where $\nabla b(x) = -x/2 + \nabla\psi(x)$. From Lemma 2.2 we have

$$x \cdot \nabla b(x) = \left(-\frac{|x|^2}{2} + x \cdot \nabla\psi(x) \right) \leq 0$$

for large $|x|$. As a consequence of Lemma 2.1, w must be a constant function. This completes the proof of Proposition 2.1. ■

3. RADIAL SYMMETRY: PROOF OF THEOREM 3

In this section we investigate the radial symmetry of solutions to (1.8) and prove Theorem 3. Namely, we show the following:

PROPOSITION 3.1. *Let $\psi \in C^2(\mathbb{R}^2)$ be a positive solution of (1.8) with (1.9). Then ψ must be radially symmetric about the origin.*

We prepare several lemmas.

LEMMA 3.1. *We have*

$$\psi(x) \leq Ce^{-\min\{\tau,1\}|x|^2/4} \quad \text{for } x \in \mathbb{R}^2 \tag{3.1}$$

with some constant $C > 0$.

Proof. Define

$$Lu = -\Delta u - \frac{\tau}{2} x \cdot \nabla u$$

and put $\kappa_\tau = \min\{1, \tau\}$. Let C be a positive constant and let $v(x) = Ce^{-\kappa_\tau|x|^2/4}$. Then

$$Lv = C\kappa_\tau \left(1 + \frac{(\tau - \kappa_\tau)}{4} |x|^2 \right) e^{-\kappa_\tau|x|^2/4} \geq C\kappa_\tau e^{-\kappa_\tau|x|^2/4}.$$

Since $L\psi = \sigma e^{-|x|^2/4} e^\psi$, if we choose C so large that $C\kappa_\tau > \sigma e^{\|\psi\|_{L^\infty(\mathbb{R}^2)}}$, then $Lv > L\psi$ in \mathbb{R}^2 . Since $v, \psi \rightarrow 0$ as $|x| \rightarrow \infty$, by the maximum principle we have $v \geq \psi$ in \mathbb{R}^2 . This implies (3.1). ■

We define $w(x, t)$ by

$$w(x, t) = t^{-\alpha} \psi \left(\frac{x}{\sqrt{t}} \right), \quad \text{where } \alpha = \frac{\sigma e^{\|\psi\|_{L^\infty(\mathbb{R}^2)}}}{\tau}. \tag{3.2}$$

LEMMA 3.2. (i) For every $T > 0$ we have $\sup_{0 < t < T} w(x, t) \rightarrow 0$ as $|x| \rightarrow \infty$.

(ii) For every $\mu > 0$ we have $\sup_{|x| > \mu} w(x, t) \rightarrow 0$ as $t \rightarrow 0$.

Proof. From Lemma 3.1 we have $|y|^{2\alpha}\psi(y) \rightarrow 0$ as $|y| \rightarrow \infty$, that is, for all $\epsilon > 0$ there exists $R > 0$ such that

$$|y|^{2\alpha}\psi(y) < \epsilon \quad \text{for } |y| \geq R. \tag{3.3}$$

From (3.2) we have

$$|x|^{2\alpha}w(x, t) = \left(\frac{|x|}{\sqrt{t}}\right)^{2\alpha} \psi\left(\frac{x}{\sqrt{t}}\right). \tag{3.4}$$

(i) Fix $T > 0$. From (3.3) and (3.4) it follows that

$$\sup_{0 < t < T} |x|^{2\alpha}w(x, t) < \epsilon \quad \text{for } |x| \geq R\sqrt{T}.$$

Since $\epsilon > 0$ is arbitrary, we obtain $\sup_{0 < t < T} w(x, t) \rightarrow 0$ as $|x| \rightarrow \infty$.

(ii) From (3.3) and (3.4) it follows that

$$\mu^{2\alpha} \sup_{|x| > \mu} w(x, t) \leq \sup_{|x| > \mu} |x|^{2\alpha}w(x, t) < \epsilon \quad \text{for } 0 < t < (\mu/R)^2.$$

Then we have $\sup_{|x| > \mu} w(x, t) \rightarrow 0$ as $t \rightarrow 0$. ■

For $\mu \in \mathbb{R}$ we define T_μ and Σ_μ by

$$T_\mu = \{x = (x_1, x_2) \in \mathbb{R}^2 \mid x_1 = \mu\} \quad \text{and} \quad \Sigma_\mu = \{x \in \mathbb{R}^2 \mid x_1 < \mu\},$$

respectively. For $x \in \mathbb{R}^2$ and $\mu \in \mathbb{R}$ let x^μ be the reflection of x with respect to T_μ , that is, $x^\mu = (2\mu - x_1, x_2)$. It is easy to see that if $\mu > 0$,

$$|x^\mu| > |x| \quad \text{for } x \in \Sigma_\mu \quad \text{and} \quad \{x^\mu : x \in \Sigma_\mu\} = \{x : x_1 > \mu\} \subset \{x : |x| \geq \mu\}.$$

By Lemma 3.2 we have the following:

LEMMA 3.3. (i) For every $T > 0$ we have $\sup_{0 < t < T} w(x^\mu, t) \rightarrow 0$ as $|x| \rightarrow \infty, x \in \Sigma_\mu$.

(ii) For every $\mu > 0$ we have $\sup_{x \in \Sigma_\mu} w(x^\mu, t) = 0$ as $t \rightarrow 0$.

LEMMA 3.4. Let $\mu > 0$. Define $z(x, t) = w(x, t) - w(x^\mu, t)$. Then

$$\tau z_t \geq \Delta z + c_\mu(x, t)z \quad \text{in } \Sigma_\mu \times (0, \infty) \quad \text{and} \quad z = 0 \quad \text{on } T_\mu \times (0, \infty), \tag{3.5}$$

where

$$c_\mu(x, t) = \frac{1}{t} \left(-\alpha\tau + \sigma e^{-|x|^2/(4t)} \int_0^1 e^{s\psi(x/\sqrt{t})+(1-s)\psi(x^\mu/\sqrt{t})} ds \right). \tag{3.6}$$

We have $c_\mu(x, t) \leq 0$ in $\mathbb{R}^2 \times (0, \infty)$.

Proof. By virtue of (3.2) we have

$$\tau w_t = \Delta w - \frac{\alpha\tau}{t} w + \sigma t^{-\alpha-1} e^{-|x|^2/4t} e^{t^\alpha w}.$$

Let $w^\mu(x, t) = w(x^\mu, t)$. Then w^μ satisfies

$$\tau w_t^\mu = \Delta w^\mu - \frac{\alpha\tau}{t} w^\mu + \sigma t^{-\alpha-1} e^{-|x^\mu|^2/4t} e^{t^\alpha w^\mu}.$$

Since $|x^\mu| \geq |x|$, we obtain

$$\tau w_t^\mu \leq \Delta w^\mu - \frac{\alpha\tau}{t} w^\mu + \sigma t^{-\alpha-1} e^{-|x|^2/4t} e^{t^\alpha w^\mu}.$$

Then we obtain $\tau z_t \geq \Delta z + c_\mu z$, where c_μ is the function in (3.6). Since α satisfies $\alpha\tau = \sigma e^{\|\psi\|_{L^\infty(\mathbb{R}^2)}}$, we have $\tau c_\mu(x, t) \leq -\alpha\tau + \sigma e^{\|\psi\|_{L^\infty(\mathbb{R}^2)}} = 0$ for $(x, t) \in \mathbb{R}^2 \times (0, \infty)$. ■

LEMMA 3.5. *Let $\mu > 0$. We have $w(x, t) \geq w(x^\mu, t)$ for $(x, t) \in \Sigma_\mu \times (0, \infty)$.*

Proof. Let $z(x, t) = w(x, t) - w(x^\mu, t)$. We show that $z(x, t) \geq 0$ for $(x, t) \in \Sigma_\mu \times (0, \infty)$. Assume to the contrary that there exists a $(x_0, t_0) \in \Sigma_\mu \times (0, \infty)$ such that $z(x_0, t_0) < 0$. Take $\delta > 0$ so small that $z(x_0, t_0) < -\delta$. By (ii) of Lemma 3.3 we can take $T_0 \in (0, t_0)$ so that $w(x^\mu, T_0) < \delta$ for $x \in \Sigma_\mu$. Then it follows from $w(x, t) > 0$ that

$$z(x, T_0) \geq -\delta \quad \text{for } x \in \Sigma_\mu. \tag{3.7}$$

Fix $T > t_0$. By (i) of Lemma 3.3 we can take $R > |x_0|$ so large that $w(x^\mu, t) < \delta$ for $|x| \geq R, x \in \Sigma_\mu, t \in [T_0, T]$. Then we obtain

$$z(x, t) \geq -\delta \quad \text{for } x \in \Sigma_\mu, |x| \geq R, t \in [T_0, T]. \tag{3.8}$$

Define $Q = \{x \in \Sigma_\mu : |x| < R\}$. Let Γ be a parabolic boundary of $Q \times (T_0, T)$, that is,

$$\Gamma = (Q \times \{T_0\}) \cup (\partial Q \times (T_0, T)).$$

From (3.5), (3.7), and (3.8) we have

$$\tau z_t \geq \Delta z + c(x, t)z \quad \text{in } Q \times (T_0, T) \quad \text{and} \quad z \geq -\delta \quad \text{on } \Gamma.$$

Put $Z = z + \delta$. Because $c_\mu(x, t) \leq 0$, it follows from the above inequality that

$$\tau Z_t \geq \Delta Z + c_\mu(x, t)Z \quad \text{in } Q \times (T_0, T) \quad \text{and} \quad Z \geq 0 \quad \text{on } \Gamma.$$

By the maximum principle [33] we have $Z \geq 0$ on $\bar{Q} \times [T_0, T]$, which implies that

$$z(x, t) \geq -\delta \quad \text{on } \bar{Q} \times [T_0, T]. \tag{3.9}$$

On the other hand, $(x_0, t_0) \in Q \times (T_0, T)$ and $z(x_0, t_0) < -\delta$. This contradicts (3.9). Hence, $z(x, t) \geq 0$ for $(x, t) \in \Sigma_\mu \times (0, \infty)$. ■

Proof of Proposition 3.1. From Lemma 3.5 we have $w(x, t) \geq w(x^\mu, t)$ for $\mu > 0$ and $(x, t) \in \Sigma_\mu \times (0, \infty)$. From the continuity of w we have $w(x, t) \geq w(x^0, t)$ for $(x, t) \in \Sigma_0 \times (0, \infty)$. We can repeat the previous arguments for the negative x_1 -direction to conclude that $w(x, t) \leq w(x^0, t)$ for $(x, t) \in \Sigma_0 \times (0, \infty)$. Hence $w(x, t)$ is symmetric with respect to the plane $x_1 = 0$, which implies that ψ is symmetric with respect to the plane $x_1 = 0$. Since Eq. (1.8) is invariant under the rotation, it follows that ψ is symmetric in every direction. Therefore, ψ is radially symmetric with respect to the origin. ■

Proof of Theorem 3. Let (ϕ, ψ) be a nonnegative solution of (1.4) with $\phi, \psi \in L^\infty(\mathbb{R}^2)$. Then ϕ is given by (1.7) for some constant $\sigma > 0$ from Proposition 2.1. It follows that $\phi > 0$ in \mathbb{R}^2 , and $\phi(x) = O(e^{-|x|^2/4})$ as $|x| \rightarrow \infty$. From the second equation of (1.4), ψ satisfies Eq. (1.8). By the strong maximum principle, $\psi > 0$ in \mathbb{R}^2 .

Assume furthermore that $\psi(x) \rightarrow 0$ as $|x| \rightarrow \infty$. Then, by Proposition 3.1, ψ must be radially symmetric about the origin. Hence $\psi = \psi(r)$, $r = |x|$, satisfies the ordinary differential equation

$$\psi_{rr} + \left(\frac{1}{r} + \frac{\tau}{2} r\right)\psi_r + \sigma e^{-r^2/4} e^\psi = 0$$

or

$$(r e^{\tau r^2/4} \psi_r)_r + \sigma r e^{(\tau-1)r^2/4} e^\psi = 0 \quad \text{for } r > 0.$$

From $\psi_r(0) = 0$, we have

$$r e^{\tau r^2/4} \psi_r = -\sigma \int_0^r s e^{(\tau-1)s^2/4} e^\psi ds < 0 \quad \text{for } r > 0.$$

This implies that $\psi_r(r) < 0$ for $r > 0$. From Lemma 3.1 we obtain $\psi(r) = O(e^{-\min\{\tau, 1\}r^2/4})$ as $r \rightarrow \infty$. This completes the proof of Theorem 3. ■

4. STRUCTURE OF THE SOLUTIONS SET TO (1.8) WITH (1.9)

From Theorem 3 the solution ψ of (1.8) with (1.9) must be radially symmetric about the origin. Then the study of the solutions is reduced to the problem

$$\begin{cases} \psi_{rr} + \left(\frac{1}{r} + \frac{\tau}{2}r\right)\psi_r + \sigma e^{-r^2/4}e^\psi = 0, & r > 0, \\ \psi_r(0) = 0 \quad \text{and} \quad \lim_{r \rightarrow \infty} \psi(r) = 0, \end{cases} \tag{4.1}_\sigma$$

where $\sigma > 0$. In this section we investigate the structure of the pair (σ, ψ) of a parameter and a solution. Define the set \mathcal{C} as

$$\mathcal{C} = \{(\sigma, \psi) : \sigma > 0 \text{ and } \psi \in C^2(0, \infty) \cap C^1[0, \infty) \text{ is a solution of (4.1)}_\sigma\}. \tag{4.2}$$

For $(\sigma, \psi) \in \mathcal{C}$ we have $\psi \in C^2[0, \infty)$ by Lemma 4.1.

PROPOSITION 4.1. *The set \mathcal{C} is written by one-parameter families $(\sigma(s), \psi(r; s))$ on $s \in \mathbb{R}$, that is, $\mathcal{C} = \{(\sigma(s), \psi(r; s)) : s \in \mathbb{R}\}$. The pairs $(\sigma(s), \psi(r; s))$ satisfy the following properties:*

- (i) $s \mapsto (\sigma(s), \psi(\cdot; s)) \in (0, \infty) \times C^2[0, \infty)$ is continuous;
- (ii) $\lim_{s \rightarrow -\infty} \sigma(s) = 0$ and $\lim_{s \rightarrow -\infty} \psi(\cdot; s) = 0$ in $C^2[0, \infty)$;
- (iii) $\lim_{s \rightarrow \infty} \|\psi(\cdot; s)\|_{L^\infty[0, \infty)} = \lim_{s \rightarrow \infty} \psi(0; s) = \infty$.

First we show the following:

LEMMA 4.1. *Let $\psi \in C^2(0, \infty) \cap C^1[0, \infty)$ be a solution to (4.1) $_\sigma$. Then $\psi \in C^2[0, \infty)$ and $\sup_{r \geq 0} \psi(r) = \psi(0)$. Moreover we have*

$$\sup_{r \geq 0} |\psi_r(r)| \leq \pi^{1/2} \sigma e^{\psi(0)} \quad \text{and} \quad \sup_{r \geq 0} |\psi_{rr}(r)| \leq \frac{3 + 2\tau}{2} \sigma e^{\psi(0)}. \tag{4.3}$$

Proof. From (4.1) $_\sigma$, we have $(re^{\tau r^2/4}\psi_r)_r + \sigma re^{(\tau-1)r^2/4}e^\psi = 0$ for $r > 0$. From $\psi_r(0) = 0$, it follows that

$$\psi_r(r) = -\frac{\sigma}{r} e^{-\tau r^2/4} \int_0^r \xi e^{(\tau-1)\xi^2/4} e^{\psi(\xi)} d\xi. \tag{4.4}$$

By using L'Hospital's rule we obtain

$$\lim_{r \rightarrow 0} \frac{\psi_r(r)}{r} = \lim_{r \rightarrow 0} -\frac{\sigma}{r^2 e^{\tau^2/4}} \int_0^r \xi e^{(\tau-1)\xi^2/4} e^{\psi(\xi)} d\xi = -\frac{\sigma e^{\psi(0)}}{2},$$

which implies $\psi \in C^2[0, \infty)$. Since $\psi_r(r) < 0$ for $r > 0$ from (4.4), we have $\sup_{r \geq 0} \psi(r) = \psi(0)$.

From (4.4) we have

$$|\psi_r(r)| \leq \left(\frac{1}{r} \int_0^r \xi e^{-\xi^2/4} d\xi \right) \sigma e^{\psi(0)}. \quad (4.5)$$

We see that $(1/r) \int_0^r \xi e^{-\xi^2/4} d\xi \leq \int_0^\infty e^{-\xi^2/4} d\xi = \pi^{1/2}$. Then the left-hand side of (4.3) holds.

From the equation in (4.1) _{σ} we have

$$|\psi_{rr}(r)| \leq \left(\frac{1}{r} + \frac{\tau}{2} r \right) |\psi_r(r)| + \sigma e^{-r^2/4} e^{\psi(r)} \leq \left(\frac{1}{r} + \frac{\tau}{2} r \right) |\psi_r(r)| + \sigma e^{\psi(0)}.$$

We note here that

$$\left(\frac{1}{r} + \frac{\tau}{2} r \right) \frac{1}{r} \int_0^r \xi e^{-\xi^2/4} d\xi \leq \frac{1}{r^2} \int_0^r \xi d\xi + \frac{\tau}{2} \int_0^\infty \xi e^{-\xi^2/4} d\xi = \frac{1}{2} + \tau. \quad (4.6)$$

It follows from (4.5) and (4.6) that

$$\left(\frac{1}{r} + \frac{\tau}{2} r \right) |\psi_r(r)| \leq \frac{1+2\tau}{2} \sigma e^{\psi(0)}.$$

Therefore, we obtain the right-hand side of (4.3). This completes the proof of Lemma 4.1. ■

To prove Proposition 4.1 we consider the initial value problem

$$\begin{cases} w_{rr} + \left(\frac{1}{r} + \frac{\tau}{2} r \right) w_r + e^{-r^2/4} e^w = 0, & r > 0, \\ w_r(0) = 0 \quad \text{and} \quad w(0) = s, \end{cases} \quad (4.7)_s$$

where $s \in \mathbb{R}$. We denote by $w(r; s)$ the solution of problem (4.7)_s. We easily see that $w(r; s)$ and $w_r(r; s)$ satisfy, respectively,

$$w(r; s) = s - \int_0^r \frac{1}{\xi} e^{-\tau \xi^2/4} \left(\int_0^\xi \eta e^{(\tau-1)\eta^2/4} e^{w(\eta; s)} d\eta \right) d\xi \tag{4.8}$$

and

$$w_r(r; s) = -\frac{1}{r} e^{-\tau r^2/4} \int_0^r \xi e^{(\tau-1)\xi^2/4} e^{w(\xi; s)} d\xi. \tag{4.9}$$

Define $I(\tau)$ as

$$I(\tau) = \int_0^\infty \frac{1}{\xi} e^{-\tau \xi^2/4} \left(\int_0^\xi \eta e^{(\tau-1)\eta^2/4} d\eta \right) d\xi.$$

From [25, Lemma 1] it follows that $I(\tau) = (\log \tau)/(\tau - 1)$ if $\tau \neq 1$, $I(\tau) = 1$ if $\tau = 1$. We easily obtain $w_r(r; s) < 0$ for $r > 0$ and $w(r; s) \geq s - e^s I(\tau)$ for $r \geq 0$ (see [25, Lemma 2]). Then $\lim_{r \rightarrow \infty} w(r; s)$ exists and is a finite value. Put $t(s) = \lim_{r \rightarrow \infty} w(r; s)$.

LEMMA 4.2. *For $s \in \mathbb{R}$, let $\psi(r; s) = w(r; s) - t(s)$. Then $\psi(r; s)$ is a solution to (4.1)_σ with $\sigma = e^{t(s)}$. Conversely, let $\psi(r)$ be a solution of (4.1)_σ. Then, for some $s \in \mathbb{R}$, $\psi(r) = \psi(r; s)$ and $\sigma = e^{t(s)}$.*

Proof. It is clear that $\psi(r; s)$ is a solution to (4.1)_σ with $\sigma = e^{t(s)}$. Conversely, let $\psi(r)$ be a solution of (4.1)_σ, and let $w(r) = \psi(r) + \log \sigma$. Then $w(r)$ satisfies (4.7)_s with $s = \psi(0) + \log \sigma$. By the uniqueness we obtain $w(r) = w(r; s)$ with $s = \psi(0) + \log \sigma$. We have $\lim_{r \rightarrow \infty} w(r; s) = \lim_{r \rightarrow \infty} w(r) = \log \sigma$. Then $t(s) = \log \sigma$, that is, $\sigma = e^{t(s)}$. Hence we obtain $\psi(r) = w(r) - \log \sigma = w(r; s) - t(s)$, which implies $\psi(r) = \psi(r; s)$. ■

From [25, Lemma 5(ii)] it follows that, for $s_1, s_2 \in \mathbb{R}$,

$$\sup_{r \geq 0} |w(r; s_1) - w(r; s_2)| \leq C_1 |s_1 - s_2|, \tag{4.10}$$

where $C_1 = \exp(e^m I(\tau))$ and $m = \max\{s_1, s_2\}$. Moreover, we have the following:

LEMMA 4.3. *Let $s_1, s_2 \in \mathbb{R}$, and let $m = \max\{s_1, s_2\}$. Then we have*

- (i) $\sup_{r \geq 0} |w_r(r; s_1) - w_r(r; s_2)| \leq C_2 |s_1 - s_2|$, where $C_2 = \pi^{1/2} e^m C_1$;
- (ii) $\sup_{r \geq 0} |w_{rr}(r; s_1) - w_{rr}(r; s_2)| \leq C_3 |s_1 - s_2|$,
where $C_3 = (3 + 2\tau)e^m C_1/2$.

Proof. From (4.9) we have

$$|w_r(r; s_1) - w_r(r; s_2)| \leq \frac{1}{r} e^{-\tau r^2/4} \int_0^r \xi e^{(\tau-1)\xi^2/4} |e^{w(\xi; s_1)} - e^{w(\xi; s_2)}| d\xi.$$

Note that $|e^{w(t; s_1)} - e^{w(t; s_2)}| \leq e^m |w(t; s_1) - w(t; s_2)|$ with $m = \max\{s_1, s_2\}$. Then from (4.10) we have $|e^{w(t; s_1)} - e^{w(t; s_2)}| \leq C_1 e^m |s_1 - s_2|$. Then it follows that

$$|w_r(r; s_1) - w_r(r; s_2)| \leq C_1 e^m |s_1 - s_2| \left(\frac{1}{r} \int_0^r \xi e^{-\xi^2/4} d\xi \right). \tag{4.11}$$

From $(1/r) \int_0^r \xi e^{-\xi^2/4} d\xi \leq \int_0^\infty e^{-\xi^2/4} d\xi = \pi^{1/2}$, we obtain (i).

From (4.7)_s we see that $w_{rr}(r; s) = -(1/r + \tau r/2)w_r(r; s) - e^{-r^2/4}e^{w(r; s)}$. Then we have

$$|w_{rr}(r; s_1) - w_{rr}(r; s_2)| \leq \left(\frac{1}{r} + \frac{\tau}{2} r \right) |w_r(r; s_1) - w_r(r; s_2)| + e^m |w(r; s_1) - w(r; s_2)|.$$

Then from (4.11) and (4.6) we obtain

$$\left(\frac{1}{r} + \frac{\tau}{2} r \right) |w_r(r; s_1) - w_r(r; s_2)| \leq \frac{1 + 2\tau}{2} C_1 e^m |s_1 - s_2|.$$

Therefore we obtain (ii). ■

LEMMA 4.4. *Let $s_1, s_2 \in \mathbb{R}$, and let $m = \max\{s_1, s_2\}$. Then we have*

- (i) $|t(s_1) - t(s_2)| \leq C_1 |s_1 - s_2|$, where $C_1 = \exp(e^m I(\tau))$;
- (ii) $\lim_{s \rightarrow -\infty} (s - t(s)) = 0$;
- (iii) $\sup_{s \in \mathbb{R}} t(s) \leq -\log I(\tau)$.

Proof. Letting $r \rightarrow \infty$ in (4.10), we have (i). Since $w(r; s) < s$ for $r > 0$, it follows from (4.8) that

$$0 < s - w(r; s) \leq e^s \int_0^r \frac{1}{\xi} e^{-\tau\xi^2/4} \left(\int_0^\xi \eta e^{(\tau-1)\eta^2/4} d\eta \right) d\xi.$$

Letting $r \rightarrow \infty$ we have $0 < s - t(s) \leq e^s I(\tau)$ for $s \in \mathbb{R}$. This implies that (ii) holds.

Since $w(r; s)$ is decreasing in $r > 0$, it follows from (4.9) that

$$w_r(r; s) \leq -\frac{1}{r} e^{w(r; s)} e^{-\tau r^2/4} \int_0^r \xi e^{(\tau-1)\xi^2/4} d\xi.$$

Then we obtain

$$\frac{d}{dr}(-e^{-w(r;s)}) \leq -\frac{1}{r} e^{-\tau r^2/4} \int_0^r \xi e^{(\tau-1)\xi^2/4} d\xi.$$

Integrating the above on $[0, \infty)$ we have $e^{-t(s)} - e^{-s} \geq I(\tau)$ or $e^{-t(s)} \geq I(\tau)$. This implies that (iii) holds. ■

Proof of Proposition 4.1. By Lemma 4.2 we have $\mathcal{C} = \{(\sigma(s), \psi(\cdot; s)) : s \in \mathbb{R}\}$, where $\sigma(s) = e^{t(s)}$ and $\psi(r; s) = w(r; s) - t(s)$. We see that $w(\cdot; s) \in C^2[0, \infty)$ and $t(s) \in \mathbb{R}$ are continuous for $s \in \mathbb{R}$ by Lemmas 4.3 and 4.4(i), respectively. Thus (i) holds.

By Lemma 4.4(ii) we have $\sigma(s) = e^{t(s)} \rightarrow 0$ and $\psi(0; s) = s - t(s) \rightarrow 0$ as $s \rightarrow -\infty$. Then, by Lemma 4.1 we conclude that $\psi(\cdot; s) \rightarrow 0$ in $C^2[0, \infty)$ as $s \rightarrow -\infty$. Thus (ii) holds.

From Lemma 4.1 we have $\|\psi(\cdot; s)\|_{L^\infty[0, \infty)} = \psi(0; s)$. From Lemma 4.4(iii) we have $\lim_{s \rightarrow \infty} \psi(0; s) = \lim_{s \rightarrow \infty} (s - t(s)) \geq \lim_{s \rightarrow \infty} (s + \log I(\tau)) = \infty$. Thus (iii) holds. This completes the proof of Proposition 4.1. ■

5. BLOWUP ANALYSIS TO SELF-SIMILAR SOLUTIONS

This section is concerned with case (iii) of Theorem 2. We study the asymptotic behavior of sequences $\{(\phi_k, \psi_k)\} \subset \mathcal{S}$ satisfying $\|\psi_k\|_{L^\infty(\mathbb{R}^2)} \rightarrow \infty$ as $k \rightarrow \infty$. We show the following:

PROPOSITION 5.1. *Let $(\phi_k, \psi_k) \in \mathcal{S}$, and let $\lambda_k = \|\phi_k\|_{L^1(\mathbb{R}^2)}$. Assume that*

$$\|\psi_k\|_{L^\infty(\mathbb{R}^2)} \rightarrow \infty \quad \text{as } k \rightarrow \infty \tag{5.1}$$

and that $\{\lambda_k\}$ is bounded. Then there exists a subsequence, which we call again (ψ_k, ϕ_k) and λ_k , satisfying $\lambda_k \rightarrow 8\pi$ as $k \rightarrow \infty$ and

$$\phi_k(x) dx \rightarrow 8\pi \delta_0(dx) \quad \text{as } k \rightarrow \infty \tag{5.2}$$

in the sense of measure, where $\delta_0(dx)$ is Dirac's delta function with the support in origin.

In order to prove Proposition 5.1 we make use of Theorems A and 4 in Section 1. We also need the following result by Brezis and Merle [2].

THEOREM B (Brezis and Merle [2]). *Assume $\{u_k\}$ is a sequence of solutions of (1.11) such that*

$$\|V_k\|_{L^\infty(\Omega)} \leq C, \quad \|u_k^+\|_{L^1(\Omega)} \leq C, \quad \text{and} \quad \int_{\Omega} V_k e^{u_k} dx < 4\pi,$$

for some constant $C > 0$, where $u^+ = \max\{u, 0\}$. Then $\{u_k^+\}$ is bounded in $L^\infty_{\text{loc}}(\Omega)$.

Now we prepare several lemmas.

LEMMA 5.1. *Assume that $f \in C(\mathbb{R}^2) \cap L^1(\mathbb{R}^2)$. Let $w \in C^2(\mathbb{R}^2) \cap L^1(\mathbb{R}^2)$ be a solution of*

$$-\Delta w - \frac{\tau}{2} x \cdot \nabla w = f \quad \text{for } x \in \mathbb{R}^2. \tag{5.3}$$

Then we have $\|w\|_{L^1(\mathbb{R}^2)} + \|\nabla w\|_{L^1(\mathbb{R}^2)} \leq C\|f\|_{L^1(\mathbb{R}^2)}$, for some positive constant C .

Proof. Define W and F , respectively, as

$$W(x, t) = w\left(\frac{x}{\sqrt{t}}\right) \quad \text{and} \quad F(x, t) = \frac{1}{t} f\left(\frac{x}{\sqrt{t}}\right).$$

Then W and F satisfy

$$\|W(\cdot, t)\|_{L^1(\mathbb{R}^2)} = t\|w\|_{L^1(\mathbb{R}^2)} \quad \text{and} \quad \|F(\cdot, t)\|_{L^1(\mathbb{R}^2)} = \|f\|_{L^1(\mathbb{R}^2)} \tag{5.4}$$

for $t > 0$. Furthermore, from (5.3) we have $\tau W_t = \Delta W + F$ in $\mathbb{R}^2 \times (0, \infty)$. Since $W \rightarrow 0$ in $L^1(\mathbb{R}^2)$ as $t \rightarrow 0$ from (5.4), we obtain

$$W(x, t) = \frac{1}{\tau} \int_0^t e^{((t-s)/\tau)\Delta} F(\cdot, s) ds.$$

Then it follows from (5.4) that

$$t\|w\|_{L^1(\mathbb{R}^2)} = \|W(\cdot, t)\|_{L^1(\mathbb{R}^2)} \leq \frac{1}{\tau} \int_0^t \|F(\cdot, s)\|_{L^1(\mathbb{R}^2)} ds \leq \frac{t}{\tau} \|f\|_{L^1(\mathbb{R}^2)}.$$

Therefore, we obtain $\|w\|_{L^1(\mathbb{R}^2)} \leq \tau^{-1} \|f\|_{L^1(\mathbb{R}^2)}$.

Next we show $\|\nabla w\|_{L^1(\mathbb{R}^2)} \leq C\|f\|_{L^1(\mathbb{R}^2)}$. By the L^p - L^q estimates (2.6) with $p = q = 1$ we have

$$\|\nabla e^{((t-s)/\tau)\Delta} F(\cdot, s)\|_{L^1(\mathbb{R}^2)} \leq C(t-s)^{-1/2} \|F(\cdot, s)\|_{L^1(\mathbb{R}^2)} = C(t-s)^{-1/2} \|f\|_{L^1(\mathbb{R}^2)}.$$

Then we obtain

$$\|\nabla W(\cdot, t)\|_{L^1(\mathbb{R}^2)} \leq \frac{1}{\tau} \int_0^t \|\nabla e^{((t-s)/\tau)\Delta} F(\cdot, s)\|_{L^1(\mathbb{R}^2)} ds \leq Ct^{1/2} \|f\|_{L^1(\mathbb{R}^2)}.$$

By the definition of W it follows that $\|\nabla W(\cdot, t)\|_{L^1(\mathbb{R}^2)} = t^{1/2} \|\nabla w\|_{L^1(\mathbb{R}^2)}$. Therefore, we conclude that $\|\nabla w\|_{L^1(\mathbb{R}^2)} \leq C\|f\|_{L^1(\mathbb{R}^2)}$. This completes the proof of Lemma 5.1. ■

Let $(\phi_k, \psi_k) \in \mathcal{S}$, and let $\lambda_k = \|\phi_k\|_{L^1(\mathbb{R}^2)}$. Then (λ_k, ψ_k) solves (1.10), that is,

$$\Delta\psi_k + \frac{\tau}{2} x \cdot \nabla\psi_k + \lambda_k e^{-|x|^2/4} e^{\psi_k} \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy = 0 \quad \text{for } x \in \mathbb{R}^2. \tag{5.5}$$

From Theorem 3, we have $\psi_k \in L^1(\mathbb{R}^2)$, $\psi_k = \psi_k(r)$, $r = |x|$, and $\partial\psi_k/\partial r < 0$ for $r > 0$. Assume that (5.1) holds. Then $\|\psi_k\|_{L^\infty(\mathbb{R}^2)} = \psi_k(0) \rightarrow \infty$ as $k \rightarrow \infty$. We always use B_r to denote a ball of radius r centered at origin, that is, $B_r = \{x \in \mathbb{R}^2 : |x| < r\}$.

LEMMA 5.2. (i) We have $\|\psi_k\|_{L^1(\mathbb{R}^2)} + \|\nabla\psi_k\|_{L^1(\mathbb{R}^2)} = O(1)$ as $k \rightarrow \infty$.

(ii) For all $r > 0$ we have $\sup_k \|\psi_k\|_{L^\infty(\mathbb{R}^2 \setminus B_r)} < \infty$.

Proof. (i) Put

$$f_k(x) = \lambda_k e^{-|x|^2/4} e^{\psi_k(x)} \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy.$$

Then $f_k \in C^2(\mathbb{R}^2) \cap L^1(\mathbb{R}^2)$. We have $\psi_k \in L^1(\mathbb{R}^2)$ and

$$-\Delta\psi_k - \frac{\tau}{2} x \cdot \nabla\psi_k = f_k \quad \text{for } x \in \mathbb{R}^2.$$

By Lemma 5.1 we obtain $\|\psi_k\|_{L^1(\mathbb{R}^2)} + \|\nabla\psi_k\|_{L^1(\mathbb{R}^2)} \leq C\|f_k\|_{L^1(\mathbb{R}^2)}$ for some constant $C > 0$. Since $\|f_k\|_{L^1(\mathbb{R}^2)} = \lambda_k = O(1)$ and as $k \rightarrow \infty$, the assertion of (i) holds.

(iii) Assume to the contrary that $\sup_k \|\psi_k\|_{L^\infty(\mathbb{R}^2 \setminus B_{r_0})} = \infty$ for some $r_0 > 0$. Since $\psi_k(r)$ is decreasing in $r > 0$, there exists a subsequence, which we call again $\{\psi_k\}$, such that $\inf_{y \in B_{r_0}} \psi_k(y) \rightarrow \infty$ as $k \rightarrow \infty$. Then $\|\psi_k\|_{L^1(\mathbb{R}^2)} \rightarrow \infty$ as $k \rightarrow \infty$, which contradicts assertion (i). ■

Take $R > 0$. Let g_k be a unique solution of the problem

$$-\Delta g_k = \frac{\tau}{2} x \cdot \nabla\psi_k \quad \text{in } B_R, \quad g_k = 0 \quad \text{on } \partial B_R.$$

LEMMA 5.3. *We have $\|g_k\|_{L^\infty(B_R)} = O(1)$ and $\|\nabla g_k\|_{L^\infty(B_R)} = O(1)$ as $k \rightarrow \infty$.*

Proof. We have $g_k = g_k(r)$, $r = |x|$, since $\psi_k = \psi_k(r)$. We see that $g_k(r)$ satisfies

$$-(rg'_k)' = \frac{\tau}{2} r^2 \psi_k, \quad 0 < r < R, \quad g'_k(0) = g_k(R) = 0,$$

where $' = d/dr$. We will show that

$$\|g_k\|_{L^\infty[0,R]} = O(1), \quad \|g'_k\|_{L^\infty[0,R]} = O(1) \quad \text{as } k \rightarrow \infty. \tag{5.6}$$

By integrating the equation above, we obtain

$$-rg'_k(r) = \frac{\tau}{2} \int_0^r s^2 \psi'_k(s) ds.$$

Then it follows that

$$|g'_k(r)| \leq \frac{\tau}{2r} \int_0^r s^2 |\psi'_k(s)| ds \leq \frac{\tau}{2} \int_0^r s |\psi'_k(s)| ds \quad \text{for } 0 \leq r \leq R.$$

Thus we obtain

$$\|g'_k\|_{L^\infty[0,R]} \leq \frac{\tau}{2} \int_0^R s |\psi'_k(s)| ds. \tag{5.7}$$

We note that $\int_r^R g'_k(s) ds = g_k(R) - g_k(r) = -g_k(r)$. Then

$$|g_k(r)| \leq \int_0^R |g'_k(s)| ds \leq R \|g'_k\|_{L^\infty[0,R]} \quad \text{for } 0 \leq r \leq R.$$

From (5.7) we obtain

$$\|g_k\|_{L^\infty[0,R]} \leq \frac{\tau R}{2} \int_0^R s |\psi'_k(s)| ds. \tag{5.8}$$

By Lemma 5.2(i) we have

$$2\pi \int_0^R s |\psi'_k(s)| ds = \|\nabla \psi_k\|_{L^1(B_R)} \leq \|\nabla \psi_k\|_{L^1(\mathbb{R}^2)} = O(1) \quad \text{as } k \rightarrow \infty.$$

From (5.7) and (5.8) we obtain (5.6). This completes the proof of Lemma 5.3. ■

Now define v_k as

$$v_k(x) = \psi_k(x) - g_k(x) - \log\left(\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy\right). \tag{5.9}$$

It follows from (5.5) that

$$-\Delta v_k = -\Delta \psi_k - \frac{\tau}{2} x \cdot \nabla \psi_k = \lambda_k e^{-|x|^2/4} e^{g_k} e^{v_k} \quad \text{for } x \in B_R. \tag{5.10}$$

Then we have

$$-\Delta v_k = V_k(x) e^{v_k} \quad \text{in } B_R, \tag{5.11}$$

where $V_k(x) = \lambda_k e^{-|x|^2/4} e^{g_k}$. Since $\{\lambda_k\}$ is bounded and by Lemma 5.3, we have $0 \leq V_k(x) \leq C_0$ and $\|\nabla V_k\|_{L^\infty(B_R)} \leq C_1$ for some constants C_0 and C_1 . Since v_k is radial symmetry and satisfies $-\Delta v_k \geq 0$ in B_R , $v_k(r)$ is nonincreasing in $r \in (0, R)$ by the maximum principle.

LEMMA 5.4. *There exists a subsequence, which we call again $\{v_k\}$, such that $v_k(0) \rightarrow \infty$ and $v_k(x) \rightarrow -\infty$ uniformly on compact subset of $B_R \setminus \{0\}$ as $k \rightarrow \infty$. Moreover,*

$$\int_{B_R} V_k e^{v_k} dx \rightarrow 8\pi \quad \text{as } k \rightarrow \infty \tag{5.12}$$

and

$$\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy \rightarrow \infty \quad \text{as } k \rightarrow \infty. \tag{5.13}$$

Proof. We see that

$$\int_{B_R} e^{v_k(y)} dy \leq e^{\|g_k\|_{L^\infty(B_R)}} \int_{B_R} e^{\psi_k(y)} dy \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4 + \psi_k(y)} dy \leq C$$

for some constant $C > 0$. Hence, by applying Theorem A, there exists a subsequence (still denoted by $\{v_k\}$) satisfying one of the alternatives (i)–(iii) in Theorem A.

Assume that the first alternative (i) holds. Since $\{v_k\}$ and $\{g_k\}$ are bounded in $L^\infty_{\text{loc}}(B_R)$ and $\psi_k(0) \rightarrow \infty$ as $k \rightarrow \infty$, it follows from (5.9) that

$$\log\left(\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy\right) = \psi_k(0) - g_k(0) - v_k(0) \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

Let $y_0 \in B_R \setminus \{0\}$. Then from (5.9) we have $\psi_k(y_0) \rightarrow \infty$ as $k \rightarrow \infty$. This contradicts Lemma 5.2(ii).

Assume that the second alternative (ii) holds. Since $v_k(r)$ is nonincreasing in r , we have $v_k \rightarrow -\infty$ uniformly on B_R . Then

$$\int_{B_R} e^{v_k} dx \rightarrow 0 \quad \text{as } k \rightarrow \infty. \tag{5.14}$$

Put

$$w_k = \psi_k - g_k \quad \text{and} \quad W_k(x) = V_k(x) \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy.$$

Then we have $-\Delta w_k = W_k e^{w_k}$ in B_R . Because $\psi_k \geq 0$, we have

$$W_k(x) \leq V_k(x) \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4} dy \leq C$$

for some constant $C > 0$. We find that $\|w_k\|_{L^1(B_R)} \leq \|\psi_k\|_{L^1(B_R)} + \|g_k\|_{L^1(B_R)} = O(1)$ as $k \rightarrow \infty$ by Lemmas 5.2 and 5.3. It follows from (5.14) that

$$\int_{B_R} W_k(y) e^{w_k(y)} dy = \int_{B_R} V_k(y) e^{v_k(y)} dy \leq C_0 \int_{B_R} e^{v_k(y)} dy \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Hence, by applying Theorem B we obtain $\|w_k^+\|_{L^\infty(B_R)} = O(1)$ as $k \rightarrow \infty$. This contradicts $w_k(0) = \psi_k(0) - g_k(0) \rightarrow \infty$ as $k \rightarrow \infty$.

Therefore, the third alternative (iii) must hold. By Lemma 5.2(ii) we have the blowup set $\mathcal{B} = \{0\}$. Then $v_k(0) \rightarrow \infty$ and $v_k(x) \rightarrow -\infty$ uniformly on compact subset of $B_R \setminus \{0\}$. Moreover,

$$\int_{B_R} V_k e^{v_k} dx \rightarrow \alpha \quad \text{as } k \rightarrow \infty \tag{5.15}$$

for some $\alpha \geq 4\pi$. Since v_k is radial symmetry, we have $\max_{\partial B_R} v_k - \min_{\partial B_R} v_k = 0$. By applying Theorem 4, we obtain $\alpha = 8\pi$ in (5.15).

Let $x_0 \in B_R \setminus \{0\}$. From $v_k(x_0) \rightarrow -\infty$ as $k \rightarrow \infty$ we have

$$\log \left(\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy \right) = \psi_k(x_0) - g_k(x_0) - v_k(x_0) \rightarrow \infty \quad \text{as } k \rightarrow \infty,$$

which implies that (5.13) holds. ■

Proof of Proposition 5.1. Let $\{v_k\}$ be a subsequence obtained in Lemma 5.4. First we verify that, for all $r > 0$,

$$\int_{\mathbb{R}^2 \setminus B_r} V_k e^{v_k} dy \rightarrow 0 \quad \text{as } k \rightarrow \infty. \tag{5.16}$$

From Lemma 5.2(ii) there exists a constant $M = M(r) > 0$ such that $|\psi_k(x)| \leq M$ for $|x| \geq r$. Since

$$\int_{\mathbb{R}^2 \setminus B_r} V_k(y) e^{v_k(y)} dy = \frac{\lambda_k \int_{\mathbb{R}^2 \setminus B_r} e^{-|y|^2/4} e^{\psi_k(y)} dy}{\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy} \leq \frac{\lambda_k e^M \int_{\mathbb{R}^2 \setminus B_r} e^{-|y|^2/4} dy}{\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi_k(y)} dy},$$

it follows from (5.13) that (5.16) holds.

From (5.10), (5.11), and the second equation of (1.4) we have

$$V_k e^{v_k} = -\Delta v_k = -\Delta \psi_k - \frac{\tau}{2} x \cdot \nabla \psi_k = \phi_k.$$

From (5.12) and (5.16) we have

$$\begin{aligned} \lambda_k &= \|\phi_k\|_{L^1(\mathbb{R}^2)} = \int_{\mathbb{R}^2} V_k e^{v_k} dy \\ &= \int_{B_R} V_k e^{v_k} dy + \int_{\mathbb{R}^2 \setminus B_R} V_k e^{v_k} dy \rightarrow 8\pi \quad \text{as } k \rightarrow \infty. \end{aligned}$$

Thus $\lambda_k \rightarrow 8\pi$ as $k \rightarrow \infty$. Since $\{\phi_k\}$ is bounded in $L^1(\mathbb{R}^2)$, we may extract a subsequence, which we call again $\{\phi_k\}$, such that ϕ_k converges in the sense of measures on \mathbb{R}^2 to some nonnegative bounded measure μ , i.e.

$$\int_{\mathbb{R}^2} \phi_k(x) \eta dx \rightarrow \int_{\mathbb{R}^2} \eta d\mu$$

for every $\eta \in C(\mathbb{R}^2)$ with compact support. From (5.16) we have $\int_{\mathbb{R}^2 \setminus B_r} \phi_k(x) dx \rightarrow 0$ as $k \rightarrow \infty$ for every $r > 0$. Then $\phi_k \rightarrow 0$ in $L^1_{\text{loc}}(\mathbb{R}^2 \setminus \{0\})$ and hence μ is supported on $\{0\}$. Thus we obtain $d\mu = \alpha \delta_0(dx)$ with $\alpha = 8\pi$, which implies that (5.2) holds. This completes the proof of Proposition 5.1. ■

6. L^1 -NORMS OF SELF-SIMILAR SOLUTIONS

This section is concerned with case (iv) of Theorem 2 and we investigate the upper bounds of $\|\phi\|_{L^1(\mathbb{R}^2)}$ for $(\phi, \psi) \in \mathcal{S}$.

PROPOSITION 6.1. *Let $(\phi, \psi) \in \mathcal{S}$. Then*

$$\|\phi\|_{L^1(\mathbb{R}^2)} \leq \max\left\{\frac{4}{3} \pi^3, \frac{4}{3} \pi^3 \tau^2\right\}.$$

Moreover, if $0 < \tau \leq 1/2$ then $\|\phi\|_{L^1(\mathbb{R}^2)} < 8\pi$.

We prove Proposition 6.1, following the idea of Biler [1]. By Theorem 1 the solution $(\phi, \psi) \in \mathcal{S}$ must be radially symmetric about the origin. Define

Φ and Ψ , respectively, as

$$\Phi(s) = \frac{1}{2} \int_0^s \phi(\sqrt{t}) dt \quad \text{and} \quad \Psi(s) = \frac{1}{2} \int_0^s \psi(\sqrt{t}) dt.$$

First we show the following:

LEMMA 6.1. *We have $\|\phi\|_{L^1(\mathbb{R}^2)} = 2\pi \lim_{s \rightarrow \infty} \Phi(s)$. Moreover, (Φ, Ψ) solves*

$$\begin{cases} \Phi'' + \frac{1}{4} \Phi' - 2\Phi'\Psi'' = 0, \\ 4s\Psi'' + \tau s\Psi' - \tau\Psi + \Phi = 0 \end{cases} \tag{6.1}$$

for $s > 0$, where $' = d/ds$.

Proof. We see that

$$\int_{\mathbb{R}^2} \phi(|y|) dy = 2\pi \int_0^\infty r\phi(r) dr = 2\pi \left(\frac{1}{2} \int_0^\infty \phi(\sqrt{t}) dt \right),$$

which implies $\|\phi\|_{L^1(\mathbb{R}^2)} = 2\pi \lim_{s \rightarrow \infty} \Phi(s)$.

Define u and v as

$$u(r, t) = \frac{1}{t} \phi\left(\frac{r}{\sqrt{t}}\right) \quad \text{and} \quad v(r, t) = \psi\left(\frac{r}{\sqrt{t}}\right),$$

respectively. Put U and V as

$$U(r, t) = \int_0^r su(s, t) ds \quad \text{and} \quad V(r, t) = \int_0^r sv(s, t) ds.$$

Then, by the change of variables, we obtain

$$U(r, t) = \frac{1}{2} \int_0^{r^2/t} \phi(\sqrt{s}) ds \quad \text{and} \quad V(r, t) = \frac{t}{2} \int_0^{r^2/t} \psi(\sqrt{s}) ds.$$

By the definition of Ψ and Φ we have

$$U(r, t) = \Phi\left(\frac{r^2}{t}\right) \quad \text{and} \quad V(r, t) = t\Psi\left(\frac{r^2}{t}\right). \tag{6.2}$$

Now we verify that (U, V) satisfies

$$\begin{cases} U_t = r(r^{-1}U_r)_r - U_r(r^{-1}V_r)_r, \\ \tau V_t = r(r^{-1}V_r)_r + U \end{cases} \tag{6.3}$$

for $(r, t) \in [0, \infty) \times (0, \infty)$. Since (u, v) solves (1.1), we see that

$$ru_t = (ru_r)_r - ru_rv_r - u(rv_r)_r \quad \text{and} \quad \tau rv_t = (rv_r)_r + ru.$$

Then we obtain

$$\int_0^r su_t(s, t) ds = ru_r - ruv_r \quad \text{and} \quad \tau \int_0^r sv_t(s, t) ds = rv_r + \int_0^r su(s, t) ds.$$

Thus we obtain (6.3). By virtue of (6.2) we have (6.1). ■

LEMMA 6.2. *We have*

$$-s\Psi''(s) = \frac{1}{4} e^{-\tau s/4} \int_0^s e^{\tau t/4} \Phi'(t) dt > 0 \quad \text{for } s > 0. \tag{6.4}$$

Proof. Put $W(s) = -4s\Psi''(s)$. From the second equation of (6.1), we have

$$\Phi' = (-4s\Psi'')' - \tau s\Psi'' = W' + \frac{\tau}{4} W.$$

Since $s\Psi''(s) = \sqrt{s}\psi'(\sqrt{s})/4$, we have $W(0) = \lim_{s \rightarrow 0} W(s) = 0$. Then we obtain

$$W(s) = e^{-\tau s/4} \int_0^s e^{\tau t/4} \Phi'(t) dt.$$

Since $\Phi'(s) = \phi(\sqrt{s})/2 > 0$, we obtain the assertion. ■

LEMMA 6.3. *We have $s\Psi''(s) \rightarrow 0$ as $s \rightarrow \infty$ and, for $s > 0$,*

$$0 < \Psi(s) - s\Psi'(s) \leq \begin{cases} \frac{\tau}{4} \int_0^s \frac{t}{e^{\tau t/4} - 1} dt < s & \text{if } 0 < \tau \leq 1, \\ \frac{\tau}{4} \int_0^s \frac{t}{e^{t/4} - 1} dt & \text{if } \tau > 1. \end{cases}$$

Proof. From the first equation of (6.1) and (6.4) we have

$$\Phi'' + \frac{1}{4}\Phi' + \frac{1}{2s}e^{-\tau s/4}\Phi' \int_0^s e^{\tau t/4}\Phi'(t) dt = 0.$$

We note that $\Phi'(s) = \phi(\sqrt{s})/2 > 0$. Then, for the case $0 < \tau \leq 1$, we have

$$\Phi'' + \frac{\tau}{4}\Phi' + \frac{1}{2s}e^{-\tau s/4}\Phi' \int_0^s e^{\tau t/4}\Phi'(t) dt \leq 0,$$

that is,

$$(e^{\tau s/4}\Phi')' + \frac{1}{2s}\Phi' \int_0^s e^{\tau t/4}\Phi'(t) dt \leq 0. \quad (6.5)$$

For the case $\tau > 1$ we have

$$\Phi'' + \frac{1}{4}\Phi' + \frac{1}{2s}e^{-\tau s/4}\Phi' \int_0^s e^{t/4}\Phi'(t) dt \leq 0,$$

that is,

$$(e^{s/4}\Phi')' + \frac{1}{2s}e^{(1-\tau)s/4}\Phi' \int_0^s e^{t/4}\Phi'(t) dt \leq 0. \quad (6.6)$$

First we consider the case $0 < \tau \leq 1$. Define Z as

$$Z(s) = \int_0^s e^{\tau t/4}\Phi'(t) dt.$$

From (6.5) we have

$$sZ'' + \frac{1}{2}e^{-\tau s/4}Z'Z \leq 0. \quad (6.7)$$

By integrating the above on $[0, s]$ we obtain

$$sZ' - Z + \frac{1}{4}e^{-\tau s/4}Z^2 + \frac{\tau}{16} \int_0^s e^{-\tau t/4}Z^2(t) dt \leq 0.$$

Then we have $sZ' - Z + e^{-\tau s/4}Z^2/4 \leq 0$. Dividing the inequality by Z^2 it follows that $(s/Z)' \geq e^{-\tau s/4}/4$. Therefore we obtain

$$Z(s) \leq \frac{\tau s}{1 - e^{-\tau s/4}}. \quad (6.8)$$

From (6.4) we have $-s\Psi'' = e^{-\tau s/4}Z(s)/4 > 0$. Then

$$0 < -s\Psi''(s) \leq \frac{\tau s}{4(e^{\tau s/4} - 1)} < 1 \quad \text{for } s > 0.$$

This implies $s\Psi''(s) \rightarrow 0$ as $s \rightarrow \infty$. By integrating the above on $[0, s]$ we obtain the assertion.

Next we consider the case $\tau > 1$. Define Z as

$$Z(s) = \int_0^s e^{t/4}\Phi'(t) dt.$$

Then from (6.6) we have (6.7). By the similar argument above we obtain (6.8). We see that

$$e^{-\tau s/4} \int_0^s e^{\tau t/4}\Phi'(t) dt = \int_0^s e^{-\tau(s-t)/4}\Phi'(t) dt \leq \int_0^s e^{-(s-t)/4}\Phi'(t) dt = e^{-s/4}Z(s).$$

Then from (6.4) and (6.8) we have

$$0 < -s\Psi''(s) \leq \frac{1}{4} e^{-s/\tau}Z(s) \leq \frac{\tau s}{4(e^{s/4} - e^{(1-\tau)s/4})} \leq \frac{\tau s}{4(e^{s/4} - 1)}.$$

Therefore, $s\Psi''(s) \rightarrow 0$ as $s \rightarrow \infty$. By integrating the above we obtain the assertion. ■

Proof of Proposition 6.1. First we consider the case $0 < \tau \leq 1$. From the second equation of (6.1) we have $\Phi(s) = -4s\Psi''(s) + \tau(\Psi(s) - s\Psi'(s))$. From Lemma 6.3 we obtain

$$\lim_{s \rightarrow \infty} \Phi(s) = \lim_{s \rightarrow \infty} \tau(\Psi(s) - s\Psi'(s)) \leq \frac{\tau^2}{4} \int_0^\infty \frac{s}{e^{\tau s/4} - 1} ds.$$

By the change of variable $z = \tau s/4$ it follows that

$$\lim_{s \rightarrow \infty} \Phi(s) \leq 4 \int_0^\infty \frac{z}{e^z - 1} dz = \frac{2}{3} \pi^2.$$

Since $\|\phi\|_{L^1(\mathbb{R}^2)} = 2\pi \lim_{s \rightarrow \infty} \Phi(s)$ from Lemma 6.1, we obtain the assertion.

Next we consider the case $\tau > 1$. By a similar argument we obtain

$$\lim_{s \rightarrow \infty} \Phi(s) \leq \frac{\tau^2}{4} \int_0^\infty \frac{s}{e^{s/4} - 1} ds = 4\tau^2 \int_0^\infty \frac{z}{e^z - 1} dz = \frac{2}{3} \pi^2 \tau^2,$$

which implies the assertion.

Finally, we consider the case $0 < \tau \leq 1/2$. The change of variables

$$\begin{aligned} t &= (\log s)/2, & k(t) &= \Phi(s), & \ell(t) &= 2s\Phi'(s), & m(t) &= \Psi(s), \\ n(t) &= 2s\Psi'(s) \end{aligned}$$

transforms (6.1) into

$$\begin{cases} \dot{k} = \ell, & \dot{m} = n, \\ \dot{\ell} = \left(2 - k + \tau m - \frac{\tau n}{2} - \frac{e^{2t}}{2}\right)\ell, \\ \dot{n} = 2n + e^{2t}\left(\frac{\tau n}{2} + \tau m - k\right), \end{cases}$$

where $\dot{} = d/dt$. Hence we have

$$\begin{aligned} \frac{d}{dt}((k - 2)^2 + 2\ell) &= 2\ell\left(\tau m - \frac{\tau n}{2} - \frac{e^{2t}}{2}\right) = 4s\Phi'(s)\left(\tau(\Psi(s) - s\Psi'(s)) - \frac{s}{2}\right) \\ &\leq 0 \end{aligned}$$

by Lemma 6.3. Then $(k(t) - 2)^2 + 2\ell(t)$ is decreasing for $t > -\infty$. We note that $\lim_{t \rightarrow -\infty} k(t) = \Phi(0) = 0$ and $\lim_{t \rightarrow -\infty} \ell(t) = \lim_{s \rightarrow 0} 2s\Phi'(s) = \lim_{s \rightarrow 0} s\phi(\sqrt{s}) = 0$. Then we have

$$(k(t) - 2)^2 + 2\ell(t) < 4 \quad \text{for } t > -\infty.$$

Since $\ell(t) = 2s\Phi'(s) = s\phi(\sqrt{s}) > 0$ and $\lim_{t \rightarrow \infty} ((k(t) - 2)^2 + 2\ell(t)) < 4$, we obtain $\lim_{t \rightarrow \infty} k(t) < 4$. Thus $\lim_{s \rightarrow \infty} \Phi(s) < 4$, which implies $\|\phi\|_{L^1(\mathbb{R}^2)} < 8\pi$. ■

7. PROOF OF THEOREM 2

By Theorem 3 it is shown that $(\phi, \psi) \in \mathcal{S}$ if and only if $\psi = \psi(r)$, $r = |y|$, solves $(4.1)_\sigma$ for some $\sigma > 0$ and ϕ is given by (1.7). By Proposition 4.1, the set \mathcal{C} defined by (4.2) is written by one-parameter families $(\sigma(s), \psi(r; s))$ on $s \in \mathbb{R}$. Let

$$\phi(r; s) = \sigma(s)e^{-r^2/4}e^{\psi(r; s)}. \tag{7.1}$$

Then \mathcal{S} is written by one-parameter families $(\phi(r, s), \psi(r, s))$ on $s \in \mathbb{R}$. From Proposition 4.1(i) and (ii) and (7.1) we have $s \mapsto (\phi(\cdot; s), \psi(\cdot; s)) \in C^2[0, \infty) \times C^2[0, \infty)$ is continuous and $(\phi(\cdot; s), \psi(\cdot; s)) \rightarrow (0, 0)$ in $C^2[0, \infty) \times C^2[0, \infty)$ as $s \rightarrow -\infty$. We see that

$$\lambda(s) = 2\pi \int_0^\infty r\phi(r; s) dr. \tag{7.2}$$

Then $\lambda(s)$ is continuous and satisfies $\lambda(s) \rightarrow 0$ as $s \rightarrow -\infty$. Hence, (i) and (ii) hold. By Proposition 6.1 we obtain (iv).

We have $\|\psi(\cdot, s)\|_{L^\infty[0, \infty)} = \psi(0, s) \rightarrow \infty$ as $s \rightarrow \infty$ from (iii) of Proposition 4.1. Let $\{s_k\}$ be a sequence satisfying $s_k \rightarrow \infty$ as $k \rightarrow \infty$. We note that $\{\lambda_k\}$ is bounded by Proposition 6.1. By applying Proposition 5.1, there exists a subsequence (still denoted by $\{s_k\}$) such that $\lambda(s_k) \rightarrow 8\pi$ and $\phi_k(|x|, s_k) dx \rightarrow 8\pi\delta_0(dx)$ as $k \rightarrow \infty$. Therefore, (iii) holds. This completes the proof of Theorem 2. ■

APPENDIX A. EXISTENCE OF SOLUTIONS TO (1.8) WITH (1.9)

The following theorem refines the previous results [26, Theorem 1; 25, Theorems 1 and 2; and 27, Theorem 1.1].

THEOREM A.1. *For any $\tau > 0$ there exists $\sigma^* > 0$ such that*

- (i) *if $\sigma > \sigma^*$, then (1.8) with (1.9) has no solution;*
- (ii) *if $\sigma = \sigma^*$, then (1.8) with (1.9) has at least one solution;*
- (iii) *if $0 < \sigma < \sigma^*$, then (1.8) with (1.9) has at least two distinct solutions $\psi_\sigma, \underline{\psi}_\sigma$ satisfying $\lim_{\sigma \rightarrow 0} \psi_\sigma(0) = 0$ and $\lim_{\sigma \rightarrow 0} \underline{\psi}_\sigma(0) = \infty$.*

Proof. By Theorem 1 problem (1.8) with (1.9) is reduced to problem (4.1) $_\sigma$. By Proposition 4.1 the set \mathcal{C} defined by (4.2) is written by one-parameter families $(\sigma(s), \psi(r; s))$ on $s \in \mathbb{R}$. From (7.1) and (7.2) we find that

$$\sigma(s) = \lambda(s) \Big/ \left(2\pi \int_0^\infty r e^{-r^2/4} e^{\psi(r;s)} dr \right) = \lambda(s) \Big/ \int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi(|y|;s)} dy.$$

From (5.13) in Lemma 5.4 we have

$$\int_{\mathbb{R}^2} e^{-|y|^2/4} e^{\psi(|y|;s)} dy \rightarrow \infty \quad \text{as } s \rightarrow \infty.$$

Then $\sigma(s) \rightarrow 0$ as $s \rightarrow \infty$. Therefore, from Proposition 4.1(ii), $\sigma(s)$ satisfies

$$\lim_{s \rightarrow \pm\infty} \sigma(s) = 0.$$

Let $\sigma^* = \sup_{s \in \mathbb{R}} \sigma(s)$. Then there exists $s^* \in \mathbb{R}$ such that $\sigma^* = \sigma(s^*)$. By Proposition 4.1 it is shown that $(4.1)_\sigma$ has a solution if and only if $\sigma = \sigma(s)$ for some $s \in \mathbb{R}$. Therefore, $(4.1)_\sigma$ has no solution, if $\sigma > \sigma^*$, and $(4.1)_\sigma$ has at least one solution, if $\sigma = \sigma^*$. If $\sigma \in (0, \sigma^*)$, by the mean value theorem, there exists $s_1, s_2 \in \mathbb{R}$, $s_1 < s^* < s_2$ such that $\sigma = \sigma(s_1) = \sigma(s_2)$. Then $(4.1)_\sigma$ has at least two solutions $\psi_{\sigma(s_1)}$ and $\psi_{\sigma(s_2)}$. We note that $\lim_{s \rightarrow -\infty} \psi_{\sigma(s)}(0) = 0$ and $\lim_{s \rightarrow \infty} \psi_{\sigma(s)}(0) = \infty$ by Proposition 4.1(ii) and (iii). Since $\lim_{s \rightarrow \pm\infty} \sigma(s) = 0$, we can choose solutions $\bar{\psi}_\sigma$ and $\underline{\psi}_\sigma$ satisfying $\lim_{\sigma \rightarrow 0} \underline{\psi}_\sigma(0) = 0$ and $\lim_{\sigma \rightarrow 0} \bar{\psi}_\sigma(0) = \infty$. This completes the proof of Theorem A.1. ■

APPENDIX B. PROOF OF THEOREM 4

Define $h_k \in C^2(\Omega) \cap C(\bar{\Omega})$ by

$$\Delta h_k = 0 \quad \text{in } \Omega \quad \text{and} \quad h_k = u_k \quad \text{on } \partial\Omega.$$

We may assume that $\{0\} \in \Omega$ without loss of generality.

LEMMA B.1. *Let $r > 0$ satisfying $\bar{B}_r \subset \Omega$. Then $\|\nabla h_k\|_{L^\infty(B_r)} = O(1)$ as $k \rightarrow \infty$.*

Proof. By the maximum principle, we have $\max_{\bar{\Omega}} h_k - \min_{\bar{\Omega}} h_k \leq \max_{\partial\Omega} h_k - \min_{\partial\Omega} h_k$. Then from (1.15) we obtain

$$\max_{\bar{\Omega}} h_k - \min_{\bar{\Omega}} h_k \leq C_2$$

with a positive constant C_2 . Let $\tilde{h}_k(x) = h_k(x) - \min_{\bar{\Omega}} h_k$. Then \tilde{h}_k satisfies

$$\Delta \tilde{h}_k = 0 \quad \text{in } \Omega, \quad 0 \leq \tilde{h}_k \leq C_2.$$

Since $\partial\tilde{h}_k/\partial x_i$, $i = 1, 2$, is harmonic, by the mean value theorem [11] and Gauss–Green theorem, we obtain

$$\frac{\partial \tilde{h}_k}{\partial x_i} = \frac{1}{\pi r^2} \int_{B_r} \frac{\partial \tilde{h}_k}{\partial x_i} dx = \frac{1}{\pi r^2} \int_{\partial B_r} \tilde{h}_k n_i ds$$

for $i = 1, 2$, where $n = (n_1, n_2)$ is the outer normal unit vector on ∂B_r . Then it follows that

$$\left| \frac{\partial \tilde{h}_k}{\partial x_i} \right| \leq \frac{1}{\pi r^2} \int_{\partial B_r} |\tilde{h}_k| ds \leq \frac{2C_1}{r}, \quad i = 1, 2.$$

Since $|\nabla h_k| = |\nabla \tilde{h}_k|$, we conclude that $\|\nabla h_k\|_{L^\infty(B_r)} = O(1)$ as $k \rightarrow \infty$. ■

Let $w_k(x) = u_k(x) - h_k(x)$ in Ω . Then

$$-\Delta w_k = W_k(x)e^{w_k} \quad \text{in } \Omega, \quad w_k = 0 \quad \text{on } \partial\Omega,$$

where $W_k(x) = e^{h_k(x)}V_k(x)$. Let $G(x, y)$ be Green's function of $-\Delta$ in Ω with respect to the zero boundary conditions

$$-\Delta_x G(x, y) = \delta_y, \quad x \in \Omega, \quad G(x, y) = 0, \quad x \in \partial\Omega.$$

Then we have

$$\nabla w_k(x) = \int_{\Omega} \nabla_x G(x, y) W_k(y) e^{w_k(y)} dy, \quad x \in \Omega. \tag{B.1}$$

Put $z_k(x) = W_k(x)e^{w_k(x)}$.

LEMMA B.2. For $\psi \in C_0^2(\Omega)$ we have

$$\begin{aligned} - \int_{\Omega} (\Delta\psi)z_k dx &= \int_{\Omega} (\nabla(\log W_k) \cdot \nabla\psi)z_k dx \\ &\quad + \frac{1}{2} \int \int_{\Omega \times \Omega} \rho(x, y)z_k(x)z_k(y) dx dy, \end{aligned} \tag{B.2}$$

where $\rho(x, y) = \nabla_x G(x, y) \cdot \nabla\psi(x) + \nabla_y G(x, y) \cdot \nabla\psi(y)$.

Proof. We see that

$$\nabla z_k = (\nabla W_k)e^{w_k} + W_k e^{w_k} \nabla w_k = z_k \nabla(\log W_k) + z_k \nabla w_k.$$

Then, for $\psi \in C_0^2(\Omega)$, we obtain

$$- \int_{\Omega} (\Delta\psi)z_k dx = \int_{\Omega} (\nabla(\log W_k) \cdot \nabla\psi)z_k dx + \int_{\Omega} (\nabla w_k \cdot \nabla\psi)z_k dx. \tag{B.3}$$

From (B.1) and Fubini's theorem, we find that

$$\begin{aligned} & \int_{\Omega} (\nabla w_k(x) \cdot \nabla \psi(x)) z_k(x) \, dx \\ &= \int \int_{\Omega \times \Omega} (\nabla_x G(x, y) \cdot \nabla \psi(x)) z_k(x) z_k(y) \, dx \, dy. \end{aligned} \tag{B.4}$$

By changing the role of x and y in (B.4) we obtain

$$\int_{\Omega} (\nabla w_k(y) \cdot \nabla \psi(y)) z_k(y) \, dy = \int \int_{\Omega \times \Omega} (\nabla_y G(x, y) \cdot \nabla \psi(y)) z_k(x) z_k(y) \, dx \, dy.$$

Hence, we obtain

$$\int_{\Omega} (\nabla w_k \cdot \nabla \psi) z_k \, dx = \frac{1}{2} \int \int_{\Omega \times \Omega} \rho(x, y) z_k(x) z_k(y) \, dx \, dy.$$

From (B.3) we obtain (B.2). ■

Without loss of generality, we may assume that the blowup set \mathcal{B} contains $\{0\}$, and that there exists a $R > 0$ satisfying $\{x : 0 < |x| < R\} \cap \mathcal{B} = \emptyset$. Therefore, $\{u_k\}$ satisfies

$$\max_{\bar{B}_R} u_k \rightarrow \infty \quad \text{and} \quad \max_{\bar{B}_R \setminus B_r} u_k \rightarrow -\infty \quad \text{as } k \rightarrow \infty \tag{B.5}$$

for all $r \in (0, R)$. Moreover,

$$V_k e^{u_k} \, dx \rightarrow \alpha \delta_0(dx) \tag{B.6}$$

on B_R in the sense of measure for some $\alpha \geq 4\pi$.

LEMMA B.3. *There exist constants $r_0 \in (0, R)$ and $a > 0$ such that $V_k(x) \geq a$ for $x \in B_{r_0}$.*

Proof. First we show $\liminf_{k \rightarrow \infty} V_k(0) > 0$. Assume to the contrary that

$$\liminf_{k \rightarrow \infty} V_k(0) = 0.$$

From (1.12) and (1.14), by taking a subsequence in $\{V_k\}$ (still denoted by $\{V_k\}$), there exists $V_0 \in C(\Omega)$ such that $V_k \rightarrow V_0$ in $C(\bar{B}_R)$ and $V_0(0) = 0$.

Let $x_k \in B_R$, $u_k(x_k) = \max_{x \in \bar{B}_R} u_k(x)$. It follows from (B.5) that

$$x_k \rightarrow 0 \quad \text{and} \quad u_k(x_k) \rightarrow \infty. \tag{B.7}$$

Let $\delta_k = e^{-u_k(x_k)/2}$. It follows from (B.7) that $\delta_k \rightarrow 0$. For $|x| \leq R/(2\delta_k)$, we consider the sequence of functions $v_k(x) = u_k(\delta_k x + x_k) + 2 \log \delta_k$. Then v_k satisfies

$$-\Delta v_k(x) = V_k(\delta_k x + x_k) e^{v_k(x)} \quad \text{for } x \in B_{R/(2\delta_k)}.$$

Moreover, we have $v_k(0) = 0$, $v_k(x) \leq 0$ in $B_{R/(2\delta_k)}$, and

$$\int_{B_{R/(2\delta_k)}} e^{v_k(x)} dx \leq \int_{B_R} e^{u_k(x)} dx \leq C$$

for some positive constant C .

For each $r > 0$ the sequence $\{v_k\}$ is well defined in B_r for k large enough. It follows from Theorem A that only alternative (i) may occur, hence $\{v_k\}$ is bounded in $L^\infty_{loc}(B_r)$ and, by standard elliptic estimates, also in $C^{2,\alpha}_{loc}(B_r)$, $0 < \alpha < 1$. Therefore, a subsequence in $\{v_k\}$ converges in $C^2_{loc}(B_r)$. We may do the same arguments for a sequence $r_k \rightarrow \infty$, and pass to a diagonal subsequence (which we will still denote as $\{v_k\}$) converging in $C^2_{loc}(\mathbb{R}^2)$ to v which satisfies $-\Delta v = V_0(0)e^v$ in \mathbb{R}^2 . Moreover, $v(0) = 0$, $v \leq 0$ in \mathbb{R}^2 , and

$$\int_{\mathbb{R}^2} e^v dx \leq C. \tag{B.8}$$

Since $V_0(0) = 0$, v is harmonic in \mathbb{R}^2 . Then v is a constant. This contradicts (B.8). Thus we conclude that $\liminf_{k \rightarrow \infty} V_k(0) > 0$.

From (1.14) there exist constants $r_0 \in (0, R)$ and $a > 0$ satisfying $V_k(x) \geq a$ for $x \in B_{r_0}$. ■

Proof of Theorem 4. We will show that $\alpha = 8\pi$ in (B.6). Take $\phi \in C^2_0(B_R)$ so that $0 \leq \phi \leq 1$ and $\phi \equiv 1$ for $x \in B_{r_0}$, where r_0 is a constant in Lemma B.3. Let $\psi(x) = |x|^2 \phi(x)$. Then we have $\psi \in C^2_0(B_R)$. Moreover, it follows that $\Delta \psi(x) = 4$ and $\nabla \psi(x) = 2x$ for $x \in B_{r_0}$.

We recall that $W_k(x) = e^{h_k(x)} V_k(x)$. Then we have

$$\nabla(\log W_k) = \frac{\nabla W_k}{W_k} = \nabla h_k + \frac{\nabla V_k}{V_k}.$$

From Lemmas B.1 and B.3 and (1.12) we obtain $|\nabla \log W_k(x)| \leq C$ for $x \in B_{r_0}$ with some constant C . Then we have

$$|\nabla \psi(x) \cdot \nabla(\log W_k(x))| \leq 2C|x| \quad \text{for } x \in B_{r_0}. \tag{B.9}$$

We see that $G(x, y) = -(1/2\pi)\log|x - y| + K(x, y)$, where $K(x, y)$ is a smooth function on $\Omega \times \Omega$. Then $\rho(x, y)$ defined in Lemma B.2 satisfies

$$\rho(x, y) = -\frac{1}{\pi} + 2x \cdot \nabla_x K(x, y) + 2y \cdot \nabla_y K(x, y) \quad \text{for } x \in B_{r_0}. \quad (\text{B.10})$$

We see that $z_k(x) = W_k(x)e^{w_k(x)} = V_k(x)e^{v_k(x)}$. From (B.6) we have $z_k(x) dx \rightarrow \alpha\delta_0(dx)$ on B_R in the sense of measure. Furthermore, we have

$$z_k(x)z_k(y) dx dy \rightarrow \alpha^2\delta_{x=0}(dx) \otimes \delta_{y=0}(dy) = \alpha^2\delta_{(x,y)=(0,0)}(dx dy)$$

on B_R in the sense of measure. Letting $k \rightarrow \infty$ in (B.2), from (B.9) and (B.10), we have $-4\pi\alpha = -\alpha^2/(2\pi)$. From $\alpha \geq 4\pi$, we obtain $\alpha = 8\pi$. This completes the proof of Theorem 4. ■

REFERENCES

1. P. Biler, Local and global solvability of some parabolic systems modelling chemotaxis, *Adv. Math. Sci. Appl.* **8** (1998), 715–743.
2. H. Brezis and F. Merle, Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions, *Comm. Partial Differential Equations* **16** (1991), 1223–1253.
3. X. Chen, Remarks on the existence of branch bubbles on the blowup analysis of equation $-\Delta u = e^{2u}$ in dimension two, *Comm. Anal. Geom.* **7** (1999), 295–302.
4. S. Childress, “Chemotactic Collapse in Two Dimensions,” *Lecture Notes in Biomathematics*, Vol. 55, pp. 217–237, Springer, Berlin, 1984.
5. S. Childress and J. K. Percus, Nonlinear aspects of chemotaxis, *Math. Biosci.* **56** (1981), 217–237.
6. M. Escobedo and O. Kavian, Variational problems related to self-similar solutions for the heat equation, *Nonlinear Anal. TMA* **11** (1987), 1103–1133.
7. H. Gajewski and K. Zacharias, Global behaviour of a reaction–diffusion system modelling chemotaxis, *Math. Nachr.* **195** (1998), 77–114.
8. B. Gidas, W.-M. Ni, and L. Nirenberg, Symmetry and related properties via the maximum principle, *Comm. Math. Phys.* **68** (1979), 209–243.
9. B. Gidas, W.-M. Ni, and L. Nirenberg, Symmetry of positive solutions of nonlinear elliptic equations in \mathbb{R}^n , in “Mathematical Analysis and Applications, Part A” (L. Nachbin, Ed.), *Advances in Mathematics Supplementary Studies*, Vol. 7, pp. 369–402, Academic Press, New York, 1981.
10. Y. Giga and T. Kambe, Large time behavior of the vorticity of two-dimensional viscous flow and its application to vortex formation, *Comm. Math. Phys.* **117** (1988), 549–568.
11. D. Gilbarg and N. S. Trudinger, “Elliptic Partial Differential Equations of the Second Order,” Springer-Verlag, Berlin, 1983.
12. G. Harada, T. Nagai, T. Senba, and T. Suzuki, Concentration lemma, Brezis–Merle type inequality, and a parabolic system of chemotaxis, *Adv. Differential Equations* **6** (2001), 1255–1280.
13. M. A. Herrero, E. Medina, and J. J. L. Velázquez, Self-similar blow-up for a reaction–diffusion system, *J. Comput. Appl. Math.* **97** (1998), 99–119.

14. M. A. Herrero and J. J. L. Velázquez, Singularity patterns in a chemotaxis model, *Math. Ann.* **306** (1996), 583–623.
15. M. A. Herrero and J. J. L. Velázquez, Chemotactic collapse for the Keller–Segel model, *J. Math. Biol.* **35** (1996), 177–196.
16. M. A. Herrero and J. J. L. Velázquez, A blow-up mechanism for a chemotaxis model, *Ann. Scuola Norm. Sup. Pisa IV* **35** (1997), 633–683.
17. D. Horstmann and G. Wang, Blow-up in a chemotaxis model without symmetry assumptions, *European J. Appl. Math.* **12** (2001), 159–177.
18. W. Jäger and S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, *Trans. Amer. Math. Soc.* **329** (1992), 819–824.
19. O. Kavian, Remarks on the large time behavior of a nonlinear diffusion equation, *Ann. Inst. H. Poincaré—Anal. Nonlinéaire* **4** (1987), 423–452.
20. T. Kawanago, Asymptotic behavior of solutions of a semilinear heat equation with subcritical nonlinearity, *Ann. Inst. H. Poincaré—Anal. Nonlinéaire* **13** (1996), 1–15.
21. E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an instability, *J. Theor. Biol.* **26** (1970), 399–415.
22. Y.-Y. Li, Harnack type inequality: the method of moving planes, *Comm. Math. Phys.* **200** (1999), 421–444.
23. Y.-Y. Li and I. Shafrir, Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two, *Indiana Univ. Math. J.* **43** (1994), 1255–1270.
24. N. Meyers and J. Serrin, The exterior Dirichlet problem for second order elliptic partial differential equations, *J. Math. Mech.* **9** (1960), 513–538.
25. Y. Mizutani, N. Muramoto, and K. Yoshida, Self-similar radial solutions to a parabolic system modelling chemotaxis via variational method, *Hiroshima Math. J.* **29** (1999), 145–160.
26. Y. Mizutani and T. Nagai, Self-similar radial solutions to a system of partial differential equations modelling chemotaxis, *Bull. Kyushu Inst. Tech. Math. Natur. Sci.* **42** (1995), 19–28.
27. N. Muramoto, Y. Naito, and K. Yoshida, Existence of self-similar solutions to a parabolic system modelling chemotaxis, *Jpn. J. Ind. Appl. Math.* **17** (2000), 427–451.
28. T. Nagai, Blow-up of radially symmetric solutions to a chemotaxis system, *Adv. Math. Sci. Appl.* **5** (1995), 581–601.
29. T. Nagai, Blow-up of nonradial solutions to parabolic–elliptic systems modeling chemotaxis in two-dimensional domain, *J. Inequality Appl.* **6** (2001), 37–55.
30. T. Nagai, T. Senba, and K. Yoshida, Application of the Trudinger–Moser inequality to a parabolic system of chemotaxis, *Funkcial. Ekvac.* **40** (1997), 411–433.
31. Y. Naito and T. Suzuki, Radial symmetry of self-similar solutions for semilinear heat equations, *J. Differential Equations* **163** (2000), 407–428.
32. V. Nanjundiah, Chemotaxis, signal relaying, and aggregation morphology, *J. Theor. Biol.* **42** (1973), 173–105.
33. M. Protter and H. Weinberger, “Maximum Principles in Differential Equations,” Prentice–Hall, Englewood Cliffs, NJ, 1967.
34. T. Senba and T. Suzuki, Chemotactic collapse in a parabolic–elliptic system of mathematical biology, *Adv. Differential Equations* **6** (2001), 21–50.
35. T. Senba and T. Suzuki, Parabolic system of chemotaxis: blow-up in a finite and the infinite time, *Methods Appl. Anal.* **8** (2001).
36. J. Serrin, A symmetry problem in potential theory, *Arch. Rational Mech. Anal.* **43** (1971), 304–318.