



Metric reduction and generalized holomorphic structures

Yicao Wang

Department of Mathematics, Hohai University, Nanjing 210098, China

ARTICLE INFO

Article history:

Received 8 August 2017

Received in revised form 14 May 2018

Accepted 7 August 2018

Available online xxxx

Keywords:

Reduction

Generalized complex structure

Generalized Kähler structure

Generalized holomorphic structure

Principal bundle

Curvature

MSC:

53D18

53D05

53C15

ABSTRACT

In this paper, metric reduction in generalized geometry is investigated. We show how the Bismut connections on the quotient manifold are obtained from those on the original manifold. The result facilitates the analysis of generalized Kähler reduction, which motivates the concept of metric generalized principal bundles and our approach to construct a family of generalized holomorphic line bundles over $\mathbb{C}P^2$ equipped with some non-trivial generalized Kähler structures.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

Generalized complex geometry initiated by N. Hitchin and his school is a simultaneous generalization of symplectic geometry and complex geometry. Since Marsden–Weinstein reduction is a basic construction in symplectic geometry, it is natural to explore a generalized version of symplectic reduction in generalized geometry. This topic was treated in great generality in the formalism of Courant reduction in [1]. When furthermore there is a generalized metric on the Courant algebroid to be reduced, it also descends to the reduced Courant algebroid under proper conditions. In [2], this 'metric reduction' was investigated; in particular, this procedure was checked from the angle of geometry of tangent bundles. The present paper arises from our work [3] on trying to understand metric reduction from a topological field theoretic viewpoint.

Considerations in generalized geometry are conceptually direct and useful, but the underlying structures often hide in depth and need careful analysis. For example, generalized Kähler reduction is easily understood from the general procedure of reduction of Dirac structures, but it contains some sophisticated details from the viewpoint of classical complex geometry. Some of these were included in [2]. In this paper, we will carry on this investigation.

We pay much attention on the special case of *isotropic trivially extended G -actions* in the sense of [1], where G is a compact connected Lie group. With an invariant generalized metric in place, the manifold M under consideration carries two horizontal distributions τ_{\pm} , which are central in our paper. Basically, they are used to express the Bismut connections in the reduced manifold $M_{\text{red}} := M/G$ in terms of Bismut connections in M . This is different from the case of reducing the Levi-Civita connection on M —In the latter case, a connection of the principal bundle $M \rightarrow M_{\text{red}}$ naturally arises from the G -invariant metric g , i.e. the horizontal distribution is just the orthogonal complement \mathcal{H} of the vertical distribution. The Levi-Civita connection on M_{red} can then be expressed using the Levi-Civita connection on M and the orthogonal projection

E-mail address: yicwang@hhu.edu.cn.

from TM to \mathcal{H} . As for reducing Bismut connections, it is not as directly solved as in the ordinary case and should be motivated by conceptual considerations in generalized geometry. This investigation of reducing Bismut connections is motivated by gauging a zero-dimensional supersymmetric σ -model in [3].

When the invariant generalized metric is from a generalized Kähler manifold \mathcal{M} , the situation becomes more interesting. To get a reduced generalized Kähler manifold, an invariant submanifold $M \subset \mathcal{M}$ should be carefully chosen and the reduced generalized Kähler structure will then sit on $M_{red} = M/G$. Hence M only serves as an intermediate object in this procedure. But in this paper M as a *metric generalized principal bundle* (see Section 5) proves to have its own interest: The curvatures of τ_{\pm} are of type $(1, 1)$ w.r.t. the reduced complex structures \tilde{J}_{\pm} on M_{red} respectively. Thus any associated complex vector bundle acquires simultaneously a \tilde{J}_{+} -holomorphic structure and a \tilde{J}_{-} -holomorphic structure.¹ This motivates our approach to constructing generalized holomorphic vector bundles from generalized Kähler reduction.

The paper is organized as follows. In Section 2, we review the basic content of generalized geometry. The goal of Section 3 is to lay the concrete background for later development by investigating the notion of isotropic trivially extended G -action in the presence of an invariant generalized metric. Compared with the work in [2], we hardly contain much essentially new content, but our viewpoint is slightly different. In particular, we include some details of the reduced structures which were missing in [2], and emphasize the basic role of the distributions k_{\pm} (Eq. (3.2) is essential for reducing the Bismut connections) which was not explicitly mentioned in [2]. In Section 4, we mainly tackle the problem of expressing the reduced Bismut connections in terms of Bismut connections in the original manifold (Theorem 4.1). The curvature of the reduced Bismut connection is also computed in terms of the reduction data (Theorem 4.3). These computations play a basic role in [3]. The last three sections devote to using generalized Kähler reduction to produce generalized holomorphic vector bundles. Section 5 discusses the notion of metric generalized principal G -bundle and its associated relative curvature. Section 6 revisits generalized Kähler reduction in the spirit of previous sections, and emphasis is put on structures on the intermediate metric generalized principal G -bundle, which carries a biholomorphic structure. These two sections pave the way for us to produce generalized holomorphic vector bundles via generalized Kähler reduction in Section 7. We give a sufficient condition for the biholomorphic structure to be generalized holomorphic in the Hamiltonian case. As examples, we have constructed generalized holomorphic line bundles on $\mathbb{C}P^2$ equipped with non-trivial generalized Kähler structures.

2. Basics of generalized geometry

In this section, we collect the most relevant aspects of generalized geometry. For a detailed account for it, we refer the reader to [4,5].

In generalized geometry, one considers geometric structures defined on the generalized tangent bundle $\mathbb{T}M = TM \oplus T^*M$ of a smooth manifold M , or more generally on an exact Courant algebroid over M .

A Courant algebroid E is a real vector bundle E over M , together with an anchor map π to TM , a non-degenerate inner product and a so-called Courant bracket $[\cdot, \cdot]_c$ on $\Gamma(E)$. These structures should satisfy some compatibility axioms. E is called exact, if the short sequence

$$0 \longrightarrow T^*M \xrightarrow{\pi^*} E \xrightarrow{\pi} TM \longrightarrow 0$$

is exact. In this paper, by 'Courant algebroid', we always mean an exact one. Given E , one can always find an isotropic right splitting $s: TM \rightarrow E$, which has a curvature form $H \in \Omega_{cl}^3(M)$ defined by

$$H(X, Y, Z) = \langle [s(X), s(Y)]_c, s(Z) \rangle, \quad X, Y, Z \in \Gamma(TM).$$

By the bundle isomorphism $s + \pi^*: TM \oplus T^*M \rightarrow E$, the Courant algebroid structure can be transported onto $\mathbb{T}M$. Then the inner product $\langle \cdot, \cdot \rangle$ is the natural pairing, i.e. $\langle X + \xi, Y + \eta \rangle = \xi(Y) + \eta(X)$, and the Courant bracket is

$$\langle X + \xi, Y + \eta \rangle_H = \langle X, Y \rangle + \mathcal{L}_X \eta - \iota_Y d\xi + \iota_Y \iota_X H, \quad (2.1)$$

called the H -twisted Courant bracket. Different splittings are related by B -field transforms, i.e. $e^B(X + \xi) = X + \xi + \iota_X B$, where B is a 2-form.

A maximal isotropic subbundle $L \subset E$ is called an almost Dirac structure. If L is involutive w.r.t. the Courant bracket, it is called a Dirac structure. These notions can be extended directly to the complexified setting which interests us most.

Definition 2.1. A generalized complex structure on E is a complex structure \mathbb{J} on E orthogonal w.r.t. the inner product and whose $\sqrt{-1}$ -eigenbundle $L \subset E \otimes \mathbb{C}$ is a complex Dirac structure.

Since \mathbb{J} and its $\sqrt{-1}$ -eigenbundle L are equivalent notions, we shall use them interchangeably to denote a generalized complex structure. At a point $x \in M$, the codimension of $\pi(L_x)$ in $T_x M \otimes \mathbb{C}$ is called the type of \mathbb{J} at x . Type can vary along some subset of M , which makes the local geometry of generalized complex structures rather non-trivial.

A generalized complex structure L is an example of complex Lie algebroids. Via the inner product, $\wedge^* L^*$ can be identified with $\wedge^* \bar{L}$, and we have an elliptic differential complex $(\Gamma(\wedge^* \bar{L}), d_L)$, which induces the Lie algebroid cohomology associated with the Lie algebroid L . The differential complex can be twisted by an L -module.

¹ Similar phenomenon, of course, occurs in ordinary Kähler reduction but is seldom emphasized in the literature.

Definition 2.2. Given a generalized complex structure L over M , an L -connection D in a complex vector bundle W is a differential operator $D : \Gamma(W) \rightarrow \Gamma(\bar{L} \otimes W)$ satisfying

$$D(fs) = d_L f \otimes s + fDs, \quad s \in \Gamma(W), f \in C^\infty(M).$$

If D is flat, i.e. $D^2 = 0$, D is called a generalized holomorphic structure and W an L -module or a generalized holomorphic vector bundle.

A standard non-trivial example of generalized holomorphic bundles is the canonical line bundle in the pure spinor description of a generalized complex structure. Further analysis and examples of generalized holomorphic structures can be found in [6,7].

Definition 2.3. A generalized (Riemannian) metric on E is an orthogonal, self-adjoint operator \mathcal{G} such that $\langle \mathcal{G}e, e \rangle > 0$ for nonzero $e \in E$. It is necessary that $\mathcal{G}^2 = id$. The \pm -eigenbundles V_\pm are positive and negative subbundles of maximal rank respectively.

A generalized metric induces a canonical isotropic splitting: $E = \mathcal{G}(T^*M) \oplus T^*M$. It is called the metric splitting. Given a generalized metric, we shall always choose the metric splitting to identify E with $\mathbb{T}M$. Then \mathcal{G} is of the form $\begin{pmatrix} 0 & g^{-1} \\ g & 0 \end{pmatrix}$ where g is an ordinary Riemannian metric, and vectors in V_\pm are of the form $X \pm g(X)$ respectively for $X \in TM$.

If H is the curvature of the metric splitting, sometimes we call the triple (M, g, H) a generalized Riemannian manifold, without explicitly mentioning the underlying Courant algebroid and generalized metric. For a generalized Riemannian manifold (M, g, H) , one can define the Bismut connections $\nabla^\pm = \nabla \pm \frac{1}{2}g^{-1}H$, where ∇ is the Levi-Civita connection. It was observed in [8,9] that these connections can be expressed using H -twisted Courant bracket:

$$[X \mp g(X), Y \pm g(Y)]_H^\pm = \nabla_X^\pm Y \pm g(\nabla_X^\pm Y), \quad (2.2)$$

where $(X + \xi)^\pm$ denote the V_\pm -part of $X + \xi \in \Gamma(\mathbb{T}M)$ w.r.t. the decomposition $E = V_+ \oplus V_-$.

A generalized metric is an ingredient of a generalized Kähler structure, which is the analogue of Kähler structure in complex geometry.

Definition 2.4. A generalized Kähler structure on E is a pair of commuting generalized complex structures \mathbb{J}_1 and \mathbb{J}_2 such that $\mathcal{G} = -\mathbb{J}_1\mathbb{J}_2$ is a generalized metric.

A generalized Kähler structure can also be characterized in terms of more ordinary notions: There are two complex structures J_\pm on M compatible with the Riemannian metric g induced from the generalized metric. Let $\omega_\pm = gJ_\pm$ and H be the curvature of the metric splitting. Then

$$d_+^c \omega_+ = -d_-^c \omega_- = -H, \quad (2.3)$$

where d_\pm^c are the d^c -differentials associated to J_\pm respectively. J_\pm is necessarily flat w.r.t. ∇^\pm respectively and H should be of type $(1, 2) + (2, 1)$ w.r.t. both J_+ and J_- . Let $T_{0,1}^\pm M$ be the anti-holomorphic tangent bundles w.r.t. J_\pm respectively. Then we can form two vector bundles over M :

$$L_\pm = \{X \pm \sqrt{-1}\omega_\pm(X) | X \in T_{0,1}^\pm M\}.$$

In the metric splitting, $L_1 := L_+ \oplus L_-$ and $L_2 := L_+ \oplus \bar{L}_-$ are precisely $\sqrt{-1}$ -eigenbundles of \mathbb{J}_1 and \mathbb{J}_2 respectively.

We are particularly interested in generalized holomorphic structures over a generalized Kähler manifold. In this setting, we choose L_1 to be the underlying generalized complex structure of a generalized holomorphic structure D . Due to the decomposition $L_1 = L_+ \oplus L_-$, D can be decomposed as $D = \bar{\delta}_+ + \bar{\delta}_-$ accordingly. Actually, $\bar{\delta}_\pm$ are ordinary J_\pm -holomorphic structures respectively. Additionally, it is necessary that

$$\bar{\delta}_+ \bar{\delta}_- + \bar{\delta}_- \bar{\delta}_+ = 0. \quad (2.4)$$

Conversely, given J_\pm -holomorphic structures $\bar{\delta}_\pm$, if Eq. (2.4) is also satisfied, then $D := \bar{\delta}_+ + \bar{\delta}_-$ is a generalized holomorphic structure [10].

3. Isotropic trivially extended action and metric reduction

Though there is a much more general framework in [1] to adapt an ordinary Lie algebra action to the setting of a Courant algebroid, we content ourselves here with the following more restrictive notion of isotropic trivially extended action of a Lie algebra \mathfrak{g} . Throughout the paper, we always assume that \mathfrak{g} is the Lie algebra of a compact connected Lie group G acting freely on M from the left. In the following, a Courant algebroid E over M is fixed.

Definition 3.1 ([1]). Let $\varphi_0 : \mathfrak{g} \rightarrow \Gamma(TM)$ be the infinitesimal action of G over M . An isotropic trivial extension of this action to E is a bracket-preserving morphism $\varphi : \mathfrak{g} \rightarrow \Gamma(E)$ such that the following diagram

$$\begin{array}{ccc} \mathfrak{g} & \xrightarrow{id} & \mathfrak{g} \\ \varphi \downarrow & & \varphi_0 \downarrow \\ \Gamma(E) & \xrightarrow{\pi} & \Gamma(TM) \end{array}$$

is commutative and the image of φ is isotropic pointwise in E . If furthermore this extended action integrates to a G -action on E , we call it an isotropic trivially extended G -action.

Let e_a , $a = 1, 2, \dots, \dim \mathfrak{g}$ be a basis of \mathfrak{g} and let V_a be the corresponding fundamental vector fields on M . When an isotropic splitting of E is chosen, $\varphi(e_a) = V_a + \xi_a$, where $\xi_{(\cdot)} : \mathfrak{g} \rightarrow \Gamma(T^*M)$ is equivariant and $\iota_a \xi_b + \iota_b \xi_a = 0$, where ι_a denotes contraction with V_a . If additionally the splitting is invariant, then its curvature H is invariant and $\iota_a H = d\xi_a$. A remarkable fact in [1] is that $H + \xi_{(\cdot)}$ is actually a closed equivariant 3-form in the Cartan model of equivariant de Rham cohomology.

Let $K \subset E$ be the subbundle generated by $\varphi(\mathfrak{g})$, and K^\perp its orthogonal complement in E w.r.t. the inner product. Then due to the reduction theory developed in [1], $E_{red} := K^\perp/G$ has the structure of a Courant algebroid induced from E . Now if \mathcal{G} is a G -invariant generalized metric over E , then E_{red} also acquires a generalized metric \mathcal{G}_{red} .

There is a useful way to describe \mathcal{G}_{red} . Let $K^\mathcal{G}$ be the \mathcal{G} -orthogonal complement of K in K^\perp , i.e.

$$K^\mathcal{G} = \mathcal{G}(K^\perp) \cap K^\perp.$$

By projection, $K^\mathcal{G}$ is isomorphism to K^\perp/K , and \mathcal{G}_{red} is actually the restriction of \mathcal{G} on the subbundle $K^\mathcal{G} \subset K^\perp$. Accordingly, we have the decomposition

$$K^\mathcal{G} = V_+^{red} \oplus V_-^{red},$$

where $V_\pm^{red} = V_\pm \cap K^\mathcal{G}$. Furthermore, by the abovementioned isomorphism, we can regard $K^\mathcal{G}/G$ as a Courant algebroid over M_{red} . The advantage of using $K^\mathcal{G}$ instead of K^\perp/K is that, when a lift $\hat{A} \in \Gamma(K^\perp)$ of $A \in \Gamma(E_{red})$ is needed, we can choose \hat{A} to be the unique one in $\Gamma(K^\mathcal{G})$.

Though the Courant algebroid structure of $K^\mathcal{G}/G$ is clear from the generalized reduction procedure, for later convenience, in the following we spell out some details of this structure. We do this mainly at the level of equivariant bundles.

The metric splitting of \mathcal{G} is, of course, invariant and in this splitting the Riemannian metric g and the curvature H are both invariant. Let $\varphi(e_a) = V_a + \xi_a$ in this splitting. Associated with the isotropic trivially extended \mathfrak{g} -action are two horizontal distributions on M [2]:

$$\tau_\pm := \{Y \in TM | g(Y, V_a) \pm \xi_a(Y) = 0, \quad a = 1, 2, \dots, \dim \mathfrak{g}\}. \quad (3.1)$$

They are just distributions derived by projecting V_\pm^{red} to TM and define two connections in the principal G -bundle $M \rightarrow M_{red} := M/G$. They are basic for our later considerations. It is convenient to use $V_a^\pm := V_a \pm g^{-1}\xi_a$. Denote the bundles generated by $\{V_a^\pm\}$ by $k_\pm \subset TM$ respectively. Then Eq. (3.1) can be rephrased as the orthogonal decomposition

$$TM = k_\pm \oplus \tau_\pm. \quad (3.2)$$

Let $q : M \rightarrow M_{red}$ be the natural quotient map. Let us first interpret the short exact sequence properly:

$$0 \longrightarrow q^*(T^*M_{red}) \xrightarrow{[\pi]^*} K^\mathcal{G} \xrightarrow{[\pi]} q^*(TM_{red}) \longrightarrow 0, \quad (3.3)$$

where q^* means the pull-back of vector bundles by the quotient map q , and $[\pi]$ denotes the composition $q_* \circ \pi$.

Lemma 3.2.

$$K^\mathcal{G} = \{Y + \eta \in TM | g(Y, V_a) + g(\eta, \xi_a) = 0, \xi_a(Y) + \eta(V_a) = 0\},$$

and

$$\ker([\pi]) = \{Y + \eta \in K^\mathcal{G} | Y \in \pi(K)\}.$$

Proof. This is obvious by definition of $K^\mathcal{G}$. \square

For E_{red} to be exact, $q^*(T^*M_{red})$ should be identified with $\ker([\pi])$ via the inner product on $K^\mathcal{G}$. This is realized as follows:

Lemma 3.3. Let $T_{ab} := g(V_a^+, V_b^+) = g(V_a^-, V_b^-)$ and denote its inverse by T^{ab} . Then for $\xi \in q^*(T^*M_{red})$,

$$[\pi]^*(\xi) = \xi - T^{ab}g(\xi, \xi_a)(V_b + \xi_b),$$

and the image of $[\pi]^*$ is precisely $\ker([\pi])$.

Proof. First note that

$$g(V_a^+, V_b^+) = g(V_a^-, V_b^-) = g(V_a, V_b) + g(\xi_a, \xi_b)$$

due to the fact $\iota_a \xi_b + \iota_b \xi_a = 0$. This implies that the restriction of g on either τ_+ or τ_- gives rise to the same Riemannian metric on TM_{red} , just as required.

For $\xi \in q^*(T^*M_{red})$, $[\pi]^*(\xi) \in K^G$ is characterized by

$$\langle [\pi]^*(\xi), Y + \eta \rangle = \xi([\pi](Y + \eta)) = \xi(q_*(Y)), \quad \forall Y + \eta \in K^G.$$

We can assume $[\pi]^*(\xi) = \xi + f^b(V_b + \xi_b)$ for some constants f^b to be determined. We thus have

$$g(\xi + f^b \xi_b, \xi_a) + g(f^b V_b, V_a) = 0, \quad a, b = 1, 2, \dots, \dim g,$$

i.e.

$$[g(V_a, V_b) + g(\xi_a, \xi_b)]f^b = T_{ab}f^b = -g(\xi, \xi_a), \quad a, b = 1, 2, \dots, \dim g.$$

This leads to our expression of $[\pi]^*(\xi)$.

It is obvious that $\text{Ran}([\pi]^*) \subset \ker([\pi])$. Thus $\text{Ran}([\pi]^*) = \ker([\pi])$ by dimensional reason. \square

We have already used the metric splitting of G to identify E with $\mathbb{T}M$. G , when restricted on K^G , also gives rise to an isotropic splitting $\mathcal{G}(\ker[\pi])$ of the sequence (3.3). Then $\mathcal{G}(\ker[\pi])/G$ is a splitting of K^G/G . Let $Q_{ab} = g(V_a, V_b)$ and Q^{ab} be its inverse.

Proposition 3.4. *If E_{red} is identified with K^G/G and T^*M_{red} with $\ker[\pi]/G$, then $\mathcal{G}(\ker[\pi])/G$ is the metric splitting of E_{red} .*

Proof. The following proof can only be viewed as a detailed analysis of the obvious conclusion. We only need to prove that in this splitting, V_+^{red}/G is the graph of the reduced metric \tilde{g} on TM_{red} , since other splittings will involve extra B -transforms.

Note that

$$\mathcal{G}(\ker[\pi]) = \{Y + \eta \in K^G \mid g^{-1}\eta \in \pi(K)\}.$$

The projection of $\mathcal{G}(\ker[\pi])$ to TM is a third horizontal distribution τ on M as a principal G -bundle. For $Y + \eta \in \mathcal{G}(\ker[\pi])$, η is uniquely determined by Y . We thus write η_Y instead of η .

As observed in [2], a typical element in $V_+ \cap K^G$ is of the form

$$A = Y + g^{-1}\eta_Y + g(Y) + \eta_Y, \quad Y \in \tau.$$

Note that $Y + \eta_Y \in \mathcal{G}(\ker[\pi])$ and $g^{-1}\eta_Y + g(Y) \in \ker[\pi]$. This means, in the splitting determined by $\mathcal{G}(\ker[\pi])/G$, A descends to

$$[Y] + g(Y) - Q^{ab}\eta_Y(V_b)\xi_a \in TM_{red} \oplus T^*M_{red},$$

since $g^{-1}\eta_Y = Q^{ab}\eta_Y(V_b)V_a$ and

$$g^{-1}\eta_Y = -Q^{ab}\eta_Y(V_b)\xi_a + Q^{ab}\eta_Y(V_b)(V_a + \xi_a).$$

Note that

$$\begin{aligned} g(Y, V_c) - Q^{ab}\eta_Y(V_b)\xi_a(V_c) &= g(Y, V_c) + Q^{ab}\eta_Y(V_b)\xi_c(V_a) \\ &= g(Y, V_c) + \xi_c(g^{-1}\eta_Y) \\ &= g(Y, V_c) + g(\eta_Y, \xi_c) \\ &= 0. \end{aligned}$$

So $g(Y) - Q^{ab}\eta_Y(V_b)\xi_a$ does live in T^*M_{red} . Now we only need to check that on τ we have

$$q^*\tilde{g}([Y]) = g(Y) - Q^{ab}\eta_Y(V_b)\xi_a, \quad \forall Y \in \tau.$$

Let $Z \in \tau$. Then on one side by definition of \tilde{g} we have

$$\begin{aligned} q^*\tilde{g}([Y])(Z) &= \tilde{g}([Y], [Z]) = g(Y + g^{-1}\eta_Y, Z + g^{-1}\eta_Z) \\ &= g(Y, Z) + \eta_Y(Z) + \eta_Z(Y) + g(\eta_Y, \eta_Z) \\ &= g(Y, Z) + g(\eta_Y, \eta_Z). \end{aligned}$$

On the other side,

$$\begin{aligned} g(Y, Z) - Q^{ab}\eta_Y(V_b)\xi_a(Z) &= g(Y, Z) + Q^{ab}\eta_Y(V_b)\eta_Z(V_a) \\ &= g(Y, Z) + \eta_Z(g^{-1}\eta_Y) \\ &= g(Y, Z) + g(\eta_Y, \eta_Z). \end{aligned}$$

Hence the claim follows. \square

Remark. τ is the average of τ_{\pm} in the following sense: If Y is the lift of $[Y]$ in τ , then $Y \pm g^{-1}\eta_Y$ are the lifts of $[Y]$ in τ_{\pm} respectively.

Let us compute the curvature \tilde{H} of the metric splitting $\mathcal{G}(\ker[\pi])/G$. Note that η is a map from τ to T^*M . Since $\mathcal{G}(\ker[\pi])$ is isotropic, we have $\eta_X(Y) + \eta_Y(X) = 0$. Additionally, it can be easily obtained that $\eta_X(V_a) = -\xi_a(X)$. We can look for a 2-form $\gamma : TM \rightarrow T^*M$ such that its restriction on τ is precisely η . Let θ^a be the connection form determined by τ . A choice of γ is then

$$\gamma = -\frac{1}{2}Q^{ab}[\xi_b - \xi_b(V_c)\theta^c] \wedge [g(V_a) - Q_{ad}\theta^d] - \xi_a \wedge \theta^a + \frac{1}{2}\xi_b(V_a)\theta^a\theta^b.$$

Proposition 3.5. *If TM_{red} is modelled on τ , then the curvature of the metric splitting of E_{red} is $\tilde{H} = (H + d\gamma)|_{\tau}$.*

Proof. Let $[X]$ here denote a vector field on M_{red} represented by the invariant lift $X \in \Gamma(\tau)$. Then

$$\begin{aligned}\tilde{H}([X], [Y], [Z]) &= \langle [X + \eta_X, Y + \eta_Y]_H, Z + \eta_Z \rangle \\ &= \langle [X, Y] + L_X\eta_Y - \iota_Y d\eta_X + \iota_Y \iota_X H, Z + \eta_Z \rangle \\ &= \eta_Z([X, Y]) + \eta_Y([Z, X]) + \eta_X([Y, Z]) + X\eta_Y(Z) \\ &\quad + Y\eta_Z(X) + Z\eta_X(Y) + H(X, Y, Z) \\ &= \gamma(Z, [X, Y]) + \gamma(Y, [Z, X]) + \gamma(X, [Y, Z]) \\ &\quad + X\gamma(Y, Z) + Y\gamma(Z, X) + Z\gamma(X, Y) + H(X, Y, Z) \\ &= (H + d\gamma)(X, Y, Z),\end{aligned}$$

as required. \square

Remark. The appearance of \tilde{H} depends on which connection among τ , τ_{\pm} is used to model TM_{red} . In [2], τ_+ is used to do this. In the next section, we will carry out the same computation in a way different from that of [2].

To conclude this section, we clarify some notation for our later use. If \mathcal{M} is a bigger manifold carrying an isotropic trivially extended G -action and M is an invariant submanifold of \mathcal{M} , the Courant algebroid \mathcal{E} on \mathcal{M} can be directly pulled back to M and the isotropic trivially extended G -action also descends. In fact, if $\varphi(e_a) = V_a + \xi_a$ on \mathcal{M} in some splitting, then a natural splitting arises in the pull-back of \mathcal{E} and $\varphi(e_a) = V_a|_M + i^*(\xi_a)$, where i is the inclusion map. By abuse of notation, we will only write $\varphi(e_a) = V_a + \xi_a$ either on \mathcal{M} or on M . The ambiguity will be clarified by the context.

4. Bismut connections in metric reduction

The basic context of this section is the same as that of the former one, and we continue to use the notation there. We try to figure out how the Bismut connections $\tilde{\nabla}^{\pm}$ in M_{red} are reduced from those in M . The curvature of $\tilde{\nabla}^-$ is a basic ingredient in [3] to interpret metric reduction in the formalism of balanced topological field theories.

Our starting point is Eq. (2.2) where Bismut connections are expressed using Courant bracket. Since by the reduction procedure established in [1], the Courant algebroid E_{red} on M_{red} can naturally be described in terms of the Courant algebroid E on M , one can expect that the Bismut connections on M_{red} could be described in terms of the Courant bracket on M . The two connections τ_{\pm} play a fundamental role in this investigation.

Note that \tilde{g} is in fact defined by restricting g on τ_+ (or τ_-). This is different from the ordinary case. Let $\tilde{\nabla}^-$ be the $-$ -Bismut connection on M_{red} and let $[X]$ denote a vector field on M_{red} represented by an invariant lift X on M . X^{\pm} are used to denote the unique lifts of $[X]$ in τ_{\pm} respectively. Let $\varrho_-([X], [Y])$ denote the unique lift of $\tilde{\nabla}_{[X]}^- [Y]$ in τ_- .

Theorem 4.1. $\varrho_-([X], [Y])$ is the projection of $\nabla_{X^+}^- Y^-$ to τ_- along k_- , namely

$$\varrho_-([X], [Y]) = \nabla_{X^+}^- Y^- + T^{ab}g(Y^-, \nabla_{X^+}^- V_b^-)V_a^-, \quad (4.1)$$

where T^{ab} is the inverse of $T_{ab} = g(V_a^-, V_b^-)$.

Proof. According to Eq. (2.2), in the metric splitting of E_{red} ,

$$\tilde{\nabla}_{[X]}^- [Y] - \tilde{g}(\tilde{\nabla}_{[X]}^- [Y]) = [[X] + \tilde{g}([X]), [Y] - \tilde{g}([Y])]_{\tilde{H}}^-.$$

Due to the discussion in Section 3 the R.H.S. of the above equation can be computed using the corresponding invariant sections of $K^{\mathcal{G}}$, i.e.

$$[X^+ + g(X^+), Y^- - g(Y^-)]_{\tilde{H}}^-.$$

It should be noted that $\Gamma^G(K^G)$ is not involutive under the Courant bracket. Involutivity can only hold up to addition of invariant section of K . Therefore,

$$[X^+ + g(X^+), Y^- - g(Y^-)]_H = A_+ + A_- + N,$$

where $A_\pm \in \Gamma(V_\pm^{red})$ and $N = 2c^a(V_a + \xi_a)$ for some functions c^a to be determined. Of course we want to separate A_- from the above expression because $\varrho_-([X], [Y]) = \pi_-(A_-)$ where π_- is the projection from V_- to TM .

We already have

$$[X^+ + g(X^+), Y^- - g(Y^-)]_H^- = A_- + N_-,$$

where $N_- = c^a(V_a - g(V_a) + \xi_a - g^{-1}\xi_a)$. Hence,

$$A_- + N_- = \nabla_{X^+}^- Y^- - g(\nabla_{X^+}^- Y^-).$$

Therefore,

$$\varrho_-([X], [Y]) + c^a(V_a - g^{-1}\xi_a) = \varrho_-([X], [Y]) + c^a V_a^- = \nabla_{X^+}^- Y^-.$$

Due to the orthogonal decomposition $TM = \tau_- \oplus k_-$, the above equation means that $\varrho_-([X], [Y])$ is actually the τ_- -part of $\nabla_{X^+}^- Y^-$ w.r.t. this decomposition. We then find

$$c^a = T^{ab}g(\nabla_{X^+}^- Y^-, V_b^-) = -T^{ab}g(Y^-, \nabla_{X^+}^- V_b^-).$$

We finally obtain the formula as required. \square

Remark. The result is very similar to the ordinary case except that a different orthogonal decomposition is used. In particular, if $[Z]$ is another vector field on M_{red} , then $\tilde{g}(\tilde{\nabla}_{[X]}^- [Y], [Z]) = g(\nabla_{X^+}^- Y^-, Z^-)$.

Now we can turn to the problem of expressing the curvature of $\tilde{\nabla}^-$ in terms of that of ∇^- . Let θ_\pm^a be the connection forms associated to τ_\pm respectively and let Ω_\pm^a be the associated curvatures. For later use, we want to express Ω_\pm^a in terms of V_a, ξ_a . Let $K_{ab} = g_{ab} - \xi_a(V_b)$ and K^{ba} its inverse, i.e. $K^{bc}K_{ab} = \delta_a^c$.

Lemma 4.2. Let Ω_\pm^a be the curvatures of τ_\pm . Then

$$\Omega_+^a|_{\tau_+} = K^{ba}d\xi_b^+|_{\tau_+}, \quad \Omega_-^a|_{\tau_-} = K^{ab}d\xi_b^-|_{\tau_-},$$

where $\xi_b^\pm = g(V_b^\pm)$.

Proof. We only compute Ω_+^a . The computation for Ω_-^a is similar. Note that

$$\theta_+ = \theta_+^a e_a = t^{ba}g(V_b^+)e_a,$$

where t^{ba} is to be determined. We have

$$t^{ba}g(V_b^+, V_c) = t^{ba}K_{cb} = \delta_c^a.$$

Then t^{ba} is precisely K^{ba} and $\theta_+^a = K^{ba}g(V_b^+)$. Then

$$\begin{aligned} \Omega_+^a(X^+, Y^+) &= d\theta_+^a(X^+, Y^+) = X^+\theta_+^a(Y^+) - Y^+\theta_+^a(X^+) - \theta_+^a([X^+, Y^+]) \\ &= -\theta_+^a([X^+, Y^+]) = -K^{ba}(V_b^+, [X^+, Y^+]) \\ &= K^{ba}(d\xi_b^+)(X^+, Y^+). \quad \square \end{aligned}$$

Let R^- and \tilde{R}^- be the curvatures of ∇^- and $\tilde{\nabla}^-$ respectively. We have

Theorem 4.3. The curvature \tilde{R}^- of $\tilde{\nabla}^-$ is

$$\begin{aligned} \tilde{g}(\tilde{R}^-([X], [Y])[Z], [W]) &= g(R^-(X^+, Y^+)Z^-, W^-) \\ &\quad - \frac{1}{2}K^{ab}(d\xi_a^+)(X^+, Y^+)(d\xi_b^-)(Z^-, W^-) \\ &\quad + T^{ab}[g(Z^-, \nabla_{Y^+}^- V_a^-)g(W^-, \nabla_{X^+}^- V_b^-) - (X \leftrightarrow Y)], \end{aligned}$$

where $(X \leftrightarrow Y)$ denotes a term similar to the term in front of it, only with X and Y exchanged.

Proof. Since $\tilde{\nabla}^-$ and ∇^- are metric connections, we have

$$\begin{aligned} \tilde{g}(\tilde{\nabla}_{[X]}^- \tilde{\nabla}_{[Y]}^- [Z], [W]) &= [X]\tilde{g}(\tilde{\nabla}_{[Y]}^- [Z], [W]) - \tilde{g}(\tilde{\nabla}_{[Y]}^- [Z], \tilde{\nabla}_{[X]}^- [W]) \\ &= X^+g(\nabla_{Y^+}^- Z^-, W^-) \end{aligned}$$

$$\begin{aligned}
& -g(\nabla_{Y^+}^- Z^- + T^{ab}g(Z^-, \nabla_{Y^+}^- V_b^-)V_a^-, \nabla_{X^+}^- W^- + T^{cd}g(W^-, \nabla_{X^+}^- V_d^-)V_c^-) \\
& = g(\nabla_{X^+}^- \nabla_{Y^+}^- Z^-, W^-) - T^{ab}g(Z^-, \nabla_{Y^+}^- V_a^-)g(W^-, \nabla_{X^+}^- V_b^-) \\
& - 2T^{ab}g(Z^-, \nabla_{Y^+}^- V_b^-)g(V_a^-, \nabla_{X^+}^- W^-) \\
& = g(\nabla_{X^+}^- \nabla_{Y^+}^- Z^-, W^-) + T^{ab}g(Z^-, \nabla_{Y^+}^- V_a^-)g(W^-, \nabla_{X^+}^- V_b^-),
\end{aligned}$$

where Eq. (4.1) is used.

Additionally,

$$\begin{aligned}
\tilde{g}(\tilde{\nabla}_{[[X],[Y]]}^- [Z], [W]) & = g(\nabla_{[X^+, Y^+] + \Omega_+(X^+, Y^+)}^- Z^-, W^-) \\
& = g(\nabla_{[X^+, Y^+]}^- Z^-, W^-) + g(\nabla_{\Omega_+(X^+, Y^+)}^- Z^-, W^-),
\end{aligned}$$

where we have used the identity²

$$[X^+, Y^+] - \widetilde{[[X], [Y]]} = -\Omega_+(X^+, Y^+) = -\Omega_+^a(X^+, Y^+)V_a,$$

and Ω_+^a is the curvature of τ_+ .

By Lemma 4.2, we have

$$g(\nabla_{\Omega_+(X^+, Y^+)}^- Z^-, W^-) = K^{ba}(d\xi_b^+)(X^+, Y^+)g(\nabla_{V_a}^- Z^-, W^-).$$

Note that

$$\begin{aligned}
g(\nabla_{V_a}^- Z^-, W^-) & = g(\nabla_{V_a} Z^-, W^-) - \frac{1}{2}H(V_a, Z^-, W^-) \\
& = g(\nabla_{V_a} Z^-, W^-) - \frac{1}{2}(d\xi_a)(Z^-, W^-),
\end{aligned}$$

$$\begin{aligned}
g(\nabla_{V_a} Z^-, W^-) & = g(\nabla_{Z^-} V_a, W^-) = Z^-(g(V_a)(W^-)) - g(V_a, \nabla_{Z^-} W^-) \\
& = Z^-(g(V_a)(W^-)) - g(V_a, \nabla_{W^-} Z^-) - g(V_a)([Z^-, W^-]) \\
& = Z^-(g(V_a)(W^-)) - W^-(g(V_a)(Z^-)) + g(\nabla_{W^-} V_a, Z^-) \\
& - g(V_a)([Z^-, W^-]) \\
& = dg(V_a)(Z^-, W^-) + g(\nabla_{W^-} V_a, Z^-),
\end{aligned}$$

and

$$g(\nabla_{V_a} Z^-, W^-) + g(\nabla_{W^-} V_a, Z^-) = 0.$$

Then we have

$$g(\nabla_{V_a}^- Z^-, W^-) = \frac{1}{2}d\xi_a^-(Z^-, W^-). \quad (4.2)$$

Combining all the above ingredients together, we come to the conclusion. \square

Similar formulae for $\tilde{\nabla}^+$ hold, but we just write down the counterpart of Eq. (4.1) for $\tilde{\nabla}^+$ for later use. The detail is left to the interested reader. Let $\varrho_+([X], [Y])$ be the τ_+ -lift of $\tilde{\nabla}_{[X]}^+[Y]$. Then

$$\varrho_+([X], [Y]) = \nabla_{X^+}^+ Y^+ + T^{ab}g(Y^+, \nabla_{X^+}^+ V_b^+)V_a^+. \quad (4.3)$$

As an application of our formula (4.3), we use it to compute the curvature \tilde{H} of the reduced metric splitting again.

Proposition 4.4. *The curvature \tilde{H} of the reduced metric splitting is $(H + \Omega_+^a \wedge \xi_a)|_{\tau_+}$.*

Proof. Since \tilde{H} is the torsion of $\tilde{\nabla}^+$, we have

$$\begin{aligned}
\tilde{H}([X], [Y], [Z]) & = \tilde{g}(\tilde{\nabla}_{[X]}^+[Y], [Z]) - \tilde{g}(\tilde{\nabla}_{[Y]}^+[X], [Z]) - \tilde{g}([X], [Y], [Z]) \\
& = g(\nabla_{X^+}^+ Y^+, Z^+) - g(\nabla_{Y^+}^+ X^+, Z^+) \\
& - g([X^+, Y^+] + \Omega_+^a(X^+, Y^+)V_a, Z^+).
\end{aligned}$$

Since X^- is uniquely determined by X^+ , define $\varsigma(X^+) = X^+ - X^-$. By definition, $g(X^+ - \varsigma(X^+), V_a^-) = 0$, i.e.

$$g(V_b, V_a^-)\varsigma^b(X^+) = g(X^+, V_a^-) = -2\xi_a(X^+).$$

² $\widetilde{[X]}$ denotes the horizontal lift of $[X]$ in τ_+ .

Since $g(V_b, V_a^-) = K_{ab}$, we find $\varsigma(X^+) = -2K^{ab}\xi_b(X^+)V_a$, and

$$\begin{aligned} g(\nabla_{\varsigma(X^+)}^+ Y^+, Z^+) &= -2K^{ab}\xi_b(X^+)g(\nabla_{V_a}^+ Y^+, Z^+) \\ &= -K^{ab}\xi_b(X^+)d\xi_a^+(Y^+, Z^+) \\ &= -\Omega_+^b(Y^+, Z^+)\xi_b(X^+), \end{aligned}$$

where we have used a counterpart of Eq. (4.2) for ∇^+ and Lemma 4.2. Thus we obtain

$$\begin{aligned} \tilde{H}([X], [Y], [Z]) &= H(X^+, Y^+, Z^+) - g(\nabla_{\varsigma(X^+)}^+ Y^+, Z^+) \\ &\quad + g(\nabla_{\varsigma(Y^+)}^+ X^+, Z^+) + \Omega_+^a(X^+, Y^+)\xi_a(Z^+) \\ &= H(X^+, Y^+, Z^+) + \Omega_+^b(Y^+, Z^+)\xi_b(X^+) \\ &\quad - \Omega_+^b(X^+, Z^+)\xi_b(Y^+) + \Omega_+^b(X^+, Y^+)\xi_b(Z^+) \\ &= (H + \Omega_+^b \wedge \xi_b)(X^+, Y^+, Z^+), \end{aligned}$$

which recovers the result in [2]. \square

As a conclusion, we briefly discuss the metric reduction from a bigger manifold \mathcal{M} to a submanifold M . The Courant algebroid \mathcal{E} over \mathcal{M} and the generalized metric \mathcal{G} can be directly pulled back to M . This situation can be treated in the same spirit as before but is much simplified. If M is locally defined by $\sigma^\alpha = 0$, $\alpha = 1, 2, \dots, \dim \mathcal{M} - \dim M$, one only needs to use $\{d\sigma^\alpha\}$ to generate the bundle K on M . Then K^\perp and $K^\mathcal{G}$ can be similarly defined. The metric splitting of \mathcal{G} directly gives rise to the metric splitting on the reduced Courant algebroid. The reduced metric \tilde{g} is just the restriction of g on TM and the curvature \tilde{H} on M is just the pull-back of H on \mathcal{M} by the inclusion map.

We still use $\tilde{\nabla}^-$ to denote the reduced $-$ -Bismut connection. Let $G^{\alpha\beta} = g(d\sigma^\alpha, d\sigma^\beta)|_M$ and $G_{\alpha\beta}$ be its inverse.

Proposition 4.5. *Let \tilde{X}, \tilde{Y} be vector fields on M , and X, Y be their arbitrary extensions to \mathcal{M} . Then*

$$\tilde{\nabla}_{\tilde{X}}^- \tilde{Y} = \nabla_X^- Y|_M + G_{\alpha\beta}(Y, \nabla_X^- d\sigma^\beta)|_M (g^{-1} d\sigma^\alpha)|_M. \quad (4.4)$$

Proof. We have an orthogonal decomposition $T\mathcal{M}|_M = TM \oplus Q$, where Q is the normal bundle of M in \mathcal{M} and is locally generated by $\{g^{-1} d\sigma^\alpha|_M\}$. It is easy to check that $\tilde{\nabla}_{\tilde{X}}^- \tilde{Y}$ is just the projection of $\nabla_X^- Y|_M$ to TM along Q . This is enough to lead to the conclusion. \square

As for the curvature of $\tilde{\nabla}^-$, we have

Proposition 4.6. *The curvature of $\tilde{\nabla}^-$ is*

$$\begin{aligned} \tilde{g}(\tilde{R}^-(\tilde{X}, \tilde{Y})\tilde{Z}, \tilde{W}) &= g(R^-(X, Y)Z, W)|_M \\ &\quad + G_{\alpha\beta}[(Z, \nabla_Y^- d\sigma^\beta)(W, \nabla_X^- d\sigma^\alpha) - (X \leftrightarrow Y)]|_M, \end{aligned}$$

where $(X \leftrightarrow Y)$ still denotes a term similar to the term in front of it, only with X and Y exchanged.

Proof. We leave the proof to the interested reader. A detailed argument can also be found in [3]. \square

5. Metric generalized principal G-bundles and relative curvatures

In this section, motivated by former observations and also for later use, we investigate generalized principal G -bundles in the presence of an invariant generalized metric.

The notion of generalized principal bundles was introduced in [7] to define generalized holomorphic structures in the setting of principal bundles.

Definition 5.1. A generalized principal G -bundle over M is a triple (P, E, φ) such that

- (i) $p: P \rightarrow M$ is an ordinary principal G -bundle,
- (ii) E is a Courant algebroid over P and φ is an isotropic trivially extended G -action on E .

In contrast with Definition 3.1, the notion of generalized principal bundles hardly contains any essentially new points, but emphasizes a different aspect of the same object. So E descends to M in the same way as before. In the following, we additionally assume that there is a G -invariant generalized metric \mathcal{G} on E and call P a *metric generalized principal bundle*. Then the two connections τ_\pm again arise. Let $\tilde{\nabla}^\pm$ denote the Bismut connections in the base manifold M .

Definition 5.2. Let X, Y be vector fields on M , and X^+, Y^- be their lifts in τ_\pm respectively. The relative curvature of the pair (τ_+, τ_-) is

$$R(X, Y) = (\tilde{\nabla}_X^- Y)^- - (\tilde{\nabla}_Y^+ X)^+ - [X^+, Y^-],$$

where $(\tilde{\nabla}_X^- Y)^-$ is the τ_- -lift of $\tilde{\nabla}_X^- Y$ and $(\tilde{\nabla}_Y^+ X)^+$ is the τ_+ -lift of $\tilde{\nabla}_Y^+ X$.

It is not hard to check that R is tensorial and takes values in the vertical distribution. There is a vector-bundle version of the notion of relative curvature.

Definition 5.3. On a generalized Riemannian manifold (M, g, H) , if a vector bundle W is equipped with two connections ∇^\pm , then the relative curvature of the pair (∇^+, ∇^-) is defined as

$$R(X, Y)s = \nabla_X^+ \nabla_Y^- s - \nabla_Y^- \nabla_X^+ s - \nabla_{\tilde{\nabla}_X^- Y}^- s + \nabla_{\tilde{\nabla}_Y^+ X}^+ s, \quad \forall s \in \Gamma(W),$$

where $\tilde{\nabla}^\pm$ are the Bismut connections in the base manifold M .

Remark. It can be recognized that the relative curvature for a vector bundle is actually part of the curvature of a generalized connection defined in [9]: In the formula for the latter, simply by letting the two arguments take values in V_+ and V_- respectively (recall that V_\pm are the eigenbundles of the generalized metric \mathcal{G}), one recovers a relative curvature.

If $\rho : G \rightarrow \text{End}(W_0)$ is a representation of G in a vector space W_0 , then τ_\pm in the metric generalized principal G -bundle P give rise to two connections ∇^\pm in the associated vector bundle $W_0 \times_\rho P$. It should be pointed out that since by our convention G acts on P from the left, G should act on W_0 from the right; in particular, $\rho_*([v, w]) = -[\rho_*(v), \rho_*(w)]$ for $v, w \in \mathfrak{g}$.

Proposition 5.4. If $R^a V_a$ is the relative curvature of the pair (τ_+, τ_-) in the metric generalized principal G -bundle P , then $R^a \rho_*(e_a)$ is the relative curvature of the pair (∇^+, ∇^-) in the associated vector bundle $W_0 \times_\rho P$.

Proof. Since the computation is essentially local, we can safely assume P is of the form $G \times M$. Let θ_\pm be the connection form of τ_\pm respectively. Then

$$X^+ = X - \theta_+^a(X)V_a, \quad Y^- = Y - \theta_-^a(Y)V_a.$$

Note that

$$\begin{aligned} \tilde{\nabla}_X^- Y - \tilde{\nabla}_Y^+ X &= \tilde{\nabla}_X Y - \tilde{\nabla}_Y X - \frac{1}{2}g^{-1}\tilde{H}(X, Y) - \frac{1}{2}g^{-1}\tilde{H}(Y, X) \\ &= \tilde{\nabla}_X Y - \tilde{\nabla}_Y X \\ &= [X, Y], \end{aligned}$$

where $\tilde{\nabla}$ is the Levi-Civita connection on M and \tilde{H} the curvature of the reduced metric splitting. Thus the relative curvature of the pair (τ_+, τ_-) is

$$\begin{aligned} R(X, Y) &= [-\theta_-^a(\tilde{\nabla}_X^- Y) + \theta_+^a(\tilde{\nabla}_Y^+ X)]V_a + [X, \theta_-^b(Y)V_b] \\ &\quad + [\theta_+^b(X)V_b, Y] - [\theta_+^a(X)V_a, \theta_-^b(Y)V_b] \\ &= [X\theta_-^a(Y) - Y\theta_+^a(X) - \theta_-^a(\tilde{\nabla}_X^- Y) + \theta_+^a(\tilde{\nabla}_Y^+ X)]V_a \\ &\quad - \theta_+^c(X)\theta_-^b(Y)f_{cb}^a V_a. \end{aligned}$$

Let s be the frame of $W_0 \times_\rho P$ corresponding to the trivialization of P . We have

$$\nabla_X^+ \nabla_Y^- s = \nabla_X^+ (s\rho_*(\theta_-(Y))) = s(\rho_*(X\theta_-(Y))) + s\rho_*(\theta_+(X))\rho_*(\theta_-(Y)),$$

and

$$\nabla_Y^- \nabla_X^+ s = \nabla_Y^- (s\rho_*(\theta_+(X))) = s(\rho_*(Y\theta_+(X))) + s\rho_*(\theta_-(Y))\rho_*(\theta_+(X)).$$

Therefore, the relative curvature of the pair (∇^+, ∇^-) is

$$\begin{aligned} R(X, Y) &= \rho_*(X\theta_-(Y)) - \rho_*(Y\theta_+(X)) - \rho_*(\theta_-(\tilde{\nabla}_X^- Y)) + \rho_*(\theta_+(\tilde{\nabla}_Y^+ X)) \\ &\quad + \rho_*(\theta_+(X))\rho_*(\theta_-(Y)) - \rho_*(\theta_-(Y))\rho_*(\theta_+(X)) \\ &= \rho_*(X\theta_-(Y)) - \rho_*(Y\theta_+(X)) - \rho_*(\theta_-(\tilde{\nabla}_X^- Y)) + \rho_*(\theta_+(\tilde{\nabla}_Y^+ X)) \\ &\quad - \rho_*([\theta_+(X), \theta_-(Y)]). \end{aligned}$$

The claim then follows. \square

Now let us go back to the context of Section 4 and view M as a metric generalized principal G -bundle over M_{red} . We want to derive a formula for the relative curvature of the pair (τ_+, τ_-) in terms of the data of the isotropic trivially extended G -action.

By Eq. (4.1) and Eq. (4.3),

$$\varrho_-([X], [Y]) = \nabla_{X^+}^- Y^- + T^{ab} g(Y^-, \nabla_{X^+}^- V_b^-) V_a^-,$$

and

$$\varrho_+([Y], [X]) = \nabla_{Y^-}^+ X^+ + T^{ab} g(X^+, \nabla_{Y^-}^+ V_b^+) V_a^+.$$

Therefore, the relative curvature is

$$\begin{aligned} R([X], [Y]) &= T^{ab} [g(Y^-, \nabla_{X^+}^- V_b^-) V_a^- - g(X^+, \nabla_{Y^-}^+ V_b^+) V_a^+] \\ &= T^{ab} [g(Y^-, \nabla_{X^+}^- V_b^-) - g(X^+, \nabla_{Y^-}^+ V_b^+)] V_a \\ &\quad - T^{ab} [g(Y^-, \nabla_{X^+}^- V_b^-) + g(X^+, \nabla_{Y^-}^+ V_b^+)] g^{-1} \xi_a. \end{aligned}$$

A simple calculation shows $g(Y^-, \nabla_{X^+}^- V_b^-) + g(X^+, \nabla_{Y^-}^+ V_b^+) = 0$. We finally have

$$R^a([X], [Y]) = -2T^{ab} g(\nabla_{X^+}^- Y^-, V_b^-). \quad (5.1)$$

If additionally M together with its structure of a metric generalized principal G -bundle comes as an invariant submanifold of a bigger manifold \mathcal{M} , which carries an isotropic trivially extended G -action and a G -invariant generalized metric, we can express the above result in terms of extensions \check{X}, \check{Y} of X^+, Y^- on \mathcal{M} . Let \check{g} be the metric on \mathcal{M} and $\check{\nabla}^\pm$ be Bismut connections on \mathcal{M} . If $M \subset \mathcal{M}$ is locally defined by $\mu^\alpha = 0$ for $\alpha = 1, 2, \dots, \dim \mathcal{M} - \dim M$, then by Eq. (4.4), we have

$$R^a([X], [Y]) = -2T^{ab} \check{g}(\check{\nabla}_{\check{X}}^- \check{Y}, V_b^-)|_M + 2T^{ab} G_{\alpha\beta} d\mu^\alpha(\check{\nabla}_{\check{X}}^- \check{Y})|_M d\mu^\beta(V_b^-)|_M, \quad (5.2)$$

where $G_{\alpha\beta}$ is the inverse of $G^{\alpha\beta} = \check{g}(d\mu^\alpha, d\mu^\beta)|_M$. This formula will be crucial in Section 7.

6. Generalized Kähler reduction

In the framework of [1] or [2], the reduction of a G -invariant generalized Kähler manifold \mathcal{M} involves two stages: (i) a G -invariant submanifold $M \subset \mathcal{M}$ is singled out, possibly by the zero-level set of an equivariant map $\mu : \mathcal{M} \rightarrow \mathfrak{h}^*$, where \mathfrak{h}^* is the dual of a \mathfrak{g} -module \mathfrak{h} . At the same time, the bundle K over M , locally generated by $\{V_a + \xi_a\}$ and $\{d\mu^\alpha\}$, is constructed. Then K^\perp is again defined as the orthogonal complement of K in $\mathbb{T}\mathcal{M}|_M$ and one gets the important bundle $K^\mathcal{G} = K^\perp \cap \mathcal{G}(K^\perp)$ over M . (ii) If \mathbb{J}_1 preserves $K^\mathcal{G}$, i.e.,

$$\mathbb{J}_1 K^\mathcal{G} = K^\mathcal{G}, \quad (6.1)$$

then $K^\mathcal{G}/G$ naturally acquires two complex structures. Since $K^\mathcal{G}/G$ can be identified with E_{red} , these are actually almost generalized complex structures on $M_{red} := M/G$. Integrability of these structures stems from the general reduction theory of Dirac structures.

We prefer to put things in another way: One can first realize the metric reduction from \mathcal{M} to M . With this in place, we are in the situation of Section 3 and can then realize the metric reduction from M to M_{red} . Now as before, there are two connections τ_\pm in M as a metric generalized principal G -bundle. Then Eq. (6.1) simply means $J_\pm \tau_\pm = \tau_\pm$, i.e. J_\pm preserve the two distributions on M respectively, where J_\pm are the underlying complex structures on \mathcal{M} .

Although integrability of the reduced generalized Kähler structure in M_{red} is almost obvious from the more general viewpoint, we would like to provide another proof of this fact, which fits in well with our viewpoint towards metric reduction. This approach is a bit more complicated and indirect, but may shed some new light on generalized Kähler reduction. Note that in the following, a Courant algebroid \mathcal{E} on \mathcal{M} carrying an isotropic trivially extended G -action is understood as the basic background.

Theorem 6.1 ([1,2]). *Let \mathcal{M} be a G -invariant generalized Kähler manifold and M a G -invariant submanifold such that $J_\pm \tau_\pm = \tau_\pm$. Then the generalized Kähler structure descends to the reduced manifold M_{red} .*

Proof. As the reduced generalized Kähler structure is well understood in the literature, we only pay attention to the integrability condition.

Since TM_{red} is modelled on both τ_+ and τ_- on M , thus $J_\pm \tau_\pm = \tau_\pm$ implies that M_{red} has two almost complex structures \tilde{J}_\pm . The compatibility of \tilde{J}_\pm with the reduced metric \tilde{g} is also obvious. In this situation, according to [4], to obtain the conclusion, we need to prove (i) $\tilde{\nabla}^\pm \tilde{J}_\pm = 0$ and (ii) \tilde{H} is of type $(2, 1) + (1, 2)$ w.r.t. both \tilde{J}_\pm . Note that here $\tilde{\nabla}^\pm$ are the Bismut connections on M_{red} and \tilde{H} is the curvature of the reduced metric splitting.

We use \check{X} to denote an extension of $X^+ \in \Gamma(\tau_+)$ or $X^- \in \Gamma(\tau_-)$ to \mathcal{M} and the Bismut connections in \mathcal{M} are denoted by $\check{\nabla}^\pm$. By Eq. (4.3) and an analogue of Eq. (4.4) for $\check{\nabla}^+$, we have

$$\begin{aligned}(\check{\nabla}_{[X]}^+ \check{J}_+ [Y], [Z]) &= (\nabla_{X^+}^+ J_+ Y^+, Z^+) = (\check{\nabla}_{\check{X}}^+ J_+ \check{Y}, \check{Z})|_M \\&= (J_+ \check{\nabla}_{\check{X}}^+ \check{Y}, \check{Z})|_M = -(\check{\nabla}_{\check{X}}^+ \check{Y}, J_+ \check{Z})|_M \\&= -(\nabla_{X^+}^+ Y^+, J_+ Z^+) = -(\check{\nabla}_{[X]}^+ [Y], \check{J}_+ [Z]) \\&= (\check{J}_+ \check{\nabla}_{[X]}^+ [Y], [Z]),\end{aligned}$$

where the fact $\check{\nabla}^+ J_+ = 0$ is used. We thus have proved that $\check{\nabla}^+ \check{J}_+ = 0$. $\check{\nabla}^- \check{J}_- = 0$ can be proved similarly.

To see that \check{H} is of type $(2, 1) + (1, 2)$ w.r.t. \check{J}_+ , since by Proposition 4.4

$$\check{H}([X], [Y], [Z]) = (\check{H}|_M + \Omega_+^a \wedge \xi_a)(X^+, Y^+, Z^+),$$

and the curvature \check{H} of the metric splitting of \mathcal{E} is of type $(2, 1) + (1, 2)$ w.r.t. J_+ , we only need to prove Ω_+^a is of type $(1, 1)$, namely

$$\Omega_+^a(X^+, Y^+) = 0$$

if $X^+, Y^+ \in (\tau_+ \otimes \mathbb{C}) \cap T_{0,1}^+ \mathcal{M}|_M$. Note that $\Omega_+^a = K^{ba} d\xi_b^+$, we should prove $d\xi_a^+(X^+, Y^+) = 0$. In fact,

$$\begin{aligned}d\xi_a^+(X^+, Y^+) &= [dg(V_a) + d\xi_a](X^+, Y^+) = X^+ g(V_a, Y^+) - Y^+ g(V_a, X^+) \\&\quad - g(V_a, [X^+, Y^+]) + \check{H}|_M(V_a, X^+, Y^+) \\&= \sqrt{-1} X^+ \omega_{+|M}(V_a^A, Y^+) - \sqrt{-1} Y^+ \omega_{+|M}(V_a^A, X^+) \\&\quad - \sqrt{-1} \omega_{+|M}(V_a^A, [X^+, Y^+]) + \check{H}|_M(V_a^A, X^+, Y^+),\end{aligned}$$

where V_a^A is the $T_{0,1}^+ \mathcal{M}$ -part of V_a and we have used the following two facts: (i) \check{g} and J_+ are compatible and $\omega_+ = \check{g}J_+$; (ii) \check{H} is of type $(2, 1) + (1, 2)$ w.r.t. J_+ . Consequently we have

$$\begin{aligned}d\xi_a^+(X^+, Y^+) &= -\sqrt{-1} d(\omega_{+|M})(X^+, Y^+, V_a^A) + \check{H}|_M(V_a^A, X^+, Y^+) \\&= -\sqrt{-1} (d\omega_{+|M})(X^+, Y^+, V_a^A) + \check{H}|_M(X^+, Y^+, V_a^A) \\&= -(d\omega_{+|M})(J_+ X^+, J_+ Y^+, J_+ V_a^A) + \check{H}|_M(X^+, Y^+, V_a^A) \\&= (d_+^c \omega_{+|M} + \check{H})|_M(X^+, Y^+, V_a^A) \\&= 0,\end{aligned}$$

where we have used the identity $d_+^c \omega_{+|M} + \check{H} = 0$ on \mathcal{M} . Similarly, the curvature Ω_-^a of τ_- is of type $(1, 1)$ w.r.t. \check{J}_- and \check{H} is of type $(2, 1) + (1, 2)$ w.r.t. \check{J}_- . \square

Remark. The proof has some byproducts. It implies that M as a principal G -bundle over generalized Kähler manifold M_{red} carries two connections τ_\pm whose curvatures are of type $(1, 1)$ w.r.t. \check{J}_\pm respectively. Thus any associated complex vector bundle W naturally has a biholomorphic structure, i.e. W is holomorphic simultaneously w.r.t. both of \check{J}_\pm . Such vector bundles play a basic role in the work [10] to find an analogue of Hermite–Einstein equations in the context of biHermitian manifolds. Thus generalized Kähler reduction actually provides examples of biholomorphic structures.

7. Generalized holomorphic structures from generalized Kähler reduction

Generalized holomorphic vector bundles are analogues of holomorphic vector bundles in complex geometry. Due to the observation in the end of Section 6, it is natural to ask whether generalized holomorphic vector bundles could arise as byproducts of generalized Kähler reduction. The goal of this section is mainly to provide some examples to give an affirmative answer to this question. We will continue to use notation in Section 6. Note that $M \subset \mathcal{M}$ is actually a metric generalized principal G -bundle, carrying the pair (τ_+, τ_-) of connections. Let $\tau_\pm^{0,1}$ denote the $-\sqrt{-1}$ -eigensubbundles of $\tau_\pm \otimes \mathbb{C}$ w.r.t. J_\pm respectively.

Lemma 7.1. Any associated complex vector bundle W of M as a principal G -bundle is naturally generalized holomorphic if the relative curvature R of the pair (τ_+, τ_-) satisfies

$$R([X], [Y]) = 0, \quad \forall [X] \in T_{0,1}^+ M_{red}, \quad [Y] \in T_{0,1}^- M_{red}. \quad (7.1)$$

Proof. Let ∇^{W^\pm} be the connections in W determined by τ_\pm in M . They can be combined to give a generalized connection \mathcal{D} in W in the sense of [9]. In fact, Since $A \in \Gamma(E_{red})$ can be uniquely written as

$$A = [X] + \check{g}([X]) + [Y] - \check{g}([Y])$$

for some $[X], [Y] \in \Gamma(TM_{red})$ due to the decomposition $E_{red} = V_+^{red}/G \oplus V_-^{red}/G$, we can define

$$\mathcal{D}_A s = \nabla_{[X]}^W s + \nabla_{[Y]}^W s, \quad s \in \Gamma(W).$$

\mathcal{D} can be additionally decomposed according to the decomposition $E_{red} \otimes \mathbb{C} = L_1^{red} \oplus \bar{L}_1^{red}$, where L_1^{red} is the $\sqrt{-1}$ -eigenbundle of the reduced generalized complex structure \mathbb{J}_1^{red} of \mathbb{J}_1 . Let $\bar{\partial}$ be the \bar{L}_1^{red} -part of \mathcal{D} . $\bar{\partial}$ can be further decomposed as $\bar{\partial} = \bar{\partial}_+ + \bar{\partial}_-$ where actually $\bar{\partial}_\pm$ are just the natural \bar{J}_\pm -holomorphic structures in W induced from ∇^{W^\pm} . Thus $\bar{\partial}$ is a generalized holomorphic structure iff Eq. (2.4) is satisfied. It is not hard to find this is exactly Eq. (7.1). \square

To provide concrete examples, we specify to the case of Hamiltonian generalized Kähler manifolds introduced by Lin and Tolman in [11]. Let \mathcal{M} be a G -invariant generalized Kähler manifold. The extended G -action is called Hamiltonian if there is an equivariant map $\mu : \mathcal{M} \rightarrow \mathfrak{g}^*$ (\mathfrak{g}^* carries the coadjoint action) such that

$$\mathbb{J}_2(V_a + \xi_a) = d\mu_a, \quad a = 1, 2, \dots, \dim \mathcal{M} \quad (7.2)$$

where $\mu_a = \mu(e_a)$. According to Lemma 4.2 in [12], in terms of the biHermitian data, Eq. (7.2) is equivalent to

$$J_+ V_a^+ = J_- V_a^- = -\check{g}^{-1} d\mu_a. \quad (7.3)$$

If G acts freely on $M = \mu^{-1}(0)$, then Eq. (6.1) naturally follows and $M_{red} = \mu^{-1}(0)/G$ carries a reduced generalized Kähler structure [11].

Recall that we use $\check{X} \in \Gamma(T\mathcal{M} \otimes \mathbb{C})$ to denote an extension of vector field $X^+ \in \Gamma(\tau_+ \otimes \mathbb{C})$ or $X^- \in \Gamma(\tau_- \otimes \mathbb{C})$ on M .

Theorem 7.2. Assume the extended G -action on generalized Kähler manifold \mathcal{M} is Hamiltonian. Then Eq. (7.1) is satisfied if

$$d\mu(\check{\nabla}_{\check{X}}^{\check{Y}})|_M = 0, \quad (7.4)$$

for any $X^+ \in \Gamma(\tau_+^{0,1})$ and $Y^- \in \Gamma(\tau_-^{0,1})$.

Remark. Eq. (7.4) means $\check{\nabla}_{\check{X}}^{\check{Y}}$ should be tangent to M .

Proof. As the result is obviously independent of which extensions we choose, we can safely assume that $\check{X} \in \Gamma(T_{0,1}^+ \mathcal{M})$ and $\check{Y} \in \Gamma(T_{0,1}^- \mathcal{M})$.

Due to Eq. (7.3), for any $Z \in \Gamma(T_{0,1}^- \mathcal{M})$ we have

$$\check{g}(Z, V_a^-) = \sqrt{-1} d\mu_a(Z).$$

Since $\check{\nabla}^- J_- = 0$, we have $\check{\nabla}_{\check{X}}^{\check{Y}} \in \Gamma(T_{0,1}^- \mathcal{M})$ and thus

$$\check{g}(\check{\nabla}_{\check{X}}^{\check{Y}}, V_a^-) = \sqrt{-1} d\mu_a(\check{\nabla}_{\check{X}}^{\check{Y}}).$$

Substituting this result in Eq. (5.2), we find $R^a([X], [Y])$ is of the form

$$R^a([X], [Y]) = -2\sqrt{-1} T^{ab} (\delta_b^c + \sqrt{-1} S_b^c) d\mu_c(\check{\nabla}_{\check{X}}^{\check{Y}})|_M,$$

for some real-valued functions S_b^c on M . The conclusion immediately follows. \square

We are now in a position to construct some generalized holomorphic bundles by generalized Kähler reduction. Here we content ourselves with a family of generalized holomorphic line bundles on \mathbb{CP}^2 equipped with certain non-trivial generalized Kähler structures. We follow the ideas of [4, 11] to deform the standard Kähler structure on \mathbb{C}^3 (as a generalized Kähler manifold) to new S^1 -invariant generalized Kähler structures while keeping the standard symplectic structure fixed. Applying the Marsden–Weinstein reduction to the symplectic structure then gives non-trivial generalized Kähler structures on \mathbb{CP}^2 . By choosing the deformations properly, we would obtain metric generalized principal S^1 -bundles over \mathbb{CP}^2 whose associated line bundles are generalized holomorphic.

Let us recall the deformation theory of [4] in some detail. Given a generalized Kähler structure on \mathcal{M} and $\epsilon \in \Gamma(\wedge^2 \bar{L}_1)$, define $L_1^\epsilon = \{X + \iota_X \epsilon | X \in L_1\}$. For ϵ small enough, L_1^ϵ is an almost generalized complex structure. The integrability condition of this deformation is the Maurer–Cartan equation

$$d_{L_1} \epsilon + \frac{1}{2} [\epsilon, \epsilon]_S = 0, \quad (7.5)$$

where $[\cdot, \cdot]_S$ is the Schouten bracket induced from the Lie algebroid L_1 . Note that since $L_1 = L_+ \oplus L_-$ and $L_2 = L_+ \oplus \bar{L}_-$, to keep L_2 fixed we should take $\epsilon \in \Gamma(\bar{L}_+ \otimes \bar{L}_-)$.

The standard Kähler structure on \mathbb{C}^3 . Let $\mathcal{M} = \mathbb{C}^3$ with its canonical complex structure J and Kähler structure $\omega = \sqrt{-1} \sum_{i=0}^2 dz_i \wedge d\bar{z}_i$. Let $E_i = \partial_{z_i} + d\bar{z}_i$, $F_i = \partial_{z_i} - d\bar{z}_i$, $i = 0, 1, 2$. We have two complex vector bundles $L_+ = \text{span}\{\bar{E}_i\}$ and $L_- = \text{span}\{F_i\}$ over \mathcal{M} . Viewed as a generalized Kähler manifold, \mathcal{M} has $L_1 = L_+ \oplus L_-$ and $L_2 = L_+ \oplus \bar{L}_-$ as the associated two generalized complex structures. A pure spinor of L_1 is $\varphi_1 = dz_0 dz_1 dz_2$ and a pure spinor of L_2 is $\varphi_2 = e^{-\sqrt{-1}\omega}$. Note that

in the present setting d_{L_1} is just the classical Dolbeault operator associated to J and from the biHermitian viewpoint, we have chosen $J_+ = J_- = J$.

S^1 acts on \mathcal{M} by scaling:

$$e^{i\theta} \cdot (z_0, z_1, z_2) = (e^{i\theta} z_0, e^{i\theta} z_1, e^{i\theta} z_2).$$

The infinitesimal action of S^1 is generated by the vector field

$$\partial_\theta = \sqrt{-1} \sum_{i=0}^2 (z_i \partial_{z_i} - \bar{z}_i \partial_{\bar{z}_i}).$$

Then $\mu = \sum_{i=0}^2 |z_i|^2 - 1$ is a moment map. By Kähler reduction, $\mu^{-1}(0)/S^1$ is a Kähler manifold. This is the canonical Kähler structure on the projective plane \mathbb{CP}^2 .

Deformations of the Kähler structure. We choose

$$\epsilon = \frac{1}{2} \left(\sum_{i=0}^2 f_i E_i \right) \wedge \left(\sum_{j=0}^2 g_j F_j \right),$$

where f_i, g_i are homogeneous polynomials of z_0, z_1 and z_2 to be determined. Since f_i, g_i are holomorphic, the integrability condition (7.5) amounts to the following equations:

$$\begin{cases} \sum_{p=0}^2 f_p (g_k \partial_{z_p} g_q - g_q \partial_{z_p} g_k) = 0, \\ \sum_{p=0}^2 g_p (f_k \partial_{z_p} f_q - f_q \partial_{z_p} f_k) = 0. \end{cases} \quad (7.6)$$

There are many solutions to these equations. We list two as follows: (i) $g_0 = g_1 = f_1 = f_2 = 0, g_2 = 1$ and $f_0 = z_0^2$; (ii) $g_i = 1, i = 0, 1, 2$ and

$$f_0 = (z_1 - z_0)(z_2 - z_0), \quad f_1 = (z_0 - z_1)(z_2 - z_1), \quad f_2 = (z_0 - z_2)(z_1 - z_2). \quad (7.7)$$

Note that deformations associated to the two solutions are both S^1 -invariant. Since we are only concerned with the behaviour of ϵ over $M = \mu^{-1}(0) = S^5$, we can simply multiply ϵ by a nonzero complex number λ such that $|\lambda|$ is small enough and then L_1^ϵ and L_2 together define a generalized Kähler structure on a bounded neighbourhood of M in \mathbb{C}^3 .

In the following, for simplicity, we will set $g_0 = g_1 = g_2 = 1$. In this case, the first equation of (7.6) holds trivially and the second equation can be written in a more compact form:

$$\sum_{p=0}^2 \partial_{z_p} \left(\frac{f_q}{f_k} \right) = 0.$$

Thus we can choose f_i to be functions of $z_1 - z_0$ and $z_2 - z_0$, e.g. our solution (ii) is such a choice. To make ϵ be S^1 -invariant, we additionally require f_i to be of degree 2. Now we have

$$L_+^\epsilon = \text{span}\{\bar{E}_i + f_i \sum_{p=0}^2 F_p\}, \quad L_-^\epsilon = \text{span}\{\bar{F}_i + \sum_{p=0}^2 f_p E_p\},$$

and $L_1^\epsilon = L_+^\epsilon \oplus L_-^\epsilon, L_2 = L_+^\epsilon \oplus \bar{L}_-^\epsilon$. Let J_\pm^ϵ be the underlying complex structures. Accordingly,

$$T_{0,1}^+ \mathcal{M} = \text{span}\{\partial_{\bar{z}_i} + f_i \sum_{p=0}^2 \partial_{z_p}\}, \quad T_{0,1}^- \mathcal{M} = \text{span}\{\partial_{\bar{z}_i} + \sum_{p=0}^2 f_p \partial_{z_p}\}.$$

It should be pointed out that generally with the above form of L_1^ϵ and L_2 , we are not in the metric splitting, but this will not bother us much. Though Eq. (2.2) is written in the metric splitting, other splittings are equally fine, because B -transforms will not affect the tangent part of the equation, which is essential to obtain Bismut connections.

The reduced generalized Kähler structure. We content ourselves with a glance at the reduced generalized Kähler structure, for we are more interested in the generalized holomorphic line bundles produced by the reduction procedure. The reduced generalized Kähler structure can be described conveniently in terms of pure spinors. We refer the interested reader to [13] for a detailed account of pure spinors in the setting of generalized reduction.

A pure spinor of L_1^ϵ is $\varphi_1^\epsilon = e^{-\epsilon} \cdot \varphi_1$. To find a pure spinor for its reduction $\tilde{\mathbb{J}}_1^\epsilon$ on \mathbb{CP}^2 , roughly speaking, one simply pulls back φ_1^ϵ to $M = S^5$ and then pushes it forward to the quotient M/S^1 (i.e. contraction with ∂_θ on M). The two stages can be exchanged. Thus one first contracts φ_1^ϵ with ∂_θ on \mathcal{M} and then pulls back the result to S^5 . By ‘roughly’, we mean actually before pulled back, $\iota_{\partial_\theta} \varphi_1^\epsilon$ should be scaled to be S^1 -invariant. The scaling cannot be realized globally, and when it breaks down, type-jumping of $\tilde{\mathbb{J}}_1^\epsilon$ occurs. A generic point on \mathbb{CP}^2 is of type 0 for $\tilde{\mathbb{J}}_1^\epsilon$, and for a point on the type-jumping locus, the type jumps to 2. A similar and more detailed analysis of some reduced generalized Kähler structures on \mathbb{CP}^2 can be found in [1]. Our argument above is along the same line of [1].

To determine the type-jumping locus of \mathbb{J}_1^ϵ , we take the 0-form component of $\iota_{\partial_{\bar{\theta}}} \varphi_1^\epsilon$. This produces a section ρ of the dual bundle of the pure spinor line bundle of \mathbb{J}_1^ϵ . The zero-locus of ρ is precisely the type-jumping locus, which is singled out by the following homogeneous equation of degree 3:

$$z_0(f_1 - f_2) + z_1(f_2 - f_0) + z_2(f_0 - f_1) = 0.$$

This can also be found using the type formula in [11]. For example, if we choose the solution (7.7) the type-jumping locus consists of three lines in \mathbb{CP}^2 : $z_0 = z_1$, $z_0 = z_2$ and $z_1 = z_2$ with a joint-point $[1 : 1 : 1]$. The type-jumping locus can be viewed as a degenerate elliptic curve.

Generalized holomorphic structures from reduction. We now check that $M = S^5$ as a metric generalized principal S^1 -bundle really gives its associated line bundles a generalized holomorphic structure, i.e., Eq. (7.4) does hold for the present setting.

Let us find $\tau_{\pm}^{0,1}$ first. Due to the proof of Theorem 7.2, we only need to find Z in $T_{0,1}^{\pm} \mathcal{M}$ such that $d\mu(Z) = 0$. Denote $z = z_0 + z_1 + z_2$ and $F = F_0 + F_1 + F_2$. The image of g on $\tau_{+}^{0,1}$ is spanned by

$$A_1 := -(z_1 + f_1 \bar{z})(\bar{E}_0 + f_0 F) + (z_0 + f_0 \bar{z})(\bar{E}_1 + f_1 F)$$

and

$$A_2 := -(z_2 + f_2 \bar{z})(\bar{E}_0 + f_0 F) + (z_0 + f_0 \bar{z})(\bar{E}_2 + f_2 F)$$

on S^5 . Denote $h = f_0 \bar{z}_0 + f_1 \bar{z}_1 + f_2 \bar{z}_2$ and $C = f_0 E_0 + f_1 E_1 + f_2 E_2$. The image of $-g$ on $\tau_{-}^{0,1}$ is spanned by

$$B_1 := -(z_1 + h)(\bar{F}_0 + C) + (z_0 + h)(\bar{F}_1 + C)$$

and

$$B_2 := -(z_2 + h)(\bar{F}_0 + C) + (z_0 + h)(\bar{F}_2 + C)$$

on S^5 . By abuse of notation, we also use A_i, B_i to denote their extensions on \mathcal{M} with the same expressions.

What left is to check that the tangent part of $[A_i, B_j]^-$ is tangent to M for $i, j = 1, 2$ according to Theorem 7.2 and Eq. (2.2). We will only compute $[A_1, B_1]^-$ and $[A_2, B_1]^-$ and the details are included in the appendix. The computation of $[A_1, B_2]^-$ and $[A_2, B_2]^-$ is similar and left to the interested reader.

The tangent part of $[A_1, B_1]^-$ is³

$$\begin{aligned} & (z_1 + f_1 \bar{z})(z_1 - z_0) \sum_{q=0}^2 \partial_{z_q} f_0 \partial_{\bar{z}_q} + 2f_0(z_1 + f_1 \bar{z})(\partial_{\bar{z}_0} - \partial_{\bar{z}_1}) \\ & + (z_0 + f_0 \bar{z})(z_0 - z_1) \sum_{q=0}^2 \partial_{z_q} f_1 \partial_{\bar{z}_q} + 2f_1(z_0 + f_0 \bar{z})(\partial_{\bar{z}_1} - \partial_{\bar{z}_0}). \end{aligned}$$

Its contraction with $d\mu$ is, up a common factor $z_1 - z_0$,

$$\begin{aligned} & (z_1 + f_1 \bar{z}) \sum_{q=0}^2 z_q \partial_{z_q} f_0 - 2f_0(z_1 + f_1 \bar{z}) \\ & - (z_0 + f_0 \bar{z}) \sum_{q=0}^2 z_q \partial_{z_q} f_1 + 2f_1(z_0 + f_0 \bar{z}) \\ & = (z_1 + f_1 \bar{z}) \left(\sum_{q=0}^2 z_q \partial_{z_q} f_0 - 2f_0 \right) - (z_0 + f_0 \bar{z}) \left(\sum_{q=0}^2 z_q \partial_{z_q} f_1 - 2f_1 \right) \\ & = 0, \end{aligned}$$

where the last equality is due to the fact that f_0 and f_1 are homogeneous functions of degree 2.

Similarly, the tangent part of $[A_2, B_1]^-$ is

$$\begin{aligned} & (z_2 + f_2 \bar{z})(z_1 - z_0) \sum_{q=0}^2 \partial_{z_q} f_0 \partial_{\bar{z}_q} + 2f_0(z_2 + f_2 \bar{z})(\partial_{\bar{z}_0} - \partial_{\bar{z}_1}) \\ & - (z_0 + f_0 \bar{z})(z_1 - z_0) \sum_{q=0}^2 \partial_{z_q} f_2 \partial_{\bar{z}_q} + 2f_2(z_0 + f_0 \bar{z})(\partial_{\bar{z}_1} - \partial_{\bar{z}_0}). \end{aligned}$$

³ Note that to obtain the expression, the equation $\sum_{p=0}^2 \partial_{z_p} f_i = 0$ is used.

Its contraction with $d\mu$ is, up to a common factor $z_1 - z_0$,

$$\begin{aligned} & (z_2 + f_2\bar{z}) \sum_{q=0}^2 z_q \partial_{z_q} f_0 - 2f_0(z_2 + f_2\bar{z}) \\ & - (z_0 + f_0\bar{z}) \sum_{q=0}^2 z_q \partial_{z_q} f_2 + 2f_2(z_0 + f_0\bar{z}) \\ & = (z_2 + f_2\bar{z}) \left(\sum_{q=0}^2 z_q \partial_{z_q} f_0 - 2f_0 \right) - (z_0 + f_0\bar{z}) \left(\sum_{q=0}^2 z_q \partial_{z_q} f_2 - 2f_2 \right) \\ & = 0, \end{aligned}$$

where the last equality is due to the fact that f_0 and f_2 are homogeneous functions of degree 2.

A similar computation shows the tangent parts of $[A_1, B_2]^-$ and $[A_2, B_2]^-$ are each tangent to M . Therefore, we have checked that any associated line bundle of M as a principal S^1 -bundle is generalized holomorphic.

The associated line bundle of the canonical representation of S^1 is actually the tautological line bundle of \mathbb{CP}^2 , and thus its first Chern class is $-[I]$, where $[I]$ denotes the homology class represented by a line in \mathbb{CP}^2 . But the first Chern class of the canonical line bundle of $\mathbb{P}_1^\mathbb{C}$ is $-3[I]$. Therefore our construction really gives rise to new generalized holomorphic line bundles.

It is expected that the approach illustrated here could also be applied to construct generalized holomorphic vector bundles of higher rank. We will turn to this elsewhere in the future.

Acknowledgment

This study is supported by the Natural Science Foundation of Jiangsu Province, China (Grant no. BK20150797) and by the China Scholarship Council (Grant no. 201806715027).

Appendix

This appendix contains the detailed computation of $[A_1, B_1]^-$ and $[A_2, B_1]^-$.

First a direct computation gives the following formula to be used later:

$$[\bar{E}_i + f_i F, \bar{F}_j + C] = \sum_{q=0}^2 \partial_{z_q} f_i (\bar{F}_q + C) - \sum_{q=0}^2 \partial_{z_q} f_i (\bar{E}_q + f_q F).$$

Note that the result is independent of j .

$$\begin{aligned} [A_1, B_1]^- &= [(z_1 + f_1\bar{z})(\bar{E}_0 + f_0 F), (z_1 + h)(\bar{F}_0 + C)]^- \\ &- [(z_1 + f_1\bar{z})(\bar{E}_0 + f_0 F), (z_0 + h)(\bar{F}_1 + C)]^- \\ &- [(z_0 + f_0\bar{z})(\bar{E}_1 + f_1 F), (z_1 + h)(\bar{F}_0 + C)]^- \\ &+ [(z_0 + f_0\bar{z})(\bar{E}_1 + f_1 F), (z_0 + h)(\bar{F}_1 + C)]^- \\ &= (z_1 + f_1\bar{z})(z_1 + h) \Sigma_{q=0}^2 \partial_{z_q} f_0 (\bar{F}_q + C) \\ &+ 2f_0(z_1 + f_1\bar{z})(\bar{F}_0 + C) - 2f_0(z_1 + f_1\bar{z})(\bar{F}_1 + C) \\ &- (z_1 + f_1\bar{z})(z_0 + h) \Sigma_{q=0}^2 \partial_{z_q} f_0 (\bar{F}_q + C) \\ &- (z_0 + f_0\bar{z})(z_1 + h) \Sigma_{q=0}^2 \partial_{z_q} f_1 (\bar{F}_q + C) \\ &- 2f_1(z_0 + f_0\bar{z})(\bar{F}_0 + C) + 2f_1(z_0 + f_0\bar{z})(\bar{F}_1 + C) \\ &+ (z_0 + f_0\bar{z})(z_0 + h) \Sigma_{q=0}^2 \partial_{z_q} f_1 (\bar{F}_q + C) \\ &= (z_1 + f_1\bar{z})(z_1 - z_0) \Sigma_{q=0}^2 \partial_{z_q} f_0 \bar{F}_q \\ &+ 2f_0(z_1 + f_1\bar{z})(\bar{F}_0 - \bar{F}_1) \\ &+ (z_0 + f_0\bar{z})(z_0 - z_1) \Sigma_{q=0}^2 \partial_{z_q} f_1 \bar{F}_q \\ &+ 2f_1(z_0 + f_0\bar{z})(\bar{F}_1 - \bar{F}_0). \end{aligned}$$

Similarly,

$$\begin{aligned}
 [A_2, B_1]^- &= [(z_2 + f_2 \bar{z})(\bar{E}_0 + f_0 F), (z_1 + h)(\bar{F}_0 + C)]^- \\
 &\quad - [(z_2 + f_2 \bar{z})(\bar{E}_0 + f_0 F), (z_0 + h)(\bar{F}_1 + C)]^- \\
 &\quad - [(z_0 + f_0 \bar{z})(\bar{E}_2 + f_2 F), (z_1 + h)(\bar{F}_0 + C)]^- \\
 &\quad + [(z_0 + f_0 \bar{z})(\bar{E}_2 + f_2 F), (z_0 + h)(\bar{F}_1 + C)]^- \\
 &= (z_2 + f_2 \bar{z})(z_1 + h) \Sigma_{q=0}^2 \partial_{z_q} f_0 (\bar{F}_q + C) \\
 &\quad + 2f_0(z_2 + f_2 \bar{z})(\bar{F}_0 + C) - 2f_0(z_2 + f_2 \bar{z})(\bar{F}_1 + C) \\
 &\quad - (z_2 + f_2 \bar{z})(z_0 + h) \Sigma_{q=0}^2 \partial_{z_q} f_0 (\bar{F}_q + C) \\
 &\quad - (z_0 + f_0 \bar{z})(z_1 + h) \Sigma_{q=0}^2 \partial_{z_q} f_2 (\bar{F}_q + C) \\
 &\quad - 2f_2(z_0 + f_0 \bar{z})(\bar{F}_0 + C) + 2f_2(z_0 + f_0 \bar{z})(\bar{F}_1 + C) \\
 &\quad + (z_0 + f_0 \bar{z})(z_0 + h) \Sigma_{q=0}^2 \partial_{z_q} f_2 (\bar{F}_q + C) \\
 &= (z_2 + f_2 \bar{z})(z_1 - z_0) \Sigma_{q=0}^2 \partial_{z_q} f_0 \bar{F}_q \\
 &\quad + 2f_0(z_2 + f_2 \bar{z})(\bar{F}_0 - \bar{F}_1) \\
 &\quad + (z_0 + f_0 \bar{z})(z_0 - z_1) \Sigma_{q=0}^2 \partial_{z_q} f_2 \bar{F}_q \\
 &\quad + 2f_2(z_0 + f_0 \bar{z})(\bar{F}_1 - \bar{F}_0).
 \end{aligned}$$

References

- [1] H. Bursztyn, G.R. Cavalcanti, M. Gualtieri, Reduction of Courant algebroids and generalized complex structures, *Adv. Math.* 211 (2) (2007) 726–765.
- [2] G.R. Cavalcanti, Reduction of metric structures on Courant algebroid, *J. Symplectic Geom.* 4 (3) (2006) 317–343.
- [3] Y. Wang, Metric reduction in generalized geometry and balanced topological field theories. [arXiv:1708.00567](https://arxiv.org/abs/1708.00567).
- [4] M. Gualtieri, Generalized Complex Geometry (Ph.D. thesis), Oxford University, 2003, [arXiv:math/0401221](https://arxiv.org/abs/math/0401221).
- [5] M. Gualtieri, Generalized complex geometry, *Ann. of Math.* 174 (2011) 75–123.
- [6] Y. Wang, Generalized holomorphic structures, *J. Geom. Phys.* 61 (2011) 1976–1984.
- [7] Y. Wang, generalized holomorphic structures, *J. Geom. Phys.* 86 (2014) 273–283.
- [8] N. Hitchin, Lectures on generalized geometry, in: N.-C. Leung, S.-T. Yau (Eds.), *Surveys in Differential Geometry*, vol. 16, International Press, Cambridge, Mass, 2011, pp. 79–124, [arXiv:10080973](https://arxiv.org/abs/10080973).
- [9] M. Gualtieri, Branes on Poisson varieties, in *The many facets of geometry*, Oxford Univ. Press, Oxford, 2010, pp. 368–394.
- [10] S. Hu, R. Morarua, R. Seyyedali, A Kobayashi-Hitchin correspondence for I_{\pm} -holomorphic bundles, *Adv. Math.* 287 (10) (2016) 519–566.
- [11] Y. Lin, S. Tolman, Symmetries in generalized Kaehler geometry, *Comm. Math. Phys.* 268 (2006) 199–222, [arXiv:math/0509069](https://arxiv.org/abs/math/0509069).
- [12] Y. Wang, The GIT aspect of generalized Kaehler reduction. I. arxiv.org/pdf/1803.01178.
- [13] T. Drummond, Generalized reduction and pure spinors, *J. Symplectic Geom.* 12 (3) (2014) 435–471.