



Asymptotic behavior of varying discrete Jacobi–Sobolev orthogonal polynomials



Juan F. Mañas-Mañas^a, Francisco Marcellán^b, Juan J. Moreno-Balcázar^{a,c,*}

^a Departamento de Matemáticas, Universidad de Almería, Spain

^b Instituto de Ciencias Matemáticas (ICMAT) and Departamento de Matemáticas, Universidad Carlos III de Madrid, Spain

^c Instituto Carlos I de Física Teórica y Computacional, Spain

ARTICLE INFO

Article history:

Received 16 May 2015

Received in revised form 18 November 2015

MSC:

33C47

42C05

Keywords:

Sobolev orthogonal polynomials

Jacobi polynomials

Mehler–Heine formulae

Asymptotics

Zeros

ABSTRACT

In this contribution we deal with a varying discrete Sobolev inner product involving the Jacobi weight. Our aim is to study the asymptotic properties of the corresponding orthogonal polynomials and the behavior of their zeros. We are interested in Mehler–Heine type formulae because they describe the essential differences from the point of view of the asymptotic behavior between these Sobolev orthogonal polynomials and the Jacobi ones. Moreover, this asymptotic behavior provides an approximation of the zeros of the Sobolev polynomials in terms of the zeros of other well-known special functions. We generalize some results appeared in the literature very recently.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

One of the aims of this paper is the study of the asymptotic behavior of sequences of polynomials $\{Q_n^{(\alpha, \beta, M_n)}\}_{n \geq 0}$ orthogonal with respect to the inner product

$$(f, g)_{S, n} = \int_{-1}^1 f(x)g(x)(1-x)^\alpha(1+x)^\beta dx + M_n f^{(j)}(1)g^{(j)}(1), \quad (1)$$

where $\alpha > -1$, $\beta > -1$, and $j \geq 0$.

We assume that $\{M_n\}_{n \geq 0}$ is a sequence of nonnegative real numbers satisfying

$$\lim_{n \rightarrow \infty} M_n n^\gamma = M > 0, \quad (2)$$

where γ is a fixed real number. Notice that this assumption is not very restrictive since the sequence $\{M_n\}_{n \geq 0}$ can behave asymptotically like any real power of the monomial n .

The main motivation to study this type of inner product arises from the papers [1,2]. In [1] the authors work with a measure supported on $[-1, 1]$. However, in [2] the authors deal with measures supported on an unbounded interval. In both cases the authors consider measures with nonzero absolutely continuous part, i.e., they work with the so-called continuous

* Corresponding author at: Departamento de Matemáticas, Universidad de Almería, Spain.

E-mail addresses: jmm939@ual.es (J.F. Mañas-Mañas), pacomarc@ing.uc3m.es (F. Marcellán), balcazar@ual.es (J.J. Moreno-Balcázar).

Sobolev orthogonal polynomials. The main topic in those papers is how to balance the Sobolev inner product to equilibrate the influence of the two measures in the asymptotic behavior of the corresponding orthogonal polynomials. This inspires us to consider the discrete Sobolev inner product

$$(f, g)_S = \int f g d\mu_0 + M \int f^{(j)} g^{(j)} d\mu_1 = \int f g d\mu_0 + M f^{(j)}(c) f^{(j)}(c),$$

which is a perturbation of a standard inner product. Now, making M dependent on n we can study the influence of the perturbation on the asymptotic behavior of the orthogonal polynomials. The literature on discrete Sobolev (or Sobolev-type) orthogonal polynomials is very wide, so we refer the interested readers on this topic to survey [3] and the references therein.

From here, in [4] the authors found the asymptotic behavior of a family of orthogonal polynomials with respect to a varying Sobolev inner product similar to (1), involving the Laguerre weight $w(x) = x^\alpha e^{-x}$, $\alpha > -1$. We remark that the techniques used in [4] are not useful in this case, and now we need to use more powerful techniques based on those considered in [5]. More recently, in [6] the same authors have even improved these techniques in such a way that they have obtained relevant results for the orthogonal polynomials with respect to a non-varying discrete Sobolev inner product being μ_0 a general measure.

Previously, in [7] J.J. Moreno-Balcázar obtained some results in this direction but only for the case $j = 0$. Again, the method used in that paper does not allow to tackle our problem.

We want to emphasize that our objective is to establish that the size of the sequence $\{M_n\}_{n \geq 0}$ has an essential influence on the asymptotic behavior of the orthogonal polynomials with respect to (1), but this influence is only local, that is, around the point where we have introduced the perturbation. In our case, this point is located at $x = 1$. Furthermore, we prove that this influence depends on the size of the sequence $\{M_n\}_{n \geq 0}$ and its relation with the parameter α in the Jacobi weight and the order of the derivative in (1). It is important to remark that for a sequence $\{M_n\}_{n \geq 0}$, we have a sequence of orthogonal polynomials for each n , so we have a square tableau $\{Q_k^{(\alpha, \beta, M_n)}\}_{k \geq 0}$. Here, we deal with the diagonal of this tableau, i.e. $\{Q_n^{(\alpha, \beta, M_n)}\}_{n \geq 0} = \{Q_0^{(\alpha, \beta, M_0)}(x), Q_1^{(\alpha, \beta, M_1)}(x), \dots, Q_i^{(\alpha, \beta, M_i)}(x), \dots\}$. At this point, in order to simplify the notation, we will denote $Q_n^{(\alpha, \beta, M_n)}(x) = Q_n(x)$.

A second aim of this paper is to establish a simple asymptotic relation between the zeros of the Sobolev polynomials which are orthogonal with respect to (1) and the zeros of combinations of Bessel functions of the first kind. This relation is deduced as an immediate consequence of Mehler–Heine formulae (Theorem 2) and they have a numerical interest since we provide an estimate of the zeros of these polynomials.

Since Jacobi classical orthogonal polynomials are involved in the varying inner product (1), we recall some of their basic properties. Jacobi polynomials are orthogonal with respect to the standard inner product

$$(f, g) = \int_{-1}^1 f(x)g(x)(1-x)^\alpha(1+x)^\beta dx, \quad \alpha, \beta > -1.$$

In the sequel, we will work with the sequence $\{P_n^{(\alpha, \beta)}\}_{n \geq 0}$, $\alpha > -1$ and $\beta > -1$, normalized by (see [8, f. (4.1.1)])

$$P_n^{(\alpha, \beta)}(1) = \binom{n+\alpha}{n} = \frac{\Gamma(n+\alpha+1)}{\Gamma(n+1)\Gamma(\alpha+1)}. \quad (3)$$

The derivatives of Jacobi polynomials satisfy (see, [8, f. (4.21.7)])

$$(P_n^{(\alpha, \beta)}(x))^{(k)} = \frac{1}{2^k} \frac{\Gamma(n+\alpha+\beta+k+1)}{\Gamma(n+\alpha+\beta+1)} P_{n-k}^{(\alpha+k, \beta+k)}(x), \quad k \geq 0. \quad (4)$$

Using (3) and (4), we deduce

$$(P_n^{(\alpha, \beta)}(1))^{(k)} = \frac{1}{2^k} \frac{\Gamma(n+\alpha+\beta+k+1)}{\Gamma(n+\alpha+\beta+1)} \frac{\Gamma(n+\alpha+1)}{\Gamma(n-k+1)\Gamma(\alpha+k+1)}, \quad (5)$$

where $(P_n^{(\alpha, \beta)}(1))^{(k)}$ denotes the k th derivative of $P_n^{(\alpha, \beta)}$ evaluated at $x = 1$.

We also note that the squared norm of a Jacobi polynomial is (see, [8, f. (4.3.3)]):

$$\|P_n^{(\alpha, \beta)}\|^2 = \frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+1)\Gamma(n+\alpha+\beta+1)}. \quad (6)$$

Finally, we will use the Mehler–Heine formula for classical Jacobi polynomials

Theorem 1 ([8, Th. 8.1.1]). *Let $\alpha, \beta > -1$. Then,*

$$\lim_{n \rightarrow \infty} n^{-\alpha} P_n^{(\alpha, \beta)}\left(\cos\left(\frac{x}{n}\right)\right) = \lim_{n \rightarrow \infty} \frac{1}{n^\alpha} P_n^{(\alpha, \beta)}\left(1 - \frac{x^2}{2n^2}\right) = (x/2)^{-\alpha} J_\alpha(x),$$

uniformly on compact subsets of \mathbb{C} . Here $J_\alpha(x)$ denotes the Bessel function of the first kind, i.e.,

$$J_\alpha(x) = \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(k + \alpha + 1)} \left(\frac{x}{2}\right)^{2k+\alpha}.$$

We will also use the following limit related to Stirling formula (see, for example, [9, f. (5.11.13)])

$$\lim_{n \rightarrow \infty} \frac{n^{b-a} \Gamma(n+a)}{\Gamma(n+b)} = 1. \quad (7)$$

We introduce the following notation: If a_n and b_n are two sequences of real numbers, then $a_n \approx b_n$ means that the sequence $\frac{a_n}{b_n}$ converges to 1.

The paper is organized as follows. In Section 2 we provide some properties of the varying Jacobi–Sobolev orthogonal polynomials which are essential to establish the Mehler–Heine asymptotics for these polynomials in Section 3. Furthermore, as a consequence of this asymptotic formula we deduce the asymptotic behavior of the corresponding zeros. Thus, as we have commented previously, we can see the influence of the parameter γ , related to the size of the sequence $\{M_n\}_{n \geq 0}$, on the location of these zeros. Finally, in Section 4 we illustrate the results obtained in Section 3 with some numerical experiments.

2. Varying Jacobi–Sobolev orthogonal polynomials

It is well known that the classical Jacobi orthogonal polynomials, $\{P_i^{(\alpha, \beta)}\}_{i=0}^n$, constitute a basis of the linear space $\mathbb{P}_n[x]$ of polynomials with real coefficients and degree at most n . Therefore, the Jacobi–Sobolev orthogonal polynomial of degree n , $Q_n(x)$, can be expressed as

$$Q_n(x) = P_n^{(\alpha, \beta)}(x) + \sum_{i=0}^{n-1} a_{n,i} P_i^{(\alpha, \beta)}(x).$$

Then, using well-known algebraic tools (see, for example, [10, Sect. 2]) we can deduce

$$Q_n(x) = P_n^{(\alpha, \beta)}(x) - \frac{M_n \left(P_n^{(\alpha, \beta)}(1)\right)^{(j)}}{1 + M_n K_{n-1}^{(j,j)}(1, 1)} K_{n-1}^{(j,0)}(1, x), \quad (8)$$

with

$$K_n^{(j,k)}(x, y) = \sum_{i=0}^n \frac{\left(P_i^{(\alpha, \beta)}(x)\right)^{(j)} \left(P_i^{(\alpha, \beta)}(y)\right)^{(k)}}{\|P_i^{(\alpha, \beta)}(x)\|^2}.$$

Next, we give a technical result useful for our purposes, interesting in itself though.

Lemma 1. Let $\{Q_n\}_{n \geq 0}$ be the sequence of orthogonal polynomials with respect to (1) and $0 \leq k \leq n$, then

(a)

$$\lim_{n \rightarrow \infty} \frac{(Q_n)^{(k)}(1)}{\left(P_n^{(\alpha, \beta)}(1)\right)^{(k)}} = \begin{cases} \frac{k-j}{\alpha+j+k+1}, & \text{if } \gamma < 2(\alpha+2j+1), \\ \theta_{\alpha, \beta, j, k}, & \text{if } \gamma = 2(\alpha+2j+1), \\ 1, & \text{if } \gamma > 2(\alpha+2j+1), \end{cases} \quad (9)$$

where

$$\theta_{\alpha, \beta, j, k} = \frac{M(k-j) + \Gamma^2(\alpha+j+1)2^{\alpha+\beta+2j+1}(\alpha+2j+1)(\alpha+j+k+1)}{(\alpha+j+k+1)(M + \Gamma^2(\alpha+j+1)2^{\alpha+\beta+2j+1}(\alpha+2j+1))}. \quad (10)$$

(b) $(Q_n, Q_n)_{S, n} \approx \|P_n^{(\alpha, \beta)}\|^2$.

Proof. Kernel polynomials related to classical families of orthogonal polynomials and their derivatives have been widely studied in the literature. Thus, we can claim that the following limit exists,

$$\lim_{n \rightarrow \infty} \frac{K_{n-1}^{(j,k)}(1, 1)}{n^{2\alpha+2j+2k+2}} \in \mathbb{R}. \quad (11)$$

It is very easy to check it by using Stolz's criterion, (5), (6), (7) and the fact that

$$n^{2\alpha+2j+2k+2} - (n-1)^{2\alpha+2j+2k+2} \approx (2\alpha+2j+2k+2)n^{2\alpha+2j+2k+1}.$$

Thus,

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{K_{n-1}^{(j,k)}(1, 1)}{n^{2\alpha+2j+2k+2}} &= \lim_{n \rightarrow \infty} \frac{K_{n-1}^{(j,k)}(1, 1) - K_{n-2}^{(j,k)}(1, 1)}{n^{2\alpha+2j+2k+2} - (n-1)^{2\alpha+2j+2k+2}} \\ &= \lim_{n \rightarrow \infty} \frac{\left(P_{n-1}^{(\alpha,\beta)}(1)\right)^{(k)} \left(P_{n-1}^{(\alpha,\beta)}(1)\right)^{(j)}}{\|P_{n-1}^{(\alpha,\beta)}\|^2 (2\alpha + 2j + 2k + 2) n^{2\alpha+2j+2k+1}} \\ &= \lim_{n \rightarrow \infty} \frac{C_{j,k} \Gamma(n + \alpha + \beta + j) \Gamma(n + \alpha + \beta + k) \Gamma(n + \alpha) \Gamma(n)}{\Gamma(n - j) \Gamma(n + \alpha + \beta) \Gamma(n - k) \Gamma(n + \beta) n^{2\alpha+2j+2k}} \\ &= C_{j,k} \in \mathbb{R},\end{aligned}$$

where

$$C_{j,k} = \frac{1}{\Gamma(\alpha + j + 1) \Gamma(\alpha + k + 1) 2^{\alpha+\beta+j+k+1} (\alpha + j + k + 1)}.$$

We will now prove part (a) of the lemma, by (8)

$$\begin{aligned}\lim_{n \rightarrow \infty} \frac{Q_n^{(k)}(1)}{\left(P_n^{(\alpha,\beta)}(1)\right)^{(k)}} &= \lim_{n \rightarrow \infty} \left(1 - \frac{M_n K_{n-1}^{(j,k)}(1, 1)}{1 + M_n K_{n-1}^{(j,j)}(1, 1)} \frac{\left(P_n^{(\alpha,\beta)}(1)\right)^{(j)}}{\left(P_n^{(\alpha,\beta)}(1)\right)^{(k)}} \right) \\ &= \lim_{n \rightarrow \infty} \left(1 - \frac{M_n n^\gamma \frac{1}{2^j} \frac{\Gamma(n+\alpha+\beta+j+1)}{\Gamma(n-j+1)\Gamma(\alpha+j+1)} \frac{K_{n-1}^{(j,k)}(1, 1)}{n^{2\alpha+2j+2k+2}} n^{2\alpha+2j+2k+2-\gamma}}{\frac{1}{2^k} \frac{\Gamma(n+\alpha+\beta+k+1)}{\Gamma(n-k+1)\Gamma(\alpha+k+1)} \left(1 + M_n n^\gamma \frac{K_{n-1}^{(j,j)}(1, 1)}{n^{2\alpha+4j+2}} n^{2\alpha+4j+2-\gamma} \right)} \right).\end{aligned}$$

To simplify the computations we introduce the following notation

$$a_n = M_n n^\gamma, \quad \text{by (2) we have } \lim_{n \rightarrow \infty} a_n = M,$$

$$b_{n,j,k} = \frac{K_{n-1}^{(j,k)}(1, 1)}{n^{2\alpha+2j+2k+2}}, \quad \text{by (11) we have } \lim_{n \rightarrow \infty} b_{n,j,k} = C_{j,k}.$$

Then, the above limit becomes

$$\begin{aligned}\lim_{n \rightarrow \infty} \left(1 - \frac{a_n b_{n,j,k} \frac{1}{2^j} \frac{\Gamma(n+\alpha+\beta+j+1)}{\Gamma(n-j+1)\Gamma(\alpha+j+1)} n^{2\alpha+2j+2k+2-\gamma}}{\frac{1}{2^k} \frac{\Gamma(n+\alpha+\beta+k+1)}{\Gamma(n-k+1)\Gamma(\alpha+k+1)} (1 + a_n b_{n,j,j} n^{2\alpha+4j+2-\gamma})} \right) \\ = 1 - \frac{2^{k-j} \Gamma(\alpha + k + 1)}{\Gamma(\alpha + j + 1)} \lim_{n \rightarrow \infty} a_n \lim_{n \rightarrow \infty} b_{n,j,k} \lim_{n \rightarrow \infty} \frac{n^{2\alpha+2j+2k+2-\gamma}}{n^{2k-2j} (1 + a_n b_{n,j,j} n^{2\alpha+4j+2-\gamma})} \\ = 1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \lim_{n \rightarrow \infty} \frac{n^{2\alpha+4j+2-\gamma}}{(1 + a_n b_{n,j,j} n^{2\alpha+4j+2-\gamma})} \\ = 1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{n^{2\alpha+4j+2-\gamma}} + a_n b_{n,j,j}}.\end{aligned}$$

Therefore, it is necessary to distinguish three cases according to the value of the parameter γ . The value of this limit is:

Case $\gamma > 2(\alpha + 2j + 1)$.

$$1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{n^{2\alpha+4j+2-\gamma}} + a_n b_{n,j,j}} = 1.$$

Case $\gamma < 2(\alpha + 2j + 1)$.

$$\begin{aligned}1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \lim_{n \rightarrow \infty} \frac{1}{\frac{1}{n^{2\alpha+4j+2-\gamma}} + a_n b_{n,j,j}} \\ = 1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \frac{1}{M C_{j,j}} = 1 - \frac{\alpha + 2j + 1}{\alpha + j + k + 1} = \frac{k - j}{\alpha + j + k + 1}.\end{aligned}$$

Case $\gamma = 2(\alpha + 2j + 1)$.

$$\begin{aligned} 1 - \frac{2^{k-j} \Gamma(\alpha + k + 1) C_{j,k} M}{\Gamma(\alpha + j + 1)} \frac{1}{1 + M C_{j,j}} &= 1 - \frac{M(\alpha + 2j + 1)}{(\alpha + j + k + 1) (M + \Gamma^2(\alpha + j + 1) 2^{\alpha+\beta+2j+1} (\alpha + 2j + 1))} \\ &= \frac{M(k - j) + \Gamma^2(\alpha + j + 1) 2^{\alpha+\beta+2j+1} (\alpha + 2j + 1) (\alpha + j + k + 1)}{(\alpha + j + k + 1) (M + \Gamma^2(\alpha + j + 1) 2^{\alpha+\beta+2j+1} (\alpha + 2j + 1))} \\ &= \theta_{\alpha, \beta, j, k}. \end{aligned}$$

Thus, we have proved (a). Now, we are going to prove (b). Using standard arguments for Sobolev orthogonal polynomials we can deduce

$$(Q_n, Q_n)_{S,n} = \|P_n^{(\alpha, \beta)}\|^2 + \frac{M_n \left(\left(P_n^{(\alpha, \beta)}(1) \right)^{(j)} \right)^2}{1 + M_n K_{n-1}^{(j,j)}(1, 1)}.$$

Then,

$$\lim_{n \rightarrow \infty} \frac{(Q_n, Q_n)_{S,n}}{\|P_n^{(\alpha, \beta)}\|^2} = \lim_{n \rightarrow \infty} \left(1 + \frac{\left(\left(P_n^{(\alpha, \beta)}(1) \right)^{(j)} \right)^2}{\|P_n^{(\alpha, \beta)}\|^2} \frac{M_n}{1 + M_n K_{n-1}^{(j,j)}(1, 1)} \right).$$

To establish (b) it is enough to prove that

$$\lim_{n \rightarrow \infty} \left(\frac{M_n \left(\left(P_n^{(\alpha, \beta)}(1) \right)^{(j)} \right)^2}{\|P_n^{(\alpha, \beta)}\|^2 \left(1 + M_n K_{n-1}^{(j,j)}(1, 1) \right)} \right) = 0.$$

Indeed, from (5) and (6) this limit can be expressed as

$$\begin{aligned} &\lim_{n \rightarrow \infty} \left(\frac{M_n \left(\left(P_n^{(\alpha, \beta)}(1) \right)^{(j)} \right)^2}{\|P_n^{(\alpha, \beta)}\|^2 \left(1 + M_n K_{n-1}^{(j,j)}(1, 1) \right)} \right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{M_n \frac{1}{2^{2j}} \frac{\Gamma(n+\alpha+\beta+j+1)}{\Gamma(n-j+1)\Gamma(\alpha+j+1)} \frac{\Gamma(n+\alpha+\beta+j+1)}{\Gamma(n+\alpha+\beta+1)} \frac{n^{-4j-2\alpha-\beta+\gamma}}{\Gamma(n-j+1)\Gamma(\alpha+j+1)} \frac{\Gamma(n+\alpha+1)}{n^{-4j-2\alpha-\beta+\gamma}} \frac{1}{n^{-4j-2\alpha-\beta+\gamma}}}{\frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{n^{-\beta}}{\Gamma(n+1)} \frac{1}{n^{-\beta}} \left(1 + M_n K_{n-1}^{(j,j)}(1, 1) \frac{n^{2\alpha+4j+2-\gamma}}{n^{2\alpha+4j+2-\gamma}} \right)} \right). \end{aligned}$$

Again, to simplify the computations we introduce some notation

$$\begin{aligned} a_n &= M_n n^\gamma, \quad \text{by (2) we have } \lim_{n \rightarrow \infty} a_n = M, \\ b_n &= \frac{\Gamma^2(n+\alpha+\beta+j+1) \Gamma(n+\alpha+1) n^{-4j-2\alpha-\beta}}{\Gamma(n-j+1) \Gamma^2(n+\alpha+\beta+1)}, \quad \text{then by (7) } \lim_{n \rightarrow \infty} b_n = 1, \\ c_n &= \frac{\Gamma(n+\beta+1) n^{-\beta}}{\Gamma(n+1)}, \quad \text{then by (7) } \lim_{n \rightarrow \infty} c_n = 1, \\ d_n &= M_n n^\gamma \frac{K_{n-1}^{(j,j)}(1, 1)}{n^{2\alpha+4j+2}}, \quad \text{then using (2) and (11) we get } \lim_{n \rightarrow \infty} d_n = M C_{j,j}, \\ E_{\alpha,j} &= \frac{1}{2^{2j}} \frac{1}{\Gamma^2(\alpha+j+1)}. \end{aligned}$$

In this way, for every γ , the above limit is

$$\lim_{n \rightarrow \infty} \frac{E_{\alpha,j} a_n b_n n^{4j+2\alpha+\beta-\gamma}}{c_n \frac{2^{\alpha+\beta+1} n^\beta}{2n+\alpha+\beta+1} \left(1 + d_n n^{4j+2\alpha+2-\gamma} \right)} = \lim_{n \rightarrow \infty} \frac{E_{\alpha,j} a_n b_n (2n+\alpha+\beta+1)}{2^{\alpha+\beta+1} n^2 \left(\frac{c_n}{n^{4j+2\alpha+2-\gamma}} + c_n d_n \right)} = 0,$$

and we have just proved (b). \square

Remark 1. Notice that taking into account (b) in the above lemma, (a) holds true when we consider orthonormal polynomials.

To tackle Mehler–Heine asymptotics we need to expand the Sobolev polynomials Q_n adequately. The following result gives us this expansion. In a more general framework it has been established in [6, Th. 1]. The idea is that the coefficients $b_i(n)$ in (12) can be obtained as a solution of a homogeneous linear system of $j + 1$ equations and $j + 2$ unknowns. In our concrete case, we can compute explicitly the entries of the corresponding coefficient matrix.

Proposition 1. *There exists a family of real numbers $\{b_i(n)\}_{i=0}^{j+1}$, not identically zero, such that the following connection formula holds*

$$Q_n(x) = \sum_{i=0}^{j+1} b_i(n)(1-x)^i P_{n-i}^{(\alpha+2i, \beta)}(x), \quad n \geq j+1. \quad (12)$$

Lemma 2. *Let $\{b_i(n)\}_{i=0}^{j+1}$ be the coefficients in (12). Then*

$$\lim_{n \rightarrow \infty} b_i(n) = b_i \in \mathbb{R}, \quad i \in \{0, 1, \dots, j+1\}.$$

Proof. We take the k th derivative in (12) and we evaluate the corresponding expression at $x = 1$,

$$\begin{aligned} Q_n^{(k)}(x) &= \sum_{i=0}^{j+1} b_i(n) \sum_{s=0}^k \binom{k}{s} ((1-x)^i)^{(s)} \left(P_{n-i}^{(\alpha+2i, \beta)}(x) \right)^{(k-s)} \\ &= \sum_{i=0}^{j+1} b_i(n) \sum_{s=0}^{\min\{i, k\}} \binom{k}{s} (-1)^s \frac{i!}{(i-s)!} (1-x)^{i-s} \left(P_{n-i}^{(\alpha+2i, \beta)}(x) \right)^{(k-s)}. \end{aligned}$$

Then,

$$Q_n^{(k)}(1) = \sum_{i=0}^k b_i(n) \binom{k}{i} (-1)^i i! \left(P_{n-i}^{(\alpha+2i, \beta)}(1) \right)^{(k-i)}.$$

From Lemma 1, $\lim_{n \rightarrow \infty} \frac{Q_n^{(k)}(1)}{(P_n^{(\alpha, \beta)}(1))^{(k)}}$ exists and its value depends on the value of parameter γ related to the size of the sequence $\{M_n\}_{n \geq 0}$, so

$$\frac{Q_n^{(k)}(1)}{(P_n^{(\alpha, \beta)}(1))^{(k)}} = \sum_{i=0}^k b_i(n) \binom{k}{i} (-1)^i i! A_i(k, n) \quad (13)$$

$$\text{with } A_i(k, n) = \frac{(P_{n-i}^{(\alpha+2i, \beta)}(1))^{(k-i)}}{(P_n^{(\alpha, \beta)}(1))^{(k)}}.$$

It only remains to prove that there exists $\lim_{n \rightarrow \infty} A_i(k, n) \in \mathbb{R}$ and, therefore the coefficients $\{b_i(n)\}_{i=0}^{j+1}$ are convergent. Indeed

$$\begin{aligned} \lim_{n \rightarrow \infty} A_i(k, n) &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^{k-i}} \frac{\Gamma(n-i+\alpha+2i+\beta+k-i+1)}{\Gamma(n-i+\alpha+2i+\beta+1)} \frac{\Gamma(n-i+\alpha+2i+1)}{\Gamma(n-i-k+i+1)\Gamma(\alpha+2i+k-i+1)}}{\frac{1}{2^k} \frac{\Gamma(n+\alpha+\beta+k+1)}{\Gamma(n+\alpha+\beta+1)} \frac{\Gamma(n+\alpha+1)}{\Gamma(n-k+1)\Gamma(\alpha+k+1)}} \\ &= A_i(k, \alpha), \end{aligned}$$

$$\text{where we denote } A_i(k, \alpha) = \frac{2^i \Gamma(\alpha+k+1)}{\Gamma(\alpha+i+k+1)}. \quad \square$$

Remark 2. Let $b_i = \lim_{n \rightarrow \infty} b_i(n)$ with $i \in \{0, 1, \dots, j+1\}$. (13) is a recursive algorithm to compute b_i .

- Step 1. For $k = 0$ we obtain b_0 in a straightforward way.
- Step 2. For $k = 1$ we deduce the value of b_1 from (13) using step 1. Similarly, for $k \geq 2$ we apply (13) in a recursive way.

3. Asymptotics and zeros of varying Jacobi–Sobolev

We focus our attention on the analysis of Mehler–Heine formulas for these discrete Jacobi–Sobolev orthogonal polynomials because we want to know how the discrete part in the inner product (1) influences the asymptotic behavior of the corresponding orthogonal polynomials. Furthermore, we will prove that this influence is related to the size of the sequence $\{M_n\}_{n \geq 0}$.

Theorem 2. For the sequence $\{Q_n\}_{n \geq 0}$ the following Mehler–Heine formula holds

$$\lim_{n \rightarrow \infty} \frac{Q_n(\cos(x/n))}{n^\alpha} = \lim_{n \rightarrow \infty} \frac{Q_n\left(1 - \frac{x^2}{2n^2}\right)}{n^\alpha} = \begin{cases} \phi_\alpha(x), & \text{if } \gamma > 2(\alpha + 2j + 1), \\ \psi_{\alpha,j}(x), & \text{if } \gamma = 2(\alpha + 2j + 1), \\ \varphi_{\alpha,j}(x), & \text{if } \gamma < 2(\alpha + 2j + 1), \end{cases} \quad (14)$$

uniformly on compact subsets of \mathbb{C} , where

$$\begin{aligned} \phi_\alpha(x) &= \left(\frac{x}{2}\right)^{-\alpha} J_\alpha(x), \\ \psi_{\alpha,j}(x) &= \sum_{i=0}^{j+1} b_i 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x), \end{aligned}$$

with

$$b_i = (-1)^i \frac{\frac{M(i-j) - \Gamma^2(\alpha+j+1)2^{\alpha+\beta+2j+1}(\alpha+2j+1)(\alpha+j+i+1)}{(\alpha+j+i+1)(M + \Gamma^2(\alpha+j+1)2^{\alpha+\beta+2j+1}(\alpha+2j+1))} - \Gamma(\alpha+i+1) \sum_{k=0}^{i-1} b_k \binom{i}{k} \frac{(-1)^k k! 2^k}{\Gamma(\alpha+i+k+1)}}{i! \frac{2^i \Gamma(\alpha+i+1)}{\Gamma(\alpha+2i+1)}},$$

for $0 \leq i \leq j+1$, and

$$\varphi_{\alpha,j}(x) = \sum_{i=0}^{j+1} b_i 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x),$$

where the coefficients b_i are computed as

$$b_i = (-1)^i \frac{\frac{i-j}{\alpha+j+i+1} - \Gamma(\alpha+i+1) \sum_{k=0}^{i-1} b_k \binom{i}{k} \frac{(-1)^k k! 2^k}{\Gamma(\alpha+i+k+1)}}{i! \frac{2^i \Gamma(\alpha+i+1)}{\Gamma(\alpha+2i+1)}}, \quad 0 \leq i \leq j+1.$$

Notice that in last two cases the coefficient b_0 is computed using the corresponding formula assuming $\sum_{i=0}^{-1} = 0$.

Proof. Scaling and taking limits in (12)

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{Q_n\left(1 - \frac{x^2}{2n^2}\right)}{n^\alpha} &= \lim_{n \rightarrow \infty} \frac{\sum_{i=0}^{j+1} b_i(n) \left(1 - \left(1 - \frac{x^2}{2n^2}\right)\right)^i P_{n-i}^{(\alpha+2i, \beta)} \left(1 - \frac{x^2}{2n^2}\right)}{n^\alpha} \\ &= \sum_{i=0}^{j+1} \lim_{n \rightarrow \infty} b_i(n) \lim_{n \rightarrow \infty} \frac{\left(1 - \left(1 - \frac{x^2}{2n^2}\right)\right)^i P_{n-i}^{(\alpha+2i, \beta)} \left(1 - \frac{x^2}{2n^2}\right)}{n^\alpha} \\ &= \sum_{i=0}^{j+1} b_i 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x), \end{aligned}$$

uniformly on compact subsets of \mathbb{C} . Notice that in the last inequality we have used Theorem 1 written in the following way

$$\lim_{n \rightarrow \infty} \frac{\left(\frac{x^2}{2n^2}\right)^i P_{n-i}^{(\alpha+2i, \beta)} \left(1 - \frac{x^2}{2n^2}\right)}{n^\alpha} = 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x),$$

uniformly on compact subsets of \mathbb{C} , where i is a fixed nonnegative integer number.

Now, we distinguish three cases according to the value of the parameter γ .

- If $\gamma > 2(\alpha + 2j + 1)$, we are going to prove that $b_0 = 1$ and $b_i = 0$ if $i \in \{1, 2, \dots, j + 1\}$. We can compute b_i from (13). If $k = 0$, then

$$\frac{Q_n(1)}{P_n^{(\alpha, \beta)}(1)} = b_0(n)A_0(0, n).$$

Using Lemma 1 and taking limits, we obtain $b_0 = 1$. If $k = 1$, then according to Lemma 1 we have

$$\frac{Q_n^{(1)}(1)}{(P_n^{(\alpha, \beta)}(1))^{(1)}} = b_0(n)A_0(1, n) - b_1(n)A_1(1, n).$$

Taking limits,

$$1 = 1 - b_1A_1(1, \alpha), \quad \text{then } b_1 = 0.$$

Applying a recursive procedure we get $b_i = 0$ for $i \in \{1, 2, \dots, j + 1\}$. To illustrate this procedure we consider the case $k = j + 1$. Thus, we have $b_i = 0$ for $i \in \{1, 2, \dots, j\}$. Then,

$$\begin{aligned} \frac{Q_n^{(j+1)}(1)}{(P_n^{(\alpha, \beta)}(1))^{(j+1)}} &= b_0(n)A_0(1, n) + \sum_{i=1}^j b_i(n) \binom{j+1}{i} (-1)^i i! A_i(j+1, n) \\ &\quad + b_{j+1}(n) (-1)^{j+1} (j+1)! A_{j+1}(j+1, n). \end{aligned}$$

Taking limits,

$$1 = 1 + b_{j+1}(-1)^{j+1} (j+1)! A_{j+1}(j+1, \alpha), \quad \text{then } b_{j+1} = 0.$$

- Case $\gamma = 2(\alpha + 2j + 1)$. From (13) and $k = 0$, we have

$$\frac{Q_n(1)}{P_n^{(\alpha, \beta)}(1)} = b_0(n)A_0(0, n).$$

Taking limits when n tends to infinity in the above expression, we get

$$b_0 = \frac{-jM - \Gamma^2(\alpha + j + 1)2^{\alpha+\beta+2j+1}(\alpha + 2j + 1)(\alpha + j + 1)}{(\alpha + j + 1)(M + \Gamma^2(\alpha + j + 1)2^{\alpha+\beta+2j+1}(\alpha + 2j + 1))}.$$

For $i \geq 1$, we use Lemma 1 again and take limits. Thus, we deduce the coefficients b_i in a recursive way from (13).

- Case $\gamma < 2(\alpha + 2j + 1)$. We can tackle this case in the same way as the case $\gamma = 2(\alpha + 2j + 1)$. \square

Next, we are going to study the zeros of the polynomials $\{Q_n\}_{n \geq 0}$ orthogonal with respect to (1). The following result was established for the non-varying case within a more general framework by H. G. Meijer in [11, Th. 4.1] (see also [12, Lemma 2]). Actually, that proof can be written in the same way for the varying case, so we omit it.

Proposition 2. The polynomial $Q_n(x)$, $n \geq 1$, has n real and simple zeros and at most one of them is located outside the interval $[-1, 1]$.

We can give more information about the location of the zeros. The case $j = 0$ was considered in [7]. We notice that in that case all the zeros are in the interval $(-1, 1)$. Thus, next we will assume $j > 0$ and we will denote by $y_{n,1} > y_{n,2} > \dots > y_{n,n-1} > y_{n,n}$ the zeros of $Q_n(x)$.

Proposition 3. For n large enough and $j > 0$, we have

- If $\gamma > 2(\alpha + 2j + 1)$, then all zeros of $Q_n(x)$ are located in $(-1, 1)$.
- If $\gamma < 2(\alpha + 2j + 1)$, then $y_{n,1} > 1$.
- If $\gamma = 2(\alpha + 2j + 1)$, then $y_{n,1} > 1$ if and only if

$$M > \frac{2^{\alpha+\beta+2j+1}(\alpha + j + 1)(\alpha + 2j + 1)\Gamma^2(\alpha + j + 1)}{j}.$$

Proof. We distinguish three cases, but essentially we use Lemma 1 (a) with $k = 0$, and the fact that the leading coefficient of Q_n is positive. Then,

- If $\gamma > 2(\alpha + 2j + 1)$, then by Lemma 1 $Q_n(1) > 0$ for n large enough. Therefore, taking into account Proposition 2, all the zeros are located in $(-1, 1)$.
- If $\gamma < 2(\alpha + 2j + 1)$, then $Q_n(1) < 0$ for n large enough, which implies that there is a zero of Q_n greater than 1 and by Proposition 2 it is the only one.

- If $\gamma = 2(\alpha + 2j + 1)$, then $y_{n,1} > 1$ if and only if $Q_n(1) < 0$ for n large enough, and this only happens if and only if

$$M > \frac{2^{\alpha+\beta+2j+1}(\alpha+j+1)(\alpha+2j+1)\Gamma^2(\alpha+j+1)}{j}. \quad \square$$

Now we deduce the asymptotic behavior of the zeros of $Q_n(x)$.

Proposition 4. Let $y_{n,1} > y_{n,2} > \dots > y_{n,n-1} > y_{n,n}$ be the zeros of $Q_n(x)$ and $\phi_\alpha(x)$, $\varphi_{\alpha,j}(x)$, and $\psi_{\alpha,j}(x)$ the functions defined in Theorem 2. We assume $j > 0$.

1. If $\gamma > 2(\alpha + 2j + 1)$, then

$$\lim_{n \rightarrow \infty} n\sqrt{2(1-y_{n,i})} = j_{\alpha,i}, \quad i \geq 1,$$

where $j_{\alpha,i}$ denotes the i th positive zero of the Bessel function of the first kind.

2. If $\gamma < 2(\alpha + 2j + 1)$, then

$$\lim_{n \rightarrow \infty} y_{n,1} = 1, \quad \lim_{n \rightarrow \infty} n\sqrt{2(1-y_{n,i})} = s_{\alpha,i-1}, \quad i \geq 2,$$

where $s_{\alpha,i}$ denotes the i th positive zero of the function $\varphi_{\alpha,j}(x)$.

3. If $\gamma = 2(\alpha + 2j + 1)$, we have two cases:

- (a) If $M \leq \frac{2^{\alpha+\beta+2j+1}(\alpha+j+1)(\alpha+2j+1)\Gamma^2(\alpha+j+1)}{j}$, then $y_{n,1} \leq 1$, for n large enough, and

$$\lim_{n \rightarrow \infty} n\sqrt{2(1-y_{n,i})} = t_{\alpha,i}, \quad i \geq 1,$$

where $t_{\alpha,i}$ denotes the i th positive zero of the function $\psi_{\alpha,j}(x)$.

- (b) If $M > \frac{2^{\alpha+\beta+2j+1}(\alpha+j+1)(\alpha+2j+1)\Gamma^2(\alpha+j+1)}{j}$, then

$$\lim_{n \rightarrow \infty} y_{n,1} = 1, \quad \lim_{n \rightarrow \infty} n\sqrt{2(1-y_{n,i})} = t_{\alpha,i-1}, \quad i \geq 2,$$

where $t_{\alpha,i}$ denotes the i th positive zero of the function $\psi_{\alpha,j}(x)$.

Proof. It follows from Theorem 2, Proposition 3, and Hurwitz's Theorem (see [8, Th. 1.91.3]). \square

To illustrate Theorem 2 we are going to recover the case $j = 0$ obtained in [7]. In that paper the author uses monic polynomials, and here we are considering a different normalization, i.e. the leading coefficient of Q_n is

$$\frac{\Gamma(2n + \alpha + \beta + 1)}{2^n \Gamma(n + 1) \Gamma(n + \alpha + \beta + 1)}.$$

Therefore, it is necessary to do some easy computations. We use the relations (see, [9, f.10.6.1], [13, 6.1.18])

$$J_\alpha(x) - \frac{2(\alpha+1)}{x} J_{\alpha+1}(x) = -J_{\alpha+2}(x), \quad (15)$$

as well as

$$\Gamma(2x) = \frac{\Gamma(x)\Gamma(x + \frac{1}{2})}{2^{1-2x}\sqrt{\pi}}. \quad (16)$$

First, using (7) and (16) we get

$$\begin{aligned} \frac{\Gamma(2n + \alpha + \beta + 1)}{2^n \Gamma(n + 1) \Gamma(n + \alpha + \beta + 1)} &\approx \frac{2^{n+\alpha+\beta}}{\sqrt{\pi}} \frac{\Gamma\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + \frac{1}{2}\right)}{\Gamma(n + 1)} \frac{\Gamma\left(n + \frac{\alpha}{2} + \frac{\beta}{2} + 1\right)}{\Gamma(n + \alpha + \beta + 1)} \\ &\approx \frac{2^{n+\alpha+\beta}}{n^{\frac{1}{2}}\sqrt{\pi}}. \end{aligned}$$

In [7] it was obtained

$$\lim_{n \rightarrow \infty} \frac{2^n \hat{P}_n^{(\alpha, \beta, M_n)}(\cos(x/n))}{n^{\alpha+1/2}} = \begin{cases} -2^{-\beta} \sqrt{\pi} x^2 z_{\alpha+2}(x), & \text{if } \gamma < 2\alpha + 2, \\ -2^{-\beta} \sqrt{\pi} (z_\alpha(x) + a_{\alpha, \beta, M} z_{\alpha+1}(x)), & \text{if } \gamma = 2\alpha + 2, \\ 2^{-\beta} \sqrt{\pi} z_\alpha(x), & \text{if } \gamma > 2\alpha + 2, \end{cases}$$

where

$$z_\alpha(x) = x^{-\alpha} J_\alpha(x),$$

$$a_{\alpha, \beta, M} = \frac{-2M(\alpha + 1)}{M + 2^{\alpha+\beta+1} \Gamma(\alpha + 2) \Gamma(\alpha + 1)},$$

and $\{\hat{P}_n^{(\alpha, \beta, M_n)}\}_{n \geq 0}$ denotes the sequence of monic polynomials which are orthogonal with respect to (1) with $j = 0$. This result can be written as follows

$$\lim_{n \rightarrow \infty} \frac{2^{n+\alpha+\beta} \hat{P}_n^{(\alpha, \beta, M_n)}(\cos(x/n))}{n^{\alpha+1/2} \sqrt{\pi}} = \begin{cases} -2^\alpha x^2 z_{\alpha+2}(x), & \text{if } \gamma < 2\alpha + 2, \\ -2^\alpha (z_\alpha(x) + a_{\alpha, \beta, M} z_{\alpha+1}(x)), & \text{if } \gamma = 2\alpha + 2, \\ 2^\alpha z_\alpha(x), & \text{if } \gamma > 2\alpha + 2. \end{cases} \quad (17)$$

We can observe that

$$\frac{2^{n+\alpha+\beta} \hat{P}_n^{(\alpha, \beta, M_n)}(\cos(x/n))}{n^{\alpha+1/2} \sqrt{\pi}} \approx \frac{Q_n(\cos(x/n))}{n^\alpha}.$$

Therefore, it only remains to compare the limit functions in (14) and (17). The case $\gamma > 2\alpha + 2$ is trivial. We pay attention to the other two cases.

- $\gamma < 2\alpha + 2$.

In this case $b_0 = 0$ and $b_1 = -1/2$. Thus we have

$$\varphi_{\alpha,0}(x) = -\left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2}(x) = -2^\alpha x^2 x^{-\alpha-2} J_{\alpha+2}(x) = -2^\alpha x^2 z_{\alpha+2}.$$

- $\gamma = 2\alpha + 2$.

In this case,

$$b_0 = -\frac{\Gamma^2(\alpha+1)2^{\alpha+\beta+1}(\alpha+1)}{M + \Gamma^2(\alpha+1)2^{\alpha+\beta+1}(\alpha+1)},$$

$$b_1 = \frac{M}{2(M + \Gamma^2(\alpha+1)2^{\alpha+\beta+1}(\alpha+1))}.$$

By using (15) we deduce

$$\begin{aligned} \psi_{\alpha,0}(x) &= b_0 \left(\frac{x}{2}\right)^{-\alpha} J_\alpha(x) + 2b_1 \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2}(x) \\ &= \frac{-\Gamma^2(\alpha+1)2^{\alpha+\beta+1}}{M + \Gamma^2(\alpha+1)2^{\alpha+\beta+1}(\alpha+1)} \left(\frac{x}{2}\right)^{-\alpha} J_\alpha(x) \\ &\quad + \frac{M}{M + \Gamma^2(\alpha+1)2^{\alpha+\beta+1}(\alpha+1)} \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2}(x) \\ &= -\left(\frac{x}{2}\right)^{-\alpha} J_\alpha(x) + \frac{M(\alpha+1)}{M + 2^{\alpha+\beta+1}\Gamma^2(\alpha+1)(\alpha+1)} \left(\frac{x}{2}\right)^{-\alpha-1} J_{\alpha+1}(x) \\ &= -2^\alpha (z_\alpha(x) + a_{\alpha, \beta, M} z_{\alpha+1}(x)). \end{aligned}$$

4. Numerical experiments

In this section we illustrate the previous results on the zeros of the polynomials Q_n with some numerical experiments where we have taken $j = 3$ for all of them. Thus, we are dealing with the varying Sobolev inner product

$$(f, g)_{S,n} = \int_{-1}^1 f(x)g(x)(1-x)^\alpha(1+x)^\beta dx + M_n f^{(3)}(1)g^{(3)}(1).$$

We have used the mathematical software *Mathematica*® 8.0 for the computations. In all the numerical experiments we have computed the four largest zeros of the polynomials $Q_n(x)$ and the corresponding scaled zeros for several values of n . We only show one example for each possible case. In the tables about the scaled zeros we show their asymptotic behavior such as it is described in Proposition 4.

- Case $\gamma > 2(\alpha + 2j + 1)$.

We choose the following values:

$$\alpha = 3, \quad \beta = 1, \quad \gamma = 25, \quad \text{and} \quad M_n = \frac{3e^n}{(6e^n + 4)n^\gamma}.$$

It was proved in Theorem 2 that in this case the Mehler–Heine formula for the polynomials Q_n is the same one as for the classical Jacobi polynomials. This behavior is due to the negligible influence of the sequence of masses $\{M_n\}_{n \geq 0}$ on the asymptotics. Obviously, as it was stated in Proposition 4, this determines the asymptotic behavior of the zeros which is illustrated in Tables 1 and 2.

Table 1

Case $\gamma = 25 > 2(\alpha + 2j + 1)$, $j = 3$, $\alpha = 3$, $\beta = 1$, $\gamma = 25$, $M_n = \frac{3e^n}{(6e^n + 4)n^\gamma}$.

	$y_{n,4}$	$y_{n,3}$	$y_{n,2}$	$y_{n,1}$
$n = 150$	0.994346	0.99636	0.997952	0.999125
$n = 250$	0.997937	0.998672	0.999254	0.999681
$n = 500$	0.999479	0.999665	0.999811	0.999919

Table 2

Case $\gamma = 25 > 2(\alpha + 2j + 1)$, $j = 3$, $\alpha = 3$, $\beta = 1$, $\gamma = 25$, $M_n = \frac{3e^n}{(6e^n + 4)n^\gamma}$.

	$n\sqrt{2(1 - y_{n,1})}$	$n\sqrt{2(1 - y_{n,2})}$	$n\sqrt{2(1 - y_{n,3})}$	$n\sqrt{2(1 - y_{n,4})}$
$n = 150$	6.27524	9.59956	12.7982	15.9503
$n = 250$	6.31687	9.66386	12.885	16.0602
$n = 500$	6.34839	9.71233	12.9501	16.1421
Limit	$\mathbf{j}_{3,1} = 6.38016$	$\mathbf{j}_{3,2} = 9.76102$	$\mathbf{j}_{3,3} = 13.0152$	$\mathbf{j}_{3,4} = 16.2235$

Table 3

Case $\gamma = 4 < 2(\alpha + 2j + 1)$, $j = 3$, $\alpha = 3$, $\beta = -1/2$, $\gamma = 4$, $M_n = \frac{7\ln(n+1)+5}{(3+\ln(n^2))n^\gamma}$.

	$y_{n,4}$	$y_{n,3}$	$y_{n,2}$	$y_{n,1}$
$n = 150$	0.994574	0.996593	0.998169	0.999286
$n = 250$	0.998176	0.998915	0.999497	1.0016
$n = 500$	0.999554	0.999739	0.999883	1.0014

Table 4

Case $\gamma = 4 < 2(\alpha + 2j + 1)$, $j = 3$, $\alpha = 3$, $\beta = -1/2$, $\gamma = 4$, $M_n = \frac{7\ln(n+1)+5}{(3+\ln(n^2))n^\gamma}$.

	$n\sqrt{2(1 - y_{n,2})}$	$n\sqrt{2(1 - y_{n,3})}$	$n\sqrt{2(1 - y_{n,4})}$
$n = 150$	9.07735	12.382	15.6257
$n = 250$	7.92964	11.6463	15.1011
$n = 500$	7.6415	11.4238	14.9355
Limit	$\mathbf{s}_{3,1} = 7.64622$	$\mathbf{s}_{3,2} = 11.4432$	$\mathbf{s}_{3,3} = 14.9699$

- Case $\gamma < 2(\alpha + 2j + 1)$.

According to Theorem 2 the limit function in the Mehler–Heine formula is given by $\varphi_{\alpha,3}(x) = \sum_{i=0}^4 b_i 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x)$, where the coefficients b_i , $0 \leq i \leq 4$, can be computed from Theorem 1. We choose the following values:

$$\alpha = 3, \quad \beta = 1, \quad \gamma = 4, \quad \text{and} \quad M_n = \frac{7\ln(n+1)+5}{(3+\ln(n^2))n^\gamma}.$$

In Table 3 we can see that the largest zero is greater than 1 for n large enough according to Proposition 3. Table 4 shows the asymptotic behavior of the scaled zeros given in Proposition 4.

- Case $\gamma = 2(\alpha + 2j + 1)$.

According to Theorem 2 the limit function in the Mehler–Heine formula is given by $\psi_{\alpha,3}(x) = \sum_{i=0}^4 b_i 2^i \left(\frac{x}{2}\right)^{-\alpha} J_{\alpha+2i}(x)$, where the coefficients b_i , $0 \leq i \leq 4$, can be computed again from Theorem 1. We choose the following values:

$$\alpha = \beta = -9/10, \quad \gamma = 61/5 = 12.2,$$

and we denote by V the quantity which appears in Proposition 4, i.e.

$$V = \frac{2^{\alpha+\beta+2j+1}(\alpha+j+1)(\alpha+2j+1)\Gamma^2(\alpha+j+1)}{j}.$$

Thus, with this data

$$V = 2^{1/5} \frac{15128}{75} \Gamma^2\left(\frac{31}{10}\right) \simeq 1119.0037947.$$

Now we take

$$M_n = \frac{Mn^2(n-1/2)(n+2)}{n^{\gamma+4}} = \frac{Mn^2(n-1/2)(n+2)}{n^{81/5}}.$$

Table 5

Case $\gamma = 61/5 = 2(\alpha + 2j + 1)$, $M = 5 \leq V$, $j = 3$, $\alpha = -9/10$, $\beta = -9/10$, $\gamma = 61/5$, $M_n = \frac{5n^2(n-1/2)(n+2)}{n^{\gamma+4}}$.

	$y_{n,4}$	$y_{n,3}$	$y_{n,2}$	$y_{n,1}$
$n = 150$	0.99778	0.99854	0.999585	0.999991
$n = 250$	0.999142	0.999585	0.999871	0.999997
$n = 500$	0.999786	0.99985	0.999968	0.999999

Table 6

Case $\gamma = 61/5 = 2(\alpha + 2j + 1)$, $M = 5 \leq V$, $j = 3$, $\alpha = -9/10$, $\beta = -9/10$, $\gamma = 61/5$, $M_n = \frac{5n^2(n-1/2)(n+2)}{n^{\gamma+4}}$.

	$n\sqrt{2(1-y_{n,1})}$	$n\sqrt{2(1-y_{n,2})}$	$n\sqrt{2(1-y_{n,3})}$	$n\sqrt{2(1-y_{n,4})}$
$n = 150$	0.649565	4.02672	7.20558	10.3659
$n = 250$	0.64887	4.02249	7.19831	10.3561
$n = 500$	0.64853	4.01929	7.19273	10.3484
Limit	$t_{0,1} = 0.648561$	$t_{0,2} = 4.01985$	$t_{0,3} = 7.19169$	$t_{0,4} = 10.3446$

Table 7

Case $\gamma = 61/5 = 2(\alpha + 2j + 1)$, $M = 10^6 > V$, $j = 3$, $\alpha = -9/10$, $\beta = -9/10$, $\gamma = 61/5$, $M_n = \frac{10^6 n^2(n-1/2)(n+2)}{n^{\gamma+4}}$.

	$y_{n,4}$	$y_{n,3}$	$y_{n,2}$	$y_{n,1}$
$n = 150$	0.996412	0.999306	0.999978	1.00042
$n = 250$	0.99931	0.999739	0.999991	1.00009
$n = 500$	0.999818	0.999928	0.999999	1.000001

Table 8

Case $\gamma = 61/5 = 2(\alpha + 2j + 1)$, $M = 10^6 > V$, $j = 3$, $\alpha = -9/10$, $\beta = -9/10$, $\gamma = 61/5$, $M_n = \frac{10^6 n^2(n-1/2)(n+2)}{n^{\gamma+4}}$.

	$n\sqrt{2(1-y_{n,2})}$	$n\sqrt{2(1-y_{n,3})}$	$n\sqrt{2(1-y_{n,4})}$
$n = 150$	1.77464	6.0132	9.53661
$n = 250$	1.10344	5.71202	9.35539
$n = 500$	1.00403	5.58651	9.27349
Limit	$t_{0,1} = 0.903528$	$t_{0,2} = 5.34057$	$t_{0,3} = 9.07889$




Fig. 1. Case $\gamma > 2(\alpha + 2j + 1)$. Limit function and scaled polynomials $Q_n(1 - x^2/(2n^2))$.

According to Proposition 4 we have two possible choices of M which determine two different asymptotic behaviors of the zeros. In Tables 5 and 6 we show the case $M \leq V$ where $M = 5$. We can see that the largest zero of Q_n is always lesser than 1. However, when $M > V$ then $y_{n,1} > 1$ for n large enough and this is illustrated in Table 7 for $M = 10^6$. In Table 8 the asymptotic behavior of the scaled zeros is shown.

Finally, we illustrate Theorem 2 plotting the curves corresponding to the limit functions and to the scaled polynomials $Q_n(1 - \frac{x^2}{2n^2})$ with $n = 150$ and $n = 500$. In all the figures we have used the same values for the parameters as those ones taken previously in the numerical experiments about the zeros (see Figs. 1–4).

Acknowledgments

We thank the two anonymous referees for their useful suggestions to improve the paper. In special, one of the referees provided us with some references such as [6] which have been relevant for our research.




Fig. 2. Case $\gamma < 2(\alpha + 2j + 1)$. Limit function and scaled polynomials $Q_n(1 - x^2/(2n^2))$.




Fig. 3. Case $\gamma = 2(\alpha + 2j + 1)$. Limit function and scaled polynomials $Q_n(1 - x^2/(2n^2))$ with $M < V$.




Fig. 4. Case $\gamma = 2(\alpha + 2j + 1)$ Limit function and scaled polynomials $Q_n(1 - x^2/(2n^2))$ with $M > V$.

The authors JFMM and JJMB are partially supported by Research Group FQM-0229 (belonging to Campus of International Excellence CEIMAR). The author JFMM is funded by a grant of Plan Propio de la Universidad de Almería. The author FM is partially supported by Dirección General de Investigación, Ministerio de Economía y Competitividad Innovación of Spain, Grant MTM2012-36732-C03-01. The author JJMB is partially supported by Dirección General de Investigación, Ministerio de Ciencia e Innovación of Spain and European Regional Development Fund, grants MTM2011-28952-C02-01 and MTM2014-53963-P, and Junta de Andalucía (excellence grant P11-FQM-7276).

References

- [1] M. Alfaro, A. Martínez-Finkelshtein, M.L. Rezola, Asymptotics properties of balanced extremal Sobolev polynomials: coherent case, *J. Approx. Theory* 100 (1999) 44–59.
- [2] M. Alfaro, J.J. Moreno-Balcázar, A. Peña, M.L. Rezola, Sobolev orthogonal polynomials: Balance and asymptotics, *Trans. Amer. Math. Soc.* 361 (2009) 547–560.
- [3] F. Marcellán, Y. Xu, On Sobolev orthogonal polynomials, *Expo. Math.* 33 (2015) 308–352.
- [4] J.F. Mañas-Mañas, F. Marcellán, J.J. Moreno-Balcázar, Varying discrete Laguerre–Sobolev orthogonal polynomials: asymptotic behavior and zeros, *Appl. Math. Comput.* 222 (2013) 612–618.
- [5] A. Peña, M.L. Rezola, Discrete Laguerre–Sobolev expansions: A Cohen type inequality, *J. Math. Anal. Appl.* 385 (2012) 254–263.
- [6] A. Peña, M.L. Rezola, Connection formulas for general discrete Sobolev polynomials: Mehler–Heine asymptotics, *Appl. Math. Comput.* 261 (2015) 216–230.
- [7] J.J. Moreno-Balcázar, Varying Jacobi–Kral orthogonal polynomials: local asymptotic behaviour and zeros, *Ramanujan J.* 28 (2012) 79–88.
- [8] G. Szegő, *Orthogonal Polynomials*, fourth ed., in: *Amer. Math. Soc. Colloq. Publ.*, vol. 23, Amer. Math. Soc., Providence, RI, 1975.
- [9] R.A. Askey, R. Roy, Gamma function, in: *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge, UK, 2010, pp. 135–147.
- [10] F. Marcellán, A. Ronveaux, On a class of polynomials orthogonal with respect to a discrete Sobolev inner product, *Indag. Math. (N.S.)* 1 (4) (1990) 451–464.
- [11] H.G. Meijer, Zero distribution of orthogonal polynomials in a certain discrete Sobolev space, *J. Math. Anal. Appl.* 172 (1993) 520–532.
- [12] M. Alfaro, G. López, M.L. Rezola, Some properties of zeros of Sobolev-type orthogonal polynomials, *J. Comput. Appl. Math.* 69 (1996) 171–179.
- [13] M. Abramowitz, I. Stegun, *Handbook of Mathematical Functions*, Dover, New York, 1972.