



# Cuspidal plane curves, syzygies and a bound on the MW-rank

Remke Kloosterman<sup>1</sup>

Institut für Mathematik, Humboldt-Universität zu Berlin, Unter den Linden 6, D-10099 Berlin, Germany

## ARTICLE INFO

### Article history:

Received 24 November 2011

Available online 30 November 2012

Communicated by Bernd Ulrich

### MSC:

14H30

13D02

14H50

14J30

### Keywords:

Elliptic threefolds

Mordell–Weil rank

Alexander polynomials of plane curves

## ABSTRACT

Let  $C = Z(f)$  be a reduced plane curve of degree  $6k$ , with only nodes and ordinary cusps as singularities. Let  $I$  be the ideal of the points where  $C$  has a cusp. Let  $\bigoplus S(-b_i) \rightarrow \bigoplus S(-a_i) \rightarrow S \rightarrow S/I$  be a minimal resolution of  $I$ . We show that  $b_i \leq 5k$ . From this we obtain that the Mordell–Weil rank of the elliptic threefold  $W: y^2 = x^3 + f$  equals  $2\#\{i \mid b_i = 5k\}$ . Using this we find an upper bound for the Mordell–Weil rank of  $W$ , which is  $\frac{1}{18}(125 + \sqrt{73} - \sqrt{2302 - 106\sqrt{73}})k + l.o.t.$  and we find an upper bound for the exponent of  $(t^2 - t + 1)$  in the Alexander polynomial of  $C$ , which is  $\frac{1}{36}(125 + \sqrt{73} - \sqrt{2302 - 106\sqrt{73}})k + l.o.t.$  This improves a recent bound of Cogolludo and Libgober almost by a factor 2.

© 2012 Elsevier Inc. All rights reserved.

## 1. Introduction

In this paper we study reduced plane curves  $C$  of degree  $d = 6k$  having only nodes and ordinary cusps as singularities. We allow  $C$  to be reducible. Let  $z_0, z_1, z_2$  be coordinates on  $\mathbf{P}^2$ , and let  $S = \mathbf{C}[z_0, z_1, z_2]$ . Let  $f \in S_{6k}$  be an equation for  $C$ . Let  $\Sigma$  be the set of cusps of  $C$  (we will ignore the nodes).

Consider now the elliptic threefold defined by

$$Z(-y^2 + x^3 + f) \subset \mathbf{P}(2k, 3k, 1, 1, 1).$$

<sup>1</sup> E-mail address: klooster@math.hu-berlin.de.

<sup>1</sup> The author thanks Alberto Calabri, Alexandru Dimca, David Eisenbud, Anatoly Libgober, Matthias Schütt and Orsola Tommasi for several comments on a previous versions of this paper. The author thanks the referee for many useful comments. The author acknowledges the partial support from the DFG under research grant KL 2244/2-1.

Let  $\text{MW}(\pi)$  be the Mordell–Weil group, i.e., the group of rational sections of the elliptic fibration. It is known that the rank of  $\text{MW}(\pi)$  can be expressed in terms of the geometry of  $C$  (see Proposition 3.1), namely

$$\text{rank MW}(\pi) = 2 \dim \text{coker} \left( S_{5k-3} \xrightarrow{\oplus \text{ev}_p} \bigoplus_{p \in \Sigma} \mathbf{c} \right).$$

We can also consider the fundamental group  $\pi_1(\mathbf{P}^2 \setminus C)$ . With this group we can associate the so-called Alexander polynomial of  $C$ . It turns out that the exponent of  $(t^2 - t + 1)$  in the Alexander polynomial equals

$$\dim \left( \text{coker } S_{5k-3} \xrightarrow{\oplus \text{ev}_p} \bigoplus_{p \in \Sigma} \mathbf{c} \right).$$

Hence both invariants coincide. Cogolludo and Libgober [2] noticed this and proved for a much larger class of singular plane curves that the degree of the Alexander polynomial is related with the Mordell–Weil group of an associated elliptic fibration.

In this paper we give a non-trivial upper bound  $g(k)$  for the Mordell–Weil rank, which also yields an upper bound for the exponent of  $t^2 - t + 1$  in the Alexander polynomial. Asymptotically we have that

$$\lim_{k \rightarrow \infty} \frac{g(k)}{k} = \frac{1}{18} (125 + \sqrt{73} - \sqrt{2302 - 106\sqrt{73}}) \approx 5.34. \quad (1)$$

The best known previous upper bound for the exponent of  $(t^2 - t + 1)$  in the Alexander polynomial of a cuspidal curve seems to be due to Cogolludo and Libgober [2], and equals  $5k - 1$  (this implies that the MW-rank is at most  $10k - 2$ ). This bound is an immediate consequence of the Shioda–Tate formula. The divisibility theorem for the Alexander polynomial of Libgober yields an upper bound of  $6k - 2$  for the exponent in the Alexander polynomial.

Our bound is deduced from two other bounds. Suppose we fix  $r, k$  and look for  $C_{2r,k}$  the minimal number of cusps on a degree  $6k$  curve such that the corresponding elliptic fibration has Mordell–Weil rank at least  $2r$ . We show that

$$C_{2,k} = 6k^2 \quad \text{and} \quad C_{2r,k} \geq 6k^2 + 3(r-1)k - \frac{3}{4}r(r-1) + O\left(\frac{1}{k}\right) \quad \text{for } k \rightarrow \infty. \quad (2)$$

The number of cusps on a degree  $d$  curve can be bounded by  $\frac{125+\sqrt{73}}{432}d^2 - \frac{511+11\sqrt{73}}{1752}d$  (see [10]). Combining both bounds yields the upper bound (1). This bound is very unlikely to be sharp. We expect that  $C_{2r,k}$  can be bounded from below by a function of the form  $h(r)k^2$ , where  $h$  is increasing in  $r$ , rather than constant. However, the bound for  $C_{2,k}$  is sharp. If we take general polynomials  $f_1 \in S_{2k}$  and  $f_2 \in S_{3k}$ , and set  $f = f_1^3 + f_2^2$ , then  $f$  has  $6k^2$  cusps and the Mordell–Weil rank is at least 2. The fact that  $C_{2,k} \geq 6k^2$  holds, can also be obtained by different methods, namely if  $C$  has less than  $6k^2$  cusps then  $\pi_1(\mathbf{P}^2 \setminus C)$  is abelian and therefore  $C$  has constant Alexander polynomial. In particular, the Mordell–Weil rank is zero in this case, see [14].

The main idea of the proofs is to consider the resolution of the ideal  $I$  of  $\Sigma$ :

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i) \rightarrow S \rightarrow S/I \rightarrow 0.$$

There are several restrictions on  $a_i, b_i$  coming from the fact that  $I$  is the ideal of a finite set of points in  $\mathbf{P}^2$ . These restrictions are classically known, see Proposition 2.1. We find further restrictions on the

$a_i$  and  $b_i$  by a combination of Bezout's theorem and an upper bound for the number of cusps on a degree  $d$  plane curve. See Proposition 2.4.

Using specializations to elliptic surfaces we show that  $b_i \leq 5k$  for all  $i$ . Using an expression for the difference between the Hilbert polynomial of  $I$  and the Hilbert function of  $I$  we obtain that  $\text{rank MW}(\pi) = 2\#\{i \mid b_i = 5k\}$ . This fact is proved in Proposition 3.5.

After distributing a preliminary version of this paper we learned the following: Zariski [14] proved that the Castelnuovo–Mumford regularity of the cuspidal locus of an *irreducible* plane curve is at most  $5k - 1$ . This is done by studying the cyclic degree  $6k$  cover of  $\mathbf{P}^2$  ramified along the curve  $C$ . The statement on the regularity implies that  $b_i \leq 5k$  holds in the case of irreducible curves. In the case of reducible curves Zariski proved that the regularity of the cuspidal locus is at most  $6k - 2$ .

The statement  $b_i \leq 5k$  in the case of reducible curves seems to be known to the experts, although we could not identify a proof for this statement in the literature. The techniques to extend Zariski's proof to the reducible case have been around since the beginning of the 1980s [6,11]. However, our proof is different from the existing proofs in the literature.

There are other examples where the highest degree syzygies of the ideal of the singular locus has a geometric interpretation. E.g., if we consider a minimal resolution of the ideals of the nodes, then a syzygy has degree at most the degree of the curve, and the number of highest degree syzygies is one less than the number of irreducible components of the curve (Proposition 3.6).

In Section 4 we prove the bound (2) under an extra technical assumption on the  $a_i$  and  $b_i$ : After permuting the  $a_i$  and  $b_i$  we may assume that the  $a_i$  and  $b_i$  both form a descending sequence. A priori, we know that  $b_i > a_i$ . In Section 4 we assume that  $a_i \leq b_{i+1}$  for all  $i$ .

In Section 5 we study the case when there is some  $i$  with  $a_i > b_{i+1}$ . We consider the ideal generated by all generators of  $I$  of degree less than  $a_i$ . This ideal defines a subscheme of  $\mathbf{P}^2$  which is the union of a (possibly non-reduced) curve and a zero-dimensional scheme. We can analyze this situation in similar way as above and obtain further restrictions on the  $a_i$  and the  $b_i$ . These further obstructions allow us to construct a new sequence  $a'_i, b'_i$  that corresponds with an ideal  $I'$  such that  $\#Z(I') \leq \#Z(I)$  and the  $a'_i, b'_i$  satisfy the extra condition used in Section 4. Hence also in this case the lower bound (2) holds.

## 2. Resolution of the locus of cusps

We cite first a result on the resolution of the ideal of finitely many points in  $\mathbf{P}^2$ .

**Proposition 2.1.** *Let  $I$  be the ideal of finitely many distinct points in  $\mathbf{P}^2$ . Then  $I$  has a free resolution*

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i) \rightarrow S \rightarrow S/I \rightarrow 0, \quad (3)$$

such that

- (1) for all  $i$  we have that  $a_i, b_i \in \mathbf{Z}$  and  $a_i > 0, b_i > 0$ ;
- (2)  $\sum_{i=1}^{t+1} a_i = \sum_{i=1}^t b_i$ ;
- (3) for  $i = 1, \dots, t$  we have  $b_i > a_i \geq a_{i+1}$  and for  $j = 1, \dots, t-1$  we have  $b_j \geq b_{j+1}$ ;
- (4)  $\#Z(I) = \frac{1}{2}(\sum b_i^2 - \sum a_i^2) = B(s) - \sum_{i=1}^{t+1} B(s - a_i) + \sum_{i=1}^t B(s - b_i)$  for every  $s$ , where  $B(s) = \frac{1}{2}(s + 1)(s + 2)$ .

**Proof.** This follows almost immediately from the fact that  $I$  has a free resolution of length 1 and that the Hilbert polynomial of  $I$  is constant. See [5, Section 3.1].  $\square$

**Definition 2.2.** For  $d$  a positive integer, define  $M(d)$  to be the maximal number of ordinary cusps on a (possibly reducible) degree  $d$  curve.

**Remark 2.3.** The best known asymptotic upper bound for  $M(d)$  we are aware of is a bound obtained by Langer (see [10, Section 11]). Langer states that

$$\limsup_{d \rightarrow \infty} M(d)/d^2 \leq \frac{125 + \sqrt{73}}{432}.$$

However, a closer inspection of the proof reveals that

$$M(d) \leq \frac{125 + \sqrt{73}}{432} d^2 - \frac{511 + 11\sqrt{73}}{1752} d.$$

On the other side, Hirano showed that [7, Corollary 3]

$$\limsup_{d \rightarrow \infty} \frac{M(d)}{d^2} \geq \frac{9}{32}.$$

**Proposition 2.4.** Suppose  $I$  is the ideal of the locus of cusps of a plane curve  $C$  of degree  $d$ . Then

$$\#Z(I) \leq \min\left(\frac{a_{t+1}d}{2}, M(d)\right) \leq \min\left(a_{t+1}, \frac{5}{8}d - \frac{3}{4}\right) \frac{d}{2}.$$

**Proof.** The upper bound  $\#Z(I) \leq M(d)$  is obvious. We prove now that  $\#Z(I) \leq \frac{a_{t+1}d}{2}$ .

Since the ideal  $I$  contains an element of degree  $a_{t+1}$ , there exists a curve  $C'$  of degree  $a_{t+1}$  such that all cusps of  $C$  are points of  $C'$ . By Bezout's theorem we have that if  $2\#Z(I) > a_{t+1}d$  then  $C$  and  $C'$  have a common component  $C''$  of degree  $d'' \leq a_{t+1}$ . Since cusps are irreducible singularities it follows that all the cusps of  $C$  that are also points of  $C'$  are actually cusps of  $C''$ . Hence

$$\#Z(I) \leq M(d'') + \frac{1}{2}(d - d'')(a_{t+1} - d'') \leq \frac{5}{16}(d'')^2 + \frac{1}{2}a_{t+1}d + \frac{1}{2}(d'')^2 - \frac{1}{2}d''(a_{t+1} + d).$$

Combining this with  $\#Z(I) \geq \frac{1}{2}da_{t+1}$  yields

$$0 \leq \frac{13}{16}(d'')^2 - \frac{1}{2}d''(a_{t+1} + d).$$

Dividing this inequality by  $d''/2$  and using  $2d'' \leq a_{t+1} + d$  yields

$$0 \leq \frac{13}{8}d'' - (a_{t+1} + d) \leq \frac{-3}{8}d''.$$

Hence the degree of  $C''$  is 0. Equivalently, the component  $C''$  does not exist and  $\#Z(I) \leq \frac{1}{2}a_{t+1}d$ .  $\square$

### 3. Syzygies and MW-rank

Let  $S := \mathbb{C}[z_0, z_1, z_2]$  be the polynomial ring in three variables. Let  $S_d$  be the subspace of homogeneous polynomials of degree  $d$ .

Fix an integer  $k$  and a square-free polynomial  $f \in S_{6k}$ , such that the plane curve  $C = Z(f)$  has only nodes and ordinary cusps as singularities. Let  $\Sigma$  denote the set of cusps of  $C$ . Let  $I \subset S$  be the ideal of  $\Sigma$ .

Let  $W_f \subset \mathbf{P}(2k, 3k, 1, 1, 1)$  be the hypersurface given by the vanishing of

$$-y^2 + x^3 + f.$$

The threefold  $W_f$  is birational to an elliptic threefold  $\pi : X \rightarrow R$ , where  $R$  is a rational surface and the elliptic fibration  $\pi$  is birational to the projection  $\psi : W_f \setminus \{(1 : 1 : 0 : 0 : 0)\} \rightarrow \mathbf{P}^2$  from  $(1 : 1 : 0 : 0 : 0)$  onto the plane  $\{x = y = 0\}$ . The explicit construction of  $\pi$  is slightly complicated, see [12]. For  $p \in \mathbf{P}^2$  the Zariski closure of  $\psi^{-1}(p)$  is either an elliptic curve with  $j$ -invariant 0 or a cuspidal cubic, depending on whether  $p \in C$  or not.

The Mordell–Weil group  $\text{MW}(\pi)$  of  $\pi$  is the group of rational sections of  $\pi$ . This is a finitely generated group, and if the singularities of  $C$  are “mild” then one has an algorithm to compute the rank of  $\text{MW}(\pi)$ , see [8]:

**Proposition 3.1.** *We have the following equality*

$$\text{rank MW}(\pi) = 2 \dim \left( \text{coker } S_{5k-3} \xrightarrow{\text{ev}_p} \bigoplus_{p \in \Sigma} \mathbf{C} \right). \quad (4)$$

**Proof.** For the case  $k = 1$  see [9, Section 9]. The general case follows along the same lines:

An  $A_1$  singularity of  $C$  yields an  $A_2$  singularity of  $W_f$ , whereas an  $A_2$  singularity of  $C$  yields a  $D_4$  singularity on  $W_f$ .

Let  $\Sigma$  be the singular locus. We can now compute  $H^4(W_f)_{\text{prim}}$  as the co-kernel of

$$H^4(\mathbf{P}(2k, 3k, 1, 1, 1) \setminus W_f) \cong H^3(W_f \setminus \Sigma) \rightarrow H^4_{\Sigma}(W_f).$$

An  $A_2$  singularity of  $W_f$  does not contribute to  $H^4_{\Sigma}$ , whereas a  $D_4$  singularity does [3, Example 1.9]. Actually, using the ideas from [3, Section 1] it follows that  $H^4_p(W_f) = \mathbf{C}(-2)^2$  if  $p$  is a  $D_4$  singularity of  $W_f$ , hence  $H^4_{\Sigma}(W_f)$  is of pure Hodge type  $(2, 2)$ . Then by the main results of [8] we have  $\text{rank MW}(\pi) = h^4(W_f) - 1$ .

Let  $\omega$  be a third root of unity. The map  $\omega : [x : y : z_0 : z_1 : z_2] \mapsto [\omega x : y : z_0 : z_1 : z_2]$  is an automorphism of  $W_f$  and fixes every point of  $\Sigma$ . The map  $H^4(\mathbf{P}(2k, 3k, 1, 1, 1) \setminus W_f) \rightarrow H^4_{\Sigma}(W_f)$  is  $\omega^*$ -equivariant, so we may decompose it in an  $\omega$ - and  $\omega^2$ -eigenspace, which have both the same dimension, and a 1-eigenspace, which is trivial.

The argument used in [9, Section 9] shows in this case that the  $\omega$ -eigenspace of the co-kernel of  $H^4(\mathbf{P}(2k, 3k, 1, 1, 1) \setminus W_f) \rightarrow H^4_{\Sigma}(W_f)$  has dimension

$$\dim \left( \text{coker } S_{5k-3} \xrightarrow{\text{ev}_p} \bigoplus_{p \in \Sigma} \mathbf{C} \right). \quad \square$$

**Remark 3.2.** With the fundamental group of  $\mathbf{P}^2 \setminus C$  one can associate the so-called Alexander polynomial. Cogolludo and Libgober [2] showed that the exponent of  $t^2 - t + 1$  in this polynomial equals half the rank of  $\text{MW}(\pi)$ .

Hence each statement which we make on the rank of  $\text{MW}(\pi)$  is also a statement on the exponent of  $t^2 - t + 1$  in the Alexander polynomial of a cuspidal curve.

We will calculate  $\text{rank MW}(\pi)$  using a projective resolution of  $I$  and use properties of the resolution to bound the Mordell–Weil rank in terms of  $k$ . Two more or less obvious restrictions on the resolution come from Bezout’s theorem (Proposition 2.4) and the upper bound for the maximal number of cusps on a degree  $d$  plane curve (Remark 2.3). The third restriction comes from considerations on the Mordell–Weil rank of elliptic surfaces.

**Lemma 3.3.** Let  $J \subset S$  be the ideal of a zero-dimensional scheme. Let

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i) \rightarrow S \rightarrow S/J \rightarrow 0$$

be a minimal resolution of  $S/J$ . Let  $n$  be integer and set  $b'_i := b_i - n$ ,  $a'_i := a_i - n$ . Then the defect of the linear system of degree  $n - 3$  polynomials vanishing along  $Z(J)$  equals

$$\left( \sum_{i|b'_i \geq 0} (b'_i + 1)(b'_i + 2) - \sum_{i|a'_i \geq 0} (a'_i + 1)(a'_i + 2) \right).$$

**Proof.** The defect of the linear system is precisely the difference between the Hilbert function of  $J$  in degree  $n - 3$  and the Hilbert polynomial of  $I$ . Let  $B(s)$  be as in the previous section. Then Hilbert polynomial  $P_J(s)$  equals

$$B(s) + \sum_{i=1}^t B(s - b_i) - \sum_{i=1}^{t+1} B(s - a_i).$$

The Hilbert function  $h_J(s)$  evaluated at  $s$  equals

$$B(s) + \sum_{i|b_i \leq s} B(s - b_i) - \sum_{i|a_i \leq s} B(s - a_i).$$

Since  $B(-1) = B(-2) = 0$  we may replace the condition “ $\leq s$ ” by “ $\leq s + 2$ ” in the above formula. Hence

$$p_J(s) - h_J(s) = \sum_{i|b_i \geq s+3} B(s - b_i) - \sum_{i|a_i \geq s+3} B(s - a_i).$$

Substituting  $s = n - 3$  in the above formula finishes the proof.  $\square$

**Lemma 3.4.** Let  $J \subset S$  be the ideal of a zero-dimensional scheme. Let

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i) \rightarrow S \rightarrow S/J \rightarrow 0$$

be a minimal resolution of  $S/J$ . Suppose there is an integer  $n$  and a linear function  $g(m)$  such that for all positive integers  $m$  and for a general choice of three homogeneous polynomials  $g_0, g_1, g_2 \in S_m$  the defect  $\delta_m$  of the linear system of degree  $mn - 3$  polynomials vanishing along  $Z(\varphi^*(J))$  is at most  $g(m)$ , where  $\varphi : \mathbf{P}^2 \rightarrow \mathbf{P}^2$  is given by  $(z_0 : z_1 : z_2) \mapsto (g_0 : g_1 : g_2)$ .

Then  $b_i \leq n$ ,  $a_i < n$  and  $\delta_m = \#\{i \mid b_i = n\}$ .

**Proof.** Define  $a'_i$  and  $b'_i$  as in Lemma 3.3. Suppose that for some  $j$  we have  $b'_j > 0$ . Fix a positive integer  $w$  such that

$$\begin{aligned} (b'_j w + 1)(b'_j w + 2) &> g(w) && \text{if } a'_j < 0 \quad \text{or} \\ (b'_j w + 1)(b'_j w + 2) - (a'_j w + 1)(a'_j w + 2) &> g(w) && \text{if } a'_j \geq 0. \end{aligned}$$

Let  $\varphi : \mathbf{P}^2 \rightarrow \mathbf{P}^2$  be a general map of degree  $w$ . Then the resolution of  $S/\varphi^*(J)$

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i w) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i w) \rightarrow S \rightarrow S/J \rightarrow 0.$$

In particular, by the previous lemma we obtain

$$\delta_w = \left( \sum_{i|b'_i \geq 0} (b'_i w + 1)(b'_i w + 2) - \sum_{i|a'_i \geq 0} (a'_i w + 1)(a'_i w + 2) \right) > g(w).$$

This contradicts  $\delta_w \leq g(w)$ . Hence  $b'_i \leq 0$ , for all  $i$  and  $a'_i < 0$  for all  $i$ .

Applying Lemma 3.3 again yields that for all  $m$  we have  $\delta_m = \#\{i \mid b_i = n\}$ .  $\square$

We return now to the case where  $C$  is a cuspidal curve and  $I$  is the ideal of the points where  $C$  has a cusp.

**Proposition 3.5.** *Let*

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i) \rightarrow S \rightarrow S/I \rightarrow 0$$

*be a resolution of  $I$ . Then for all  $i$  we have that  $b_i \leq 5k$ ,  $a_i < 5k$  and*

$$\text{rank MW}(\pi) = 2\#\{i \mid b_i = 5k\}.$$

**Proof.** Take a general line  $\ell \subset \mathbf{P}^2$ . The (projective) surface  $\overline{\pi^{-1}(\ell)} \subset \mathbf{P}(2k, 3k, 1, 1, 1)$  might be singular. Denote with  $\widetilde{\pi^{-1}(\ell)}$  a resolution of singularities of this surface. This surface admits a natural elliptic fibration  $\pi_\ell : \widetilde{\pi^{-1}(\ell)} \rightarrow \ell$ . From the theory of elliptic surface we obtain the following well-known inequalities:

$$\text{rank MW}(\pi) \leq \text{rank MW}(\pi_\ell) \leq h^{1,1}(\widetilde{\pi^{-1}(\ell)}) - 2 = 10k - 2. \quad (5)$$

The first inequality is a standard result on specializations. The second inequality follows from the Shioda–Tate formula. The final equality is a well-known fact for elliptic surfaces, see e.g., [13].

Now take three general polynomials  $g_0, g_1, g_2$  of degree  $w$ . Let  $\varphi : \mathbf{P}^2 \rightarrow \mathbf{P}^2$  be the map defined by  $\varphi(z_0 : z_1 : z_2) = (g_0 : g_1 : g_2)$ . Let  $\tilde{f} = \varphi^*(f) \in S_{6kw}$  and let  $\pi_w : X_w \rightarrow R'$  be the pullback of the elliptic fibration  $\pi : X \rightarrow R$ . For general  $g_i$  the curve defined by  $\tilde{f}$  has only nodes and ordinary cusps as singularities and the locus  $\tilde{\Sigma}$  consisting of the cusps of  $Z(\tilde{f})$  equals  $\varphi^{-1}(\Sigma)$ . In particular, the corresponding ideal  $\tilde{I}$  has the following minimal free resolution

$$0 \rightarrow \bigoplus_{i=1}^t S(-b_i w) \rightarrow \bigoplus_{i=1}^{t+1} S(-a_i w) \rightarrow S \rightarrow S/\tilde{I} \rightarrow 0.$$

From Proposition 3.1 it follows that the rank of  $\text{MW}(\pi)$  is twice the dimension of the co-kernel of the evaluation map  $S_{5k-3} \xrightarrow{\text{ev}_p} \bigoplus_{p \in \Sigma} \mathbf{C}$ , which equals the defect of the linear system of degree  $5k-3$  polynomials through  $Z(I)$ . Similarly the Mordell–Weil rank of  $\pi_w$  equals the defect of the linear system of degree  $5kw-3$  polynomials through  $Z(\varphi^*I)$ . This defect is bounded by  $10kw-2$ , which is

linear in  $w$ . Hence we can apply Lemma 3.4 and we obtain that  $b_i \leq 5k$ ,  $a_i < 5k$  and  $\text{rank MW}(\pi) = 2\#\{i \mid b_i = 5k\}$ .  $\square$

If  $C'$  is an irreducible curve of degree  $d$  and  $I'$  the ideal of the points of  $C'$  where  $C'$  has a node then each syzygy of  $I'$  has degree at most  $d - 1$  (this is implied by the exercises 24 and 31 of [1, Appendix A]). Actually, a statement analogous to Proposition 3.5 holds for the locus of nodes of a plane curve, but we could not find this particular result in the literature.

**Proposition 3.6.** *Let  $C' \subset \mathbf{P}^2$  be a reduced plane curve of degree  $d$  with only nodes and ordinary cusps as singularities. Let  $c$  be the number of irreducible components of  $C'$ . Define  $\mathcal{N}$  to be the locus of nodes of  $C'$ . Let  $I'$  be the ideal of  $\mathcal{N}$  and*

$$0 \rightarrow \bigoplus S(-b_i) \rightarrow \bigoplus S(-a_i) \rightarrow S \rightarrow S/I' \rightarrow 0$$

be a minimal resolution of  $I'$ . Then  $b_i \leq d$  and

$$\#\{i \mid b_i = d\} = c - 1.$$

**Proof.** From the Mayer–Vietoris sequence it follows easily that  $h^2(C') = c$ . We would like to use Dimca’s method [3] to calculate  $h^2(C')$  in terms of the defect of a linear system. However, the previous discussion was based on the results from [3], and this text considers only hypersurfaces in weighted projective spaces of dimension at least 3. A variant that works for plane curves is presented in [4, p. 201].

Let  $\Sigma' = C'_{\text{sing}}$ . If  $\Sigma'$  is empty then there is nothing to prove, so assume that  $\Sigma'$  is non-empty. Let  $C^* = C' \setminus \Sigma'$ , let  $\mathbf{P}^* = \mathbf{P}^2 \setminus \Sigma'$  and  $U = \mathbf{P}^* \setminus C^* = \mathbf{P}^2 \setminus C'$ .

Let  $p \in \Sigma'$ ,  $V_p \subset \mathbf{P}^2$  be a small neighborhood of  $p$  and  $D_p = C' \cap V_p$ . Then in [4, p. 201] it is shown that  $H^2(C')_{\text{prim}}$  equals the co-kernel of

$$H^2(U)(1) \rightarrow \bigoplus_{p \in \Sigma'} H^2(V_p \setminus D_p).$$

Following the discussion after [3, Formula (1.7)] we obtain that  $H^2(V_p \setminus D_p)$  is zero if  $p$  is a cusp and is one-dimensional if  $p$  is a node. In the nodal case we have that  $H^2(V_p \setminus D_p)$  is spanned by  $\frac{1}{xy} dx \wedge dy$ .

The map  $H^2(U)(1) \rightarrow \bigoplus_{p \in \mathcal{N}} \mathbf{C} \frac{1}{xy} (dx \wedge dy)$  can be given explicitly as follows. Let  $f \in S_d$  be a polynomial defining  $C'$ . Set

$$\Omega := z_0 z_1 z_2 \left( \frac{dz_1}{z_1} \wedge \frac{dz_2}{z_2} - \frac{dz_0}{z_0} \wedge \frac{dz_2}{z_2} + \frac{dz_0}{z_0} \wedge \frac{dz_1}{z_1} \right).$$

Following [3, Section 1] we have  $F^3 H^2(U) = 0$ ,

$$F^2 H^2(U) \subset \left\{ \frac{g}{f} \Omega \mid g \in S_{d-3} \right\}$$

and  $H^2(U) = F^0 H^2(U) = F^1 H^2(U)$ . Moreover, the latter space is spanned by

$$\left\{ \frac{g}{f} \Omega \mid g \in S_{d-3} \right\} \cup \left\{ \frac{h}{f^2} \Omega \mid h \in S_{2d-3} \right\}.$$



Since  $H^2(U)(1) \rightarrow H^2_{\Sigma'}(C)$  is a morphism of Hodge structures and the image has only classes of type  $(1, 1)$ , it follows that  $\text{Gr}_1^F H^2(U)$  is mapped to zero in  $H^2_{\Sigma'}(C')$ . Hence to determine the co-kernel of  $H^2(U)(1) \rightarrow \bigoplus \mathbf{C} \frac{1}{xy} (dx \wedge dy)$  we can restrict this map to  $F^2 H^2(U)$ . From the local construction of this map it follows directly that  $\frac{g}{f} \Omega$  is mapped to  $g(p) \frac{1}{xy} (dx \wedge dy)$ . Combining everything we obtain that

$$c - 1 = h^2(C')_{\text{prim}} = \dim \text{coker} \left( S_{d-3} \rightarrow \bigoplus_{p \in \mathcal{N}} \mathbf{C} \right). \quad (6)$$

Hence  $c - 1$  equals the defect of the linear system of degree  $d - 3$  polynomial through the set of nodes of  $C$ . For a general degree  $m$  base change, the pullback  $C'_m$  of  $C'$  is a nodal curve, and the ideal of the nodes of  $C'$  is the pullback of  $I$ . The number of irreducible components of  $C_m$ , and hence the defect of the linear system of degree  $md - 3$  polynomials through the nodes, can be bounded by  $md$ . Hence we may apply Lemma 3.4 to the minimal resolution of  $I'$ , and obtain that  $b_i \leq d$ ,  $a_i < d$  and  $c - 1 = \#\{i \mid b_i = d\}$ .  $\square$

#### 4. Upper bound for rank $\text{MW}(\pi)$ (strongly admissible case)

As discussed in the proof of Proposition 3.5 we have

$$\text{rank MW}(\pi) \leq 10k - 2.$$

This upper bound is a corollary from the Shioda–Tate formula for the Mordell–Weil rank of elliptic surfaces. Cogolludo and Libgober [2] used this bound to bound the degree of the Alexander polynomial.

In this and the next section we will give an upper bound  $g(k)$  for  $\text{rank MW}(\pi)$  such that

$$\lim_{k \rightarrow \infty} \frac{g(k)}{k} = \frac{1}{18} (125 + \sqrt{73} + \sqrt{2302 - 106\sqrt{73}}) \approx 5.34.$$

**Definition 4.1.** Fix a positive integer  $t$ . Let  $a_1, \dots, a_{t+1}, b_1, \dots, b_t$  be a sequence of positive integers. If no confusion arises we write  $\mathbf{a}, \mathbf{b}$  for this sequence. We call  $\mathbf{a}, \mathbf{b}$   $k$ -admissible for rank  $2r$  and lowest degree  $D_0$  if:

- (1)  $\sum a_i = \sum b_i$ .
- (2)  $a_i < b_i$  for  $i = 1, \dots, t$ .
- (3)  $a_i \geq a_{i+1}$ ,  $b_i \geq b_{i+1}$ .
- (4)  $a_{t+1} = D_0$ .
- (5)  $b_i \leq 5k$  for  $i = 1, \dots, t$ .
- (6)  $\#\{i \mid b_i = 5k\} \geq r$ .
- (7)  $c(\mathbf{a}, \mathbf{b}) := \frac{1}{2} (\sum_{i=1}^{t+1} b_i^2 - \sum_{i=1}^t a_i^2) \leq \min(M(6k), 3ka_{t+1})$ .

We call a sequence *strongly  $k$ -admissible* if  $a_i \leq b_{i+1}$  for all  $i = 1, \dots, t - 1$ .

We call a (strongly)  $k$ -admissible sequence *reduced* if  $\{a_i\} \cap \{b_i\} = \emptyset$ .

**Remark 4.2.** Given a (strongly)  $k$ -admissible sequence  $\mathbf{a}, \mathbf{b}$  we can construct a reduced  $k$ -admissible sequence by repeatedly throwing out  $b_i$  and  $a_j$  in case they are equal. The new sequence is easily to be seen  $k$ -admissible, since this operation does not change  $c(\mathbf{a}, \mathbf{b})$ . Moreover, this reduction transforms a  $k$ -admissible sequence into a  $k$ -admissible sequence and it transforms a strongly  $k$ -admissible sequence into a strongly  $k$ -admissible sequence.

The results of Sections 2 and 3 show that:

**Lemma 4.3.** Suppose that  $y^2 = x^3 + f$  has Mordell–Weil rank  $2r$ . Let  $D_0$  be the degree of a generator of minimal degree of  $I$ . Then there exists a  $k$ -admissible sequence  $(\mathbf{a}, \mathbf{b})$  of rank  $2r$  and lowest degree  $D_0$ , such that  $c(\mathbf{a}, \mathbf{b}) = \#Z(I)$ .

**Remark 4.4.** In this section we will study strongly  $k$ -admissible sequences. In the next section we will show that if  $y^2 = x^3 + f$  has rank  $2r$  then there exists a strongly  $k$ -admissible sequence for rank  $2r$  and lowest degree  $D'_0 \leq D_0$ . Hence to prove the desired bounds we only have to consider strongly  $k$ -admissible sequences.

For technical reasons, we have to discuss the following type of sequences separately:

**Lemma 4.5.** Suppose that  $t = r$ ,  $a_1 = \dots = a_r = 5k - 1$ ,  $a_{r+1} = r$  and  $k \geq 2$ . Then

$$c(\mathbf{a}, \mathbf{b}) \leq \min(M(6k), 3kD_0)$$

implies  $k = 1$  and  $r = 3$ .

**Proof.** The first two assumptions imply that

$$c(\mathbf{a}, \mathbf{b}) = 5rk - \frac{1}{2}r(r+1)$$

and  $D_0 = r$ .

If  $r < 4k - 1$  then

$$c(\mathbf{a}, \mathbf{b}) > 3kr = 3kD_0$$

hence we can exclude this case.

If  $r \geq 4k$  then the bound for  $M(6k)$  from Remark 2.3 and the inequality  $5rk - \frac{1}{2}r(r+1) \leq M(6k)$  yield

$$r \leq 5k - 1/2 - \frac{1}{438} \sqrt{799350k^2 - 287766k + 47961 - 31974\sqrt{73}k^2 + 14454\sqrt{73}k}.$$

A straightforward calculation shows that the right hand side is strictly smaller than  $4k$ .

It remains to check the case  $r = 4k - 1$ . Then  $c(\mathbf{a}, \mathbf{b}) = 12k^2 - 3k$ . Now  $c(\mathbf{a}, \mathbf{b}) \leq M(6k)$  implies that  $k < 2$ . Hence the only case that might occur is  $k = 1$ ,  $r = 3$ .  $\square$

**Remark 4.6.** This exceptional case  $k = 1$ ,  $t = r = 3$  does occur: Let  $C$  be the dual of a smooth cubic curve. This is a sextic curve with 9 cusps and no further singularities. Since  $c(\mathbf{a}, \mathbf{b}) \leq 3kD_0$  it follows that  $D_0 \geq 3$ . Hence  $b_i \in \{4, 5\}$  and the  $a_i \in \{3, 4\}$ .

Let  $r$  be the number of  $b_i$  that equals 5, let  $A_4$  be the difference between the number of  $i$  such that  $b_i = 4$  and the number of  $i$  such that  $a_i = 4$ , let  $A_3$  be minus the number of  $i$  such that  $a_i = 3$ . Then we have the following three equalities

$$r + A_4 + A_3 + 1 = 0, \quad 5r + 4A_4 + 3A_3 = 0, \quad 25r + 16A_4 + 9A_3 = 18.$$

These equalities come from the following facts: there is one more  $a_i$  than  $b_i$ ; we have  $\sum a_i = \sum b_i$  and we have  $\sum b_i^2 - \sum a_i^2 = 2c(\mathbf{a}, \mathbf{b})$ .

The only solution to this system of equations is  $r = 3$ ,  $A_4 = -3$ ,  $A_3 = -1$ . In order to actually determine the minimal resolution we need to determine  $t$ . If  $t$  were strictly larger than  $r$  then both  $a_4$

and  $b_4$  equal 4 and in particular the resolution is not minimal. Hence  $t = r = 3$ . From this it follows that  $b_1 = b_2 = b_3 = 5$ ,  $a_1 = a_2 = a_3 = 4$  and  $a_{t+1} = 3$ . Hence the exceptional case  $k = 1$ ,  $r = 3$  of the above lemma does actually occur.

**Proposition 4.7.** Suppose  $(\mathbf{a}', \mathbf{b}')$  is strongly  $k$ -admissible for rank  $2r$  and degree  $D_0$ . Then there exists a strongly  $k$ -admissible sequence  $(\mathbf{a}, \mathbf{b})$  for rank  $2r$  and degree  $D_0$  such that  $c(\mathbf{a}, \mathbf{b}) \leq c(\mathbf{a}', \mathbf{b}')$  and

- (1)  $k = 1$ ,  $r = t = 3$ ,  $a_1 = a_2 = a_3 = 4$ ,  $a_4 = 3$ ;
- (2)  $r = t$ , there exists an integer  $w$  between 0 and  $r - 1$  such that  $a_1 = \dots = a_w = 5k - 1$ ;  $a_w > a_{w+1} \geq a_{w+2}$ ;  $a_{w+2} = a_{w+3} = \dots = a_{r+1} = D_0$  and  $b_1 = \dots = b_r = 5k$ ; or
- (3) there exists an integer  $w$  between 0 and  $r - 1$  such that  $a_1 = \dots = a_w = 5k - 1$ ;  $a_{w+1} = \dots = a_{r+1} = D_0$ ;  $b_1 = \dots = b_r = 5k$  and  $b_i = a_i + 1$  for  $r < i \leq t$ .

**Proof.** We start by setting  $\mathbf{a} := \mathbf{a}'$  and  $\mathbf{b} := \mathbf{b}'$ . We apply a series of modifications to  $\mathbf{a}$ ,  $\mathbf{b}$  in order to end up in one of the three above mentioned forms.

First of all we may reduce  $\mathbf{a}$ ,  $\mathbf{b}$ , i.e. there are no pairs  $i, j$  such that  $a_i = b_j$ . Moreover, from Lemma 4.5 it follows that  $D_0 < 5k - 1$ .

Recall that  $c(\mathbf{a}, \mathbf{b}) = \frac{1}{2}(\sum b_i^2 - \sum a_i^2)$ . We apply several operations on  $\mathbf{a}$ ,  $\mathbf{b}$  that fix  $r$  and  $D_0$ , keep the sequence strongly  $k$ -admissible and reduced and lower the function  $c$ :

- (1) Let  $i$  be the smallest index such that  $a_i < 5k - 1$ , let  $j > i$  such that  $a_j > D_0$ . Assume that  $i < r$ , hence  $b_{i+1} = 5k > a_i + 1$ . Replace in  $\mathbf{a}$ ,  $a_i$  by  $a_i + 1$  and  $a_j$  by  $a_j - 1$ . The new sequence is clearly strongly  $k$ -admissible (here one uses  $i < r$ ) and has a lower value of  $c(\mathbf{a}, \mathbf{b})$  (here one uses  $a_i > a_j$ ).
- (2) If for some  $r < i < t$  we have that  $b_i - b_t \geq 2$  and  $b_i - a_{i-1} \geq 2$  then we can decrease  $b_i$  by one and increase  $b_t$  by one.
- (3) If for some  $r \leq i < t$  we have that  $a_i > D_0$  then we can decrease both  $a_i$  and  $b_{i+1}$ .

It might be that one has to reorder the  $a_i$  and  $b_i$  after applying one of the above operations or that one has to reduce the sequence.

**Step 1: Adjust  $\mathbf{a}$  such that at most one  $a_i$  is different from  $5k - 1$  and  $D_0$ .** Applying the first operation several times brings us in the situation that at most one of the  $a_i$  is different from  $5k - 1$ ,  $D_0$ , or that  $a_{r-1}$  equals  $5k - 1$ . If we are in the latter case and at least two of the  $a_i$  are different from  $5k - 1$ ,  $D_0$  then  $t > r$ . In this case we apply the third operation (combined with reducing and sorting if necessary) until either  $t = r$  or  $a_r = D_0$  holds. Hence we are now in the situation that at most one  $a_i$  is different from  $5k - 1$ ,  $D_0$ .

**Step 2: Case  $t = r$ .** If  $t = r$  then all the  $b_i$  equal  $5k$ . From Lemma 4.5 it follows that either at least two of the  $a_i$  are different from  $5k - 1$  or  $k = 1$ ,  $t = r = 3$  holds. Hence we are either in the first or in the second case of the proposition.

**Step 3: Case  $t \neq r$ .** Suppose now that  $t > r$ . Applying the third operation several times brings us in the situation where all the  $a_i$  are either  $5k - 1$  or  $D_0$ .

Suppose now that  $a_r = 5k - 1$ . Let  $i$  be the largest index such that  $a_i = 5k - 1$ , let  $j$  be the largest such that  $b_j = 5k$ . Since  $\mathbf{a}$ ,  $\mathbf{b}$  is strongly  $k$ -admissible we have that  $j > i \geq r$ . Replace in  $\mathbf{a}$ ,  $\mathbf{b}$   $a_i$  by  $D_0$  and  $b_j$  by  $D_0 + 1$ , and sort  $\mathbf{b}$ . Then the new sequence has a lower value of  $c$ . Iterating this allows us to assume that  $b_{r+1} < 5k$  and hence that  $a_r = D_0$ .

Let  $i$  be largest index such that  $b_i \neq D_0 + 1$ . If  $i = r$  then our sequence is of the third form. Suppose now that  $i > r$ . From  $\mathbf{a}$ ,  $\mathbf{b}$  we obtain a new strongly  $k$ -admissible sequence of length  $t + 1$ , by decreasing  $b_i$  by one and by setting  $b_{t+1} = D_0 + 1$ ,  $a_{t+2} = D_0$ . The new sequence has a lower value of  $c$ . Iterating this yields a sequence  $\mathbf{a}$ ,  $\mathbf{b}$  such that  $a_i$  is either  $5k - 1$  or  $D_0$  and  $b_i$  is either  $5k$  or  $D_0 + 1$ , and such that  $a_r = D_0$ .  $\square$

**Remark 4.8.** If  $(\mathbf{a}, \mathbf{b})$  is  $k$ -admissible then  $c(\mathbf{a}, \mathbf{b}) \leq \min(M(6k), 3ka_{t+1})$  holds. Let  $m(d)$  be the smallest integer bigger or equal than  $2M(d)/d$ . Then the above mentioned condition can be rephrased as  $c(\mathbf{a}, \mathbf{b}) \leq 3ka_{t+1}$  if  $a_{t+1} \leq m(6k)$  and  $c(\mathbf{a}, \mathbf{b}) \leq M(6k)$  if  $a_{t+1} \geq m(6k)$ .

**Proposition 4.9.** Suppose that  $\mathbf{a}, \mathbf{b}$  is a strongly  $k$ -admissible sequence of rank  $2r$ . Then  $r < m(6k)$  if  $k > 1$  and  $r \leq m(6) = 3$  if  $k = 1$ .

**Proof.** Suppose we have a strongly  $k$ -admissible sequence  $\mathbf{a}, \mathbf{b}$  with  $r \geq m(6k)$ . Without loss of generality we may assume that  $\mathbf{a}, \mathbf{b}$  is reduced.

If  $D_0 = r$  then by Lemma 4.5 it follows that  $k = 1$  and  $r = 3$ . If  $(k, r) \neq (1, 3)$  then we have that  $D_0 > r$ , hence at least two of the  $a_i$  are different from  $5k - 1$ .

By decreasing  $a_{t+1}$  by one and increasing  $a_t$  by one, we obtain a new sequence, that is again strongly  $k$ -admissible: the value of  $c$  decreases by this operation, and since the new  $D_0$  is still larger or equal than  $m(6k)$  we have that  $\min(M(6k), 3ka_{t+1}) = M(6k)$ . Since the value of  $c$  for the old sequence was already smaller than this quantity the value of  $c$  for this new sequence is that again. If necessary replace  $\mathbf{a}, \mathbf{b}$  by its reduction. By iterating this we end up in the case that  $a_t = 5k - 1$  and  $D_0 = r$ . This is impossible by Lemma 4.5.  $\square$

**Proposition 4.10.** Suppose  $\mathbf{a}, \mathbf{b}$  is a strongly  $k$ -admissible sequence of rank  $2r$ . Then

$$c(\mathbf{a}, \mathbf{b}) \geq \frac{3k}{2}(r - 1 + 2k + \sqrt{-r^2 + 4kr + 1 - 4k + 4k^2}).$$

**Proof.** For fixed  $r, D_0$  the minimum value of  $c(\mathbf{a}, \mathbf{b})$  is attained by a sequence of the form described in Proposition 4.7.

The first case of Proposition 4.7 occurs only for  $(k, r) = (1, 3)$ . In this case  $c(\mathbf{a}, \mathbf{b}) = 9$  holds and we have

$$c(\mathbf{a}, \mathbf{b}) = 9 = \frac{3k}{2}(r - 1 + 2k + \sqrt{-r^2 + 4kr + 1 - 4k + 4k^2}).$$

In the second and third case we will vary  $D_0$  and the additional parameter  $w$  to determine the minimum value of  $c(\mathbf{a}, \mathbf{b})$  for fixed  $r$ .

Consider now sequences of the form (2).

Suppose first that  $D_0 > \frac{1}{2}(5k + r - 1)$ . Then  $w < r - 1$ . Set  $w' = (5rk - rD_0 - D_0)/(5k - 1 - D_0)$ . From

$$B(s - a_{w+1}) \leq (w - w')B(s - 5k + 1) + (1 - w + w')B(s - D_0)$$

it follows that

$$\begin{aligned} c(\mathbf{a}, \mathbf{b}) &= rB(s - 5k) + B(s) - wB(s - 5k + 1) - B(s - a_{w+1}) - (r - w)B(s - D_0) \\ &\geq rB(s - 5k) + B(s) - w'B(s - 5k + 1) - (r + 1 - w')B(s - D_0) \\ &= \frac{1}{2}(5kD_0 + 5rk - rD_0 - D_0). \end{aligned}$$

The right hand side is decreasing in  $D_0$ . Moreover, the right hand side is precisely  $c(\mathbf{a}, \mathbf{b})$  for  $D_0 = \frac{1}{2}(5k + r - 1)$ . Hence the minimal value for  $c(\mathbf{a}, \mathbf{b})$  is attained at  $D_0 = \frac{1}{2}(5k + r - 1)$ . For this value one has that

$$c(\mathbf{a}, \mathbf{b}) = \frac{1}{4}(1 - 10k + 10rk + 25k^2 - r^2)$$

which is smaller than

$$3kD_0 = \frac{3}{2}k(r + 5k - 1).$$

Hence if for any  $D_0 > \frac{1}{2}(5k+r-1)$  there exists a strongly admissible sequence of type (2) then there exists one with  $D_0 = \lceil \frac{1}{2}(5k+r-1) \rceil$ .

Suppose now that  $D_0 < \lceil \frac{1}{2}(5k+r-1) \rceil$  and hence that  $w = r-1$  holds. Then

$$c(\mathbf{a}, \mathbf{b}) = \frac{1}{2}r(1-r) - D_0 + 5kD_0 - D_0^2 + D_0r.$$

We look now for the smallest  $D_0$  such that  $c(\mathbf{a}, \mathbf{b}) \leq 3kD_0$ . Call this value  $D_{0,\min}$ . For  $D_0 < D_{0,\min}$  our sequence  $\mathbf{a}, \mathbf{b}$  is not  $k$ -admissible, for  $D_0 > D_{0,\min}$  we find a higher value of  $c(\mathbf{a}, \mathbf{b})$ . Solving

$$3kD_0 = \frac{1}{2}r(1-r) - D_0 + 5kD_0 - D_0^2 + D_0r$$

yields

$$D_{0,\min} = \frac{1}{2}(2k-1+r+\sqrt{4k^2+4kr-r^2-4k+1}).$$

One easily checks that  $D_{0,\min} < \frac{1}{2}(5k+r-1)$ . Hence the minimal possible value of  $c(\mathbf{a}, \mathbf{b})$  is then  $3kD_{0,\min}$  which is precisely the bound mentioned in the statement.

Consider now case (3) of Proposition 4.7. We have  $b_1 = \dots = b_r = 5k$ ,  $b_{r+1} = \dots = b_t = D_0 + 1$ ,  $a_1 = \dots = a_w = 5k-1$  and  $a_{w+1} = \dots = a_{t+1} = D_0$ ,  $w \leq r < D_0$ . In order to have  $\sum a_i = \sum b_i$  we need

$$t = (5k-1-D_0)(w-r) + D_0.$$

Note that  $t$  is increasing in  $w$  and the function  $2c(\mathbf{a}, \mathbf{b})$  equals

$$(D_0+1-5k)(5k-2-D_0)w + D_0 - 5rk + D_0^2 + 25rk^2 - 10rkD_0 + D_0r + D_0^2r.$$

This function is either constant or is decreasing in  $w$ , hence we may take  $w$  as large as possible, namely  $w = r-1$ , and therefore

$$t = 1 - 5k + 2D_0.$$

Note that  $D_0 \geq \frac{1}{2}(r+5k-1)$ . Differentiating  $c$  with respect to  $D_0$  yields that  $c$  increases as function of  $D_0$  for  $D_0 \geq \frac{1}{2}(r+5k-1)$ . Hence the minimal value for  $c$  is attained at  $D_0 = \frac{1}{2}(r+5k-1)$ . In that case we have

$$c(\mathbf{a}, \mathbf{b}) = \frac{1}{4}(25k^2 + 2rk - 10k - r^2 + 1).$$

A straightforward calculation shows that  $3kD_{0,\min} - \frac{1}{4}(25k^2 + 2rk - 10k - r^2 + 1)$  has a maximum in  $r = 2k$ , and that for  $r = 2k$  this function is negative. Hence for each  $D_0$  we have that  $c(\mathbf{a}, \mathbf{b}) \geq 3kD_{0,\min}$  which finishes the proof.  $\square$

**Remark 4.11.** For  $r = 1$  we find that  $C$  has at least  $6k^2$  cusps. If we expand the right hand side of the inequality as a function for  $k \rightarrow \infty$  we obtain

$$\frac{3k}{2}(r-1+2k+\sqrt{-r^2+4kr+1-4k+4k^2}) = 6k^2 + 3(r-1)k + \frac{3}{4}r(1-r) + O\left(\frac{1}{k}\right).$$

**Remark 4.12.** We can specialize to the case  $k = 1$ . Then we find that to have  $r = 1$  one needs at least 6 cusps, to have  $r = 2$  one needs at least 8 cusps, to have  $r = 3$  one needs at least 9 cusps and for  $r > 3$  one would need at least 10 cusps. Since a sextic has at most 9 cusps, this is not possible. The above mentioned bounds for the minimal number of cusps are sharp, see [9, Theorem 9.2].

**Corollary 4.13.** *There is a function  $g : \mathbf{Z}_{>0} \rightarrow \mathbf{Z}_{>0}$  such that for each  $r > g(k)$  there does not exist a strongly  $k$ -admissible sequence  $\mathbf{a}, \mathbf{b}$  of rank  $2r$ . Moreover one has for  $k \geq 2$  that*

$$g(k) \leq \frac{1}{2628}(-219 - 33\sqrt{73} + (9125 + 73\sqrt{73})k - \sqrt{\alpha}),$$

where

$$\alpha = 3\,325\,734 - 14\,454\sqrt{73} + (-11\,766\,432 + 287\,328\sqrt{73})k + (12\,267\,358 - 564\,874\sqrt{73})k^2.$$

In particular,

$$\limsup_{k \rightarrow \infty} \frac{g(k)}{k} \leq \frac{1}{36}(125 + \sqrt{73} - \sqrt{2302 - 106\sqrt{73}}) \approx 2.67.$$

**Proof.** We know that for fixed  $r$  and  $k$  we have by the previous proposition that

$$c(\mathbf{a}, \mathbf{b}) \geq \frac{3k}{2}(r - 1 + 2k + \sqrt{-r^2 + 4kr + 1 - 4k + 4k^2}).$$

If  $\mathbf{a}, \mathbf{b}$  is  $k$ -admissible then

$$c(\mathbf{a}, \mathbf{b}) \leq M(k) \leq \frac{125 + \sqrt{73}}{12}k^2 - \frac{511 + 11\sqrt{73}}{292}k.$$

Combining both inequalities yields that

$$r \leq \frac{1}{2628}(-219 - 33\sqrt{73} + (9125 + 73\sqrt{73})k - \sqrt{\alpha})$$

or

$$r \geq \frac{1}{2628}(-219 - 33\sqrt{73} + (9125 + 73\sqrt{73})k + \sqrt{\alpha}).$$

In the latter case  $r \geq 4k - 1 \geq m(6k)$ . This case is excluded by Proposition 4.9.  $\square$

## 5. Admissible case

We consider now the case where the resolution of  $I$ , the ideal of the cusps, yields a sequence  $\mathbf{a}', \mathbf{b}'$  that is  $k$ -admissible but not strongly  $k$ -admissible. We show that in this case there is a curve of small degree containing many of the cusps. This yields additional numerical constraints on  $\mathbf{a}', \mathbf{b}'$ . We will use these extra constraints to construct a strongly  $k$ -admissible sequence  $\mathbf{a}, \mathbf{b}$  for the same rank such that  $c(\mathbf{a}, \mathbf{b}) \leq c(\mathbf{a}', \mathbf{b}')$ :

**Proposition 5.1.** *Suppose  $\mathbf{a}', \mathbf{b}'$  form a sequence coming from the resolution of the ideal of cusps of a cuspidal curve for rank  $2r$ . Suppose that  $\mathbf{a}', \mathbf{b}'$  is not strongly  $k$ -admissible. Then there exists a strongly  $k$ -admissible sequence  $\mathbf{a}, \mathbf{b}$  for rank  $2r$  such that  $c(\mathbf{a}, \mathbf{b}) \leq c(\mathbf{a}', \mathbf{b}')$ .*

**Proof.** Set  $\mathbf{a} = \mathbf{a}'$  and  $\mathbf{b} = \mathbf{b}'$ . We are going to modify  $\mathbf{a}$  and  $\mathbf{b}$  such that they become strongly admissible.

**Step 1: Set-up and goal.** To study non-strongly  $k$ -admissible sequence we need to introduce some further notation. Without loss of generality we may assume that  $\mathbf{a}, \mathbf{b}$  is reduced, in particular  $a_i \neq b_{i+1}$ .

Since the  $a_i, b_i$  are not strongly  $k$ -admissible, there is an  $i$  such that  $a_i > b_{i+1}$ . Let  $i_0$  be the smallest index such that  $a_{i_0} > b_{i_0+1}$  holds. Let  $A = a_{i_0}$  and  $D_2 = \sum_{i=1}^{i_0} b_i - a_i$ .

Define

$$h_1 := \sum_{i=1}^{i_0} B(s - b_i) - B(s - a_i)$$

and

$$h_2 := B(s) - B(s - a_{t+1}) + \sum_{i=i_0+1}^t B(s - b_i) - B(s - a_i).$$

Then  $h_1 + h_2$  is the Hilbert polynomial of  $I$ . In particular,  $\# \Sigma = h_1 + h_2$ .

Consider now the ideal  $I'$  generated by the elements of  $I$  of degree strictly less than  $A$ . Let  $f_1$  be the greatest common divisor of the elements of  $I'$ . Let  $I''$  be the ideal generated by the elements of  $I'$  divided by  $f_1$ . Let  $D_1 := \deg(f_1)$ .

Now  $I''$  defines a zero-dimensional scheme, which is possibly non-reduced, moreover  $I''$  might be not saturated. If  $I''$  were saturated then its resolution has length 1, but otherwise the resolution might be of length 2. From this it follows that  $I'$  has a resolution of length at most 2. Hence there might exist an  $s_0 \in \mathbb{Z}_{\geq 0}$  and  $c_j, d_j \in \mathbb{Z}$  for  $j = 1, \dots, s_0$  such that  $A \leq c_j < d_j$  and

$$0 \rightarrow \bigoplus_{j=1}^{s_0} S(-d_j) \rightarrow \bigoplus_{j=1}^{s_0} S(-c_j) \oplus \bigoplus_{i>i_0} S(-b_i) \rightarrow \bigoplus_{i>i_0} S(-a_i) \rightarrow S \rightarrow S/I'$$

is a minimal resolution of  $I'$ . Let  $h_3 = \sum_{j=1}^{s_0} (B(s - c_j) - B(s - d_j))$ .

With this notation we have that the Hilbert polynomial of  $I$  equals  $h_1 + h_2$ , the Hilbert polynomial of  $I'$  equals  $h_2 + h_3$  and the Hilbert polynomial of  $I''$  equals  $h_2 + h_3 - h_C$ , with  $h_C(s) = D_1 s - \frac{1}{2} D_1 (D_1 - 3)$ , the Hilbert polynomial of the ideal  $(f_1)$ .

We will find some additional restrictions.

Note that the Hilbert polynomial of  $I''$  is constant. The coefficient of  $s$  in  $h_2(s) + h_3(s)$  equals  $D_2 + \sum d_j - c_j$ . Hence  $D_1 = D_2 + \sum (d_j - c_j)$ . Since  $d_j > c_j$  it follows that  $D_2 \leq D_1$ . Using  $i_0 \geq r$  and  $D_2 = \sum_{i=1}^{i_0} (b_i - a_i)$  we obtain that  $D_2 \geq r$ .

Suppose the curve  $f_1 = 0$  contains more than  $3kD_1$  cusps of  $C$ . Then  $C$  and  $Z(f_1)$  have a common component. Using the same reasoning as in Proposition 2.4 one can show that the common component has non-positive degree. From this it follows that  $Z(f_0)$  contains at most  $3kD_1$  points of  $\Sigma$ . Therefore  $Z(I'')$  contains at least  $c(\mathbf{a}, \mathbf{b}) - 3kD_1$  points of  $\Sigma$  and

$$3kD_1 \geq \#Z(I) - \#Z(I'') \geq h_1 + h_2 - h_2 - h_3 + h_C = h_1 - h_3 + h_C.$$

Note that up to degree  $A - 1$  the generators and syzygies of  $I$  and  $I'$  agree. Hence  $h_I(A - 1) = h_{I'}(A - 1)$ . Now  $h_I(A - 1) = h_2(A - 1)$  and  $h_{I'} = h_{I''} + h_C$ . From this we get

$$0 \leq h_{I''}(A - 1) = h_{I'}(A - 1) - h_C(A - 1) = h_2(A - 1) - h_C(A - 1).$$

Summarizing we found sequences **a**, **b**, **c**, **d** and integers  $i_0, D_0, D_1, D_2, A$  such that (with  $h_1, h_2, h_3, h_C$  as above):

- (1)  $a_i \geq A$  for  $i = 1, \dots, i_0$ .
- (2)  $b_i < A$  for  $i = i_0 + 1, \dots, t$ .
- (3)  $D_0 \leq a_i < b_i \leq 5k$  for  $i = 1, \dots, t$ .
- (4)  $b_i = 5k$  for  $i = 1, \dots, r$ .
- (5)  $D_0 = a_{t+1} = \sum_{i=1}^t (b_i - a_i)$ .
- (6)  $A \leq c_j < d_j$  for  $j = 1, \dots, s_0$ .
- (7)  $a_i < b_{i+1}$  for  $i = 1, \dots, i_0 - 1$ .
- (8)  $D_2 = \sum_{i=1}^{i_0} (b_i - a_i)$ .
- (9)  $r \leq D_2 \leq D_1 \leq D_0 \leq A \leq 5k - 1$ .
- (10)  $h_1 + h_2 \leq 3kD_0$ .
- (11)  $h_1 + h_2 \leq \frac{1}{4}(45k^2 - 9k)$ .
- (12)  $h_1 - h_3 + h_C \leq 3kD_1$ .
- (13)  $h_2(A - 1) \geq h_C(A - 1)$ .

We want to show that for given  $r$ , a sequence **a**, **b**, **c**, **d** and integers  $D_1, D_2, A$  satisfying the above conditions there exists a sequence **a'**, **b'** with same rank  $r$ , but that is strongly  $k$ -admissible and  $c(\mathbf{a}, \mathbf{b}) \geq c(\mathbf{a}', \mathbf{b}')$ . We do this by changing the above mentioned parameters in such a way that  $c(\mathbf{a}, \mathbf{b})$  decreases and such that in the end we have either  $D_0 = A$  or  $D_0 = D_1 = D_2$  holds. In the former case we clearly have a strongly  $k$ -admissible sequence. In the latter case we use that  $D_0 = D_2$  implies  $i_0 = t$ , hence  $a_i < b_{i+1}$  for  $i = 1, \dots, i_0 - 1 = t - 1$ , which in turn implies that the sequence is strongly  $k$ -admissible.

**Step 2: Optimization of  $h_1, h_2, h_3$  without changing  $D_0, D_1, D_2, A$  and  $r$ .** We first optimize our **a**, **b**, **c**, **d** without changing  $D_0, D_1, D_2, A$  and  $r$ . Specifically, we aim at decreasing the constant coefficients of  $h_1$  and  $h_2$  and at increasing the constant coefficient of  $h_3$ . Hence at this stage we only have to consider the conditions (1)–(8).

The sequence  $a_1, \dots, a_{i_0}, D_1; b_1, \dots, b_{i_0}$  is a strongly  $k$ -admissible sequence. This means that we can apply the same transformations as in the proof of Proposition 4.7, only that we need to impose  $a_i \geq A$  for  $i = 1, \dots, i_0$ .

If for some  $i < j \leq i_0$  we have  $5k - 1 > a_i \geq a_j > A$ , then we can increase  $a_i$  by one and decrease  $a_j$  by one then reduce and sort. The new sequence still satisfies the above mentioned conditions, but  $c(\mathbf{a}, \mathbf{b})$  decreases. So  $a_i \in \{5k - 1, A\}$  for all but at most one  $i \leq i_0$ .

Suppose  $i_0 > r$  and for some  $i < i_0$  we have that  $A < a_i < 5k - 1$ . Let  $j$  be such that  $b_j \neq 5k$ . If  $b_j < a_i$  then we can increase both  $a_i$  and  $b_j$  by  $5k - 1 - a_i$ , and hence all the  $a_i \in \{5k - 1, A\}$ . If  $b_j > a_i$  then we can lower them both with  $a_i - A$  and sort the  $b_i$  if necessary. From this it follows that  $a_i \in \{5k - 1, A\}$  for all  $i$ .

Suppose  $i_0 > r$  and  $b_{i_0} > A + 1$ , then we lower  $b_{i_0}$  by one and increase the length of our original sequence **a**, **b** by adding  $A$  to **a** and adding  $A + 1$  to **b**. Now sort and reduce.

The optimization of  $h_1$  allows us to assume for  $i \leq i_0$  that either

- (1)  $i_0 = r, b_i = 5k, i = 1, \dots, r$  and there exists a  $w \leq r - 1$  such that  $a_1 = \dots = a_w = 5k - 1 > a_{w+1} \geq a_{w+2} = \dots = a_r = A$ . (If  $w = r - 1$  then  $a_{w+1} = A$ . In this case we might disregard  $a_{w+2}$ , since  $w + 2 > i_0$ .)
- (2)  $i_0 > r, b_i = 5k, i = 1, \dots, r, b_i = A + 1$  for  $i = r + 1, \dots, i_0$  and there exists a  $w \leq r - 1$  such that  $a_1 = \dots = a_w = 5k - 1, a_{w+1} = \dots = a_{i_0} = A$ .

This description of **a**, **b** implies  $D_2 \geq 5k - A + r - 1$  or, equivalently,  $A \geq 5k + r - 1 - D_2$ .

Suppose we are in case (1). Then  $A$  and  $D_2$  determine both  $w$  and  $a_{w+1}$  hence the function  $h_1$  depends only on  $A$  and  $D_2$ . Denote this function by  $h_{1,a}(A, D_2)$ . If we are in case (2) then  $A, D_2$  and  $w$  determine  $i_0$ . Hence we have a function  $h_{1,b}(A, D_2, w)$ . Now  $h_{1,b}$  is decreasing in  $w$ , hence we may assume that  $w = r - 1$ . Since



$$h_{1,b}(A, D_2, r-1) \leq h_{1,a}(A, D_2) + \frac{1}{2}(A+2-5k)(A-5k-r+d_1+1)$$

we have that the constant polynomial  $h_{1,b} - h_{1,a}$  is negative. Hence we may assume that  $h_1 = h_{1,b}(A, D_2, r-1)$ , i.e.,

$$h_1(s) = -D_2s - r + 1 + 5kr - rA - D_2 + D_2A - \frac{15}{2}k + \frac{3}{2}A + \frac{1}{2}A^2 + \frac{25}{2}k^2 - 5kA.$$

The optimization of  $h_3$  is relatively easy. First lowering  $c_j$  and  $d_j$  simultaneously increases the constant coefficient of  $h_3$ , so we may assume that  $c_j = A$  for all  $j$ . Suppose that for some  $j$  we have  $d_j \neq A+1$ . We can lower  $d_j$  by one as follows: we increase the length of  $\mathbf{c}$  by one by setting  $c_{s_0+1} = A$ . We increase the length of  $\mathbf{d}$  by setting  $d_{s_0+1} = A+1$ , and decreasing  $d_j$  by one. Then the constant coefficient of  $h_3$  increases under this operation. This allows us to assume that  $d_j = A+1$  and

$$h_3 = (D_1 - D_2)s + (D_1 - D_2)(1 - A).$$

We can optimize  $h_2$  as follows. If for some  $j > i_0$  we decrease  $b_j$  and  $a_j$  simultaneously by one then the constant coefficient of  $h_2$  decreases. If we extend  $\mathbf{a}, \mathbf{b}$  by setting  $a_{t+2} = D_0$ ,  $b_{t+1} = D_0 + 1$ , and lowering one of the  $b_j$  for some  $j > i_0$  then  $c(\mathbf{a}, \mathbf{b})$  decreases. However, we have to stop as soon as  $h_2(A-1) = h_C(A-1)$ . I.e., this allows us to assume that  $h_2 = \max(h_{2,a}, h_{2,b})$  with

- $h_{2,a} = B(s) - B(s - D_0) + (D_0 - D_2)(B(s - D_0 - 1) - B(s - D_0)) = D_2s + \frac{1}{2}D_0^2 + \frac{1}{2}D_0 - D_2(D_0 + 1)$ ,
- $h_{2,b} = D_2s + \frac{1}{2}(D_1 - D_1^2) + D_1A - D_2(A - 1)$ .

In the next step we are going to vary  $A, D_0, D_1, D_2$  in such a way that if we start with a  $k$ -admissible sequence, not strongly  $k$ -admissible and satisfying the conditions (1)–(13) then the new sequence is still  $k$ -admissible, satisfies (1)–(13), but might have a lower value of  $c(\mathbf{a}, \mathbf{b}) = h_1 + \max(h_{2,a}, h_{2,b})$ . It turns out that we end up either in the case  $A = D_0$  or  $D_2 = D_0$ .

The only remaining variables are  $A, D_0, D_1, D_2$ , hence we may disregard the conditions (1)–(8). Also we want to minimize  $h_1 + h_2$ , hence condition (11) is a priori fulfilled. By replacing  $h_2$  by  $\max(h_{2,a}, h_{2,b})$  we forced condition (13) to hold. Hence we need only to consider the conditions (9), (10) and (12). We consider (9) as describing a domain in which the parameters  $A, D_0, D_1, D_2$  may vary, and try to minimize  $h_1 + h_2$  in such a way that (10) and (12) hold.

**Step 3: Elimination of  $A$ .** Note that the main conditions we are considering are

$$h_1 + h_{2,a} \leq 3kD_0, h_1 + h_{2,b} \leq 3kD_0, h_1 + h_C - h_3 \leq 3kD_1.$$

Since  $h_{2,b} + h_3 - h_C$  is a constant polynomial,  $h_3(A-1) = 0$  and  $h_{2,b}(A-1) = h_C(A-1)$  it follows that  $h_{2,b} = h_C - h_3$ . So the left hand side of the second and third inequality agree. Since  $D_0 > D_1$  we might ignore the second inequality.

Both  $h_1 + h_{2,a}$  and  $h_1 + h_{2,b}$  are increasing as a function of  $A$ , for  $A \geq 5k + r - 1 - D_2$ . Hence we might take  $A$  as small as possible, which means that either  $A = D_0$  or  $A = 5k + r - 1 - D_2$ . In the first case we are done. So from now on we assume that

$$A = 5k + r - 1 - D_2.$$

**Step 4: Elimination of  $D_1$ .** Substituting  $A = 5k + r - 1 - D_2$  yields new functions  $h_1, h_{2,a}, h_{2,b}$  and  $h_3$ . The function  $h_1 + h_{2,a}$  increases with  $D_0$ , whereas  $h_1 + h_{2,b}$  is independent of  $D_0$ . So we might decrease  $D_0$  until one of the following three cases occurs:

$$D_0 = D_1, \quad h_{2,b} = h_{2,a} \quad \text{or} \quad h_1 + h_{2,a} = 3kD_0.$$

We claim now that even in the second and in the third case we may also assume that  $D_0 = D_1$  holds.

If  $D_0$  is such that  $h_{2,b} = h_{2,a}$  then the only interesting inequalities are

$$h_1 + h_{2,b} \leq \min\left(3kD_0, \frac{1}{4}(45k^2 - 9k)\right), \quad h_1 + h_{2,b} \leq 3kD_1.$$

Now  $h_1$  and  $h_{2,b}$  are independent of  $D_0$ . Since  $D_1 < D_0$  we may simplify these bounds to  $h_1 + h_{2,b} \leq \min(3kD_1, 45k^2 - 9k)$ , which is completely independent of  $D_0$ . Hence we may decrease  $D_0$  such that  $D_1 = D_0$  since  $c(\mathbf{a}, \mathbf{b})$  is increasing in  $D_0$  and  $D_0$  is not involved in any of the further bounds except  $D_1 \leq D_0$ .

Suppose now that  $D_0$  is such that  $h_1 + h_{2,a} = 3kD_0$  and  $h_{2,a} > h_{2,b}$ . Then  $3kD_1 - h_1 - h_{2,b}$  is increasing in  $D_1$  if  $D_2 + D_1 \geq r + 2k$ . Now  $D_1 \geq D_2 \geq r$ , hence this condition is automatically satisfied as soon as  $r \geq 2k$ .

If  $D_1 + D_2 \geq r + 2k$  then this implies that we may increase  $D_1$  until we reach  $D_1 = D_0$  or  $h_{2,a} = h_{2,b}$ . The latter case we can apply the above argument to obtain  $D_0 = D_1$ . So if  $D_1 + D_2 \geq r + 2k$  we may assume  $D_0 = D_1$ .

If  $D_1 + D_2 \leq r + 2k$  and  $h_{2,a} > h_{2,b}$  then  $c(\mathbf{a}, \mathbf{b})$  is independent of  $D_1$ . Hence we might decrease  $D_1$ , since this decreases  $h_{2,b} + h_1$  and increases  $h_{2,a} - h_{2,b}$ . Hence we get in the situation that  $D_1 = D_2$ . The new function  $3kD_2 - h_{2,b} + h_1$  decreases with  $D_2$ , whereas  $c(\mathbf{a}, \mathbf{b})$  increases with  $D_2$ . So we may decrease  $D_2 = D_1$  as long as all lower bounds for  $D_2$  are satisfied. However, the only bound for  $D_2$  is  $D_2 \geq r$ . Hence we are in the situation that  $D_1 = D_2 = r$ . From this it follows that  $A = 5k - 1$ . Substituting this in  $h_1 + h_{2,b} \leq 3kD_1$  yields

$$-2kn + (1/2)k + (1/2)k^2 \geq 0.$$

In particular  $r \geq 4k - 1$ . Since  $D_2 + D_1 \geq r + 2k$  implies  $r \leq 2k$  this case does not occur.

**Step 5.** Assume now  $D_0 = D_1$ . Consider the following bounds

$$h_1 + h_{2,a} \leq 3kD_0, \quad h_1 + h_{2,b} \leq 3kD_0 \quad \text{and} \quad h_1 + h_c - h_3 \leq 3kD_1.$$

The second and third inequality coincide: as remarked before we have that  $h_{2,b} = h_c - h_3$ , moreover we assumed now that  $D_0 = D_1$ .

Suppose that  $h_{2,b} \leq h_{2,a}$ . Then we only need to consider the first inequality  $h_1 + h_{2,a} \leq 3kD_0$ . Now  $h_1 + h_{2,a}$  increases with  $D_2$  hence in this case we might decrease  $D_2$  until either  $D_2 = r$  holds, or  $h_{2,a} = h_{2,b}$  holds.

Suppose that  $h_{2,b} \geq h_{2,a}$ . Then we only need to consider the first inequality  $h_1 + h_{2,b} \leq 3kD_0$ . Now  $h_1 + h_{2,b}$  decreases with  $D_2$  hence in this case we might increase  $D_2$  until either  $D_2 = D_1 = D_0$  holds or  $h_{2,a} = h_{2,b}$  holds.

So we have either  $h_{2,a} = h_{2,b}$ ,  $D_2 = r$  or  $D_0 = D_1 = D_2$ . In the latter case we are done.

**Subcase  $D_2 = r$ .** If  $D_2 = r$ , then

$$h_1 + h_{2,b} = 5kD_0 - \frac{1}{2}D_0(D_0 + 1).$$

One of the conditions we need to check is that  $h_1 + h_{2,b} \leq 3kD_0$ . This inequality implies that  $D_0 \geq 4k - 1$ . Now,  $h_1 + h_{2,b}$  is increasing in  $D_0$  for  $D_0 < 5k$ . In particular,

$$h_1 + h_{2,b} \geq 5k(4k - 1) - \frac{1}{2}(4k - 1)4k = 12k^2 - 3k.$$

This contradicts  $h_1 + h_{2,b} \leq \frac{45}{4}k^2 - \frac{9}{4}k$ , unless  $k = 1$  and  $D_0 \geq 3$ . Now if  $k = 1$  and  $D_0 \geq 3$  then  $a_i \in \{3, 4\}$  and  $b_i \in \{4, 5\}$ . An easy computation shows that  $4 - r$  of the  $a_i$  equal 3. In particular,

we have  $b_1 = 5$ ,  $b_2 = 4$ ,  $a_1 = a_2 = a_3 = 3$ ,  $b_1 = b_2 = 5$ ,  $a_1 = 4$ ,  $a_2 = a_3 = 3$  or  $b_1 = b_2 = b_3 = 5$ ,  $a_1 = a_2 = a_3 = 4$ ,  $a_4 = 3$ . These are all strongly  $k$ -admissible.

**Subcase  $h_{2,a} = h_{2,b}$ .** Hence the remaining case is  $h_{2,a} = h_{2,b}$ . Using that  $D_0 = D_1$ ,  $A = 5k + r - 1 - D_2$  it follows that  $h_{2,a} = h_{2,b}$  implies

$$0 = (D_0 - D_2)(D_0 + 1 - r - 5k + D_1).$$

Hence we have either  $D_0 = D_2$  or  $D_0 = r - 1 + 5k - D_2 = A$ . In both cases we are done.  $\square$

**Theorem 5.2.** Suppose  $C = Z(f)$  is a reduced degree  $6k$  curve with only nodes and ordinary cusps as singularities. Then both twice the exponent of  $t^2 - t + 1$  in the Alexander polynomial of  $C$  and the Mordell–Weil rank of the elliptic 3-fold given by  $y^2 = x^3 + f$  are at most

$$\frac{1}{1314}(-219 - 33\sqrt{73} + (9125 + 73\sqrt{73})k - \sqrt{\alpha}),$$

where

$$\alpha = 3\,325\,734 - 14\,454\sqrt{73} + (-11\,766\,432 + 287\,328\sqrt{73})k + (12\,267\,358 - 564\,874\sqrt{73})k^2.$$

**Proof.** Let  $r$  be the Mordell–Weil rank. By Proposition 5.1 there exists a strongly  $k$ -admissible sequence of rank  $r$ . From Corollary 4.13 it follows that  $r$  can be bounded by the above mentioned quantity.  $\square$

## References

- [1] E. Arbarello, M. Cornalba, P.A. Griffiths, J. Harris, *Geometry of Algebraic Curves*, vol. I, Grundlehren Math. Wiss. (Fundamental Principles of Mathematical Sciences), vol. 267, Springer-Verlag, New York, 1985.
- [2] J.I. Cogolludo-Agustín, A. Libgober, Mordell–Weil groups of elliptic threefolds and the Alexander module of plane curves, Preprint, available at arXiv:1008.2018v1, 2010.
- [3] A. Dimca, Betti numbers of hypersurfaces and defects of linear systems, *Duke Math. J.* 60 (1990) 285–298.
- [4] A. Dimca, *Singularities and Topology of Hypersurfaces*, Universitext, Springer-Verlag, New York, 1992.
- [5] D. Eisenbud, *The Geometry of Syzygies*, Grad. Texts in Math., vol. 229, Springer-Verlag, New York, 2005.
- [6] H. Esnault, Fibre de Milnor d'un cône sur une courbe plane singulière, *Invent. Math.* 68 (1982) 477–496.
- [7] A. Hirano, Construction of plane curves with cusps, *Saitama Math. J.* 10 (1992) 21–24.
- [8] K. Hulek, R. Kloosterman, Calculating the Mordell–Weil rank of elliptic threefolds and the cohomology of singular hypersurfaces, *Ann. Inst. Fourier (Grenoble)* 61 (2011) 1133–1179.
- [9] R. Kloosterman, On the classification of rational elliptic threefolds with constant  $j$ -invariant, Preprint, 2008.
- [10] A. Langer, Logarithmic orbifold Euler numbers of surfaces with applications, *Proc. Lond. Math. Soc.* 3 (86) (2003) 358–396.
- [11] A. Libgober, Alexander invariants of plane algebraic curves, in: *Singularities, Part 2*, Arcata, CA, 1981, in: *Proc. Sympos. Pure Math.*, vol. 40, Amer. Math. Soc., Providence, RI, 1983, pp. 135–143.
- [12] R. Miranda, Smooth models for elliptic threefolds, in: *The Birational Geometry of Degenerations*, Cambridge, MA, 1981, in: *Progr. Math.*, vol. 29, Birkhäuser, Boston, MA, 1983, pp. 85–133.
- [13] R. Miranda, *The basic theory of elliptic surfaces*, Dottorato di Ricerca in Matematica, ETS Editrice, Pisa, 1989.
- [14] O. Zariski, On the irregularity of cyclic multiple planes, *Ann. of Math.* (2) 32 (1931) 485–511.