



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



Asymmetry of Ext-groups

Izuru Mori

Department of Mathematics, Faculty of Science, Shizuoka University, Shizuoka 422-8529, Japan

ARTICLE INFO

Article history:

Received 6 January 2009

Available online 16 April 2009

Communicated by Michel Van den Bergh

Dedicated to Luchezar Avramov on his 60th birthday

Keywords:

Ext-groups

AS-Gorenstein algebras

Frobenius Koszul algebras

Noncommutative projective geometry

ABSTRACT

In this paper, we will use techniques of noncommutative projective geometry to construct examples of algebras R over a field k not satisfying the following two types of symmetric behaviors of Ext-groups: **(EE)** For any pair of finitely generated R -modules (M, N) , $\dim_k \text{Ext}_R^i(M, N) < \infty$ for all $i \in \mathbb{N}$ if and only if $\dim_k \text{Ext}_R^i(N, M) < \infty$ for all $i \in \mathbb{N}$. **(ee)** For any pair of finitely generated R -modules (M, N) , $\text{Ext}_R^i(M, N) = 0$ for all $i \gg 0$ if and only if $\text{Ext}_R^i(N, M) = 0$ for all $i \gg 0$. In particular, and contrary to the commutative case, we give a simple example of a noncommutative noetherian Gorenstein (Frobenius) local algebra satisfying **(uac)** (uniform Auslander condition) but not satisfying **(ee)**.

© 2009 Elsevier Inc. All rights reserved.

1. Introduction

1.1. Motivation

Throughout, we fix a base field k . Let R be an algebra over k . We denote by $\text{mod } R$ the category of finitely generated right R -modules. For $M, N \in \text{mod } R$, we defined in [16] the intersection multiplicity of M and N by $M \cdot N := (-1)^{\text{GKdim } R - \text{GKdim } M} \xi(M, N)$ where $\text{GKdim } M$ is the Gelfand–Kirillov dimension of M and

$$\xi(M, N) := \sum_{i \in \mathbb{N}} (-1)^i \dim_k \text{Ext}_R^i(M, N)$$

is the Euler form of M and N . In order for this definition to yield a good intersection theory, we must have the property $M \cdot N = N \cdot M$ for reasonably nice R, M, N . Although, for each i , $\text{Ext}_R^i(M, N)$ and $\text{Ext}_R^i(N, M)$ are very different in general (even if R is commutative), the property $M \cdot N = N \cdot M$

E-mail address: simouri@ipc.shizuoka.ac.jp.

holds in many situations. For example, over a commutative noetherian Gorenstein local algebra, this property is equivalent to the Serre’s vanishing conjecture [11]. Moreover, for a noetherian connected graded algebra R , and finitely generated graded right R -modules M and N , it is often the case that $M \cdot N = N \cdot M$ as long as both $M \cdot N$ and $N \cdot M$ are well defined [9]. So the natural question is to ask whether $M \cdot N$ is well defined if and only if $N \cdot M$ is well defined (in the graded case). Since $M \cdot N$ is well defined if and only if (i) $\dim_k \text{Ext}_R^i(M, N) < \infty$ for all $i \in \mathbb{N}$, and (ii) $\text{Ext}_R^i(M, N) = 0$ for all $i \gg 0$, we ask whether R satisfies the following two types of symmetric behaviors of Ext-groups:

- **(EE)**: for all $M, N \in \text{mod } R$, $\dim_k \text{Ext}_R^i(M, N) < \infty$ for all $i \in \mathbb{N}$ if and only if $\dim_k \text{Ext}_R^i(N, M) < \infty$ for all $i \in \mathbb{N}$.
- **(ee)**: for all $M, N \in \text{mod } R$, $\text{Ext}_R^i(M, N) = 0$ for all $i \gg 0$ if and only if $\text{Ext}_R^i(N, M) = 0$ for all $i \gg 0$.

It is known that every commutative noetherian local algebra satisfies **(EE)** [11, Corollary 3.2]. On the other hand, it is easy to see that if R is a local algebra satisfying **(ee)**, then R must be Gorenstein, so we focus on studying Gorenstein algebras in this paper. Presumably, Avramov and Buchweitz were the first people who studied the condition **(ee)**, and they proved that every commutative complete intersection ring satisfies **(ee)** in [2]. The first example of a noetherian Gorenstein algebra not satisfying **(ee)** was given by Jorgensen and Sega in [6], which is even a commutative Frobenius local algebra. Later, a simpler but noncommutative (Frobenius) algebra not satisfying **(ee)** was given in [4]. (We thank Petter Andreas Bergh for pointing this out.) In this paper, we will use techniques of non-commutative projective geometry to construct noncommutative algebras not satisfying **(EE)** and those not satisfying **(ee)**.

Related to the condition **(ee)**, there is another condition **(uac)** (uniform Auslander condition) on a ring R :

- **(uac)**: there is an integer $d_R \in \mathbb{N}$ such that, for all $M, N \in \text{mod } R$, if $\text{Ext}_R^i(M, N) = 0$ for all $i \gg 0$, then $\text{Ext}_R^i(M, N) = 0$ for all $i > d_R$.

It was shown [5, Theorem 4.1] that every commutative noetherian Gorenstein local ring satisfying **(uac)** satisfies **(ee)**. Contrary to the commutative case, we give a simple example of a noncommutative noetherian Gorenstein (Frobenius) local algebra satisfying **(uac)** but not satisfying **(ee)**. (In fact, we will see that this happens quite frequently.)

Along the way toward the main results, we also prove that, for every FBN (fully bounded noetherian) AS-Gorenstein Koszul algebra, there is a bijection between isomorphism classes of point modules over A and those over A^0 , extending [15, Theorem 6.3].

1.2. Noncommutative projective geometry

In this subsection, we review some of the language of noncommutative projective geometry which will be needed in this paper. We refer to [1] for details. Let A be a graded algebra. We denote by $\text{GrMod } A$ the category of graded right A -modules, and by $\text{grmod } A$ the full subcategory of finitely generated graded right A -modules. Morphisms in $\text{GrMod } A$ are A -module homomorphisms preserving degrees. The category of (finitely generated) graded left A -modules can be identified with $\text{GrMod } A^0$ ($\text{grmod } A^0$) where A^0 is the opposite graded algebra of A .

For a vector space V over k , we denote by V^* the dual vector space of V . For a graded vector space $V \in \text{GrMod } k$, we define the graded vector space dual $V^* \in \text{GrMod } k$ by $(V^*)_i := (V_{-i})^*$ for $i \in \mathbb{Z}$. Moreover, for an integer $n \in \mathbb{Z}$, we define the truncation $V_{\geq n} := \bigoplus_{i \geq n} V_i \in \text{GrMod } k$ and the shift $V(n) \in \text{GrMod } k$ by $V(n)_i := V_{n+i}$ for $i \in \mathbb{Z}$. We say that V is right bounded if $V_{\geq n} = 0$ for some $n \in \mathbb{Z}$. We say that V is locally finite if $\dim_k V_i < \infty$ for all $i \in \mathbb{Z}$. In this case, we define the Hilbert series of V by

$$H_V(t) := \sum_{i \in \mathbb{Z}} (\dim_k V_i) t^i \in \mathbb{Z}[[t, t^{-1}]].$$

For $M, N \in \text{GrMod } A$, we denote by $\text{Hom}_A(M, N) = \text{Hom}_{\text{GrMod } A}(M, N)$ the set of all A -module homomorphisms $f : M \rightarrow N$ preserving degrees, and by $\text{Ext}_A^i(M, N) = \text{Ext}_{\text{GrMod } A}^i(M, N)$ its right derived functors. We further define the graded k -vector space

$$\underline{\text{Ext}}_A^i(M, N) := \bigoplus_{n \in \mathbb{Z}} \text{Ext}_A^i(M, N(n)).$$

Note that if A is right noetherian and $M \in \text{grmod } A$, then it is easy to see that $\underline{\text{Ext}}_A^i(M, N) \cong \text{Ext}_{\text{Mod } A}^i(M, N)$ as vector spaces where we forget graded structures in the left-hand side.

We say that a module $M \in \text{GrMod } A$ is torsion if it is a direct limit of right bounded modules. We denote by $\text{Tors } A$ the full subcategory of $\text{GrMod } A$ consisting of torsion modules, and $\text{Tails } A := \text{GrMod } A / \text{Tors } A$ the quotient category. The natural functor $\pi : \text{GrMod } A \rightarrow \text{Tails } A$ is exact and has a right adjoint $\omega : \text{Tails } A \rightarrow \text{GrMod } A$. For $M \in \text{GrMod } A$, we often write $\mathcal{M} := \pi M \in \text{Tails } A$. The set of morphisms in $\text{Tails } A$ is denoted by $\text{Hom}_{\mathcal{A}}(\mathcal{M}, \mathcal{N}) = \text{Hom}_{\text{Tails } A}(\mathcal{M}, \mathcal{N})$, and we define

$$\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N}) := \bigoplus_{n \in \mathbb{Z}} \text{Ext}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N}(n))$$

as before.

Let A be a connected graded algebra with the unique maximal homogeneous ideal $\mathfrak{m} := A_{\geq 1}$. We will define two notions of Cohen–Macaulay. The local cohomology modules of $M \in \text{GrMod } A$ are defined by

$$\underline{H}_{\mathfrak{m}}^i(M) := \lim_{n \rightarrow \infty} \underline{\text{Ext}}_A^i(A/A_{\geq n}, M) \in \text{GrMod } A.$$

For $M \in \text{GrMod } A$, we define the following numbers:

$$\begin{aligned} j(M) &:= \inf\{i \mid \underline{\text{Ext}}_A^i(M, A) \neq 0\}, \\ \text{depth } M &:= \inf\{i \mid \underline{H}_{\mathfrak{m}}^i(M) \neq 0\}, \\ \text{ldim } M &:= \sup\{i \mid \underline{H}_{\mathfrak{m}}^i(M) \neq 0\}. \end{aligned}$$

We say that A satisfies Cohen–Macaulay property with respect to GK-dimension if

$$j(M) + \text{GKdim } M = \text{GKdim } A < \infty$$

for all $M \in \text{grmod } A$. We say that M is Cohen–Macaulay if $\text{depth } M = \text{ldim } M < \infty$. We say that A satisfies the condition χ if $\underline{H}_{\mathfrak{m}}^i(M)$ are right bounded for all $M \in \text{grmod } A$ and all $i \in \mathbb{N}$.

The following classes of algebras are important in noncommutative projective geometry.

Definition 1.1. Let A be a noetherian connected graded algebra.

- (1) We say that A is AS Cohen–Macaulay if
 - A and A^0 satisfy the condition χ , and
 - A and A^0 are Cohen–Macaulay as graded modules over themselves.
- (2) We say that A is AS-Gorenstein if
 - A is AS Cohen–Macaulay, and
 - $\text{id } A = \text{id } A^0 < \infty$.
- (3) We say that A is a quantum polynomial algebra if
 - A is AS-Gorenstein,
 - $\text{gldim } A = \text{gldim } A^0 = d < \infty$,

- $H_A(t) = (1 - t)^{-d}$, and
- A satisfies Cohen–Macaulay property with respect to GK-dimension.

If A is a noetherian AS Cohen–Macaulay algebra of depth $A = d$, then the graded A - A bimodule $\omega_A := \underline{H}_m^d(A)^*$ is called the balanced dualizing module [10]. It is known that a noetherian AS Cohen–Macaulay algebra is AS-Gorenstein if and only if there are a graded algebra automorphism $\nu \in \text{Aut } A$ called the generalized Nakayama automorphism, and an integer $\ell \in \mathbb{Z}$ called the Gorenstein parameter, such that $\omega_A \cong A_\nu(-\ell)$ as graded A - A bimodules. Note that every artinian connected graded algebra is AS Cohen–Macaulay with the balanced dualizing module $\omega_A = A^*$. It follows that every Frobenius connected graded algebra is AS-Gorenstein and the generalized Nakayama automorphism is the usual Nakayama automorphism in this case (see [15]).

A linear resolution of $M \in \text{grmod } A$ is a free resolution of M of the form

$$\dots \rightarrow A(-3)^{\oplus r_3} \rightarrow A(-2)^{\oplus r_2} \rightarrow A(-1)^{\oplus r_1} \rightarrow A^{\oplus r_0} \rightarrow M \rightarrow 0$$

for some $r_i \in \mathbb{N}$. The full subcategory of $\text{grmod } A$ consisting of modules having linear resolutions is denoted by $\text{lin } A$. We say that A is Koszul if $k := A/\mathfrak{m} \in \text{lin } A$. It is known that A is Koszul if and only if A^0 is Koszul. Every Koszul algebra A is quadratic, that is, $A = T(V)/(R)$ where $R \subset V \otimes_k V$ is a subspace and (R) is the two-sided ideal of $T(V)$ generated by R . A useful fact about a Koszul algebra A is that there is a duality $E_A : \text{lin } A \leftrightarrow \text{lin } A^1 : E_{A^1}$ where A^1 is the quadratic dual of A . It is known that A^1 is also Koszul, called the Koszul dual of A . We refer to [18] for details on Koszul algebras.

For the rest of this paper, we only consider graded algebras finitely generated in degree 1 over a base field k . Such an algebra can be written as $A = T(V)/I$ where V is a finite dimensional vector space over k , $T(V)$ is the tensor algebra of V over k , and I is a homogeneous ideal of $T(V)$. We often fix this presentation of an algebra A .

2. The condition (EE) over AS-Gorenstein algebras

The purpose of this section is to produce a simple example of an algebra not satisfying (EE).

2.1. The condition (PC)

First, we will define the notion of complete point module and the condition (PC).

Definition 2.1. Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. A point module over A is a cyclic module $M \in \text{grmod } A$ such that

$$H_M(t) = \sum_{i=0}^{\infty} t^i = 1 + t + t^2 + t^3 + t^4 + \dots$$

The full subcategory of $\text{grmod } A$ consisting of point modules over A is denoted by $\text{pmod } A$. The point module sequence of a point module $M \in \text{pmod } A$ is a sequence of points $\{p_0, p_1, p_2, p_3, \dots\}$ where $p_n := \mathcal{V}(\{f \in V = A_1 \mid M_n f = 0\}) \subset \mathbb{P}(V^*)$.

A point module $M \in \text{pmod } A$ is called complete if $(\omega\pi M)_{\geq n}(n) \in \text{pmod } A$ for all $n \in \mathbb{Z}$. The complete point module sequence of a complete point module $M \in \text{pmod } A$ is a sequence of points $\{\dots, p_{-2}, p_{-1}, p_0, p_1, p_2, p_3, \dots\}$ where $p_n := \mathcal{V}(\{f \in V = A_1 \mid (\omega\pi M)_n f = 0\}) \subset \mathbb{P}(V^*)$. We say that A satisfies (PC) if every point module is complete.

Every quantum polynomial algebra satisfies (PC) by [15, Proposition 6.6]. (Recall that if A is a quantum polynomial algebra, then every point module over A has a linear resolution, that is, $\text{pmod } A = \text{plin } A := \text{pmod } A \cap \text{lin } A$ [13, Corollary 5.7].) We will find below a larger class of algebras satisfying (PC).

Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. Note that if $M \in \text{pmod } A$, then $M_{\geq n}(n) \in \text{pmod } A$ for all $n \in \mathbb{N}$. For a point $p \in \mathbb{P}(V^*)$, we define a module $M_p := A/p^\perp A \in \text{grmod } A$ where $p^\perp = \{f \in A_1 = V \mid f(p) = 0\}$. By symmetry, we define ${}_p M := A/Ap^\perp \in \text{grmod } A^0$. We say that A is quasi-geometric if every point module over A is isomorphic to M_p for some $p \in \mathbb{P}(V^*)$. In this case, we define the geometric pair $\mathcal{P}(A) = (E, \sigma)$ where $E = \{p \in \mathbb{P}(V^*) \mid M_p \in \text{pmod } A\}$ and $\sigma : E \rightarrow E$ is the map defined by $(M_p)_{\geq 1}(1) \cong M_{\sigma(p)}$ (see [13]).

Lemma 2.2. *Let A be a graded algebra, and $M \in \text{GrMod } A$. For $f \in A_i$, if $M_{j-i} f = 0$, then $f(M_j)^* = 0$.*

Proof. Let $\phi \in (M_j)^*$ so that $\phi : M_j \rightarrow k$ is a linear map. For $f \in A_i$ and $m \in M_{j-i}$, $mf \in M_j$ and $(f\phi)(m) = \phi(mf)$. So if $mf = 0$ for all $m \in M_{j-i}$, then $f\phi = 0$ for all $\phi \in (M_j)^*$. \square

Theorem 2.3. *Let A be a noetherian AS Cohen–Macaulay algebra of depth $A = d$. If*

$$\text{Ext}_A^{d-1}(-, \omega_A)(1) : \text{pmod } A \leftrightarrow \text{pmod } A^0 : \text{Ext}_{A^0}^{d-1}(-, \omega_A)(1)$$

is a duality, then A and A^0 satisfy (PC). In addition, if A and A^0 are quasi-geometric, then $\mathcal{P}(A) = (E, \sigma)$ and $\mathcal{P}(A^0) = (E, \sigma^{-1})$ for some bijection $\sigma : E \rightarrow E$. Moreover, for a point $p \in E$, the complete point module sequence of $M_p \in \text{pmod } A$ is $\{\sigma^n(p)\}_{n \in \mathbb{Z}}$.

Proof. Since A is a noetherian AS Cohen–Macaulay algebra, for $M \in \text{GrMod } A$ and $i \in \mathbb{Z}$, we have $H_m^i(M)^* \cong \text{Ext}_A^{d-i}(M, \omega_A)$ in $\text{GrMod } A^0$ by local duality [21, Theorem 5.1]. By assumption, if $M \in \text{pmod } A$, then $\text{Ext}_A^{d-1}(M, \omega_A)(1) \in \text{pmod } A^0$, so

$$H_{H_m^1(M)}(t) = H_{\text{Ext}_A^{d-1}(M, \omega_A)}(t^{-1}) = t^{-1} H_{\text{Ext}_A^{d-1}(M, \omega_A)(1)}(t^{-1}) = t^{-1} + t^{-2} + t^{-3} + \dots$$

By [1, Proposition 7.2(2)], there is an exact sequence

$$0 \rightarrow H_m^0(M) \cong 0 \rightarrow M \rightarrow \omega\pi M \rightarrow H_m^1(M) \rightarrow 0,$$

so

$$(\omega\pi M)_i \cong \begin{cases} M_i \cong k & \text{if } i \geq 0, \\ H_m^1(M)_i \cong k & \text{if } i < 0. \end{cases}$$

By [1, Proposition 7.2(2)] again, there is an exact sequence

$$0 \rightarrow H_m^0(\omega\pi M) \rightarrow \omega\pi M \rightarrow \omega\pi(\omega\pi M) \cong \omega\pi M \rightarrow H_m^1(\omega\pi M) \rightarrow 0,$$

so $H_m^0(\omega\pi M) = 0$, hence $\omega\pi M$ is torsion-free. It follows that $(\omega\pi M)_{\geq n}$ is cyclic, so $(\omega\pi M)_{\geq n}(n) \in \text{pmod } A$ for all $n \in \mathbb{Z}$, hence A satisfies (PC). By symmetry, A^0 satisfies (PC).

If A is quasi-geometric with $\mathcal{P}(A) = (E, \sigma)$, define a map $\tau : E \rightarrow E$ by $(\omega\pi M_p)_{\geq -1}(-1) = M_{\tau(p)} \in \text{pmod } A$. Since

$$M_{\sigma\tau(p)} \cong (M_{\tau(p)})_{\geq 1}(1) = ((\omega\pi M_p)_{\geq -1}(-1))_{\geq 1}(1) \cong (\omega\pi M_p)_{\geq 0} \cong M_p,$$

$\sigma\tau = \text{id}_E$. Since

$$\begin{aligned} M_{\tau\sigma(p)} &= (\omega\pi M_{\sigma(p)})_{\geq -1}(-1) \cong (\omega\pi((M_p)_{\geq 1}(1)))_{\geq -1}(-1) \\ &\cong (\omega\pi M_p)(1)_{\geq -1}(-1) \cong (\omega\pi M_p)_{\geq 0} \cong M_p, \end{aligned}$$

$\tau\sigma = \text{id}_E$, so $\sigma : E \rightarrow E$ is a bijection and $\tau = \sigma^{-1} : E \rightarrow E$. It follows that, for a point $p \in E$, the complete point module sequence of $M_p \in \text{pmod } A$ is $\{\sigma^n(p)\}_{n \in \mathbb{Z}}$.

By symmetry, if A^0 is quasi-geometric with $\mathcal{P}(A^0) = (E^0, \sigma^0)$, then $\sigma^0 : E^0 \rightarrow E^0$ is a bijection. Define a map $\phi : E \rightarrow E^0$ by $\underline{\text{Ext}}_A^{d-1}(M_p, \omega_A)(1) \cong_{\phi(p)} M \in \text{pmod } A^0$. For $M = M_p \in \text{pmod } A$,

$$\begin{aligned} \phi(p) &= \mathcal{V}(\{f \in V = A_1 \mid f \underline{\text{Ext}}_A^{d-1}(M_p, \omega_A)(1)_0 = 0\}) \\ &= \mathcal{V}(\{f \in V = A_1 \mid f \underline{\text{Ext}}_A^{d-1}(M_p, \omega_A)_1 = 0\}) \\ &= \mathcal{V}(\{f \in V = A_1 \mid f(\underline{H}_m^1(M)_{-1})^* = 0\}) \\ &\subset \mathcal{V}(\{f \in V = A_1 \mid \underline{H}_m^1(M)_{-2}f = 0\}) \\ &= \mathcal{V}(\{f \in V = A_1 \mid (\omega\pi M)_{-2}f = 0\}) \\ &= \sigma^{-2}(p) \end{aligned}$$

by Lemma 2.2, so $E = E^0$ and $\phi = \sigma^{-2} : E \rightarrow E$. A short exact sequence

$$0 \rightarrow M_{\geq 1} \rightarrow M \rightarrow M/M_{\geq 1} \rightarrow 0$$

induces an exact sequence

$$\underline{H}_m^1(M_{\geq 1}) \rightarrow \underline{H}_m^1(M) \rightarrow \underline{H}_m^1(M/M_{\geq 1}) \cong 0,$$

so there is an injection of shifted point modules

$$\begin{aligned} \sigma^{-2}(p)M(-1) &= \phi(p)M(-1) = \underline{\text{Ext}}_A^{d-1}(M, \omega_A) \cong \underline{H}_m^1(M)^* \\ &\rightarrow \underline{H}_m^1(M_{\geq 1})^* \cong \underline{\text{Ext}}_A^{d-1}(M_{\geq 1}, \omega_A) \cong \underline{\text{Ext}}_A^{d-1}(M_{\geq 1}(1), \omega_A)(1) \\ &\cong \underline{\text{Ext}}_A^{d-1}(M_{\sigma(p)}, \omega_A)(1) = \phi_{\sigma(p)}M = \sigma^{-1}(p)M. \end{aligned}$$

It follows that

$$\sigma^0\sigma^{-1}(p)M = \sigma^{-1}(p)M_{\geq 1}(1) \cong_{\sigma^{-2}(p)} M(-1)_{\geq 1}(1) \cong_{\sigma^{-2}(p)} M$$

for all $p \in E$, so $\sigma^0 = \sigma^{-1} : E \rightarrow E$. \square

The following corollary extends [15, Theorem 6.3].

Corollary 2.4. *If A is an AS Cohen–Macaulay Koszul algebra which is a graded quotient algebra of a quantum polynomial algebra, or an FBN (fully bounded noetherian) AS-Gorenstein Koszul algebra, then A and A^0 satisfy (PC), and $\mathcal{P}(A) = (E, \sigma)$ and $\mathcal{P}(A^0) = (E, \sigma^{-1})$ for some bijection $\sigma : E \rightarrow E$. In particular, there is a bijection between isomorphism classes of point modules over A and those over A^0 .*

Proof. If A is an AS Cohen–Macaulay Koszul algebra which is a graded quotient algebra of a quantum polynomial algebra, or an FBN AS-Gorenstein Koszul algebra, then A and A^0 are quasi-geometric by [13, Corollary 5.7, Theorems 5.8 and 3.8], and $\underline{\text{Ext}}_A^{d-1}(-, \omega_A)(1) : \text{pmod } A \leftrightarrow \text{pmod } A^0 : \underline{\text{Ext}}_{A^0}^{d-1}(-, \omega_{A^0})(1)$ is a duality by [15, Lemma 4.6], so the result follows from Theorem 2.3. \square

2.2. The condition (EE)

In this subsection, we will give a geometric way of constructing algebras not satisfying (EE). The basic idea is to reduce the problem in the category $\text{GrMod } A$ to that in the category $\text{Tails } A$. So we first compare Ext -groups in $\text{GrMod } A$ with those in $\text{Tails } A$.

Lemma 2.5. *Let A be a right noetherian locally finite connected graded algebra satisfying χ , and $M, N \in \text{grmod } A$. Then, for each $i \in \mathbb{N}$, $\dim_k \underline{\text{Ext}}_A^i(M, N) < \infty$ if and only if $\underline{\text{Ext}}_A^i(\mathcal{M}, \mathcal{N})$ is right bounded.*

Proof. For each $i \in \mathbb{N}$,

- (1) $\dim_k \underline{\text{Ext}}_A^i(M, N)_n < \infty$ for all $n \in \mathbb{Z}$ by [1, Proposition 3.1(3)],
- (2) $\underline{\text{Ext}}_A^i(M, N)_n = 0$ for all $n \ll 0$ by [1, Proposition 3.1(1)(c)], and
- (3) $\underline{\text{Ext}}_A^i(M, N)_{\geq n} \cong \underline{\text{Ext}}_A^i(\mathcal{M}, \mathcal{N})_{\geq n}$ as graded vector spaces for all $n \gg 0$ by [1, Corollary 7.3(2)],

hence the result. \square

We will see below that Ext -groups in $\text{Tails } A$ are also controlled by geometry. For an abelian category \mathcal{C} , we denote by $\mathcal{D}(\mathcal{C})$ ($\mathcal{D}^b(\mathcal{C})$) the (bounded) derived category of \mathcal{C} . For $X \in \mathcal{D}(\mathcal{C})$ and $n \in \mathbb{Z}$, $X[n] \in \mathcal{D}(\mathcal{C})$ is defined by $(X[n])^i := X^{n+i}$ for $i \in \mathbb{Z}$. The following lemma is well known. We will include the proof for the convenience of the reader.

Lemma 2.6. *Let A be a right noetherian connected graded algebra. For $K \in \text{GrMod } A$, we have an isomorphism of functors*

$$\underline{\text{RHom}}_A(\mathcal{K}, -) \cong \underline{\text{RHom}}_A(K, \text{R}\omega(-)) : \mathcal{D}^b(\text{Tails } A) \rightarrow \mathcal{D}(\text{GrMod } k).$$

Proof. Note that $\text{GrMod } A, \text{Tails } A, \text{GrMod } k$ are all abelian categories having enough injectives. Since $\underline{\text{Hom}}_A(K, -) : \text{GrMod } A \rightarrow \text{GrMod } k$ is a left exact functor, and $\omega : \text{Tails } A \rightarrow \text{GrMod } A$ is a left exact functor preserving injectives by [1, Proposition 7.1(1)],

$$\begin{aligned} \underline{\text{RHom}}_A(\mathcal{K}, -) &\cong \text{R}(\underline{\text{Hom}}_A(K, \omega(-))) = \text{R}(\underline{\text{Hom}}_A(K, -) \circ \omega(-)) \\ &\cong \underline{\text{RHom}}_A(K, -) \circ \text{R}\omega(-) = \underline{\text{RHom}}_A(K, \text{R}\omega(-)) \end{aligned}$$

as functors $\mathcal{D}^b(\text{Tails } A) \rightarrow \mathcal{D}(\text{GrMod } k)$ by [19, Corollary 10.8.3]. \square

Definition 2.7. Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. A cyclic graded module of the form $K = A/fA \in \text{grmod } A$ where $f \in A_1 = V$ is a homogeneous regular element of degree 1 is called a hyperplane module over A (associated to the hyperplane $H = \mathcal{V}(f) \subset \mathbb{P}(V^*)$).

Proposition 2.8. *Let $A = T(V)/I$ be a noetherian AS-Gorenstein algebra satisfying Cohen–Macaulay property with respect to GK-dimension. If $M \in \text{pmod } A$ is a complete point module with the complete point module sequence $\{p_n\}_{n \in \mathbb{Z}}$, and $K = A/fA \in \text{GrMod } A$ is a hyperplane module where $f \in A_1 = V$ is a homogeneous regular element of degree 1, then*

$$\dim_k \underline{\text{Ext}}_A^i(\mathcal{K}, \mathcal{M})_n = \begin{cases} k & \text{if } i = 0, 1 \text{ and } p_n \in \mathcal{V}(f), \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Since A is a noetherian AS-Gorenstein algebra satisfying Cohen–Macaulay property with respect to GK-dimension, $\text{GKdim } A = j(k) + \text{GKdim } k = \text{depth } A$, so $\text{depth } M \leq \text{l dim } M = \text{GKdim } M = 1$

by [12, Remark 2.4]. By [1, Proposition 7.2(2)], $h^i(\mathbf{R}\omega\mathcal{M}) \cong \underline{H}_m^{i+1}(M) = 0$ for all $i \geq 1$, so $\mathbf{R}\omega\mathcal{M} \cong h^0(\mathbf{R}\omega\mathcal{M}) \cong \omega\mathcal{M}$ in $\mathcal{D}(\text{GrMod } A)$. By Lemma 2.6, $\underline{\mathbf{RHom}}_A(\mathcal{K}, \mathcal{M}) \cong \underline{\mathbf{RHom}}_A(K, \mathbf{R}\omega\mathcal{M}) \cong \underline{\mathbf{RHom}}_A(K, \omega\mathcal{M})$. Consider the long exact sequence

$$\begin{aligned} \cdots \rightarrow \underline{\text{Ext}}_A^{i-1}(K, \omega\mathcal{M}/(\omega\mathcal{M})_{\geq n}) &\rightarrow \underline{\text{Ext}}_A^i(K, (\omega\mathcal{M})_{\geq n}) \\ &\rightarrow \underline{\text{Ext}}_A^i(K, \omega\mathcal{M}) \rightarrow \underline{\text{Ext}}_A^i(K, \omega\mathcal{M}/(\omega\mathcal{M})_{\geq n}) \rightarrow \cdots \end{aligned}$$

Since $\underline{\text{Ext}}_A^i(K, \omega\mathcal{M}/(\omega\mathcal{M})_{\geq n})_{\geq n} = 0$ for all $i \in \mathbb{N}$ and all $n \in \mathbb{Z}$ by [1, Proposition 3.1(2)], $\underline{\text{Ext}}_A^i(K, (\omega\mathcal{M})_{\geq n})_{\geq n} \cong \underline{\text{Ext}}_A^i(K, \omega\mathcal{M})_{\geq n}$, so

$$\underline{\text{Ext}}_A^i(\mathcal{K}, \mathcal{M})_n \cong \underline{\text{Ext}}_A^i(K, \omega\mathcal{M})_n \cong \underline{\text{Ext}}_A^i(K, (\omega\mathcal{M})_{\geq n})_n \cong \underline{\text{Ext}}_A^i(K, (\omega\mathcal{M})_{\geq n}(n))_0$$

for all $i \in \mathbb{N}$ and all $n \in \mathbb{Z}$. Since $(\omega\mathcal{M})_{\geq n}(n)$ is a point module with the point module sequence $\{p_n, p_{n+1}, p_{n+2}, p_{n+3}, \dots\}$, the result follows from the proof of [13, Theorem 6.1]. \square

Next, we will see below that an asymmetric behavior of Ext in the category $\text{Tails } A$ is controlled by the generalized Nakayama automorphism.

Definition 2.9. We say that an abelian category \mathcal{C} has finite global dimension if there is an integer $n \in \mathbb{N}$ such that $\text{Ext}_{\mathcal{C}}^i(\mathcal{M}, \mathcal{N}) = 0$ for all $\mathcal{M}, \mathcal{N} \in \mathcal{C}$ and all $i > n$.

Example 2.10. The following are examples of graded algebras A such that $\text{Tails } A$ has finite global dimension.

- (1) If A is a commutative graded algebra finitely generated in degree 1 and $X = \text{Proj } A$ is the associated projective scheme, then $\text{Tails } A \cong \text{Mod } X$ where $\text{Mod } X$ is the category of quasi-coherent sheaves on X , so $\text{Tails } A$ has finite global dimension if and only if $\text{Proj } A$ is smooth. So we may think of the condition “ $\text{Tails } A$ has finite global dimension” as the condition “ $\text{Proj } A$ is smooth” even if A is not commutative.
- (2) If A has finite global dimension, then $\text{Tails } A$ has finite global dimension.
- (3) If A is a right noetherian graded algebra generated in degree 1, then $\text{Tails } A^{(r)} \cong \text{Tails } A$ for any $r \in \mathbb{N}^+$ by [1, Proposition 5.10(3)] where $A^{(r)}$ is the r th Veronese subalgebra of A , so if A has finite global dimension, then $\text{Tails } A^{(r)}$ has finite global dimension.
- (4) If A is an FBN AS Cohen–Macaulay algebra of finite CM-representation type, then, for any $M, N \in \text{grmod } A$, $\text{Ext}_A^i(\mathcal{M}, \mathcal{N}) = 0$ for all $i \geq \text{depth } A$ by [7, Theorem 2.5], so $\text{tails } A$ has finite global dimension.

Let A be a graded algebra and $\tau \in \text{Aut } A$ a graded algebra automorphism. For a graded module $M \in \text{GrMod } A$, we define a new graded module $M_\tau \in \text{GrMod } A$ by $M_\tau = M$ as a graded vector space with the new right action $m * a := m\tau(a)$ for $m \in M, a \in A$.

Lemma 2.11. Let A be a noetherian AS-Gorenstein algebra of injective dimension d and Gorenstein parameter ℓ , and let $\nu \in \text{Aut } A$ be the generalized Nakayama automorphism of A . If $\text{Tails } A$ has finite global dimension, then, for $M, N \in \text{grmod } A$ and all $i \in \mathbb{Z}$,

$$\underline{\text{Ext}}_A^i(\mathcal{M}, \mathcal{N}) \cong \underline{\text{Ext}}_A^{d-1-i}(\mathcal{N}, \mathcal{M}_\nu)^*(\ell)$$

as graded vector spaces. In particular, for all $n \in \mathbb{Z}$,

$$\underline{\text{Ext}}_A^i(\mathcal{M}, \mathcal{N})_n \cong \underline{\text{Ext}}_A^{d-1-i}(\mathcal{N}, \mathcal{M}_\nu)_{-n-\ell}$$

as vector spaces.

Proof. Since A is a noetherian AS-Gorenstein algebra, $A_v(-\ell)[d]$ is the balanced dualizing complex for A in the sense of [20]. Since $\text{Tails } A$ has finite global dimension,

$$-\otimes_{\mathcal{A}}^L \mathcal{A}_v(-\ell)[d-1] : \mathcal{D}^b(\text{tails } A) \rightarrow \mathcal{D}^b(\text{tails } A)$$

is the Serre functor by [17, Theorem A.4], so, for $M, N \in \text{grmod } A$,

$$\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N}) \cong \underline{\text{Ext}}_{\mathcal{A}}^{-i}(\mathcal{N}, \mathcal{M} \otimes_{\mathcal{A}}^L \mathcal{A}_v(-\ell)[d-1])^* \cong \underline{\text{Ext}}_{\mathcal{A}}^{d-1-i}(\mathcal{N}, \mathcal{M}_v)^*(\ell)$$

as graded vector spaces. \square

Let $A = T(V)/I$ be a graded algebra. If $\tau \in \text{Aut } A$ is a graded algebra automorphism, then it restricts to an automorphism $\tau : V \rightarrow V$, so its dual induces an automorphism $\tau^* : \mathbb{P}(V^*) \rightarrow \mathbb{P}(V^*)$. If A is quasi-geometric with $\mathcal{P}(A) = (E, \sigma)$, then it is easy to see that τ^* restricts to a bijection $\tau^* : E \rightarrow E$ (see [15]).

Theorem 2.12. *Let $A = T(V)/I$ be a quasi-geometric FBN AS-Gorenstein algebra satisfying (PC) with $\mathcal{P}(A) = (E, \sigma)$ such that $\text{Tails } A$ has finite global dimension, and let $v \in \text{Aut } A$ be the generalized Nakayama automorphism of A . If there are a point $p \in E$ and a hyperplane $\mathcal{V}(f) \subset \mathbb{P}(V^*)$ where $f \in A_1 = V$ is a homogeneous regular element of degree 1 such that $\#\{n \in \mathbb{N} \mid \sigma^n(p) \in \mathcal{V}(f)\} = \infty$ but $\#\{n \in \mathbb{N} \mid \sigma^{-n}v^*(p) \in \mathcal{V}(f)\} < \infty$, then A does not satisfy (EE).*

Proof. By [22, Theorem 3.1(1)], A satisfies Cohen–Macaulay property with respect to GK-dimension. Let $M = M_p$ be a point module and $K = A/fA$ a hyperplane module. Since $\{\sigma^n(p)\}_{n \in \mathbb{Z}}$ is a complete point module sequence of M by Theorem 2.3, if $\#\{n \in \mathbb{N} \mid \sigma^n(p) \in \mathcal{V}(f)\} = \infty$, then, for $i = 0, 1$, $\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{K}, \mathcal{M})_n \neq 0$ for infinitely many $n \in \mathbb{N}$ by Proposition 2.8, so $\dim_k \underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{K}, \mathcal{M}) = \infty$ by Lemma 2.5. On the other hand, since $\{\sigma^n v^*(p)\}_{n \in \mathbb{Z}}$ is a complete point module sequence of M_v by [15, Lemma 3.2(1)] and Theorem 2.3, if $\#\{n \in \mathbb{N} \mid \sigma^{-n}v^*(p) \in \mathcal{V}(f)\} < \infty$, then $\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{K}, \mathcal{M}_v)_{-n} = 0$ for all $i \in \mathbb{N}$ and all $n \gg 0$ by Proposition 2.8. It follows that $\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{K})$ are right bounded for all $i \in \mathbb{N}$ by Lemma 2.11, hence $\dim_k \underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{K}) < \infty$ for all $i \in \mathbb{N}$ by Lemma 2.5. \square

Remark 2.13. In the above arguments, techniques of noncommutative projective geometry play an essential role in the following sense: since $\mathcal{D}^b(\text{grmod } A)$ almost never has Serre functor, an asymmetric behavior of Ext in the category $\text{GrMod } A$ is not controlled by the generalized Nakayama automorphism. In fact, although we know that $\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N})_{\geq n} \cong \underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{M}, \mathcal{N})_{\geq n}$ and $\underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{N}, \mathcal{M}_v)_{\geq n} \cong \underline{\text{Ext}}_{\mathcal{A}}^i(\mathcal{N}, \mathcal{M}_v)_{\geq n}$ for all $n \gg 0$ by [1, Corollary 7.3(2)], $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{M}, \mathcal{N})_{\gg 0}$ and $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{N}, \mathcal{M}_v)_{\gg 0}$ are not related in general, so $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{M}, \mathcal{N})_{\gg 0}$ and $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{N}, \mathcal{M}_v)_{\gg 0}$ are not related at all. What we know is that $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{M}, \mathcal{N})_{\gg 0}$ and $\underline{\text{Ext}}_{\mathcal{A}}^{\bullet}(\mathcal{N}, \mathcal{M}_v)_{\ll 0}$ are related by Lemma 2.11. (We would like to emphasize this point, because some of the readers might think that the above result is trivial by a simple application of the Serre functor.) That is why we must pass through the category $\text{Tails } A$ and use rather special modules, namely, complete point modules in the above arguments.

Corollary 2.14. *Let $A = T(V)/I$ be a quasi-geometric FBN AS-Gorenstein domain satisfying (PC) with $\mathcal{P}(A) = (E, \sigma)$ such that $\text{Tails } A$ has finite global dimension, and let $v \in \text{Aut } A$ be the generalized Nakayama automorphism of A . If*

- k is an infinite field,
- $\dim_k V \geq 3$,
- $|\sigma| < \infty$, and
- there is a point $p \in E$ such that $\sigma^i(p) \neq v^*(p)$ for any $i \in \mathbb{Z}$,

then A does not satisfy (EE).

Proof. Since A is a domain, any homogeneous element $f \in A_1$ of degree 1 is regular. Since k is infinite, $\dim_k V \geq 3$, and the orbit S of $v^*(p)$ under σ is a finite set not containing p , there is a hyperplane $\mathcal{V}(f) \subset \mathbb{P}(V^*)$ such that $p \in \mathcal{V}(f)$ but $\mathcal{V}(f) \cap S = \emptyset$. Since $|\sigma| < \infty$, $\sigma^n(p) \in \mathcal{V}(f)$ for infinitely many $n \in \mathbb{N}$ but $\sigma^{-n}v^*(p) \notin \mathcal{V}(f)$ for any $n \in \mathbb{N}$, hence the result follows from Theorem 2.12. \square

Example 2.15. Let $A = k\langle x, y, z \rangle / (zy - \alpha yz, xz - \beta zx, yx - \gamma xy)$ where $\alpha, \beta, \gamma \in k$ such that $\alpha\beta\gamma \neq 0, 1$. It is well known that $\mathcal{P}(A) = (E, \sigma)$ where $E = \mathcal{V}(xyz) \subset \mathbb{P}(V^*)$ and

$$\begin{aligned} \sigma(0, b, c) &= (0, \alpha b, c), \\ \sigma(a, 0, c) &= (a, 0, \beta c), \\ \sigma(a, b, 0) &= (\gamma a, b, 0). \end{aligned}$$

Since

$$\begin{aligned} (xyz)x &= (\gamma/\beta)x(xyz), \\ (xyz)y &= (\alpha/\gamma)y(xyz), \\ (xyz)z &= (\beta/\alpha)z(xyz) \end{aligned}$$

in A , the bijection $v^* : E \rightarrow E$ induced by the generalized Nakayama automorphism $v \in \text{Aut } A$ of A is given by

$$v^*(a, b, c) = ((\gamma/\beta)a, (\alpha/\gamma)b, (\beta/\alpha)c)$$

by [20, Theorem 7.18]. Suppose that k is an infinite field, and α, β, γ are roots of unity so that $|\sigma| < \infty$. If $\alpha^i \neq \beta\gamma$ for any $i \in \mathbb{Z}$, then $\sigma^i(0, 1, 1) = (0, \alpha^i, 1) \neq (0, \alpha/\gamma, \beta/\alpha) = v^*(0, 1, 1)$ for any $i \in \mathbb{Z}$, so A does not satisfy (EE) by Corollary 2.14. By symmetry, if $\beta^i \neq \gamma\alpha$ for any $i \in \mathbb{Z}$, or $\gamma^i \neq \alpha\beta$ for any $i \in \mathbb{Z}$, then A does not satisfy (EE).

3. The condition (ee) over Frobenius Koszul algebras

The purpose of this section is to produce a simple example of an algebra not satisfying (ee). We will see that there are many analogies between the arguments in this section and the previous one.

3.1. The condition (pc)

First, we will define the notion of complete co-point module and the condition (pc).

Definition 3.1. Let A be a noetherian graded algebra. A graded module $N \in \text{grmod } A$ is called totally reflexive if $\text{Ext}_A^i(N, A) = 0 = \text{Ext}_{A^0}^i(\text{Hom}_A(N, A), A)$ for all $i \geq 1$, and there is a canonical isomorphism $N \cong \text{Hom}_{A^0}(\text{Hom}_A(N, A), A)$ in $\text{grmod } A$.

In other words, $N \in \text{grmod } A$ is totally reflexive if and only if $\text{RHom}_A(N, A) \cong \text{Hom}_A(N, A)$ in $\mathcal{D}^b(\text{grmod } A^0)$ and $\text{RHom}_{A^0}(\text{RHom}_A(N, A), A) \cong N$ in $\mathcal{D}^b(\text{grmod } A)$. Note that if A is Frobenius, then every finitely generated graded module is totally reflexive.

Definition 3.2. Let A be a noetherian graded algebra. An acyclic complex of finitely generated free modules T is called totally acyclic if $\text{Hom}_A(T, A)$ is also acyclic.

For a complex of free modules

$$T : \dots \xrightarrow{\partial^{n-2}} T^{n-1} \xrightarrow{\partial^{n-1}} T^n \xrightarrow{\partial^n} T^{n+1} \xrightarrow{\partial^{n+1}} \dots,$$

we define $\Omega^n T := \text{Coker } \partial^{n-1}$. For $N \in \text{GrMod } A$, we define $\Omega^n N := \Omega^n F$ where F is the minimal free resolution of N .

Lemma 3.3. (See [3, Theorem 3.1].) *Let A be a noetherian graded algebra. A graded module $N \in \text{grmod } A$ is totally reflexive if and only if there is a totally acyclic complex T such that $\Omega^0 T \cong N$.*

We call T in the above lemma a complete resolution of N . A complete resolution can be defined for more general modules (see [3]).

Definition 3.4. Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. A co-point module over A having a linear resolution is a module $N \in \text{grmod } A$ having a linear resolution of the form

$$\dots \xrightarrow{v_3} A(-3) \xrightarrow{v_2} A(-2) \xrightarrow{v_1} A(-1) \xrightarrow{v_0} A \rightarrow N \rightarrow 0$$

where $v_i \in A_1 = V$. The full subcategory of $\text{grmod } A$ consisting of co-point modules over A having linear resolutions is denoted by $\text{clin } A$.

A co-point module $N \in \text{clin } A$ having a linear resolution is called complete if N has a complete resolution of the form

$$T : \dots \xrightarrow{v_2} A(-2) \xrightarrow{v_1} A(-1) \xrightarrow{v_0} A \xrightarrow{v_{-1}} A(1) \xrightarrow{v_{-2}} A(2) \xrightarrow{v_{-3}} \dots$$

where $v_i \in A_1 = V$. The complete co-point module sequence of a complete co-point module N is a sequence of points $\{\dots, p_{-2}, p_{-1}, p_0, p_1, p_2, \dots\}$ where $p_i = [v_i] \in \mathbb{P}(V)$ are the images of $v_i \in V$. We say that A satisfies **(pc)** if every co-point module having a linear resolution is complete.

If A is a Frobenius Koszul algebra such that A^1 is a quantum polynomial algebra, then A satisfies **(pc)** by the proof of [15, Theorem 6.1(1)]. We will find below a larger class of algebras satisfying **(pc)**.

Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. Note that if $N \in \text{clin } A$, then $\Omega^n N(n) \in \text{clin } A$ for all $n \in \mathbb{N}$. For a point $p \in \mathbb{P}(V)$, we define a module $N_p := A/vA \in \text{grmod } A$ where $v \in A_1 = V$ such that $p = [v] \in \mathbb{P}(V)$. By definition, every co-point module over A having a linear resolution is isomorphic to N_p for some $p \in \mathbb{P}(V)$. We define the co-geometric pair $\mathcal{P}^1(A) = (E, \sigma)$ where $E = \{p \in \mathbb{P}(V) \mid N_p \in \text{clin } A\}$ and $\sigma : E \rightarrow E$ is the map defined by $\Omega N_p(1) \cong N_{\sigma(p)}$ (see [13]).

We now prepare two easy lemmas.

Lemma 3.5. *Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1, and $\tau \in \text{Aut } A$ a graded algebra automorphism. If $N \in \text{clin } A$ is a complete co-point module with the complete co-point module sequence $\{\dots, p_{-2}, p_{-1}, p_0, p_1, p_2, \dots\}$, then $N_\tau \in \text{clin } A$ is also a complete co-point module with the complete co-point module sequence $\{\dots, \tau^{-1}(p_{-2}), \tau^{-1}(p_{-1}), \tau^{-1}(p_0), \tau^{-1}(p_1), \tau^{-1}(p_2), \dots\}$.*

Proof. If

$$\dots \xrightarrow{v_2} A(-2) \xrightarrow{v_1} A(-1) \xrightarrow{v_0} A \xrightarrow{v_{-1}} A(1) \xrightarrow{v_{-2}} \dots$$

is a complete resolution of N where $p_i = [v_i] \in \mathbb{P}(V)$, then

$$\dots \xrightarrow{\tau^{-1}(v_2)} A(-2) \xrightarrow{\tau^{-1}(v_1)} A(-1) \xrightarrow{\tau^{-1}(v_0)} A \xrightarrow{\tau^{-1}(v_{-1})} A(1) \xrightarrow{\tau^{-1}(v_{-2})} \dots$$

is a complete resolution of N_τ where $\tau^{-1}(p_i) = [\tau^{-1}(v_i)] \in \mathbb{P}(V)$. \square

Lemma 3.6. *Let A be a graded algebra. If $\text{id } A < \infty$, then every acyclic complex of finitely generated free modules is totally acyclic.*

Proof. Let $d = \text{id } A < \infty$ and T an acyclic complex of finitely generated free modules. Since $T^{\leq d+1-n}[d+1-n]$ is a free resolution of $\Omega^{d+1-n}T$,

$$\begin{aligned} \text{Ext}_A^n(T, A) &\cong \text{Ext}_A^n(T^{\leq d+1-n}, A) \cong \text{Ext}_A^{d+1}(T^{\leq d+1-n}[d+1-n], A) \\ &\cong \text{Ext}_A^{d+1}(\Omega^{d+1-n}T, A) = 0 \end{aligned}$$

for all $n \in \mathbb{Z}$, hence the result. \square

Theorem 3.7. Let A be a noetherian AS-Gorenstein algebra. If $\mathcal{P}^1(A) = (E, \sigma)$ for some bijection $\sigma : E \rightarrow E$, then, for each $p \in E$, $N_p \in \text{cln } A$ is a complete co-point module with the complete co-point module sequence $\{\sigma^n(p)\}_{n \in \mathbb{Z}}$. In particular, A satisfies **(pc)**.

Proof. Since $\sigma : E \rightarrow E$ is a bijection, for every point $p \in E$, $\sigma^n(p) \in E$ for all $n \in \mathbb{Z}$, so

$$T : \dots \xrightarrow{\sigma^2(p)\cdot} A(-2) \xrightarrow{\sigma(p)\cdot} A(-1) \xrightarrow{p\cdot} A \xrightarrow{\sigma^{-1}(p)\cdot} A(1) \xrightarrow{\sigma^{-2}(p)\cdot} A(2) \xrightarrow{\sigma^{-3}(p)\cdot} \dots$$

is an acyclic complex of finitely generated free modules such that $\Omega^0 T \cong N_p$. Since T is totally acyclic by Lemma 3.6, it is a complete resolution of $N_p \in \text{cln } A$. \square

We know so far no example of an algebra such that $\sigma : E \rightarrow E$ is not a bijection, so we hope that a large class of noetherian AS-Gorenstein algebras satisfy **(pc)**. For example, the following corollary provides a class of algebras satisfying **(pc)**.

Corollary 3.8. Let A be a noetherian AS-Gorenstein Koszul algebra. If its Koszul dual $A^!$ is a graded quotient algebra of a quantum polynomial algebra, or an FBN AS-Gorenstein Koszul algebra, then A and A^0 satisfy **(pc)**.

Proof. By [13, Corollary 5.7, Theorems 5.8 and 3.8] and Corollary 2.4, $\mathcal{P}^1(A) = \mathcal{P}(A^!) = (E, \sigma)$ for some bijection $\sigma : E \rightarrow E$, so A satisfies **(pc)** by Theorem 3.7. By symmetry, A^0 satisfies **(pc)**. \square

The following are characterizations of algebras satisfying **(pc)**.

Theorem 3.9. Let A be a noetherian AS-Gorenstein algebra. Then the following are equivalent:

- (1) Every co-point module over A and A^0 having linear resolution is totally reflexive and

$$\text{Hom}_A(-, A)(1) : \text{cln } A \quad \leftrightarrow \quad \text{cln } A^0 : \text{Hom}_{A^0}(-, A)(1)$$

is a duality.

- (2) $\mathcal{P}^1(A) = (E, \sigma)$ and $\mathcal{P}^1(A^0) = (E, \sigma^{-1})$ for some bijection $\sigma : E \rightarrow E$.
- (3) A and A^0 satisfy **(pc)**.

Proof. (1) \Rightarrow (2): This follows from the proof of [15, Theorem 6.1(1)].

(2) \Rightarrow (3): This follows from Theorem 3.7.

(3) \Rightarrow (1): Suppose that every co-point module $N_{p_0} \in \text{cln } A$ has a complete resolution of the form

$$\dots \xrightarrow{v_2\cdot} A(-2) \xrightarrow{v_1\cdot} A(-1) \xrightarrow{v_0\cdot} A \xrightarrow{v_{-1}\cdot} A(1) \xrightarrow{v_{-2}\cdot} A(2) \xrightarrow{v_{-3}\cdot} \dots$$

Applying the functor $\text{Hom}_A(-, A)$, we have an acyclic complex

$$\dots \xleftarrow{v_2\cdot} A(2) \xleftarrow{v_1\cdot} A(1) \xleftarrow{v_0\cdot} A \xleftarrow{v_{-1}\cdot} A(-1) \xleftarrow{v_{-2}\cdot} A(-2) \xleftarrow{v_{-3}\cdot} \dots,$$

so it is a complete resolution of $\underline{\text{Hom}}_A(N_{p_0}, A)$. It follows that $\underline{\text{Hom}}_A(N_{p_0}, A)(1) \cong_{p-2} N \in \text{clin } A^0$, so there is a functor $\underline{\text{Hom}}_A(-, A)(1) : \text{clin } A \rightarrow \text{clin } A^0$. Since N_{p_0} is totally reflexive by Lemma 3.3,

$$\underline{\text{Hom}}_{A^0}(\underline{\text{Hom}}_A(N_{p_0}, A)(1), A)(1) \cong \underline{\text{Hom}}_{A^0}(\underline{\text{Hom}}_A(N_{p_0}, A), A) \cong N_{p_0}.$$

By symmetry, every co-point module over A^0 having a linear resolution is totally reflexive, and there is a functor $\underline{\text{Hom}}_{A^0}(-, A)(1) : \text{clin } A^0 \rightarrow \text{clin } A$ such that $\underline{\text{Hom}}_A(\underline{\text{Hom}}_{A^0}(N, A)(1), A)(1) \cong N$ for all $N \in \text{clin } A^0$. It follows that $\underline{\text{Hom}}_A(-, A)(1) : \text{clin } A \leftrightarrow \text{clin } A^0 : \underline{\text{Hom}}_{A^0}(-, A)(1)$ is a duality. \square

3.2. The condition (ee)

We will use a co-hyperplane module defined below to find algebras not satisfying (ee).

Definition 3.10. Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1, and $H = \mathbb{P}(W) \subset \mathbb{P}(V)$ a hyperplane where $W \subset V$ is a co-dimension 1 subspace. A co-hyperplane module over A associated to H is a cyclic graded right A -module defined by $L_H := A/(WA + A_{\geq 2}) \in \text{grmod } A$.

There is a correspondence between hyperplane modules and co-hyperplane modules via Koszul duality.

Lemma 3.11. Let $A = T(V)/(R)$ be a Koszul algebra where $R \subset V \otimes_k V$ is a subspace, and $H = \mathcal{V}(f) \subset \mathbb{P}(V^*)$ a hyperplane where $f \in A_1 = V$ is a homogeneous regular element of degree 1. If $K = A/f \in \text{lin } A$ is a hyperplane module over A , then $E_A(K) = L_H \in \text{lin } A^1$ is a co-hyperplane module over A^1 associated to the hyperplane H .

Proof. If $kf \subset V$ is a 1-dimensional subspace generated by $f \in A_1 = V$, then $(kf)^\perp := \{\lambda \in V^* \mid \lambda(f) = 0\} \subset V^*$ is a co-dimension 1 subspace, and

$$\begin{aligned} \mathcal{V}(f) &= \{p \in \mathbb{P}(V^*) \mid f(p) = 0\} \\ &= \{[\lambda] \in \mathbb{P}(V^*) \mid \lambda(f) = 0\} \\ &= \{[\lambda] \in \mathbb{P}(V^*) \mid \lambda \in (kf)^\perp\} \\ &= \mathbb{P}((kf)^\perp). \end{aligned}$$

Since $f \in A_1 = V$ is a homogeneous regular element of degree 1,

$$0 \rightarrow A(-1) \xrightarrow{f} A \rightarrow K \rightarrow 0$$

is a linear free resolution of K , so $H_K(t) = H_A(t) - tH_A(t) = (1 - t)H_A(t)$. Since

$$0 \rightarrow kf \otimes_k A \rightarrow k \otimes_k A \rightarrow K \rightarrow 0$$

is the minimal free resolution of $K \in \text{lin } A$, the minimal free resolution of $E_A(K) \in \text{lin } A^1$ begins

$$(kf)^\perp \otimes_k A^1 \rightarrow k^* \otimes_k A^1 \rightarrow E_A(K) \rightarrow 0$$

by [18, Theorem 6.3(3)], so $E_A(K) = A^1/(kf)^\perp A^1 \in \text{lin } A^1$. By [18, Theorem 6.3(2)], $H_{E_A(K)}(t) = H_K(-t)/H_A(-t) = 1 + t$, so $E_A(K) = A^1/(kf)^\perp A^1 = A^1/((kf)^\perp A^1 + A^1_{\geq 2}) \in \text{lin } A^1$ is a co-hyperplane module associated to $H = \mathcal{V}(f)$. \square

Let A be a noetherian graded algebra and $M, N \in \text{grmod } A$. Recall that if N has a complete resolution T , then we define the Tate cohomologies by $\widehat{\text{Ext}}_A^i(N, M) := h^i(\underline{\text{Hom}}_A(T, M))$. The Tate cohomologies can be defined for more general modules (see [3,8]).

Lemma 3.12. *Let $A = T(V)/I$ be a graded algebra finitely generated in degree 1. If $N \in \text{clin } A$ is a complete co-point module with the complete co-point module sequence $\{p_n\}_{n \in \mathbb{Z}}$, and $L = L_H \in \text{grmod } A$ is a co-hyperplane module associated to a hyperplane $H \subset \mathbb{P}(V)$, then $\widehat{\text{Ext}}_A^i(N, L) \neq 0$ if and only if $p_i \in H$ or $p_{i-1} \in H$.*

Proof. Since the complete resolution of $N \in \text{clin } A$ is

$$\dots \xrightarrow{v_2} A(-2) \xrightarrow{v_1} A(-1) \xrightarrow{v_0} A \xrightarrow{v_{-1}} A(1) \xrightarrow{v_{-2}} A(2) \xrightarrow{v_{-3}} \dots$$

where $p_i = [v_i] \in \mathbb{P}(V)$, $\widehat{\text{Ext}}_A^i(N, L)$ are the homologies of the complex

$$\dots \xrightarrow{v_{-3}} L(-2) \xrightarrow{v_{-2}} L(-1) \xrightarrow{v_{-1}} L \xrightarrow{v_0} L(1) \xrightarrow{v_1} L(2) \xrightarrow{v_2} \dots$$

Since $L = A/(WA + A_{\geq 2}) = k \oplus (V/W)$ where $W \subset V$ is a codimension 1 subspace such that $H = \mathbb{P}(W)$,

$$\text{Ker}\{L(i) \xrightarrow{v_i} L(i+1)\} = \begin{cases} L(i) & \text{if } v_i \in W, \\ L(i)_{-i+1} & \text{if } v_i \notin W, \end{cases}$$

and

$$\text{Im}\{L(i-1) \xrightarrow{v_{i-1}} L(i)\} = \begin{cases} 0 & \text{if } v_{i-1} \in W, \\ L(i)_{-i+1} & \text{if } v_{i-1} \notin W, \end{cases}$$

hence the result. \square

Proposition 3.13. *Let $A = T(V)/I$ be a noetherian AS-Gorenstein algebra with $\mathcal{P}^1(A) = (E, \sigma)$ such that $\sigma : E \rightarrow E$ is a bijection. If $N = N_p \in \text{clin } A$ is a co-point module over A having a linear resolution associated to a point $p \in E$, and $L = L_H \in \text{grmod } A$ is a co-hyperplane module associated to a hyperplane $H \subset \mathbb{P}(V)$, then $\widehat{\text{Ext}}_A^i(N, L) \neq 0$ if and only if $\sigma^i(p) \in H$ or $\sigma^{i-1}(p) \in H$.*

Proof. By Theorem 3.7, for every $p \in E$, $N_p \in \text{clin } A$ is a complete co-point module with the complete co-point module sequence $\{\sigma^n(p)\}_{n \in \mathbb{Z}}$, so the result follows from Lemma 3.12. \square

The following theorem provides more examples of algebras not satisfying **(uac)**, extending [13, Corollary 6.2].

Theorem 3.14. *Let $A = T(V)/I$ be a noetherian AS-Gorenstein algebra with $\mathcal{P}^1(A) = (E, \sigma)$ such that $\sigma : E \rightarrow E$ is a bijection. If there are a point $p \in \mathbb{P}(V)$ and a hyperplane $H \subset \mathbb{P}(V)$ such that $0 < \#\{i \in \mathbb{Z} \mid \sigma^i(p) \in H\} < \infty$, then A does not satisfy **(uac)**.*

Proof. Let $d = \text{id } A < \infty$. We fix $n \in \mathbb{Z}$ such that $\sigma^n(p) \in H$. By [14, Lemma 3.1], $\widehat{\text{Ext}}_A^i(N_{\sigma^{n-d-1}(p)}, L_H) \cong \widehat{\text{Ext}}_A^i(N_{\sigma^{n-d-1}(p)}, L_H)$ for $i > d$. Since $\sigma^i(\sigma^{n-d-1}(p)) = \sigma^{i+n-d-1}(p) \notin H$ for any $i \gg 0$ by the assumption, $\widehat{\text{Ext}}_A^i(N_{\sigma^{n-d-1}(p)}, L_H) = 0$ for all $i \gg 0$ by Proposition 3.13. However, since $\sigma^{d+1}(\sigma^{n-d-1}(p)) = \sigma^n(p) \in H$,

$$\widehat{\text{Ext}}_A^{d+1}(N_{\sigma^{n-d-1}(p)}, L_H) \cong \widehat{\text{Ext}}_A^{d+1}(N_{\sigma^{n-d-1}(p)}, L_H) \neq 0$$

by Proposition 3.13 again, so A does not satisfy **(uac)** by [14, Corollary 3.3]. \square

Note that in order for the geometric condition in the above theorem to be satisfied, we must have $|\sigma| = \infty$.

The following is the main theorem of this section.

Theorem 3.15. *Let $A = T(V)/(R)$ be a Frobenius Koszul algebra with $\mathcal{P}^1(A) = (E, \sigma)$ such that A^1 is a quantum polynomial algebra, and let $\nu \in \text{Aut } A$ be the Nakayama automorphism of A . If there are a point $p \in E$ and a hyperplane $H \subset \mathbb{P}(V)$ such that $\#\{i \in \mathbb{N} \mid \sigma^i(p) \in H\} = \infty$ but $\#\{i \in \mathbb{N} \mid \sigma^{-i}\nu(p) \in H\} < \infty$, then A does not satisfy (ee).*

Proof. If $\#\{i \in \mathbb{N} \mid \sigma^i(p) \in H\} = \infty$, then $\widehat{\text{Ext}}_A^i(N, L) \neq 0$ for infinitely many $i \in \mathbb{N}$ by Proposition 3.13, so $\underline{\text{Ext}}_A^i(N, L) \neq 0$ for infinitely many $i \in \mathbb{N}$ by [14, Lemma 3.1]. On the other hand, suppose that $\#\{i \in \mathbb{N} \mid \sigma^{-i}\nu(p) \in H\} < \infty$. Since

$$\widehat{\text{Ext}}_A^i(L, N) \cong \widehat{\text{Ext}}_A^{-1-i}((N_p)_{\nu^{-1}}(-d), L)^* \cong \widehat{\text{Ext}}_A^{-1-i}(N_{\nu(p)}, L)^*(-d)$$

as graded vector spaces where $d = \text{gldim } A^1$ by [8, Proposition 8] and [15, Lemma 3.2(2)], $\widehat{\text{Ext}}_A^i(L, N) \neq 0$ for finitely many $i \in \mathbb{N}$ by Proposition 3.13 again, hence $\underline{\text{Ext}}_A^i(L, N) \neq 0$ for finitely many $i \in \mathbb{N}$ by [14, Lemma 3.1] again. \square

Corollary 3.16. *Let $A = T(V)/(R)$ be a Frobenius Koszul algebra with $\mathcal{P}^1(A) = (E, \sigma)$ such that A^1 is a quantum polynomial algebra, and let $\nu \in \text{Aut } A$ be the Nakayama automorphism of A . If*

- k is an infinite field,
- $\dim_k V \geq 3$,
- $|\sigma| < \infty$, and
- there is a point $p \in E$ such that $\sigma^i(p) \neq \nu(p)$ for any $i \in \mathbb{Z}$,

then A does not satisfy (ee).

Proof. The proof is similar to that of Corollary 2.14. \square

Remark 3.17. Recall that the map $\sigma : E \rightarrow E$ corresponds to the syzygy functor $\Omega : \underline{\text{gmod}} A \rightarrow \underline{\text{gmod}} A$, and the map $\nu : E \rightarrow E$ corresponds to the Nakayama functor $-\otimes_A A^* : \underline{\text{gmod}} A \rightarrow \underline{\text{gmod}} A$, so the condition $\sigma^i(p) \neq \nu(p)$ for any $i \in \mathbb{Z}$ in the above corollary implies that $\underline{\text{gmod}} A$ is not Calabi-Yau (up to degree shifts) or A is not stably symmetric in the sense of [14].

Example 3.18. Let $A = k\langle x, y, z \rangle / (\alpha zy + yz, \beta xz + zx, \gamma yx + xy, x^2, y^2, z^2)$ where $\alpha, \beta, \gamma \in k$ such that $\alpha\beta\gamma \neq 0, 1$. Since $A^1 \cong k\langle x, y, z \rangle / (zy - \alpha yz, xz - \beta zx, yx - \gamma xy)$, $\mathcal{P}^1(A) = \mathcal{P}(A^1) = (E, \sigma)$ where $E = \mathcal{V}(xyz) \subset \mathbb{P}(V)$ and

$$\begin{aligned} \sigma(0, b, c) &= (0, \alpha b, c), \\ \sigma(a, 0, c) &= (a, 0, \beta c), \\ \sigma(a, b, 0) &= (\gamma a, b, 0) \end{aligned}$$

by [13, Theorem 3.8], and the map $\nu : E \rightarrow E$ induced by the Nakayama automorphism $\nu \in \text{Aut } A$ of A is given by

$$\nu(a, b, c) = ((\gamma/\beta)a, (\alpha/\gamma)b, (\beta/\alpha)c)$$

by [21, Theorem 9.2]. Suppose that k is an infinite field, and α, β, γ are roots of unity so that $|\sigma| < \infty$. If $\alpha^i \neq \beta\gamma$ for any $i \in \mathbb{Z}$, $\beta^i \neq \gamma\alpha$ for any $i \in \mathbb{Z}$, or $\gamma^i \neq \alpha\beta$ for any $i \in \mathbb{Z}$, then A does not satisfy (ee) by the same argument as in Example 2.15.

Remark 3.19. In the commutative case, the condition **(uac)** implies the condition **(ee)** over noetherian commutative Gorenstein local rings [5, Theorem 4.1]. However, in the above example, if α, β, γ are roots of unity, then A satisfies **(uac)** by [13, Theorem 6.5]. So, for example, if $\alpha = 1$ and $\beta = \gamma$ is the primitive n th root of unity for $n \geq 3$, then A is a noetherian AS-Gorenstein (Frobenius) algebra satisfying **(uac)** but not satisfying **(ee)**. It follows that the condition **(uac)** does not imply the condition **(ee)** over noncommutative Gorenstein rings. In [4], there is given a simple example of a noncommutative (Frobenius) algebra not satisfying **(ee)**, however, that example does not satisfy **(uac)** either by [13, Theorem 6.5], so Example 3.18 may provide the first example of an algebra satisfying **(uac)** but not satisfying **(ee)**. However, Example 3.18 also suggests that, for a Frobenius, non-stably symmetric Koszul algebra A with $\mathcal{P}^1(A) = (E, \sigma)$, if $|\sigma| < \infty$, then A tends to satisfy **(uac)** but A tends not to satisfy **(ee)**, so, contrary to the commutative situation, **(uac)** and **(ee)** are rather exclusive conditions.

References

- [1] M. Artin, J.J. Zhang, Noncommutative projective schemes, *Adv. Math.* 109 (1994) 228–287.
- [2] L.L. Avramov, R. Buchweitz, Support varieties and cohomology over complete intersections, *Invent. Math.* 142 (2000) 285–318.
- [3] L.L. Avramov, A. Martsinkovsky, Absolute, relative, and Tate cohomology of modules of finite Gorenstein dimension, *Proc. London Math. Soc.* 85 (2002) 393–440.
- [4] P.A. Bergh, On the Hochschild (co)homology of quantum exterior algebras, *Comm. Algebra* 35 (2007) 3440–3450.
- [5] C. Huneke, D.A. Jorgensen, Symmetry in the vanishing of Ext over Gorenstein rings, *Math. Scand.* 93 (2003) 161–184.
- [6] D.A. Jorgensen, L.M. Sega, Asymmetric complete resolutions and vanishing of Ext over Gorenstein rings, *Int. Math. Res. Not.* 56 (2005) 3459–3477.
- [7] P. Jorgensen, Finite Cohen–Macaulay type and smooth non-commutative schemes, *Canad. J. Math.* 60 (2008) 379–390.
- [8] R. Martinez-Villa, A. Martsinkovsky, Cohomology of tails, Tate–Vogel cohomology, and noncommutative Serre duality over Koszul quiver algebras, *J. Algebra* 280 (2004) 58–83.
- [9] I. Mori, Intersection multiplicity over noncommutative algebras, *J. Algebra* 252 (2002) 241–257.
- [10] I. Mori, Homological properties of balanced Cohen–Macaulay algebras, *Trans. Amer. Math. Soc.* 355 (2003) 1025–1042.
- [11] I. Mori, Serre’s vanishing conjecture for Ext-groups, *J. Pure Appl. Algebra* 187 (2004) 207–240.
- [12] I. Mori, Noncommutative projective schemes and point schemes, in: *Algebras, Rings and Their Representations*, World Sci. Publ., 2006, pp. 215–239.
- [13] I. Mori, Co-point modules over Koszul algebras, *J. London Math. Soc.* 74 (2006) 639–656.
- [14] I. Mori, Symmetry in the vanishing of Ext over stably symmetric algebras, *J. Algebra* 310 (2007) 708–729.
- [15] I. Mori, Co-point modules over Frobenius Koszul algebras, *Comm. Algebra* 36 (2008) 4659–4677.
- [16] I. Mori, S.P. Smith, Bézout’s theorem for non-commutative projective spaces, *J. Pure Appl. Algebra* 157 (2001) 279–299.
- [17] K. de Naeghel, M. Van den Bergh, Ideal classes of three-dimensional Sklyanin algebras, *J. Algebra* 276 (2004) 515–551.
- [18] S.P. Smith, Some finite dimensional algebras related to elliptic curves, in: *Representation Theory of Algebras and Related Topics*, Mexico City, 1994, in: *CMS Conf. Proc.*, vol. 19, Amer. Math. Soc., Providence, RI, 1996, pp. 315–348.
- [19] C. Weibel, *An Introduction to Homological Algebra*, Cambridge Stud. Adv. Math., vol. 38, Cambridge University Press, Cambridge, 1995.
- [20] A. Yekutieli, Dualizing complexes over noncommutative graded algebras, *J. Algebra* 153 (1992) 41–84.
- [21] M. Van den Bergh, Existence theorems for dualizing complexes over non-commutative graded and filtered rings, *J. Algebra* 195 (1997) 662–679.
- [22] J.J. Zhang, Connected graded Gorenstein algebras with enough normal elements, *J. Algebra* 189 (1997) 390–405.