



ELSEVIER

Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



An insertion algorithm on multiset partitions with applications to diagram algebras



Laura Colmenarejo^a, Rosa Orellana^b, Franco Saliola^c,
Anne Schilling^{d,*}, Mike Zabrocki^{e,*}

^a Department of Mathematics and Statistics, UMass Amherst, 710 N Pleasant St, Amherst, USA

^b Mathematics Department, Dartmouth College, 6188 Kemeny Hall, Hanover, NH 03755, USA

^c Département de mathématiques, Université du Québec à Montréal, Canada

^d Department of Mathematics, University of California, One Shields Avenue, Davis, CA 95616-8633, USA

^e Department of Mathematics and Statistics, York University, 4700 Keele Street, Toronto, Ontario M3J 1P3, Canada

ARTICLE INFO

Article history:

Received 9 May 2019

Available online 20 April 2020

Communicated by Volodymyr Mazorchuk

Keywords:

Partition algebras

RSK algorithm

Multiset tableaux

Representation theory

ABSTRACT

We generalize the Robinson–Schensted–Knuth algorithm to the insertion of two row arrays of multisets. This generalization leads to new enumerative results that have representation theoretic interpretations as decompositions of centralizer algebras and the spaces they act on. In addition, restrictions on the multisets lead to further identities and representation theory analogues. For instance, we obtain a bijection between words of length k with entries in $[n]$ and pairs of tableaux of the same shape with one being a standard Young tableau of size n and the other being a standard multiset tableau of content $[k]$. We also obtain an algorithm from partition diagrams to pairs of a standard tableau and a standard multiset tableau of the same shape, which has the remarkable property

* Corresponding authors.

E-mail addresses: laura.colmenarejo.hernando@gmail.com (L. Colmenarejo), Rosa.C.Orellana@dartmouth.edu (R. Orellana), saliola.franco@uqam.ca (F. Saliola), anne@math.ucdavis.edu (A. Schilling), zabrocki@mathstat.yorku.ca (M. Zabrocki).

URLs: <https://sites.google.com/view/l-colmenarejo/home> (L. Colmenarejo), <https://math.dartmouth.edu/~orellana/> (R. Orellana), <http://lacim.uqam.ca/~saliola/> (F. Saliola), <http://www.math.ucdavis.edu/~anne> (A. Schilling), <http://garsia.math.yorku.ca/~zabrocki/> (M. Zabrocki).

<https://doi.org/10.1016/j.jalgebra.2020.04.010>

0021-8693/© 2020 Elsevier Inc. All rights reserved.

that it is well-behaved with respect to restricting a representation to a subalgebra. This insertion algorithm matches recent representation-theoretic results of Halverson and Jacobson [15].

© 2020 Elsevier Inc. All rights reserved.

1. Introduction

We explore a variant of the Robinson–Schensted–Knuth (RSK) algorithm, where we insert multisets instead of integer entries. If we restrict the multisets to all have size one, the algorithm we are using is the usual RSK algorithm. Applying this insertion to different arrays of multisets gives rise to a purely enumerative result that is a combinatorial manifestation of a double centralizer theorem from representation theory. Although representation theory serves as a principal motivation for studying these algorithms, no familiarity is assumed in our exposition of the enumerative and combinatorial results.

The RSK algorithm evolved over the last century from a procedure defined on permutations (in the work of Robinson [31]) to a procedure defined on finite sequences of integers (in the work of Schensted [35]) and finally to a procedure defined on “generalized permutations” by Knuth [22]. In each of these versions, the algorithm establishes a correspondence between the initial input and pairs of combinatorial objects called tableaux subject to certain constraints (see Section 2 for definitions).

Each of the above procedures reflects a classical direct-sum decomposition result in representation theory. While the reader will find more details in Section 4, we present here an overview. Broadly speaking, we start with two families of operators, say \mathcal{A} and \mathcal{B} , acting on a vector space V , and we determine the finest decomposition of V into a direct sum of subspaces that are invariant for all the operators, say $V = \bigoplus_{\lambda \in \Lambda} V^\lambda$ for some indexing set Λ . Under certain circumstances, the actions of \mathcal{A} and \mathcal{B} neatly separate the subspaces V^λ ; more precisely, V^λ can be expressed as a tensor product $U^\lambda \otimes W^\lambda$, where the action of \mathcal{A} only affects U^λ and the action of \mathcal{B} only affects W^λ . Thus, we obtain the decomposition:

$$V \cong \bigoplus_{\lambda \in \Lambda} (U^\lambda \otimes W^\lambda).$$

At the combinatorial level, this decomposition implies that there is a bijection between \mathcal{V} and $\bigcup_{\lambda \in \Lambda} (U^\lambda \times W^\lambda)$, where \mathcal{V} , U^λ and W^λ denote bases of V , U^λ and W^λ , respectively.

One example of this is when $GL(V)$ and S_k both act on $V^{\otimes k}$ (see Section 4 for details). In this case, we deduce the existence of a bijection between the set of finite sequences of length k with entries in $\{1, 2, \dots, \dim(V)\}$ (*i.e.*, a basis of $V^{\otimes k}$) and the union of the set of pairs consisting of a semistandard tableau of shape λ with entries in $\{1, 2, \dots, \dim(V)\}$ (*i.e.*, a basis of U^λ) and a standard tableau of shape λ and size k (*i.e.*, a basis of W^λ). This is precisely what the RSK algorithm does (see Section 3).

The above situation also holds for many other pairs of families of operators acting on $V^{\otimes k}$; for instance: the partition algebra and the symmetric group; the Brauer algebra and the orthogonal group; and the Hecke algebra and the quantum group of type A .

In this paper, we adapt the RSK algorithm to the insertion of arrays of multisets. This adaptation gives combinatorial descriptions of other direct-sum decomposition results in representation theory. Furthermore, restrictions on the multisets result in a bijection and an enumerative result relating sets of combinatorial objects. For instance, by considering a vector space on which both the symmetric group and the partition algebra act, we obtain a bijection between words of length k with entries in $[n]$ and pairs of tableaux of the same shape with one being a standard Young tableau of size n and the other being a standard multiset tableau of content $[k]$. We also obtain an algorithm from monomials in a polynomial ring to pairs of a standard tableau and a standard multiset tableau of the same shape and from elements of diagram algebras to pairs of standard multiset tableaux.

Note that algorithms that relate partition diagrams and pairs of paths in the Bratteli diagram for the partition algebras have been known since the late 1990s [16,26]. These paths are referred as “vacillating tableaux” and they are analogues of a path in the Young’s lattice, which is the Bratteli diagram for the symmetric groups. Paths in the Young lattice are encoded by standard Young tableaux.

Recently, a new combinatorial interpretation for the dimensions of the irreducible representations for the partition algebra has appeared in the literature [2,3,29,15,13]. In particular, Benkart and Halverson [2] presented a bijection between vacillating tableaux and “set-partition tableaux” (tableaux whose entries are sets of positive integers). There are two main advantages to working with set-partition tableaux instead of vacillating tableaux. Firstly, they are closer in spirit to the ubiquitous Young tableaux. Secondly, the definition extends naturally to the notion of multiset tableaux (tableaux whose entries are multisets of positive integers) and working with multiset tableaux leads to new enumerative and algebraic results that are not obvious by other means (see Proposition 5.1, Corollary 5.4, and Theorem 6.3).

Our insertion algorithm from partition diagrams to pairs of a standard tableau and a standard multiset tableau of the same shape has the remarkable property that it is well-behaved with respect to the subalgebra structure of the partition algebra. One surprising consequence is that we are able to provide explicit combinatorial descriptions of the sets of tableaux that give the dimensions of the irreducible representations associated to the prominent subalgebras of the partition algebras, such as the symmetric group, the Brauer algebra, the rook algebra, the rook-Brauer algebra, the Temperley–Lieb algebra, the Motzkin algebra, the planar rook algebra, and the planar algebra (see Lemma 6.6). This gives rise to analogues of the famous identity $n! = \sum_{\lambda \vdash n} (f^\lambda)^2$ for the symmetric group, where f^λ is the number of standard tableaux of shape λ , to all of the above mentioned algebras (see Corollary 6.8). We prove that the dimensions of the irreducible representations of the various algebras is equal to the number of our combinatorially-defined tableaux by establishing that the branching rules are encoded in the tableaux (see

Section 6.3 and Corollary 6.16). Our insertion is different from combining the insertion of Halverson and Lewandowski [16] from partition diagrams to paths in the Bratteli diagram with the bijection of Halverson and Benkart [2] from paths in the Bratteli diagram to set-partition tableaux (see Section 6.3).

The paper is organized as follows. In Section 2, we define the principal combinatorial objects used throughout this paper: multiset tableaux. In Section 3, we review the RSK algorithm for associating a pair of tableaux to a generalized permutation. The above discussion is expanded in Section 4 by providing more details on how RSK on permutations, words, and generalized permutations, reflects decomposition results in representation theory. In Section 5, the RSK algorithm is adapted to the multiset tableaux setting and corresponding enumerative results are obtained. The section opens with a description of the enumerative and combinatorial results and closes by connecting these results with representation theory. In Section 6 the algorithm is applied to partition algebra diagrams and it is shown that the new insertion algorithm is well-behaved when restricted to sub-algebras. Finally, the connection to the representation theory recently developed in [15] is established.

Acknowledgments

The authors would like to thank BIRS and AIM for the opportunity to collaborate in Banff in April 2018 and in San José in November 2018, which greatly facilitated the work on this project. We thank Tom Halverson for sharing an early version of [13] with us.

The second author was partially supported by NSF grant DMS-1700058 and the fourth author was partially supported by NSF grants DMS-1760329 and DMS-1764153. The third and fifth authors were supported by NSERC Discovery Grants.

2. Multiset tableaux

Throughout this paper, we work with tableaux whose entries are multisets. Note that any Young tableau—that is, a tableau with integer entries—can be viewed as a multiset tableau by considering each entry to be a multiset of cardinality 1. In this section, we fix notation and define the total orders on multisets that we use in order to extend the property of being (semi)standard to multiset tableaux.

2.1. Partitions

A *partition* of $n \in \mathbb{N}$ is a sequence of positive integers $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_r)$ with $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_r > 0$ whose sum is n . Note that the empty sequence $()$ is a partition of 0. The notation $\lambda \vdash n$ is used to indicate that λ is a partition of n . The *length* of the partition is denoted by $\ell(\lambda) = r$. As is customary, we depict partitions as diagrams; see

Example 2.1. The *cells* of the partition are the coordinates of the boxes in the diagram; that is, $\text{cells}(\lambda) = \{(i, j) \mid 1 \leq i \leq \lambda_j, 1 \leq j \leq \ell(\lambda)\}$. The operation of removing the first row of the partition λ is denoted by $\bar{\lambda} = (\lambda_2, \lambda_3, \dots, \lambda_r)$.

2.2. Set partitions

A *set partition* π of a set S is a collection $\{\pi_1, \dots, \pi_k\}$ of non-empty subsets of S that are mutually disjoint, *i.e.*, $\pi_i \cap \pi_j = \emptyset$ for all $i \neq j$, and $\bigcup_i \pi_i = S$. The subsets π_i are called the *blocks* of the set partition. We write $\pi \vdash S$ to mean that π is a set partition of the set S .

2.3. Multisets

Let (A, \leq_A) be a totally ordered set, which we refer to as an *(ordered) alphabet*. A *multiset* $S = \{\{a_1, a_2, \dots, a_r\}\}$ over A is an unordered collection of elements of A , allowing repeats. The collection of multisets forms an associative monoid with operation

$$\{\{a_1, a_2, \dots, a_r\}\} \uplus \{\{b_1, b_2, \dots, b_d\}\} = \{\{a_1, a_2, \dots, a_r, b_1, b_2, \dots, b_d\}\} .$$

To simplify notation, we let $\{\{a^{m_a}, b^{m_b}, c^{m_c}, \dots\}\}$ denote the multiset that contains m_a occurrences of a , m_b occurrences of b , and so on; for example $\{\{1^2, 4^3, 5\}\} = \{\{1, 1, 4, 4, 4, 5\}\}$.

A *multiset partition*¹ of a multiset S is a multiset of multisets, $\pi = \{\{S^{(1)}, S^{(2)}, \dots, S^{(r)}\}\}$, such that $S = S^{(1)} \uplus S^{(2)} \uplus \dots \uplus S^{(r)}$. We indicate this by the notation $\pi \Vdash S$.

2.4. Ordering multisets

We will use two different methods to totally order the collection of all multisets over an ordered alphabet A . In Section 5, we use *graded lexicographic order*. If $S = \{\{a_1, \dots, a_r\}\}$ with $a_1 \leq_A \dots \leq_A a_r$ and $S' = \{\{a'_1, \dots, a'_t\}\}$ with $a'_1 \leq_A \dots \leq_A a'_t$, then we say $S < S'$ in the *graded lexicographic order* if:

- $r < t$; or
- $r = t$ and there exists $1 \leq i \leq t$ such that $a_1 = a'_1, \dots, a_{i-1} = a'_{i-1}$, and $a_i <_A a'_i$.

This is a total order [5], with minimum element the empty multiset.

In Section 6, where we need only compare disjoint sets, we use the *last letter order*. Given two disjoint sets S and S' with elements in an ordered set A , we say $S < S'$ in the *last letter order* if $\max(S) <_A \max(S')$, where $\max(S)$ is the largest element in S . For

¹ Multisets are in bijection with integer vectors and multiset partitions are in bijection with objects known as vector partitions [6,9,23,32]. Since integer vectors can be identified with sequences of monomials in a set of variables, another interpretation for multisets is as monomials in the variables $\{x_{c_1}, x_{c_2}, \dots\}$.

example, $\{1, 3, 5\} < \{2, 7\}$. (This order can be realized as the restriction of a total order on multisets, for example *reverse lexicographic order*, but this is not necessary here.)

2.5. Multiset tableaux

Let λ be a partition, A an ordered alphabet, and a fixed total order $<$ on multisets (such as the graded lexicographic order or the last letter order if the multisets are all disjoint sets). A *semistandard multiset tableau* of *shape* λ is a function T that associates with each cell $(i, j) \in \text{cells}(\lambda)$ a multiset over A such that:

- $T(i, j) \leq T(i, j + 1)$ whenever (i, j) and $(i, j + 1)$ both belong to $\text{cells}(\lambda)$; and
- $T(i, j) < T(i + 1, j)$ whenever (i, j) and $(i + 1, j)$ both belong to $\text{cells}(\lambda)$.

The *shape* of a multiset tableau T is the partition λ , and the *cells* of T are the cells of its shape. If $T(i, j) = S$, then we say that S *labels* the cell (i, j) , and that S is an *entry* of T .

When drawing multiset tableaux, the multisets are abbreviated as words without the surrounding multiset delimiters $\{\!\!\{, \}\!\!\}$, and empty sets are depicted by blank cells.

2.6. Content of multiset tableaux

The *content* of a semistandard multiset tableau T is the (disjoint) union of the entries of T . More precisely, the content of T is the multiset

$$\text{content}(T) = \bigsqcup_{(i,j) \in \text{cells}(T)} T(i, j).$$

A semistandard multiset tableau is said to be

- a *standard multiset tableau* if its content is the set $[k] := \{1, 2, \dots, k\}$ for some $k \in \mathbb{N}$; in other words, each letter $1, 2, \dots, k$ appears exactly once in the tableau;
- a *semistandard Young tableau* if all its entries are multisets of size 1;
- a *standard Young tableau* if it is both standard and a semistandard Young tableau.

Finally, for a multiset S , let

- $\text{SSMT}(\lambda, S)$ be the set of *semistandard* multiset tableaux of shape λ and content S ;
- $\text{SMT}(\lambda, k)$ be the set of *standard* multiset tableaux of shape λ and content $[k]$;
- $\text{SSYT}(\lambda, S)$ be the set of semistandard Young tableaux of shape λ and content S ;
- and
- $\text{SYT}(\lambda)$ be the set of standard Young tableaux of shape λ .

The *set-partition tableaux* of [2, Definition 3.14] are closely related to our standard multiset tableaux.

Example 2.1. Let $A = \{1, 2, 3, 4, 5\}$ with the usual order on integers. Then

114		
2	14	
	2	5

3		
2	3	
1	1	4

134		
5	26	
	7	8

are three semistandard multiset tableaux of shape $(3, 2, 1)$. The leftmost tableau has content $\{\{1^3, 2^2, 4^2, 5\}\}$. The middle tableau has content $\{\{1^2, 2, 3^2, 4\}\}$ and is also a semistandard Young tableau. The rightmost multiset tableau is standard since its content is $\{1, \dots, 8\}$.

3. The RSK correspondence

We present here the Robinson–Schensted–Knuth (RSK) algorithm for certain finite sequences in $A \times B$, where A and B are totally ordered sets. Our presentation is slightly more general than the original [22], where it was defined on certain finite sequences in $\mathbb{N} \times \mathbb{N}$. The proofs in [22] still hold in this context because they only make use of the fact that \mathbb{N} is a totally ordered set.

Let A and B be two ordered alphabets. A *generalized permutation* from A to B is a two-line array of the form $\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}$ satisfying: $a_1, \dots, a_r \in A$; $b_1, \dots, b_r \in B$; $a_i \leq_A a_{i+1}$ for $1 \leq i \leq r - 1$; and $b_i \leq_B b_{i+1}$ whenever $a_i = a_{i+1}$.

The RSK algorithm constructs two semistandard tableaux of the same shape from a generalized permutation $\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}$. The cells of one of the tableaux are labelled by a_1, \dots, a_r and the cells of the other tableau are labelled by b_1, \dots, b_r . The key step in the algorithm is the following insertion procedure.

Definition 3.1 (*RSK insertion procedure*). Let T be a semistandard tableau with entries in an ordered set X and $x \in X$. The *(row) insertion* of x into T is the tableau defined recursively as follows, starting with $r = 1$.


- (1) If x is greater than or equal to all entries of the r -th row of T (this includes the case where the r -th row has no entries), then append x to the end of the row; otherwise
- (2) let \hat{x} be the leftmost entry of the row that is greater than x , replace \hat{x} with x , and insert the \hat{x} into row $r + 1$ (i.e., repeat step (1) with $x = \hat{x}$ and r replaced by $r + 1$).

Iterating this insertion procedure allows us to associate two semistandard tableaux with any generalized permutation $\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}$ as follows (cf. Example 3.2).

- Start with a pair of empty tableaux; *i.e.*, tableaux with no rows and no columns.
- Insert b_1 into the first tableau using the insertion procedure; this introduces a new cell, say in position (i_1, j_1) ; add a cell labelled a_1 in position (i_1, j_1) of the second tableau.
- Insert b_2 into the first tableau, which introduces a new cell to the first tableau; label the corresponding cell in the second tableau by a_2 .
- Repeat with $(a_3, b_3), (a_4, b_4), \dots, (a_r, b_r)$.

The result is two semistandard tableaux of the same shape, the first has entries b_1, \dots, b_r , and the second has entries a_1, \dots, a_r .

Example 3.2. Consider the alphabets $A = \{a < b < c < d\}$ and $B = \{w < x < y < z\}$. The insertion procedure applied to the generalized permutation $\begin{pmatrix} a & b & b & c & c & d & d & d \\ x & y & z & w & y & w & y & y \end{pmatrix}$, gives the following sequence of tableaux.



Theorem 3.3 (Knuth). *There is a one-to-one correspondence between generalized permutations from A to B and pairs of tableaux (P, Q) satisfying: P and Q are of the same shape; P is semistandard with entries in B ; and Q is semistandard with entries in A .*

Two special cases of this correspondence coincide with the correspondences discovered by Robinson [31] and Schensted [35].

- (1) By encoding a permutation σ of size n as the generalized permutation $\begin{pmatrix} 1 & 2 & \dots & n \\ \sigma_1 & \sigma_2 & \dots & \sigma_n \end{pmatrix}$, where $\sigma_i = \sigma(i)$, we obtain a bijection between permutations of size n and pairs of standard Young tableaux of the same shape of size n .
- (2) By encoding a word $w_1 \dots w_k$ as $\begin{pmatrix} 1 & 2 & \dots & k \\ w_1 & w_2 & \dots & w_k \end{pmatrix}$, we obtain a bijection between words of length k with entries in $[n]$ and pairs of Young tableaux of the same shape, the first semistandard with entries in $[n]$ and the second standard with entries in $[k]$.

4. RSK and representation theory

As described in Section 1, the RSK algorithm is a combinatorial manifestation of certain direct-sum decompositions from representation theory. Below we consider the

bijections induced by the RSK algorithm on permutations, on words, and on generalized permutations. For each bijection, we describe a vector space equipped with an action by two families of operators, and the decomposition of the space into a direct sum of invariant subspaces.

4.1. RSK and the group algebra $\mathbb{C}S_n$

When considered as a procedure on permutations, RSK establishes a correspondence between permutations of $[n]$ and pairs of standard tableaux of the same shape of size n . If $f^\lambda := \# \text{SYT}(\lambda)$ denotes the number of standard tableaux of shape λ , then this correspondence gives the enumerative result:

$$n! = \sum_{\lambda \vdash n} (f^\lambda)^2. \tag{4.1}$$

This formula reflects the following classic decomposition result in representation theory. Let $\mathbb{C}S_n$ denote the group algebra of the symmetric group S_n with coefficients in \mathbb{C} . This is both a left and a right module over S_n , and so it admits a decomposition into simple two-sided S_n -modules. This decomposition takes the form

$$\mathbb{C}S_n \cong \bigoplus_{\lambda \vdash n} (S^\lambda)^* \otimes S^\lambda, \tag{4.2}$$

where $\{S^\lambda \mid \lambda \vdash n\}$ is a complete set of non-isomorphic simple right S_n -modules, and $(S^\lambda)^* = \text{Hom}_{S_n}(S^\lambda, \mathbb{C})$. Note that since S^λ is a *right* S_n -module, its dual $\text{Hom}_{S_n}(S^\lambda, \mathbb{C})$ is a *left* S_n -module. One recovers Equation (4.1) from the decomposition in Equation (4.2) by computing the dimension of the vector spaces and noting that $\dim(S^\lambda) = \dim((S^\lambda)^*) = f^\lambda$.

4.2. RSK and the $GL(V) \times S_k$ -structure on $V^{\otimes k}$

When considered as a procedure on finite sequences, RSK establishes a correspondence between finite sequences of length k with entries in $[n]$ and pairs of tableaux (P, Q) satisfying: P and Q have the same shape; P is a semistandard tableau with entries in $[n]$; and Q is a standard tableau with entries $[k]$. Enumeratively,

$$n^k = \sum_{\lambda \vdash k} \# \text{SSYT}_n(\lambda) \cdot \# \text{SYT}(\lambda), \tag{4.3}$$

where $\text{SSYT}_n(\lambda)$ is the set of semistandard tableaux of shape λ with entries in $[n]$, and where $\text{SYT}(\lambda)$ is the set of standard tableaux of shape λ .

This formula also reflects a classic decomposition result in representation theory. Let $V = \mathbb{C}^n$ and consider the tensor product $V^{\otimes k}$ of V with itself k times. This is a left GL_n -module, with $g \in GL_n$ acting on simple tensors by

$$g \cdot (v_1 \otimes v_2 \otimes \cdots \otimes v_k) = g(v_1) \otimes g(v_2) \otimes \cdots \otimes g(v_k);$$

as well as a right S_k -module, with $\sigma \in S_k$ acting on simple tensors by

$$(v_1 \otimes v_2 \otimes \cdots \otimes v_k) \cdot \sigma = v_{\sigma(1)} \otimes v_{\sigma(2)} \otimes \cdots \otimes v_{\sigma(k)};$$

and these two actions commute:

$$g \cdot \left((v_1 \otimes \cdots \otimes v_k) \cdot \sigma \right) = \left(g \cdot (v_1 \otimes \cdots \otimes v_k) \right) \cdot \sigma.$$

Therefore, $V^{\otimes k}$ admits a decomposition into simple $GL_n \times S_k$ -modules, and this decomposition takes the form

$$V^{\otimes k} \cong \bigoplus_{\lambda \vdash k} W_n^\lambda \otimes S^\lambda,$$

where W_n^λ is a simple left GL_n -module and S^λ is a simple right S_k -module. Since $\dim W_n^\lambda = \# \text{SSYT}_n(\lambda)$ and $\dim S^\lambda = \# \text{SYT}(\lambda)$, one immediately recovers Equation (4.3).

The fact that the simple left GL_n -modules are also indexed by partitions λ is a special case of Schur–Weyl duality. It was first observed by Schur [33,34] in his thesis and later promoted by Weyl [40] in his book on the representation theory of the classical groups. The idea extends to a much more general result known as the *double commutant theorem* (see, for instance, [10]). Roughly speaking, it states: if a vector space is acted on by two algebras of operators whose actions mutually centralize each other, then the space decomposes into a direct sum of tensors of two simple modules (for instance, $W_n^\lambda \otimes S^\lambda$), and the two simple modules determine each other.

4.3. RSK and the $GL_n \times GL_k$ -structure on $\mathbb{C}\{x_{ij}\}$

Knuth’s generalization establishes a correspondence between monomials of degree r in n sets of k variables $\{x_{ij} \mid i \in [n], j \in [k]\}$ and pairs of semistandard tableaux (P, Q) of size r satisfying: P and Q have the same shape; P is semistandard with entries in $[n]$; Q is semistandard with entries in $[k]$. This is achieved by encoding monomials as generalized permutations from $[k]$ to $[n]$: each occurrence of x_{ij} is encoded by the column $\binom{j}{i}$, for example $x_{12}^3 x_{14} x_{23}^2$ becomes $\begin{pmatrix} 2 & 2 & 2 & 3 & 3 & 4 \\ 1 & 1 & 1 & 2 & 2 & 1 \end{pmatrix}$.

This correspondence reflects a decomposition of the polynomial ring generated by the variables $\{x_{ij} \mid i \in [n], j \in [k]\}$ when it is viewed as a $GL_n \times GL_k$ -module. To define the module structure, we first arrange the variables in the form of an $n \times k$ matrix:

$$X = \begin{bmatrix} x_{11} & x_{12} & \cdots & x_{1k} \\ x_{21} & x_{22} & \cdots & x_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ x_{n1} & x_{n2} & \cdots & x_{nk} \end{bmatrix}.$$

The left action of $A \in GL_n$ on the polynomial ring corresponds to multiplying X on the left by A . Explicitly, the variable x_{ij} is replaced with $(AX)_{ij} = \sum_l a_{il}x_{lj}$. The right action of $B \in GL_k$ corresponds to multiplying X on the right by B (explicitly, $x_{ij} \mapsto (XB)_{ij}$). Since these actions correspond to multiplying X on the left and right by matrices, the fact that the two actions commute is a consequence of the associativity of matrix multiplication.

Consequently, the polynomial ring admits a direct sum decomposition whose summands are tensors of pairs of simple modules (see, for instance, [17]):

$$\mathbb{C}[X] \cong \bigoplus_{\lambda} W_n^{\lambda} \otimes (W_k^{\lambda})^*,$$

where λ runs over all partitions, W_n^{λ} is the simple left GL_n -module indexed by λ , and $(W_k^{\lambda})^*$ is the simple right GL_k -module indexed by λ . Since the dimension of W_m^{λ} is the number of semistandard tableaux of shape λ with entries in $[m]$, one sees that the set of monomials in $\{x_{ij} \mid i \in [n], j \in [k]\}$ is in bijection with the set of pairs (P, Q) of semistandard tableaux of the same shape with P having entries in $[k]$ and Q having entries in $[n]$.

In particular, for every multiset $\{1^{\alpha_1}, \dots, k^{\alpha_k}\}$, the monomials $\prod_{i=1}^n \prod_{j=1}^k x_{ij}^{b_{ij}}$ satisfying $\sum_{i=1}^n b_{ij} = \alpha_j$ for all $j \in [k]$ span a GL_n -invariant subspace of $\mathbb{C}[X]$ whose dimension is equal to $\prod_{i=1}^k \binom{n+\alpha_i-1}{\alpha_i}$. These monomials are in bijection with the pairs (P, Q) of semistandard tableaux of the same shape with P having entries in $[n]$ and Q having content $\{1^{\alpha_1}, \dots, k^{\alpha_k}\}$. From this we immediately obtain

$$\prod_{i=1}^k \binom{n + \alpha_i - 1}{\alpha_i} = \sum_{\lambda \vdash r} \sum_S \# \text{SSYT}(\lambda, S) \cdot \# \text{SSYT}(\lambda, \{1^{\alpha_1}, 2^{\alpha_2}, \dots, k^{\alpha_k}\}), \quad (4.4)$$

where the inner sum is over multisets S of size $r = \sum_{i=1}^k \alpha_i$ with entries in $[n]$.

5. Application: a new insertion on generalized permutations

Throughout this section, semistandard multiset tableaux are defined using **graded lexicographic order**; see Sections 2.4 and 2.5 for details.

Section 4 illustrated how RSK parallels the direct-sum decomposition of three distinct vector spaces. In each setting, the correspondence was facilitated by parameterizing a basis of the vector space by generalized permutations. More specifically, permutations, then words, and finally monomials were encoded as generalized permutations to which the RSK insertion procedure was applied (*cf.* Section 4.1–Section 4.3). In this section, we begin with an alternative encoding, which produces new combinatorial and enumerative results that parallel a different decomposition of the polynomial ring of Section 4.3.

5.1. *The correspondence*

Let $\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}$ be a generalized permutation from $[k]$ to $[n]$. We transform this into a generalized permutation from *multisets over $[k]$ to $[n]$* as follows. The columns of the generalized permutation are $\begin{pmatrix} M_i \\ i \end{pmatrix}$, where $M_i = \{a_j \mid b_j = i\}$ and $i \in [n]$. The columns are ordered so that the entries of the top row are weakly-increasing in graded lexicographic order and $i < i'$ whenever $M_i = M_{i'}$.

This encoding, together with Theorem 3.3, establishes the following result.

Proposition 5.1. *There is a one-to-one correspondence between generalized permutations $\begin{pmatrix} a_1 & a_2 & \dots & a_n \\ b_1 & b_2 & \dots & b_n \end{pmatrix}$, where a_1, \dots, a_n are multisets over $[k]$ and b_1, \dots, b_n are distinct elements of $[n]$, and pairs (S, T) satisfying: S and T are tableaux of the same shape; S is a standard Young tableau of size n ; and T is a semistandard multiset tableau of content $a_1 \uplus \dots \uplus a_n$.*

Example 5.2. Consider the generalized permutation from $[6]$ to $[5]$,

$$\begin{pmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 3 & 3 & 3 & 4 & 6 & 6 & 6 \\ 1 & 5 & 5 & 2 & 3 & 1 & 3 & 5 & 5 & 1 & 1 & 2 & 3 \end{pmatrix},$$

with which we associate the following generalized permutation whose top row consists of (possibly empty) multisets over $[6]$ and whose bottom row are the elements $\{1, 2, 3, 4, 5\}$, each appearing exactly once:

$$\left(\begin{array}{ccccc} \{\} & \{2, 6\} & \{2, 3, 6\} & \{1, 1, 3, 3\} & \{1, 3, 4, 6\} \\ 4 & 2 & 3 & 5 & 1 \end{array} \right).$$

This generalized permutation in turn corresponds to the following pair of tableaux:

$$\left(\begin{array}{|c|} \hline 4 \\ \hline 2 \\ \hline 1 & 3 & 5 \\ \hline \end{array}, \begin{array}{|c|} \hline 1346 \\ \hline 26 \\ \hline & 236 & 1133 \\ \hline \end{array} \right).$$

Example 5.3. Consider the case where $n = 3, k = 2$ and the generalized permutations are of the form $\begin{pmatrix} 1 & 1 & 2 \\ a & b & c \end{pmatrix}$. The pairs of tableaux under the correspondence of Proposition 5.1 are depicted in Fig. 1, whereas the pairs of tableaux under the usual RSK correspondence are depicted in Fig. 2.

5.2. *Special case: insertion on words*

In the special case where the top row of $\begin{pmatrix} a_1 & a_2 & \dots & a_r \\ b_1 & b_2 & \dots & b_r \end{pmatrix}$ satisfies $a_j = j$ for all j , we obtain the following correspondence.

	c = 1	c = 2	c = 3
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 1 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 2 & 112 & \\ \hline 1 & 3 & \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 11 \\ \hline 2 & 2 \\ \hline 1 & \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 2 & & 11 \\ \hline 1 & 3 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 3 & 12 \\ \hline 2 & 1 \\ \hline 1 & \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & & 1 \\ \hline 1 & 2 & 12 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 2 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 3 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 2 & 12 & \\ \hline 1 & 3 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 2 & \\ \hline 1 & 2 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 2 & & 1 & \\ \hline 1 & 3 & & 12 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 2 & 2 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 3 & 2 & \\ \hline 1 & 2 & 11 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 112 & \\ \hline 1 & 2 & \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 3 & & 11 & \\ \hline 1 & 2 & & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 2 & 3 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 2 & 2 & \\ \hline 1 & 3 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 12 & \\ \hline 1 & 2 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c c } \hline 1 & 2 & 3 & & 1 & 12 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 3 & 3 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 2 & 2 & \\ \hline 1 & 3 & 11 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 2 & 3 & 2 & 11 \\ \hline \end{array}$	$\begin{array}{ c c c c c } \hline 1 & 2 & 3 & & 112 \\ \hline \end{array}$

Fig. 1. The associated pair of tableaux by the correspondence of Proposition 5.1.

	c = 1	c = 2	c = 3
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 1 & c \end{pmatrix}$	$\begin{array}{ c c c } \hline 1 & 1 & 1 \\ \hline 1 & 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 1 & 1 & 2 \\ \hline 1 & 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 1 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 2 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline 1 & 1 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 1 & 2 & 2 \\ \hline 1 & 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 2 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 1 & 3 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 1 & 1 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 2 & \\ \hline 1 & 2 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 3 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 2 & 2 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline 1 & 2 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 2 & 2 & 2 \\ \hline 1 & 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 2 & 2 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 2 & 3 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline 1 & 3 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 2 & \\ \hline 2 & 2 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 2 & 3 & 3 & 1 & 1 & 2 \\ \hline \end{array}$
$\begin{pmatrix} 1 & 1 & 2 \\ 3 & 3 & c \end{pmatrix}$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 1 & 3 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 3 & 2 & \\ \hline 2 & 3 & 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 3 & 3 & 3 & 1 & 1 & 2 \\ \hline \end{array}$

Fig. 2. The associated pair of tableaux by the usual RSK correspondence.

Corollary 5.4. *There is a one-to-one correspondence between words of length k with entries in $[n]$ and pairs (S, T) satisfying: S and T are tableaux of the same shape; S is a standard Young tableau of size n ; T is a standard multiset tableau of content $[k]$.*

Example 5.5. The generalized permutation associated with the word 155231315 over the alphabet $[6]$ is

$$\left(\begin{array}{cccccc} \{\} & \{\} & \{4\} & \{5, 7\} & \{1, 6, 8\} & \{2, 3, 9\} \\ 4 & 6 & 2 & 3 & 1 & 5 \end{array} \right),$$

which corresponds to the pair

	$b = 1$	$b = 2$	$b = 3$	$b = 4$
$a = 1$	$\begin{array}{ c c } \hline 1 & 1 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 1 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$
$a = 2$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 2 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 2 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$
$a = 3$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 2 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$
$a = 4$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 2 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 1 & 1 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$

Fig. 3. The pair of tableaux associated with the word ab by the usual RSK correspondence.

	$b = 1$	$b = 2$	$b = 3$	$b = 4$
$a = 1$	$\begin{array}{ c c c } \hline 2 & & 12 \\ \hline 1 & 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 4 \\ \hline 1 & 2 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 2 & 4 \\ \hline 1 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c c } \hline 2 & & 1 \\ \hline 1 & 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$
$a = 2$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 2 & 1 \\ \hline 1 & 4 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 12 \\ \hline 1 & 2 & 4 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 1 \\ \hline 1 & 2 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 1 \\ \hline 1 & 2 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$
$a = 3$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 2 & 1 \\ \hline 1 & 3 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 3 & 1 \\ \hline 1 & 2 \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 12 \\ \hline 1 & 2 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 2 & 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$
$a = 4$	$\begin{array}{ c c } \hline 2 & 2 \\ \hline 1 & 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c } \hline 3 & 2 \\ \hline 1 & 2 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c } \hline 4 & 2 \\ \hline 1 & 2 & 3 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$	$\begin{array}{ c c c c } \hline 1 & 2 & 3 & 4 \\ \hline \end{array} \quad \begin{array}{ c c } \hline & \\ \hline \end{array}$

Fig. 4. The pair of tableaux associated with the word ab by the correspondence of Corollary 5.4.

$$\left(\begin{array}{|c|c|c|} \hline 4 & & \\ \hline 2 & 6 & \\ \hline 1 & 3 & 5 \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline 168 & & \\ \hline 4 & 57 & \\ \hline & & 239 \\ \hline \end{array} \right).$$

Example 5.6. Consider the (relatively small) example where $n = 4$ and $k = 2$ so that there are 16 words of length 2 with entries in $\{1, 2, 3, 4\}$. Fig. 3 depicts the pairs of tableaux associated with these words by the usual RSK algorithm; and Fig. 4 depicts the pairs of tableaux associated with these words by the correspondence of Corollary 5.4.

5.3. Enumerative results

By restricting the correspondence of Proposition 5.1 to generalized permutations $\begin{pmatrix} a_1 & a_2 & \dots & a_n \\ b_1 & b_2 & \dots & b_n \end{pmatrix}$ satisfying $a_1 \uplus \dots \uplus a_n = \{1^{\alpha_1}, 2^{\alpha_2}, \dots, k^{\alpha_k}\}$, we obtain the following enumerative statement:

$$\prod_{i=1}^k \binom{n + \alpha_i - 1}{\alpha_i} = \sum_{\lambda \vdash n} \# \text{SYT}(\lambda) \cdot \# \text{SSMT} \left(\lambda, \{1^{\alpha_1}, 2^{\alpha_2}, \dots, k^{\alpha_k}\} \right).$$

Compare this with Equation (4.4), which is the enumerative statement that accompanies Theorem 3.3.

The analogous enumerative statement that accompanies Corollary 5.4 is the special case where $\alpha_i = 1$ for all i :

$$n^k = \sum_{\lambda \vdash n} \# \text{SYT}(\lambda) \cdot \# \text{SMT}(\lambda, k) . \tag{5.1}$$

Compare this with the enumerative statement obtained from the usual application of the RSK correspondence to words (see Section 4.2 and Equation (4.3)):

$$n^k = \sum_{\mu \vdash k} \sum_C \# \text{SSYT}(\mu, C) \cdot \# \text{SYT}(\mu), \tag{5.2}$$

where the inner sum is over all multisets C of size k with entries in $\{1, 2, \dots, n\}$.

5.4. Connections with representation theory

Consider the vector space $V^{\otimes k}$, where $V = \mathbb{C}^n$. This space admits an action of the symmetric group S_n as well as an action of the partition algebra $P_k(n)$. The S_n -action is obtained by identifying the symmetric group S_n with the subgroup of GL_n consisting of the permutation matrices and restricting the GL_n -action on $V^{\otimes k}$ defined in Section 4.2. The precise definition of the $P_k(n)$ -action is not necessary here, so we refer the interested reader to [12, Eq. (1.3.4)]. More information on the partition algebra $P_k(n)$ is presented in Section 6.

These two actions are closely related when $n \geq 2k$: the algebra of linear endomorphisms of $V^{\otimes k}$ that commute with the S_n -action is isomorphic to $P_k(n)$; and conversely, the algebra of linear endomorphisms of $V^{\otimes k}$ that commute with the $P_k(n)$ -action is isomorphic to $\mathbb{C}S_n$ [21]. That is, provided that $n \geq 2k$,

$$\text{End}_{S_n}(V^{\otimes k}) \cong P_k(n) \quad \text{and} \quad \text{End}_{P_k(n)}(V^{\otimes k}) \cong \mathbb{C}S_n.$$

Consequently, $V^{\otimes k}$ admits a direct sum decomposition whose summands are tensors of a simple S_n -module and a simple $P_k(n)$ -module (see, for instance, [18, Theorem 3.2.2] or [7, Theorem 8.3.18]):

$$V^{\otimes k} \cong \bigoplus_{\substack{\lambda \vdash n \\ |\bar{\lambda}| \leq k}} S^\lambda \otimes V_{P_k(n)}^{\bar{\lambda}},$$

where S^λ is the simple S_n -module indexed by λ and $V_{P_k(n)}^{\bar{\lambda}}$ is the simple $P_k(n)$ -module indexed by $\bar{\lambda}$ (recall from Section 2.1 that $\bar{\lambda}$ is obtained from λ by deleting the first row). Since the dimension of S^λ is the number of standard tableaux of shape λ and the dimension of $V_{P_k(n)}^{\bar{\lambda}}$ is the number of standard multiset tableaux of shape λ and content $[k]$, we immediately obtain the enumerative result in Equation (5.1).

As pointed out in the introduction, this combinatorial interpretation for the dimension is different from that of [16,26], which makes use of vacillating tableaux instead of multiset tableaux. It is more closely aligned with the results in [2,3,29,15].

6. Application: diagram algebras

Throughout this section, standard multiset tableaux are defined using **last letter order**; see Sections 2.4 and 2.5 for details.


For any parameter n and positive integer k , the partition algebra, $P_k(n)$, is defined as the complex vector space with basis given by the set partitions on two disjoint sets $[k] \cup [\bar{k}] = \{1, 2, \dots, k\} \cup \{\bar{1}, \bar{2}, \dots, \bar{k}\}$:

$$P_k(n) = \text{span}_{\mathbb{C}}\{\pi \mid \pi \vdash [k] \cup [\bar{k}]\}.$$

Although we do not define the product here, as we will not use it explicitly, we remark that the dependency of the algebra on n arises when we multiply the set partitions [18].

A *diagram* is a graphical representation of a set partition of the set $[k] \cup [\bar{k}]$: the vertex set of the graph is $[k] \cup [\bar{k}]$ arranged in two horizontal rows, where the top row is labelled by $1, 2, \dots, k$ and the bottom row are labelled by $\bar{1}, \bar{2}, \dots, \bar{k}$; and there is a path connecting two vertices if and only if they belong to the same block of the set partition. Note that there is more than one graph that represents a set partition, but this is immaterial to the following. In our examples, we will connect vertices in the same block with a cycle.

Example 6.1. The set partition $\pi = \{\{1, 2, 4, \bar{2}, \bar{5}\}, \{3\}, \{5, 6, 7, \bar{3}, \bar{4}, \bar{6}, \bar{7}\}, \{8, \bar{8}\}, \{\bar{1}\}\}$ is represented by the following diagram:



The partition algebra $P_k(n)$ is semisimple whenever the parameter $n \notin \{0, 1, \dots, 2k - 2\}$ in which case the irreducible representations are indexed by partitions λ with $0 \leq |\lambda| \leq k$ [27]. We assume throughout that $n \geq 2k$, so that $P_k(n)$ is semisimple and isomorphic to $\text{End}_{S_n}(V^{\otimes k})$ as explained in Section 5.4.

In [16,26], the authors introduce RSK-type algorithms between partition algebra diagrams and pairs of paths in the Bratteli diagram of the partition algebras; in [16] these paths are called vacillating tableaux. In [2], the authors define a bijection between vacillating tableaux and standard multiset tableaux. In this section we provide a different bijection from partition algebra diagrams to standard multiset tableaux.

This algorithm not only encodes the representation theory of the partition algebra, in the sense that the tableaux of shape λ index an irreducible representation associated with λ , but it also encodes the representation theory of subalgebras of the partition algebra when we restrict the set of diagrams considered. This allows us to obtain enumerative results for representations of various diagram algebras using standard multiset tableaux.

In this section, we only consider centralizer algebras acting on $V^{\otimes k}$, but this construction indicates that more general diagram algebras are also of interest. If instead one considers the centralizer algebras acting on the polynomial ring $\mathbb{C}[X]$ (as described in Section 4.3), then the corresponding diagrams would have repeated entries and the dimensions of the irreducible representations will be subsets of semistandard multiset tableaux. Currently little is known about these centralizer algebras, see for instance [28,30].

6.1. The correspondence

A block in a set partition π is called *propagating* if it contains vertices in both $[k]$ and $[\bar{k}]$. For example, $\{1, 2, 4, \bar{2}, \bar{5}\}$ is a propagating block. A block is called *non-propagating* otherwise. The number of propagating blocks in π is called the *propagating number*. We denote the propagating number by $\text{pr}(\pi)$. For example, the set partition π in Example 6.1 has $\text{pr}(\pi) = 3$.

Let $\pi = \{\pi_1, \pi_2, \dots, \pi_r\}$ be a set partition of $[k] \cup [\bar{k}]$. We associate with π a pair (T, S) of standard multiset tableaux as follows. To begin,

- let $\pi_{j_1}, \pi_{j_2}, \dots, \pi_{j_p}$ denote the propagating blocks of π ordered so that $\pi_{j_1}^+ < \dots < \pi_{j_p}^+$ in the last letter order, where $\pi_j^+ = \pi_j \cap [k]$;
- let $\sigma_{i_1}, \dots, \sigma_{i_a} \subseteq [k]$ denote the non-propagating blocks contained in $[k]$ and ordered so that $\sigma_{i_1} < \dots < \sigma_{i_a}$ in the last letter order;
- let $\tau_{i_1}, \dots, \tau_{i_b} \subseteq [\bar{k}]$ denote the non-propagating blocks contained in $[\bar{k}]$ and ordered so that $\tau_{i_1} < \dots < \tau_{i_b}$ in the last letter order.

Let (P, Q) denote the pair of standard multiset tableaux obtained by applying the RSK algorithm to the generalized permutation

$$\begin{pmatrix} \pi_{j_1}^+ & \pi_{j_2}^+ & \cdots & \pi_{j_p}^+ \\ \pi_{j_1}^- & \pi_{j_2}^- & \cdots & \pi_{j_p}^- \end{pmatrix},$$

where $\pi_j^+ = \pi_j \cap [k]$ and $\pi_j^- = \pi_j \cap [\bar{k}]$. Let T be the tableau obtained from P by adjoining a row containing $n - p - b$ empty cells followed by cells labelled $\tau_{i_1}, \dots, \tau_{i_b}$. Let S be the tableau obtained from Q by adjoining a row containing $n - p - a$ empty cells followed by cells labelled $\sigma_{i_1}, \dots, \sigma_{i_a}$.


Example 6.2. Let $\pi = \{\{2, 3, 4, \overline{4}, \overline{5}\}, \{5, \overline{2}, \overline{3}\}, \{1, 6, \overline{7}, \overline{8}\}, \{7, 8\}, \{9, \overline{6}\}, \{\overline{1}\}, \{\overline{9}\}\} \in P_9(18)$. The non-propagating blocks are $\{\overline{1}\}, \{\overline{9}\}$ and $\{7, 8\}$, and the generalized permutation constructed from the propagating blocks is

$$\left(\begin{array}{cccc} \{2, 3, 4\} & \{5\} & \{1, 6\} & \{9\} \\ \{\overline{4}, \overline{5}\} & \{\overline{2}, \overline{3}\} & \{\overline{7}, \overline{8}\} & \{\overline{6}\} \end{array} \right).$$

Apply the RSK algorithm to obtain the following pair of multiset tableaux:

$$P = \begin{array}{|c|c|} \hline \overline{45} & \overline{78} \\ \hline \overline{23} & \overline{6} \\ \hline \end{array}, \quad Q = \begin{array}{|c|c|} \hline 5 & 9 \\ \hline 234 & 16 \\ \hline \end{array}.$$

Finally, adjoin a new row to P and a new row to Q containing the non-propagating blocks so that the resulting tableaux are of size $n = 18$:



Theorem 6.3. Let $n \geq 2k$. The set partitions of $[k] \cup [\overline{k}]$ are in bijection with pairs (T, S) of standard multiset tableaux satisfying: T and S are of the same shape λ , where λ is a partition of n ; T has content $[\overline{k}]$; and S has content $[k]$.

Proof. Let $\pi = \{\pi_1, \pi_2, \dots, \pi_r\} \vdash [k] \cup [\overline{k}]$ and let (T, S) denote the tableaux constructed by the above procedure. Notice that the propagation number of π is at most k , and since $n \geq 2k$, there will be at least k empty cells in the first row of T and in the first row of S guaranteeing that both are semistandard multiset tableaux. In addition, the cells of T are filled with the blocks of a set partition of $[\overline{k}]$ and the cells of S with the blocks of a set partition of $[k]$, and so there are no repetitions in T or S . Hence both tableaux are standard multiset tableaux.

Observe that we can reconstruct the set partition π from (T, S) : non-propagating blocks are the elements of the first rows; and the inverse of the RSK algorithm recovers the generalized permutation defined by the propagating blocks of π . \square

Since the number of set partitions of a set of cardinality $2k$ is equal to the Bell number $B(2k)$ [19, A000110, A020557], Theorem 6.3 implies that for $n \geq 2k$,

$$B(2k) = \sum_{\lambda \vdash n} \# \text{SMT}(\lambda, k)^2. \tag{6.1}$$

Example 6.4. Fig. 5 depicts the correspondence of Theorem 6.3 for the 15 diagrams for $P_2(4)$.





Fig. 5. The correspondence from Theorem 6.3 for the 15 diagrams for $P_2(4)$.

For any diagram $\pi \vdash [k] \cup [\bar{k}]$, we define $\text{flip}(\pi)$ to be the reflection of π along its horizontal axis. If



The properties in the next proposition follow directly from the RSK algorithm.

Proposition 6.5. Let $\pi \vdash [k] \cup [\bar{k}]$.

- (a) If π inserts to (T, S) with S and T of shape λ , then $|\bar{\lambda}| = \text{pr}(\pi)$.
- (b) If π inserts to (T, S) , then $\text{flip}(\pi)$ inserts to (S, T) .

6.2. Restriction to subalgebras

There are other bijections between partition algebra diagrams and pairs of standard multiset tableaux, but an important aspect of the algorithm in this paper is that it is compatible with the (representation theory) restriction to many prominent subalgebras of $P_k(n)$. More precisely, we will see that this single procedure captures the combinatorics of the representation theory of all these subalgebras.

For instance, for an integer r with $0 \leq r \leq k$ the subspace spanned by the set partitions with propagating number at most r is a subalgebra of $P_k(n)$ and the irreducible representations of this subalgebra are indexed by partitions of size less than or equal to r . By Proposition 6.5, a refinement of Equation (6.1) states




Fig. 6. Examples of types of set partition diagrams.

Table 1
Subalgebras of the partition algebra $P_k(n)$.

Subalgebra	Diagrams spanning the subalgebra	Dimension
Partition algebra $P_k(n)$	all diagrams	$B(2k)$
Group algebra of symmetric group $\mathbb{C}S_k$	permutations	$k!$
Brauer algebra $B_k(n)$ [4,38]	perfect matchings	$(2k - 1)!!$
Rook algebra $R_k(n)$ [36]	partial permutations	$\sum_{i=0}^k \binom{k}{i}^2 i!$
Rook-Brauer algebra $RB_k(n)$ [14,25]	matchings	$\sum_{i=0}^k \binom{2k}{2i} (2i - 1)!!$
Temperley–Lieb algebra $TL_k(n)$ [37,20,39,24]	planar perfect matchings	$\frac{1}{k+1} \binom{2k}{k}$
Motzkin algebra $M_k(n)$ [1]	planar matchings	$\sum_{i=0}^k \frac{1}{i+1} \binom{2i}{i} \binom{2k}{2i}$
Planar rook algebra $PR_k(n)$ [8]	planar partial permutations	$\binom{2k}{k}$
Planar algebra $PP_k(n)$ [21]	planar diagrams	$\frac{1}{2k+1} \binom{4k}{2k}$

$$\#\left\{ \pi \vdash [k] \cup [\bar{k}] \mid \text{pr}(\pi) \leq r \right\} = \sum_{\substack{\lambda \vdash n \\ |\bar{\lambda}| \leq r}} \# \text{SMT}(\lambda, k)^2 . \tag{6.2}$$

6.2.1. Definition of the subalgebras

We introduce some terminology that will make it easier to define the subalgebras. See Fig. 6 for examples of the types of diagrams that we define below. A set partition π is called *planar* if it can be represented as a graph without edge crossings inside the rectangle formed by its vertices. A set partition is called a *matching* if all its blocks are of size at most 2. We call a set partition a *perfect matching* if all its blocks are of size 2. The number of perfect matchings of $2n$ elements is equal to $(2n - 1)!! = (2n - 1)(2n - 3) \cdots (1)$. A perfect matching, where each block contains an element in $[k]$ and an element in $[\bar{k}]$ is a *permutation*. A set partition is a *partial permutation* if all its blocks have size one or two and every block of size two is propagating.

Table 1 summarizes the definitions of the subalgebras that we work with. In [15], the authors construct the irreducible representations of these subalgebras using standard multiset tableaux (which they call *set-partition tableaux*) and compute their characters.

Table 2

Dimensions and index sets for irreducible representations of certain subalgebras A_k of the partition algebra $P_k(n)$ [15]. We highlight that the bottom four subalgebras are all planar and that their irreducible representations are indexed by partitions having a single part.

A_k	Index set for irreducibles	Dimension of irreducible $V_{A_k}^\lambda$
$P_k(n)$	$\{\lambda \mid \lambda \vdash m, 0 \leq m \leq k\}$	$f^\lambda \sum_{i= \lambda }^k \binom{k}{i} \left\{ \begin{matrix} i \\ \lambda \end{matrix} \right\} B(k-i)$
$\mathbb{C}S_k$	$\{\lambda \mid \lambda \vdash k\}$	f^λ
$B_k(n)$	$\{\lambda \mid \lambda \vdash k-2r, 0 \leq 2r \leq k\}$	$f^\lambda \binom{k}{ \lambda } (k- \lambda -1)!!$
$R_k(n)$	$\{\lambda \mid \lambda \vdash m, 0 \leq m \leq k\}$	$f^\lambda \binom{k}{ \lambda }$
$RB_k(n)$	$\{\lambda \mid \lambda \vdash m, 0 \leq m \leq k\}$	$f^\lambda \binom{k}{ \lambda } \sum_{i=0}^{(k- \lambda)/2} \binom{k- \lambda }{2i} (2i-1)!!$
$TL_k(n)$	$\{(k-2r) \mid 0 \leq 2r \leq k\}$	$\binom{k}{(k-m)/2} - \binom{k}{(k-m)/2-1}$
$M_k(n)$	$\{(m) \mid 0 \leq m \leq k\}$	$\sum_{i=0}^{\lfloor (k-m)/2 \rfloor} \binom{k}{m+2i} \left(\binom{m+2i}{i} - \binom{m+2i}{i-1} \right)$
$PR_k(n)$	$\{(m) \mid 0 \leq m \leq k\}$	$\binom{k}{m}$
$PP_k(n)$	$\{(m) \mid 0 \leq m \leq k\}$	$\binom{2k}{k-m} - \binom{2k}{k-m-1}$

Their results provide a detailed study of the representation theory of these subalgebras from which we extract the information in Table 2.

6.2.2. Restricting the correspondence to the subalgebras

We characterize the standard multiset tableaux produced by the correspondence of Section 6.1 when restricted to the diagrams spanning one of the subalgebras A_k in Table 1. We denote this set by $SMT_{A_k}(\lambda)$.

A standard multiset tableau is *matching* if the first row contains sets of size less than or equal to 2 and all other rows contain only sets of size 1. In Lemma 6.6, we show these are the multiset tableaux that correspond to matching diagrams by our insertion algorithm.

Two sets S and S' are *non-crossing* if there do not exist elements $a, b \in S$ and $c, d \in S'$ such that either $a < c < b < d$ or $c < a < d < b$. We say that $c \in [k]$ is *between* a set S if there exist $a, b \in S$ such that $a < c < b$. We call a standard multiset tableau *planar* if it has two rows, if the sets in the first row are pairwise non-crossing, and if no element belonging to one of the sets in the second row is between any set in the tableau (apart from the set containing the element). In Lemma 6.6, we show these are the multiset tableaux that correspond to planar diagrams by our insertion algorithm.

Lemma 6.6. *Let k be any positive integer, λ a partition of an integer n with $n \geq 2k$, and A_k one of the subalgebras of $P_k(n)$ defined in Table 1. If we apply the insertion procedure of Theorem 6.3 to the diagrams spanning A_k , then the resulting standard multiset tableaux are characterized by the properties listed in Table 3.*

Proof. The case $A_k = P_k(n)$ follows from Theorem 6.3.

Table 3

Properties characterizing the standard multiset tableaux that belong to $SMT_{A_k}(\lambda)$. These are the tableaux produced by the correspondence of Section 6.1 when restricted to the diagrams spanning A_k .

A_k	Diagrams spanning A_k	Properties characterizing SMT_{A_k}	
		Sizes of entries in first row	Other properties
$P_k(n)$	all diagrams	—	—
$PP_k(n)$	planar diagrams	—	planar
CS_k	permutations	0	matching
$B_k(n)$	perfect matchings	0, 2	matching
$R_k(n)$	partial permutations	0, 1	matching
$RB_k(n)$	matchings	0, 1, 2	matching
$TL_k(n)$	planar perfect matchings	0, 2	matching & planar
$M_k(n)$	planar matchings	0, 1, 2	matching & planar
$PR_k(n)$	planar partial permutations	0, 1	matching & planar

For $A_k = PP_k(n)$, observe that if a set partition π is planar, then the propagating blocks $\pi_{i_1}, \dots, \pi_{i_p}$ are inserted in order as the blocks are non-crossing. This means that the shape of the insertion tableau T (as well as that of the recording tableau S) has at most two rows. Also notice that non-propagating blocks have to be non-crossing (since the diagram is planar) and these entries constitute the entries in the first row of T and S . Furthermore, propagating blocks in π correspond to entries in the second row of T and S . If a letter in the second row of T (respectively, S) is between another set in T (respectively, S), then the diagram π is not planar.

By the correspondence described in Section 6.1, a propagating block of a diagram π is a set of size 2 with one element in $[k]$ and one element in $[\bar{k}]$ if and only if the corresponding entries in the pair of standard multiset tableaux (T, S) have size one and appear in the second row or above in both T and S . The non-propagating blocks are all in the first row and, in a matching diagram, all of the blocks are of size less than or equal to 2. This implies that, if A_k is spanned by diagrams that are matching, then these diagrams insert to tableaux which are matching. Similarly, if the subalgebra is spanned by diagrams that are planar, then these diagrams insert to tableaux which are planar.

Since the non-empty sets that appear in the first row of T and the first row of S correspond to the non-propagating blocks of the set partition, we obtain the restrictions on sizes of the sets appearing in the first row. For instance, if $A_k = CS_k$, then there are no non-propagating blocks, and so the first row of S and of T contain only empty sets. If A_k is $R_k(n)$ or $PR_k(n)$, then the non-propagating blocks are all of size 1. If A_k is $B_k(n)$ or $TL_k(n)$, then the blocks (and hence the non-propagating blocks) are all of size 2. If A_k is $RB_k(n)$ or $M_k(n)$, then non-propagating blocks are of size at most 2. \square

Example 6.7. In Table 4, we give examples of the tableaux described in Lemma 6.6 for $k = 9$ and n sufficiently large.

In addition, we obtain the following corollary of Theorem 6.3 and Lemma 6.6.

Corollary 6.8. *If $n \geq 2k$, then for each subalgebra A_k of the partition algebra $P_k(n)$ described in Table 1, we have*

Table 4
Examples of the tableaux in Lemma 6.6 for $k = 9$ and $n \geq 2k$.

Algebra	Diagram	Insertion tableaux (T)	Recording tableaux (S)
$P_k(n)$		$\begin{array}{cccccccc} \overline{26} & & & & & & & \\ 345 & \overline{9} & & & & & & \\ \hline & & & \cdots & & \overline{1} & \overline{78} & \end{array}$	$\begin{array}{cccccccc} 567 & & & & & & & \\ 124 & 89 & & & & & & \\ \hline & & & \cdots & & & & 3 \end{array}$
CS_k		$\begin{array}{cccccccc} \overline{9} & & & & & & & \\ 4 & \overline{6} & \overline{7} & & & & & \\ \hline \overline{1} & \overline{2} & \overline{3} & \overline{5} & \overline{8} & & & \cdots \end{array}$	$\begin{array}{cccccccc} 9 & & & & & & & \\ 3 & 4 & 8 & & & & & \\ \hline 1 & 2 & 5 & 6 & 7 & & & \cdots \end{array}$
$B_k(n)$		$\begin{array}{cccccccc} \overline{8} & & & & & & & \\ 4 & \overline{7} & & & & & & \\ \hline \overline{1} & \overline{5} & & & \cdots & & \overline{23} & \overline{69} \end{array}$	$\begin{array}{cccccccc} 5 & & & & & & & \\ 3 & 9 & & & & & & \\ \hline 2 & 8 & & & \cdots & & 14 & 67 \end{array}$
$R_k(n)$		$\begin{array}{cccccccc} \overline{8} & & & & & & & \\ 4 & & & & & & & \\ \hline 2 & \overline{3} & \overline{7} & & & \cdots & \overline{1} & \overline{5} & \overline{6} & \overline{9} \end{array}$	$\begin{array}{cccccccc} 6 & & & & & & & \\ 5 & & & & & & & \\ \hline 3 & 7 & 9 & & & \cdots & 1 & 2 & 4 & 8 \end{array}$
$RB_k(n)$		$\begin{array}{cccccccc} \overline{3} & & & & & & & \\ \overline{1} & \overline{5} & & & & & & \\ \hline & & & \cdots & & \overline{46} & \overline{27} & \overline{8} & \overline{9} \end{array}$	$\begin{array}{cccccccc} 4 & & & & & & & \\ 1 & 7 & & & & & & \\ \hline & & & \cdots & & \overline{23} & \overline{6} & \overline{5} & \overline{8} & 9 \end{array}$
$TL_k(n)$		$\begin{array}{cccccccc} \overline{1} & \overline{2} & \overline{7} & & & & & \\ \hline & & & \cdots & & \overline{45} & \overline{36} & \overline{89} \end{array}$	$\begin{array}{cccccccc} 5 & 6 & 9 & & & & & \\ \hline & & & \cdots & & \overline{12} & \overline{34} & \overline{78} \end{array}$
$M_k(n)$		$\begin{array}{cccccccc} \overline{4} & \overline{5} & \overline{6} & \overline{8} & & & & \\ \hline & & & \cdots & & \overline{2} & \overline{13} & \overline{7} & \overline{9} \end{array}$	$\begin{array}{cccccccc} 5 & 7 & 8 & 9 & & & & \\ \hline & & & \cdots & & \overline{12} & \overline{34} & \overline{6} \end{array}$
$PR_k(n)$		$\begin{array}{cccccccc} \overline{1} & \overline{2} & \overline{4} & \overline{5} & \overline{7} & \overline{8} & & \\ \hline & & & \cdots & & \overline{3} & \overline{6} & \overline{9} \end{array}$	$\begin{array}{cccccccc} 2 & 4 & 5 & 7 & 8 & 9 & & \\ \hline & & & \cdots & & \overline{1} & \overline{3} & \overline{6} \end{array}$
$PP_k(n)$		$\begin{array}{cccccccc} \overline{1} & \overline{5} & \overline{6} & \overline{7} & \overline{9} & & & \\ \hline & & & \cdots & & \overline{234} & \overline{8} & \end{array}$	$\begin{array}{cccccccc} 134 & 5 & 7 & & & & & \\ \hline & & & \cdots & & \overline{2} & \overline{6} & \overline{89} \end{array}$

$$\dim(A_k) = \sum_{\lambda \vdash n} (\# \text{SMT}_{A_k}(\lambda))^2,$$

where the dimension of A_k is also given in Table 1.

6.3. From standard multiset tableaux to Bratteli diagrams

Let A_k denote one of the subalgebras from Lemma 6.6. We establish a bijection between the standard multiset tableaux for A_k and the paths in the Bratteli diagram for $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$.

A *Bratteli diagram* associated to a tower of algebras $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$ is an infinite \mathbb{N} -graded graph defined as follows. The vertices at level $k \in \mathbb{N}$ are in bijection

Table 5
Branching rules for various subalgebras of the partition algebra.

A_k	Branching rule for λ of A_k to μ of A_{k-1}	Reference
$P_k(n)$ $PP_k(n)$	remove 1 or 0 cells from λ to get τ , then add 1 or 0 cells to τ to get μ	[11, Equation (1.4.1)] [18, Section 1]
CS_k	remove a cell from λ to get μ	
$B_k(n)$ $TL_k(n)$	add or remove a cell from λ to get μ	[38, p. 192] [20, p. 19]
$R_k(n)$ $PR_k(n)$	remove 1 or 0 cells from λ to get μ	[12, Sec. 3.1] [8, Equation (4)]
$RB_k(n)$ $M_k(n)$	add or remove 1 or 0 cells from λ to get μ	[14, Equation (3.4)] [1, Equation (3.12)]

with the isomorphism classes of the irreducible representations of A_k ; if the irreducible representations are parameterized by some index set, then we label the vertices by the elements of the index set. Note that it is possible that vertices at different levels carry the same label (this happens for some of the index sets listed in Table 2), but the associated representations are different. The edges in the graph connect vertices of level k with vertices at level $k + 1$: the number of edges from the vertex associated with an irreducible A_k -representation V to the vertex associated with an irreducible A_{k+1} -representation V' is the multiplicity of V in the restriction of V' to A_k .

In all the examples we consider, there is exactly one irreducible A_0 -representation and it is of dimension 1. It follows from an induction argument that the dimension of an irreducible representation V is equal to the number of paths in the Bratteli diagram from the unique level-0 vertex to the vertex associated with V .

Example 6.9. Young’s lattice is an example of a Bratteli diagram for the tower of symmetric group algebras $\mathbb{C}S_0 \subseteq \mathbb{C}S_1 \subseteq \mathbb{C}S_2 \subseteq \dots$. Indeed, recall that there is exactly one edge in Young’s lattice $\mu \rightarrow \lambda$ if and only if μ is obtained from λ by removing a corner cell. And, the multiplicity of the irreducible S_k -representation indexed by μ in the restriction to $\mathbb{C}S_k$ of the irreducible S_{k+1} -representation indexed by λ is equal to 1 if $\mu \rightarrow \lambda$ in Young’s lattice and is equal to 0 otherwise.

By *branching rule*, we mean any combinatorial description of the edge multiplicities in the Bratteli diagram in terms of the index sets of the irreducible representations. Table 5 summarizes the branching rule for various subalgebras of the partition algebra, where the index sets for the irreducible representations are given in Table 2.

Remark 6.10. A proof that the planar algebra $PP_k(n)$ is isomorphic to the Temperley–Lieb algebra $TL_{2k}(n)$ can be found in [18, Section 1]. Consequently, the branching rule for $PP_k(n)$ is obtained by a repeated application of the branching rule for $TL_{2k}(n)$.

Paths in the Brauer algebra Bratteli diagram are often called updown tableaux or oscillating tableaux in the literature; see [16] and the references therein.

Proposition 6.11. *Let k be a positive integer and λ a partition of $n \geq 2k$. There is a bijection*

$$\phi: \text{SMT}(\lambda, k) \rightarrow \left\{ (S, \tau) \mid S \in \text{SMT}(\mu, k-1), \tau \rightarrow \mu, \tau \rightarrow \lambda \right\}.$$

Proof. Let T be an element of $\text{SMT}(\lambda, k)$. Let A denote the unique set appearing in T containing k . Because we are using last letter order, the cell labelled by A is a corner cell. Let T' be the tableau obtained from T by deleting A . If $A \neq \{k\}$, then let S be the tableau obtained by inserting $A \setminus \{k\}$ in the second row of T' using the RSK insertion procedure. If $A = \{k\}$, then let S be the tableau obtained from T' by adding a blank cell at the beginning of its first row. Set $\phi(T) = (S, \tau)$, where τ is the shape of T' . Note that $\tau \rightarrow \lambda$.

Conversely, let (S, τ) be such that $S \in \text{SMT}(\mu, k-1)$, $\tau \rightarrow \mu$ and $\tau \rightarrow \lambda$. If the unique cell in μ/τ is in the first row, then let T be the tableau obtained from S by removing a blank cell from the first row of S , and then adding a cell labelled $\{k\}$ at the cell in λ/τ .

Otherwise, let S' denote the tableau obtained from S by deleting its first row. Reverse the RSK insertion procedure starting with the cell μ/τ to produce a tableau T' and a set A' such that inserting A' into T' produces S' . Let T be the tableau obtained from T' by adjoining the first row of S and adding a new cell labelled $A' \cup \{k\}$ at the cell in λ/τ . \square

This correspondence is particularly useful because it respects the properties characterizing the tableaux in $\text{SMT}_{A_k}(\lambda)$ (see Table 3).

Example 6.12. Consider the following tableau in $\text{SMT}((n-3, 2, 1), 9)$,

$$T = \begin{array}{|c|c|c|c|c|c|c|c|} \hline 7 & & & & & & & \\ \hline 5 & 8 & & & & & & \\ \hline & & & \dots & & 12 & 46 & 39 \\ \hline \end{array},$$

in particular, it is an element of $\text{SMT}_{B_9(n)}((n-3, 2, 1))$. To compute $\phi(T) = (S, \tau)$, we remove the cell labelled $\{3, 9\}$ and insert $\{3\}$ in the second row, obtaining

$$S = \begin{array}{|c|c|c|c|c|c|c|} \hline 7 & & & & & & \\ \hline 5 & & & & & & \\ \hline 3 & 8 & & & & & \\ \hline & & & \dots & & 12 & 46 \\ \hline \end{array}$$

where $S \in \text{SMT}_{B_8(n)}((n-4, 2, 1, 1))$ and $\tau = (n-4, 2, 1)$. Proposition 6.11 also states that T can be recovered from S and the partitions $\tau = (n-4, 2, 1)$ and $\lambda = (n-3, 2, 1)$.

Now we are ready for the main result of this section, which states that the standard multiset tableaux in $\text{SMT}_{A_k}(\lambda)$ encode the branching rule for the subalgebra A_k .

Theorem 6.13. *Let A_k be any of the subalgebras of $P_k(n)$ defined in Table 1 with $n \geq 2k$, and $\lambda, \mu \vdash n$.*

- (1) *If $T \in \text{SMT}_{A_k}(\lambda)$ and $\phi(T) = (S, \tau)$, then $S \in \text{SMT}_{A_{k-1}}(\mu)$.*
- (2) *For each $S \in \text{SMT}_{A_{k-1}}(\mu)$, the number of $T \in \text{SMT}_{A_k}(\lambda)$ such that $\phi(T) = (S, \tau)$ for some partition τ is equal to the number of edges from $\bar{\mu}$ to $\bar{\lambda}$ in the Bratteli diagram for the tower of algebras $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$.*

Proof. (1) We first verify that if T is planar (respectively, matching) and $\phi(T) = (S, \tau)$, then S is also planar (respectively, matching).

Let T be a planar tableau and let A denote the set in T that contains k . If $A = \{k\}$, then S is obtained from T by deleting the cell labelled A and adding a blank cell to the first row. Since T is planar, all the sets appearing in T satisfy the conditions in the definition of planar, and so S is also planar.

Suppose $A \neq \{k\}$ and that A appears in the second row of T . Let a be the largest element in $A \setminus \{k\}$. Then $a + 1, a + 2, \dots, k - 1$ must be in the first row of T (otherwise, these elements are between A , contradicting that T is planar). Therefore, $A \setminus \{k\}$ is greater than all the sets in the second row of T in the last letter order. Thus, S is obtained from T by deleting k , and it follows that S is planar.

Suppose $A \neq \{k\}$ and that A appears in the first row of T . If one of the sets in the second row of T contains $c \in [k]$ satisfying $\max(A \setminus \{k\}) < c < k$, then c is between A , which contradicts the hypothesis that T is planar. Hence, $A \setminus \{k\}$ is greater than all sets appearing in the second row of T , and so S is obtained from T by deleting the cell labelled A and appending $A \setminus \{k\}$ to the second row. To prove that S is planar, it remains to show that no element of $A \setminus \{k\}$ is between any other set in S .

Suppose there exists $b \in A \setminus \{k\}$ that is between some set B . Then there exist $a, c \in B$ such that $a < b < c < k$. If B is in the first row of S , then A and B are crossing, which contradicts the fact that the sets in the first row of T are pairwise non-crossing. If B is in the second row of S , then c is between A , which contradicts the fact that no element belonging to the second row of T is between any set in the tableau. Hence, S is also planar.

Let T be a matching tableau and let A denote the set in T that contains k . If $A = \{k\}$, then S is obtained from T by deleting the cell labelled A and adding a blank cell to the first row. Since T is matching, all the sets appearing in T satisfy the conditions in the definition of matching, and so S is also matching.

Suppose $A \neq \{k\}$. Then A is a set of size 2, say $A = \{a, k\}$, and it appears in the first row of T . Then S is obtained from T by deleting A and inserting $\{a\}$ into the second row using the RSK algorithm. Thus, all sets in S are of size at most 2, and the sets of size 2 belong to its first row. Hence, S is matching.

Finally, note that if the sizes of the sets in the first row of T are constrained to be in some set Σ that contains 0, then the same is true for the sets in the first row of S :

indeed, S is obtained from T by deleting a set and either adding a blank cell in the first row or by adding a non-empty cell in some row besides the first row.

(2) Next we check that for a fixed $S \in \text{SMT}_{A_{k-1}}(\mu)$,

$$\#\{T \in \text{SMT}_{A_k}(\lambda) \mid \phi(T) = (S, \tau) \text{ for some } \tau\}$$

is the multiplicity of $V_{A_{k-1}}^{\bar{\mu}}$ in the restriction of $V_{A_k}^{\bar{\lambda}}$ to A_{k-1} described in Table 5. We will do this on a case by case basis for each of the four pairs of subalgebras. Throughout this proof, let $S \in \text{SMT}_{A_{k-1}}(\mu)$ and let $T \in \text{SMT}_{A_k}(\lambda)$ be such that $\phi(T) = (S, \tau)$ for some τ .

Let A_k be either $R_k(n)$ or $PR_k(n)$. Note that all the sets appearing in S are of size at most 1, and S is obtained from T by deleting the cell labelled k and adding an empty cell to the first row. The cell labelled k is removed from the first row if and only if $\bar{\lambda} = \bar{\mu}$. And if the cell is removed from some other row, then $\bar{\mu}$ is obtained from $\bar{\lambda}$ by deleting a cell. This agrees with the branching rule in Table 5 with λ replaced by $\bar{\lambda}$ and μ replaced by $\bar{\mu}$.

Let A_k be $B_k(n)$ or $TL_k(n)$. The non-empty sets in the first row of S are all of size 2 and the sets in the other rows are all of size 1. If the set of T containing k is $\{k\}$, then it appears in the second row or above of T . In this case, S is obtained from T by deleting the cell labelled $\{k\}$ and adding an empty cell in the first row. Thus, μ is obtained from λ by moving a cell to the first row, or in other words, $\bar{\mu}$ is obtained from $\bar{\lambda}$ by deleting a cell.

Otherwise, the set containing k is $\{a, k\}$, for some a , and it appears at the end of the first row. Then S is obtained from T by deleting $\{a, k\}$ and inserting $\{a\}$ in the second row. Thus, μ is obtained from λ by removing a cell from the first row and adding a cell to some other row. In other words, $\bar{\mu}$ is obtained from $\bar{\lambda}$ by adding a cell. This agrees with the branching rule in Table 5 with λ replaced by $\bar{\lambda}$ and μ replaced by $\bar{\mu}$.

Let A_k be $RB_k(n)$ or $M_k(n)$. The non-empty sets in the first row of S are all of size 1 or 2 and those in the other rows are all of size 1. Hence, the set in T containing k is either $\{k\}$ or $\{a, k\}$ for some a . In the first case, S is obtained from T by deleting the cell labelled $\{k\}$ and adding an empty cell to the first row, from which it follows that we have $\bar{\lambda} = \bar{\mu}$ (when $\{k\}$ is in the first row of T) or $\bar{\mu} \rightarrow \bar{\lambda}$ (otherwise). In the second case, S is obtained from T by deleting the cell labelled $\{a, k\}$ at the end of the first row and using the RSK insertion procedure to insert $\{a\}$ into the second row. Thus, μ is obtained from λ by moving a cell from the first row to some other row. In other words, $\bar{\mu}$ is obtained from $\bar{\lambda}$ by adding a cell. This agrees with the branching rule in Table 5 with λ replaced by $\bar{\lambda}$ and μ replaced by $\bar{\mu}$.

Let A_k be either $P_k(n)$ or $PP_k(n)$. Let τ denote the shape of the tableau obtained from T by deleting the set containing k . If the set containing k is $\{k\}$, then S is obtained from T by deleting the cell labelled $\{k\}$ and adding an empty cell to the first row. In

this case, μ is obtained from λ by moving a cell to the first row. If the moved cell came from the first row, then $\overline{\mu} = \overline{\lambda}$, and otherwise $\overline{\mu}$ is obtained from $\overline{\lambda}$ by deleting a cell.

If the set containing k is not $\{k\}$, then S is the tableau obtained from T by deleting the cell containing k and inserting a set in the second row using the RSK insertion procedure. Thus, μ is obtained from λ by deleting a cell and adding a cell in a row that is not the first row. If the deleted cell belonged to the first row, then $\overline{\mu}$ is obtained from $\overline{\lambda}$ by adding a cell. Otherwise, $\overline{\mu}$ is obtained from $\overline{\lambda}$ by removing a cell and then adding a cell. This is precisely the branching rule in Table 5, with λ replaced by $\overline{\lambda}$ and μ replaced by $\overline{\mu}$. \square

The map ϕ from Proposition 6.11 allows us to establish a bijection between standard multiset tableaux and *vacillating tableaux*. A *vacillating tableau* is a sequence of partitions satisfying the condition $\lambda^{(r)} \vdash n$ and $\lambda^{(r+\frac{1}{2})} \vdash n - 1$ with $\lambda^{(r)} \leftarrow \lambda^{(r+\frac{1}{2})}$ and $\lambda^{(r+\frac{1}{2})} \rightarrow \lambda^{(r+1)}$ for $0 \leq r < k$ [16,2]. A different bijection appears in [2]. The bijection we provide here is compatible with the families of tableaux for each of the subalgebras and the Bratteli diagrams for those subalgebras.

Proposition 6.14. *For each family of subalgebras A_k in Table 1 and for each λ a partition of $n \geq 2k$, there is a bijection between $\text{SMT}_{A_k}(\lambda)$ and the set of vacillating tableaux of the form*

$$\left((n) = \lambda^{(0)}, \lambda^{(\frac{1}{2})}, \lambda^{(1)}, \lambda^{(1\frac{1}{2})}, \dots, \lambda^{(k-\frac{1}{2})}, \lambda^{(k)} = \lambda \right),$$

where

$$\overline{\lambda^{(0)}} \Rightarrow \overline{\lambda^{(1)}} \Rightarrow \dots \Rightarrow \overline{\lambda^{(k)}}$$

is a path in the Bratteli diagram for the tower of algebras $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$.

Proof. Let $T^{(k)}$ be a tableau in $\text{SMT}_{A_k}(\lambda)$. If $\phi(T^{(k)}) = (S, \tau)$, then set $T^{(k-1)} = S$, $\lambda^{(k-1)} = \text{shape}(S)$, and $\lambda^{(k-\frac{1}{2})} = \tau$. Repeat this process on $T^{(k-r)}$ for k steps until $T^{(0)}$ is the unique empty tableau in $\text{SMT}_{A_0}((n))$. Record at each step of this process $\lambda^{(k-1)}$ and $\lambda^{(k-\frac{1}{2})}$. Now given the sequence of partitions

$$\left((n) = \lambda^{(0)}, \lambda^{(\frac{1}{2})}, \lambda^{(1)}, \lambda^{(1\frac{1}{2})}, \dots, \lambda^{(k-\frac{1}{2})}, \lambda^{(k)} = \lambda \right)$$

we can reverse the steps and recover the standard multiset tableau from the sequence.

A consequence of Theorem 6.13 is that the sequence of partitions

$$\overline{\lambda^{(0)}} \Rightarrow \overline{\lambda^{(1)}} \Rightarrow \dots \Rightarrow \overline{\lambda^{(k)}}$$

is a path in the Bratteli diagram for the tower of algebras $A_0 \subseteq A_1 \subseteq A_2 \subseteq \dots$; furthermore, for a partition $\lambda \vdash n$ with $n \geq 2k$, the number of these paths that end on the partition $\overline{\lambda}$ is equal to the number of standard multiset tableaux in $\text{SMT}_{A_k}(\lambda)$. \square




Fig. 7. An example of the bijection from Proposition 6.14; see Example 6.15 for details.

Example 6.15. Let T and S be the two tableaux from Example 6.12. Start with $T^{(9)} = T$. It follows from Example 6.12 that $T^{(8)} = S$ and so we record

$$\lambda^{(9)} = (n - 3, 2, 1), \lambda^{(8\frac{1}{2})} = (n - 4, 2, 1), \text{ and } \lambda^{(8)} = (n - 4, 2, 1, 1).$$

The remaining steps of the bijection are given in Fig. 7. The corresponding sequence $\{\overline{\lambda^{(i)}}\}_{i=1}^9$ is the following path in the Bratteli diagram for the Brauer algebra.

$$\emptyset \Rightarrow (1) \Rightarrow \emptyset \Rightarrow (1) \Rightarrow (2) \Rightarrow (2, 1) \Rightarrow (1, 1) \Rightarrow (1, 1, 1) \Rightarrow (2, 1, 1) \Rightarrow (2, 1).$$

What we have presented in this section completes the connection between the results in [15] and those in [16]. The insertion presented in Theorem 6.3 is a correspondence between diagrams and pairs of standard multiset tableaux that motivates the tableaux that arise in the paper [15]. Theorem 6.13 then provides a correspondence between standard multiset tableaux and paths in the Bratteli diagram.

Since the dimensions of the irreducibles are equal to the number of paths in the Bratteli diagram, it follows that the number of tableaux of a given shape is equal to the dimension of the irreducible representation. This establishes the following result, which can also be proven by enumerating the tableaux in Lemma 6.6 by a purely combinatorial argument and verifying that the values agree with Table 2.

Corollary 6.16. *Let $n \geq 2k$ and $\lambda \vdash n$. For each of the algebras A_k described in Table 1, let $V_{A_k}^{\bar{\lambda}}$ be the irreducible A_k -representation indexed by $\bar{\lambda}$. Then*

$$\dim \left(V_{A_k}^{\bar{\lambda}} \right) = \# \text{SMT}_{A_k}(\lambda).$$

Remark 6.17. Benkart and Halverson [2] give a bijection between standard multiset tableaux and vacillating tableaux that is different from the correspondence that we have just described. Their bijection does not behave well under restriction to all of the subalgebras A_k and the corresponding standard multiset tableaux in $\text{SMT}_{A_k}(\lambda)$. This can be demonstrated via an example. In $TL_2(4)$, there are two diagrams and two pairs of tableaux (see the third diagram in the second row and the first diagram in the fourth row of Example 6.4), and the Benkart–Halverson produces the following vacillating tableau

$$\begin{array}{|c|c|c|c|} \hline & & & 12 \\ \hline \end{array} \xrightarrow{\text{Benkart–Halverson}} \left(\begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array} \right),$$

however $(\emptyset, \emptyset, \emptyset)$ is not a path in the Temperley–Lieb Bratteli diagram (see [20, p. 19]). On the other hand, our correspondence produces the following vacillating tableau

$$\begin{array}{|c|c|c|c|} \hline & & & 12 \\ \hline \end{array} \xrightarrow{\text{Proposition 6.14}} \left(\begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \\ \hline \end{array}, \begin{array}{|c|c|c|} \hline & & \\ \hline \end{array}, \begin{array}{|c|c|} \hline & \\ \hline \end{array}, \begin{array}{|c|c|c|c|} \hline & & & \\ \hline \end{array} \right),$$

and $(\emptyset, \square, \emptyset)$ is a path in the Temperley–Lieb Bratteli diagram.

References

- [1] Georgia Benkart, Tom Halverson, Motzkin algebras, *Eur. J. Comb.* 36 (2014) 473–502, MR 3131911.
- [2] Georgia Benkart, Tom Halverson, Partition algebras and the invariant theory of the symmetric group, in: *Recent Trends in Algebraic Combinatorics*, in: *Assoc. Women Math. Ser.*, vol. 16, Springer, Cham, 2019, pp. 1–41, MR 3969570.
- [3] Georgia Benkart, Tom Halverson, Nate Harman, Dimensions of irreducible modules for partition algebras and tensor power multiplicities for symmetric and alternating groups, *J. Algebraic Comb.* 46 (1) (2017) 77–108, MR 3666413.
- [4] Richard Brauer, On algebras which are connected with the semisimple continuous groups, *Ann. Math.* (2) 38 (4) (1937) 857–872, MR 1503378.
- [5] David A. Cox, John Little, Donal O’Shea, *Ideals, Varieties, and Algorithms: An Introduction to Computational Algebraic Geometry and Commutative Algebra*, fourth ed., *Undergraduate Texts in Mathematics*, Springer, Cham, 2015, MR 3330490.
- [6] Louis Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, enlarged ed., D. Reidel Publishing Co., Dordrecht, 1974, MR 0460128.

- [7] Tullio Ceccherini-Silberstein, Fabio Scarabotti, Filippo Tolli, Representation Theory of the Symmetric Groups: The Okounkov-Vershik Approach, Character Formulas, and Partition Algebras, Cambridge Studies in Advanced Mathematics, vol. 121, Cambridge University Press, Cambridge, 2010, MR 2643487.
- [8] Daniel Flath, Tom Halverson, Kathryn Herbig, The planar rook algebra and Pascal's triangle, Enseign. Math. (2) 55 (1–2) (2009) 77–92, MR 2541502.
- [9] Ira M. Gessel, Enumerative applications of a decomposition for graphs and digraphs, in: Formal Power Series and Algebraic Combinatorics, Montreal, PQ, 1992, Discrete Math. 139 (1–3) (1995) 257–271, MR 1336842.
- [10] Roe Goodman, Nolan R. Wallach, Representations and Invariants of the Classical Groups, Encyclopedia of Mathematics and Its Applications, vol. 68, Cambridge University Press, Cambridge, 1998, MR 1606831.
- [11] Tom Halverson, Characters of the partition algebras, J. Algebra 238 (2) (2001) 502–533, MR 1823772.
- [12] T. Halverson, Representations of the q -rook monoid, J. Algebra 273 (2004) 227–251.
- [13] Tom Halverson, Set-partition tableaux, symmetric group multiplicities, and partition algebra modules, Sémin. Lothar. Comb. 82B (2019) 58.
- [14] Tom Halverson, Elise deMas, Representations of the rook-Brauer algebra, Commun. Algebra 42 (1) (2014) 423–443, MR 3169580.
- [15] Tom Halverson, Theodore N. Jacobson, Set-partition tableaux and representations of diagram algebras, arXiv e-prints, arXiv:1808.08118, 2018.
- [16] Tom Halverson, Tim Lewandowski, RSK insertion for set partitions and diagram algebras, Electron. J. Comb. 11 (2) (2004/2006) R24, MR 2195430.
- [17] Roger Howe, Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond, in: The Schur Lectures, Tel Aviv, 1992, in: Israel Math. Conf. Proc., vol. 8, Bar-Ilan Univ., Ramat Gan, 1995, pp. 1–182, MR 1321638.
- [18] Tom Halverson, Arun Ram, Partition algebras, Eur. J. Comb. 26 (6) (2005) 869–921, MR 2143201.
- [19] OEIS Foundation Inc., The on-line encyclopedia of integer sequences, 2019 (online).
- [20] V.F.R. Jones, Index for subfactors, Invent. Math. 72 (1) (1983) 1–25, MR 696688.
- [21] V.F.R. Jones, The Potts model and the symmetric group, in: Subfactors, Kyuzeso, 1993, World Sci. Publ., River Edge, NJ, 1994, pp. 259–267, MR 1317365.
- [22] Donald E. Knuth, Permutations, matrices, and generalized Young tableaux, Pac. J. Math. 34 (1970) 709–727, MR 0272654.
- [23] Percy A. MacMahon, Combinatory Analysis, vols. I, II (bound in one volume), Dover Phoenix Editions, Dover Publications, Inc., Mineola, NY, 2004, Reprint of *An Introduction to Combinatory Analysis* (1920) and *Combinatory Analysis, vols. I, II* (1915, 1916), MR 2417935.
- [24] Paul P. Martin, Representations of graph Temperley-Lieb algebras, Publ. Res. Inst. Math. Sci. 26 (3) (1990) 485–503, MR 1068862.
- [25] Paul Martin, Volodymyr Mazorchuk, On the representation theory of partial Brauer algebras, Q. J. Math. 65 (1) (2014) 225–247, MR 3179659.
- [26] P.P. Martin, G. Rollet, The Potts model representation and a Robinson-Schensted correspondence for the partition algebra, Compos. Math. 112 (2) (1998) 237–254, MR 1626017.
- [27] Paul Martin, Hubert Saleur, On an algebraic approach to higher-dimensional statistical mechanics, Commun. Math. Phys. 158 (1) (1993) 155–190, MR 1243720.
- [28] Sridhar Narayanan, Digjoy Paul, Shraddha Srivastava, The multiset partition algebra, arXiv e-prints, arXiv:1903.10809, 2019.
- [29] Rosa Orellana, Mike Zabrocki, Symmetric group characters as symmetric functions, arXiv e-prints, arXiv:1605.06672, 2016.
- [30] Rosa Orellana, Mike Zabrocki, A combinatorial model for the decomposition of multivariate polynomials rings as an S_n -module, arXiv e-prints, arXiv:1906.01125, 2019.
- [31] G. de B. Robinson, On the representations of the symmetric group, Am. J. Math. 60 (3) (1938) 745–760, MR 1507943.
- [32] Mercedes Helena Rosas, A Combinatorial Overview of the Theory of MacMahon Symmetric Functions and a Study of the Kronecker Product of Schur Functions, ProQuest LLC, Ann Arbor, MI, 2000, Thesis (Ph.D.), Brandeis University, MR 2700302.
- [33] I. Schur, Über eine Klasse von Matrizen, die sich einer gegebenen Matrix zuordnen lassen, Dissertation, Berlin, 1901, JMF 32.0165.04.
- [34] I. Schur, Über die rationalen Darstellungen der allgemeinen linearen Gruppe, Sitz.ber. Akad. Berlin (1927) 58–75, JMF 53.0108.05.

- [35] C. Schensted, Longest increasing and decreasing subsequences, *Can. J. Math.* 13 (1961) 179–191, MR 0121305.
- [36] Louis Solomon, Representations of the rook monoid, *J. Algebra* 256 (2) (2002) 309–342, MR 1939108.
- [37] H.N.V. Temperley, E.H. Lieb, Relations between the “percolation” and “colouring” problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the “percolation” problem, *Proc. R. Soc. Lond. Ser. A* 322 (1549) (1971) 251–280, MR 0498284.
- [38] Hans Wenzl, On the structure of Brauer’s centralizer algebras, *Ann. Math. (2)* 128 (1) (1988) 173–193, MR 951511.
- [39] B.W. Westbury, The representation theory of the Temperley-Lieb algebras, *Math. Z.* 219 (4) (1995) 539–565, MR 1343661.
- [40] Hermann Weyl, *The Classical Groups: Their Invariants and Representations*, fifteenth printing, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1997, Princeton Paperbacks, MR 1488158.