

Deligne–Lusztig varieties and period domains over finite fields

S. Orlik^{a,*}, M. Rapoport^b

^a *Universität Leipzig, Fakultät für Mathematik und Informatik, Postfach 100920, 04009 Leipzig, Germany*

^b *Mathematisches Institut der Universität Bonn, Beringstrasse 1, 53115 Bonn, Germany*

Received 10 September 2007

Available online 8 May 2008

Communicated by Michel Broué

Abstract

We prove that the Drinfeld space is essentially the only Deligne–Lusztig variety which is at the same time a period domain over a finite field. We also discuss an affineness criterion for Deligne–Lusztig varieties and its relation to cohomology vanishing theorems.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Deligne–Lusztig varieties; Period domains; Minimal length elements in F -conjugacy classes of Weyl groups

1. Introduction

Let G_0 be a reductive group over \mathbb{F}_q . There are two classes of algebraic varieties over an algebraic closure \mathbb{F} of \mathbb{F}_q attached to G_0 . Let us recall their definition. We set $G = G_0 \times_{\mathbb{F}_q} \mathbb{F}$.

To G there is associated *the* maximal torus, *the* Weyl group W and *the* set of fundamental reflections in W , cf. [DL], 1.1. Let $X = X_G$ be the set of all Borel subgroups of G . Then X is a smooth projective algebraic variety homogeneous under G . The set of orbits of G on $X \times X$ can be identified with W , and this defines *the relative position map* $\text{inv} : X \times X \rightarrow W$ (associate to an element of $X \times X$ the G -orbit containing it). Let $w \in W$. The *Deligne–Lusztig variety associated to* (G_0, q, w) is the locally closed subset of X given by

* Corresponding author.

E-mail addresses: orlik@mathematik.uni-leipzig.de (S. Orlik), rapoport@math.uni-bonn.de (M. Rapoport).

$$X(w) = X_{G_0}(w) = \{x \in X \mid \text{inv}(x, Fx) = w\}.$$

Here $F : X \rightarrow X$ denotes the Frobenius map over \mathbb{F}_q . It is known ([DL], 1.4) that $X(w)$ is a smooth quasi-projective variety of dimension $\ell(w)$, which is equipped with an action of $G_0(\mathbb{F}_q)$. If F^e is the minimal power of F with $F^e(w) = w$, then $X(w)$ is defined over \mathbb{F}_{q^e} .

For the other class of varieties, fix a conjugacy class \mathcal{N} of cocharacters $\nu : \mathbb{G}_m \rightarrow G$. Any such ν defines a parabolic subgroup P_ν of G and all parabolics obtained from elements $\nu \in \mathcal{N}$ are conjugate. Let $X_G(\mathcal{N})$ be the set of these conjugates, a smooth projective algebraic variety homogeneous under G . Any $\nu \in \mathcal{N}$ defines via the adjoint representation a \mathbb{Z} -filtration \mathcal{F}_ν on $\text{Lie}(G)$, and ν is called *semi-stable* if $(\text{Lie}(G_0), \mathcal{F}_\nu)$ is semi-stable as an \mathbb{F}_q -vector space equipped with a \mathbb{Z} -filtration on the corresponding \mathbb{F} -vector space, cf. [R,F]. This condition only depends on the point in $X(\mathcal{N})$ corresponding to ν and defines an open subset $X(\mathcal{N})^{\text{ss}} = X_{G_0}(\mathcal{N})^{\text{ss}}$ of $X(\mathcal{N})$, called the *period domain associated with* (G_0, q, \mathcal{N}) , cf. [R]. Hence $X(\mathcal{N})^{\text{ss}}$ is a smooth connected quasi-projective variety of dimension $\dim X(\mathcal{N})$. It is equipped with an action of $G_0(\mathbb{F}_q)$. If the conjugacy class \mathcal{N} is defined over the extension \mathbb{F}_{q^e} , then $X(\mathcal{N})^{\text{ss}}$ is defined over \mathbb{F}_{q^e} .

The Drinfeld space Ω^n (relative to \mathbb{F}_q) is a DL-variety, as well as a period domain. More precisely, let $G_0 = \text{GL}_n$. Let $w = s_1 s_2 \cdots s_{n-1} = (1, 2, \dots, n)$ be the standard Coxeter element. Then $X_{G_0}(w)$ can be identified with the Drinfeld space

$$\Omega^n = \Omega_{\mathbb{F}_q}^n = \mathbb{P}^{n-1} \setminus \bigcup_{H/\mathbb{F}_q} H$$

(complement of all \mathbb{F}_q -rational hyperplanes in the projective space of lines in \mathbb{F}^n), cf. [DL], Section 2. For any Coxeter element w for GL_n , the corresponding DL-variety $X(w)$ is universally homeomorphic to Ω^n , cf. [L], Prop. 1.10. On the other hand, let us identify as usual the set of conjugacy classes \mathcal{N} for GL_n with

$$(\mathbb{Z}^n)_+ = \{(x_1, \dots, x_n) \in \mathbb{Z}^n \mid x_1 \geq x_2 \geq \cdots \geq x_n\}.$$

Let $(x, y^{(n-1)}) \in (\mathbb{Z}^n)_+$ with $x > y$ (here $y^{(n-1)}$ indicates that the entry y is repeated $n-1$ times). Then the corresponding period domain is equal to Ω^n , cf. [R]. Similarly, if $(x^{(n-1)}, y) \in (\mathbb{Z}^n)_+$ with $x > y$, then the corresponding period domain is isomorphic to Ω^n (it is equal to the dual $\hat{\Omega}^n$, the set of hyperplanes of \mathbb{F}^n not containing any \mathbb{F}_q -rational line). In [R], Section 3, it is shown on examples that the Drinfeld space has various special features that are not shared by more general period domains. In the present paper we exhibit another such feature: the Drinfeld space is essentially the only period domain which is at the same time a DL-variety. Before formulating the result, we note that both $X_{G_0}(w)$ and $X_{G_0}(\mathcal{N})^{\text{ss}}$ only depend on the adjoint group $G_{0\text{ad}}$. Also, if G_0 is the direct product of groups, then the corresponding Deligne–Lusztig varieties and period domains also split into products. Hence we may assume that G_0 is \mathbb{F}_q -simple and adjoint. Then G_0 is of the form $G_0 = \text{Res}_{\mathbb{F}_{q'}/\mathbb{F}_q}(G'_0)$ for an absolutely simple group G'_0 over $\mathbb{F}_{q'}$. Then \mathcal{N} is of the form $(\mathcal{N}_1, \dots, \mathcal{N}_t)$ corresponding to the \mathbb{F}_q -embeddings of $\mathbb{F}_{q'}$ into \mathbb{F} . Here $t = |\mathbb{F}_{q'} : \mathbb{F}_q|$ and $\mathcal{N}_1, \dots, \mathcal{N}_t$ are conjugacy classes of G' .

The main result of this note is the following theorem.

Theorem 1.1. *Let G_0 be absolutely simple of adjoint type over \mathbb{F}_q . A Deligne–Lusztig variety $X_{G_0}(w)$ is never universally homeomorphic to a period domain $X_{G_0}(\mathcal{N})^{\text{ss}}$, unless $G_0 = \text{PGL}_n$, w is a Coxeter element and \mathcal{N} corresponds to $\nu \in (\mathbb{Z}^n)_+$ of the form $\nu = (x, y^{(n-1)})$ or*

$v = (x^{(n-1)}, y)$ with $x > y$, in which case $X_{G_0}(w)$ and $X_{G_0}(\mathcal{N})^{\text{ss}}$ are both universally homeomorphic to $\Omega_{\mathbb{F}_q}^n$.

More generally, let $G_0 = \text{Res}_{\mathbb{F}_{q'}/\mathbb{F}_q}(G'_0)$ be simple of adjoint type, and let $t = |\mathbb{F}_{q'} : \mathbb{F}_q|$. Then a Deligne–Lusztig variety $X_{G_0}(w)$ is never universally homeomorphic to a period domain $X_{G_0}(\mathcal{N})^{\text{ss}}$, unless $G'_0 = \text{PGL}_n$, w is a Coxeter element in the sense of [L], 1.7, and \mathcal{N} is of the form $(v_1, \dots, v_t) \in ((\mathbb{Z}^n)_+)^t$ with v_i scalar for all indices $i = 1, \dots, t$, except one index where the entry is of the form $(x, y^{(n-1)})$ or $(x^{(n-1)}, y)$ with $x > y$. In this case $X_{G_0}(w)$ and $X_{G_0}(\mathcal{N})^{\text{ss}}$ are both universally homeomorphic to $\Omega_{\mathbb{F}_{q'}}^n$.

This theorem comes about by comparing a cohomology vanishing theorem for the DL-varieties with a cohomology non-vanishing theorem for period domains. In the sequel we denote for any variety X over \mathbb{F} by $H_c^i(X)$ the ℓ -adic cohomology group with compact supports $H_c^i(X, \overline{\mathbb{Q}}_\ell)$.

The vanishing theorem for DL-varieties is the following statement.

Proposition 1.2.

$$H_c^i(X_{G_0}(w)) = 0 \quad \text{for } 0 \leq i < l(w).$$

This vanishing property is due to Digne, Michel and Rouquier [DMR], Cor. 3.3.22. When $q \geq h$ (where h denotes the Coxeter number of G) then all DL-varieties $X_{G_0}(w)$ are affine, cf. [DL], Thm. 9.7. In this case, the vanishing statement follows by Poincaré duality from a general vanishing theorem for the étale cohomology of affine varieties. Before we became aware of the paper [DMR], we pursued a strategy for proving Proposition 1.2, which relates its statement to the general problem of determining which DL-varieties are affine. Since we believe that our approach has its own merits, we give it in Section 2. It seems more elementary than the approach in [DMR], and is also applicable to the Deligne–Lusztig local systems on DL-varieties. However, we did not succeed completely, since we have to base ourselves on the following hypothesis.

Aff(G_0, q, w): For every w' of minimal length in the F -conjugacy class of w , the corresponding DL-variety $X_{G_0}(w')$ is affine.

It seems to us quite likely that this condition is always satisfied.¹ Lusztig's result [L], Cor. 2.8, that $X_{G_0}(w)$ is affine when w is a Coxeter element may be viewed as supporting this belief. In any case, we show that **Aff**(G_0, q, w) is satisfied when G_0 is a split classical group (cf. Section 5). It is also satisfied when G_0 is of type G_2 , cf. [H'], 4.18. On the other hand, we believe that the hypothesis that w be of minimal length in its conjugacy class cannot be totally dropped, i.e., we believe it may happen for small q that there are DL-varieties which are not affine, although we have no example for this (but a concrete candidate over the field with 2 elements, cf. Remark 5.1). On the other hand, there is the following non-vanishing result [O], Cor. 1.2, for period domains. Let $r_0 = \text{rk}_{\mathbb{F}_q}(G_0)$ denote the \mathbb{F}_q -rank of G_0 (dimension of a maximal \mathbb{F}_q -split torus of G_0).

Proposition 1.3. Let G_0 be a simple group of adjoint type over \mathbb{F}_q . If \mathcal{N} is non-trivial, then

$$H_c^{r_0}(X_{G_0}(\mathcal{N})^{\text{ss}}) \neq 0;$$

¹ X. He [He] has recently given a proof of this conjecture which is inspired by our method in Section 5. Yet another proof is due to C. Bonnafé and R. Rouquier [BR'].

in fact, the representation of $G_0(\mathbb{F}_q)$ on this cohomology group is irreducible and is equivalent to the Steinberg representation.

In order to carry out the comparison between these two results, we use the following observation.

Proposition 1.4. *Let G be a simple group of adjoint type over an algebraically closed field k . For any proper parabolic subgroup P , the following inequality holds,*

$$\mathrm{rk}(G) \leq \dim G/P,$$

with strict inequality, except when $G = \mathrm{PGL}_n$ and P is a parabolic subgroup of type $(n-1, 1)$ or $(1, n-1)$.

Our approach to Proposition 1.2 is given in Section 2, and the proof of Proposition 1.4 in Section 3. The main theorem is proved in Section 4. In the final Section 5, we verify the condition $\mathbf{Aff}(G_0, q, w)$ for split classical groups by checking the criterion of Deligne and Lusztig [DL], 9.6.

2. A vanishing theorem

Let \mathcal{F} be a smooth $\bar{\mathbb{Q}}_\ell$ -sheaf on a connected normal variety X over \mathbb{F} . We say that \mathcal{F} is a *smooth prime-to- p $\bar{\mathbb{Q}}_\ell$ -sheaf*, if it is defined by a constant toric sheaf and the corresponding representation of the fundamental group $\pi_1(X) = \pi_1(X, x)$ on the fiber \mathcal{F}_x at a geometric point x of X factors through the prime-to- p part $\pi_1(X)^{(p)}$ of $\pi_1(X)$. This is independent of the choice of x . The extension of this definition to non-connected normal schemes is immediate.

We will use the following stability property of smooth prime-to- p $\bar{\mathbb{Q}}_\ell$ -sheaves. Let S be a normal scheme and let $f : X \rightarrow S$ be a smooth morphism of relative dimension one, with all fibers affine curves. We assume that f factors as $f = \bar{f} \circ j$, where $j : X \hookrightarrow \bar{X}$ is an open immersion, and where $\bar{f} : \bar{X} \rightarrow S$ is proper and smooth, and such that $D = \bar{X} \setminus X$ is a smooth relative divisor over S . Let \mathcal{F} be a prime-to- p smooth $\bar{\mathbb{Q}}_\ell$ -sheaf on X . Then $R^i f_!(\mathcal{F})$ is a smooth prime-to- p $\bar{\mathbb{Q}}_\ell$ -sheaf on S and is trivial for $i \neq 1, 2$. Indeed, \mathcal{F} is tamely ramified along D , so that the smoothness of $R^i f_!(\mathcal{F})$ follows from [SGA4'], app. to Th. finitude, esp. 1.3.3 and 2.7. Also, the vanishing of $R^i f_!(\mathcal{F})$ for $i \neq 1, 2$ follows from the proper base change theorem, and the calculation of the cohomology of affine curves. Alternatively, one may use Poincaré duality to reduce the question to the analogous statement concerning $R^i f_*(\mathcal{F})$ (for $i \neq 0, 1$ and for the dual sheaf), and refer to [SGA1], XIII, Prop. 1.14 and Remark 1.17 for the smoothness of $R^i f_*(\mathcal{F})$, and to [SGA1], Thm. 2.4, (1) for the commutation of $R^i f_*(\mathcal{F})$ with base change. For $i = 0, 1$, $(R^i f_*(\mathcal{F}))_s = H^i(X_s, \mathcal{F})$ is equal to the Galois cohomology group $H^i(\pi_1(X_s, x), \mathcal{F}_x)$, cf. [M], Thm. 14.14. Under this identification, the action of $\pi_1(S, s)$ is obtained from the action of $\pi_1(X, x)$ on \mathcal{F}_x in the sense of [S], I.2.6, (b).

[Illusie pointed out to us that this requires justification. For this, it suffices to prove the analogous statement for a smooth torsion sheaf \mathcal{F} . By restricting f to smaller and smaller open subsets of S , we may pass to the generic fiber and are then in the following situation. Let X be an affine smooth curve over a field k and let \mathcal{F} be a smooth torsion sheaf on X . Consider the exact sequence of fundamental groups

$$1 \rightarrow \pi_1(X_{\bar{k}}, x) \rightarrow \pi_1(X, x) \rightarrow \mathrm{Gal}(\bar{k}/k) \rightarrow 1.$$

The étale cohomology groups $H^i(X_{\bar{k}}, \mathcal{F})$ may be identified with the Galois cohomology groups $H^i(\pi_1(X_{\bar{k}}, x), \mathcal{F}_x)$ since the inverse image of \mathcal{F} to the universal covering of $X_{\bar{k}}$ is acyclic [M], Thm. 14.14. There are two actions of $\text{Gal}(\bar{k}/k)$ on these cohomology groups: one on the Galois cohomology group coming from the fact that the action of $\pi_1(X_{\bar{k}}, x)$ on \mathcal{F}_x comes by restricting the action of the bigger group $\pi_1(X, x)$ on \mathcal{F}_x , and the action of $\text{Gal}(\bar{k}/k)$ on the étale cohomology group $H^i(X_{\bar{k}}, \mathcal{F})$ by functoriality. It is obvious that these two actions coincide for $i = 0$. Since the two functors arise as derived functors, the two actions coincide then for all i .

Now the homomorphism $\pi_1(X, x) \rightarrow \pi_1(S, s)$ is surjective [SGA1], IX, 5.6, hence this action factors through $\pi_1(S, s)^{(p)}$.

After these preliminaries, we may state the vanishing theorem.

Theorem 2.1. *Assume $\text{Aff}(G_0, q, w)$. Let \mathcal{F} be a smooth prime-to- p $\bar{\mathbb{Q}}_\ell$ -sheaf on $X(w)$. Then*

$$H_c^i(X(w), \mathcal{F}) = 0 \quad \text{for } 0 \leq i < \ell(w).$$

For the constant sheaf $\mathcal{F} = \bar{\mathbb{Q}}_\ell$, we obtain the statement of Proposition 1.2, except that here we have to make the hypothesis $\text{Aff}(G_0, q, w)$.

Let T_0 be a maximal torus in G_0 , with corresponding maximal torus T of G . We identify the Weyl group with the Weyl group of T . Then to every $w \in W$ and every character $\theta : T(\mathbb{F})^{wF} \rightarrow \bar{\mathbb{Q}}_\ell^\times$, Deligne and Lusztig have associated a smooth prime-to- p sheaf \mathcal{F}_θ on $X(w)$, cf. [DL], p. 111 (when θ is trivial, then $\mathcal{F}_\theta = \bar{\mathbb{Q}}_\ell$). As an application of Theorem 2.1 we have the following result.

Corollary 2.2. *Assume $\text{Aff}(G_0, q, w)$. For any θ*

$$H_c^i(X(w), \mathcal{F}_\theta) = 0 \quad \text{for } 0 \leq i < \ell(w).$$

If θ is nonsingular, then

$$H_c^i(X(w), \mathcal{F}_\theta) = 0 \quad \text{for } i \neq \ell(w).$$

Proof. The first statement follows from Theorem 2.1. For the second statement, we use the fact [DL], Thm. 9.8, that if θ is nonsingular, then the natural homomorphism from $H_c^i(X(w), \mathcal{F}_\theta)$ to $H^i(X(w), \mathcal{F}_\theta)$ is an isomorphism. Therefore the assertion follows from Poincaré duality. \square

Remark 2.3. The previous statement for nonsingular θ is due to Haastert [H], Satz 3.2, as an application of his result that $X(w)$ is quasi-affine, cf. [H], Satz 2.3. He does not have to assume the hypothesis $\text{Aff}(G_0, q, w)$. Of course, when $X(w)$ is affine, this statement is proved in [DL].

For the proof of Theorem 2.1 we first recall the following result of Geck, Kim and Pfeiffer. Denote by S the set of simple reflections in W . Let $w, w' \in W$ and $s \in S$. Set $w \xrightarrow{s}_F w'$ if $w' = s w F(s)$ and $\ell(w') \leq \ell(w)$. We write $w \rightarrow_F w'$ if $w = w'$ or if there exist elements $s_1, \dots, s_r \in S$ and $w = w_1, \dots, w_r = w' \in W$ with $w_i \xrightarrow{s_i}_F w_{i+1}$, $i = 1, \dots, r - 1$.

Theorem 2.4. (See [GKP], Thm. 2.6.) *Let C be an F -conjugacy class of W and let C_{\min} be the set of elements in C of minimal length. For any $w \in C$, there exists some $w' \in C_{\min}$ such that $w \rightarrow_F w'$.*

We also recall the following lemma (Case 1 of Thm. 1.6 in [DL]).

Lemma 2.5. *Let w and w' be F -conjugate. Suppose that there are $w_1, w_2 \in W$ with $w_1 w_2 = w$, $w_2 F(w_1) = w'$ and $\ell(w) = \ell(w_1) + \ell(w_2) = \ell(w')$. Then $X(w)$ and $X(w')$ are universally homeomorphic and hence $H_c^*(X(w), \mathcal{F}) \cong H_c^*(X(w'), \mathcal{F})$ for any \mathbb{Q}_ℓ -sheaf \mathcal{F} .*

As is well known, this lemma has the following consequence.

Lemma 2.6. *Let $s \in S$ and let $w, w' \in W$ with $w' = s w F(s)$. Suppose that $\ell(w) = \ell(w')$. Then $X(w)$ and $X(w')$ are universally homeomorphic and hence $H_c^*(X(w), \mathcal{F}) \cong H_c^*(X(w'), \mathcal{F})$ for any \mathbb{Q}_ℓ -sheaf \mathcal{F} .*

Proof. We consider the following three cases.

Case 1. $\ell(sw) = \ell(w) - 1$. Then we put $w_1 = s$, $w_2 = sw$ and apply the previous lemma.

Case 2. $\ell(wF(s)) = \ell(w) - 1$. We put $w_1 = s$, $w_2 = sw'$. Again we apply the previous lemma, with the roles of w and w' interchanged.

Case 3. $\ell(sw) = \ell(w) + 1$ and $\ell(wF(s)) = \ell(w) + 1$. Then we apply Lemma 1.6.4 of [DL] to deduce that $w = w'$. \square

Proof of Theorem 2.1. We prove the claim by induction on $\ell(w)$. The case $\ell(w) = 0$ is trivial.

Let $w \in W$ and suppose that the vanishing property holds for all elements in W with length smaller than $\ell(w)$. If w is minimal within its F -conjugacy class, then the vanishing follows by our assumption **Aff**(G_0, q, w) from Poincaré duality and a general vanishing property of affine schemes. If w is not minimal, there is by Theorem 2.4 a chain of simple reflections $s_1, \dots, s_r \in S$ and $w = w_1, \dots, w_r \in W$ with $w_i \xrightarrow{s_i} w_{i+1}$, $i = 1, \dots, r-1$ such that w_r is minimal. By Lemma 2.6 and by induction, we may assume that $w' = s w F(s)$ where $s \in S$ and $\ell(swF(s)) < \ell(w)$, i.e., $\ell(w') = \ell(w) - 2$. As in the proof of Theorem 1.6 in [DL], we may write $X(w)$ as a (set-theoretical) disjoint union

$$X(w) = X_1 \cup X_2$$

where X_1 is closed in $X(w)$ and X_2 is its open complement. By applying the long exact cohomology sequence, it suffices to show that $H_c^i(X_1, \mathcal{F}|_{X_1}) = 0$ and $H_c^i(X_2, \mathcal{F}|_{X_2}) = 0$ for $i < \ell(w)$. Note that the restrictions $\mathcal{F}|_{X_1}$ and $\mathcal{F}|_{X_2}$ are also prime-to- p , since the corresponding representations of their fundamental groups are induced by the canonical maps $\pi_1(X_i) \rightarrow \pi_1(X(w))$, $i = 1, 2$. Now X_1 has the structure of an \mathbb{A}^1 -fibering over $X(w')$. Let $f : X_1 \rightarrow X(w')$ be the \mathbb{A}^1 -fibering. Consider the Leray spectral sequence

$$H_c^i(X(w'), R^j f_* \mathcal{F}|_{X_1}) \Rightarrow H_c^{i+j}(X_1, \mathcal{F}|_{X_1}).$$

The stalks of $R^j f_* \mathcal{F}$ are isomorphic to $H_c^j(\mathbb{A}^1, \mathcal{F}|_{\mathbb{A}^1})$. Now $\pi_1(\mathbb{A}^1)^{(p)} = 0$, cf. [SGA1], XIII, Cor. 2.12. Since \mathcal{F} is prime-to- p , $\mathcal{F}|_{\mathbb{A}^1}$ is constant and $H_c^1(\mathbb{A}^1, \mathcal{F}) = 0$. We deduce that

$$H_c^i(X_1, \mathcal{F}) = H_c^{i-2}(X(w'), R^2 f_* \mathcal{F}).$$

Since $\mathcal{F}' = R^2 f_! \mathcal{F}$ is a smooth prime-to- p $\bar{\mathbb{Q}}_\ell$ -sheaf on $X(w')$, the induction hypothesis applies to it and it follows that $H^{i-2}(X(w'), \mathcal{F}') = 0$ for all $i - 2 < \ell(w')$. Thus $H_c^i(X_1, \mathcal{F}_{|X_1}) = 0$ for all $i < \ell(w)$.

The vanishing of $H_c^i(X_2, \mathcal{F}_{|X_2})$ is even easier. In the proof of [DL], Thm. 1.6, it is shown that X_2 is universally homeomorphic to a line bundle over $X(sw')$ with the zero section removed. Let $g : X_2 \rightarrow X(sw')$ be the corresponding morphism. Then the Leray spectral sequence gives a long exact sequence

$$\cdots \rightarrow H_c^{i-1}(X(sw'), R^1 g_! \mathcal{F}) \rightarrow H_c^i(X_2, \mathcal{F}_{|X_2}) \rightarrow H_c^{i-2}(X(sw'), R^2 g_! \mathcal{F}) \rightarrow \cdots.$$

We have $\ell(sw') = \ell(w') + 1$. By induction $H_c^i(X(sw'), R^j g_! \mathcal{F}) = 0$ for all $i < \ell(sw') = \ell(w) - 1$ and all j . Thus $H_c^{i-1}(X(sw'), R^1 g_! \mathcal{F}) = 0$ and $H_c^{i-2}(X(sw'), R^2 g_! \mathcal{F}) = 0$ for $i < \ell(w)$. The claim follows. \square

3. Proof of Proposition 1.4

We retain the notation of the statement of the proposition. It obviously suffices to prove the statement for a *maximal* parabolic subgroup P . Let B be a Borel subgroup contained in P and let T be a maximal torus in B . Let M be the Levi subgroup of P containing T . Then

$$\begin{aligned} \dim G/P &= \dim G/B - \dim M/M \cap B \\ &= |\Phi^+| - |\Phi_M^+|, \end{aligned}$$

where $\Phi^+ = \Phi_G^+$ resp. Φ_M^+ denotes the set of positive roots of G resp. of M . The assertion is now reduced to a purely combinatorial statement that can be checked mechanically for each type in the tables [Bou]. We adopt the notation used in these tables.

Type A_ℓ : Here $|\Phi^+| = \frac{\ell(\ell+1)}{2}$. If Δ_M is obtained by deleting the root α_i , then Φ_M is of type $A_{i-1} \times A_{\ell-i}$ (with the convention $A_0 = \emptyset$). Hence $|\Phi_M^+| = \frac{i(i-1)}{2} + \frac{(\ell-i)(\ell-i+1)}{2}$. Hence $|\Phi^+| - |\Phi_M^+| \geq \ell$, with equality iff $i = 1$ or $i = \ell$.

Type B_ℓ ($\ell \geq 2$): Here $|\Phi^+| = \ell^2$. If Δ_M is obtained by deleting α_i , then Φ_M is of type $A_{i-1} \times B_{\ell-i}$ (with the convention $B_0 = \emptyset$, $B_1 = A_1$) and $|\Phi_M^+| = \frac{i(i-1)}{2} + (\ell-i)^2$. Hence $|\Phi^+| - |\Phi_M^+| > \ell$ in all cases. The type C_ℓ is identical.

Type D_ℓ ($\ell \geq 4$): Here $|\Phi^+| = (\ell-1)\ell$. If Δ_M is obtained by deleting α_i , then Φ_M is of type $A_{i-1} \times D_{\ell-i}$ except when $i = \ell-1$ or $i = \ell$ in which case Φ_M is of type $A_{\ell-1}$, and except when $i = \ell-2$ in which case Φ_M is of type $A_{\ell-3} \times A_1 \times A_1$, and except when $i = \ell-3$ in which case Φ_M is of type $A_{\ell-4} \times A_3$. For $1 \leq i \leq \ell-4$, $|\Phi_M^+| = \frac{i(i-1)}{2} + (\ell-i)(\ell-i-1)$ and hence $|\Phi^+| - |\Phi_M^+| > \ell$. For $i = \ell-1$ or $i = \ell$, $|\Phi_M^+| = \frac{(\ell-1)\ell}{2}$ and for $i = \ell-2$, $|\Phi_M^+| = \frac{(\ell-3)(\ell-2)}{2} + 2$, and for $i = \ell-3$, $|\Phi_M^+| = \frac{(\ell-4)(\ell-3)}{2} + 6$. In all cases $|\Phi^+| - |\Phi_M^+| > \ell$.

For the exceptional types one gets for the differences $|\Phi^+| - |\Phi_M^+|$, as Δ_M is obtained by deleting $\alpha_1, \dots, \alpha_\ell$, the following integers:

$$E_6: 16, 21, 25, 29, 25, 16,$$

$$E_7: 33, 42, 47, 53, 50, 42, 27,$$

$$E_8: 78, 92, 98, 106, 104, 97, 83, 57,$$

$$F_4: 15, 20, 20, 15,$$

$$G_2: 5, 5.$$

In each case the numbers are strictly larger than the rank. \square

4. Proof of Theorem 1.1

Let us first treat the case when G_0 is absolutely simple. Let us assume that $X = X_{G_0}(w)$ is universally homeomorphic to $X_{G_0}(\mathcal{N})^{\text{ss}}$. By Proposition 1.2 we have $H_c^i(X) = 0$ for $i < \ell(w) = \dim X$. Comparing with Proposition 1.3 we obtain

$$\dim X_{G_0}(w) \leq r_0.$$

Now the relative rank r_0 of G_0 is at most the absolute rank r . From Proposition 1.4 we obtain the chain of inequalities

$$\dim X_{G_0}(w) \leq r_0 \leq r \leq \dim X(\mathcal{N}). \quad (4.1)$$

Hence all inequalities are equalities and by Proposition 1.4, we have that $G = \text{PGL}_n$ and that \mathcal{N} corresponds to $(x, y^{(n-1)})$ or $(x^{(n-1)}, y)$ with $x > y$. Indeed, the case where \mathcal{N} corresponds to $(x^{(n)})$ is excluded, since this would imply that $\ell(w) = \dim X(\mathcal{N}) = 0$, hence $X(w) = X(\mathbb{F}_q)$ would not be connected. Also the equality $r_0 = r$ implies that $G_0 = \text{PGL}_n$. It follows that $X_{G_0}(\mathcal{N}) \cong \Omega^n$ and $\ell(w) = n - 1$. On the other hand, since $X(w)$ is connected, w has to be an elliptic element in S_n , i.e., every fundamental reflection has to appear in a minimal expression of w , cf. [L], p. 26, and [BR] (the converse is also true, but more difficult to prove). Hence every fundamental reflection appears exactly once, i.e. w is a Coxeter element. Now, the assertion follows from the remarks in the introduction.

Now let G_0 be of the form $G_0 = \text{Res}_{\mathbb{F}_{q'}/\mathbb{F}_q}(G'_0)$, where G'_0 is absolutely simple of adjoint type, and let $t = |\mathbb{F}_{q'} : \mathbb{F}_q|$. As in the introduction we write $\mathcal{N} = (\mathcal{N}_1, \dots, \mathcal{N}_t)$, where the \mathcal{N}_i are conjugacy classes of G'_0 . Let r be the absolute rank of G'_0 . Let t_1 be the number of indices i , where \mathcal{N}_i is non-trivial. The inequality (4.1) is replaced by

$$\dim X_{G_0}(w) \leq r_0 \leq r t_1 \leq \dim X(\mathcal{N}). \quad (4.2)$$

Since $r_0 \leq r$, we deduce from the fact all inequalities in (4.2) are equalities, that $r_0 = r$ and $t_1 = 1$ (as before the case $t_1 = 0$ is excluded). As in the absolutely simple case we deduce that $G'_0 = \text{PGL}_n$, and that for the one index i with non-trivial \mathcal{N}_i this conjugacy class of PGL_n corresponds to $(x, y^{(n-1)})$ or $(x^{(n-1)}, y)$ with $x > y$. Reasoning as before, this implies that w is a Coxeter element in the sense of [L'], i.e., every F -orbit of simple reflections appears precisely once in a minimal expression of w as a product of simple reflections. All these Coxeter elements define universally homeomorphic DL-varieties, cf. [L'], Prop. 1.10. To identify the variety $X = X_{G_0}(w) = X_{G_0}(\mathcal{N})^{\text{ss}}$ with $\Omega_{\mathbb{F}_{q'}}^n$, one may use either incarnation of X . On the DL-side, one can use the Coxeter element $w = (w_1, \dots, w_t)$ with $w_1 = s_1 s_2 \dots s_{n-1}$ and $w_2 = \dots = w_t = 1$. Since the action of F on the flag variety of G_0 is given by $F(B_1, \dots, B_t) = (F^t B_t, B_1, \dots, B_{t-1})$, one sees easily that $X_{G_0}(w) \simeq \Omega_{\mathbb{F}_{q'}}^n$. \square

5. The condition $\text{Aff}(G_0, q, w)$

In this section we show that the condition $\text{Aff}(G_0, q, w)$ is satisfied for classical split groups.

We shall use the following criterion of Deligne and Lusztig [DL], 9.6. Let $C \subset X_*(T)_{\mathbb{R}}$ be the (open) Weyl chamber. For $w \in W$, let

$$D(C, -w^{-1}C) = \{x \in X_*(T)_{\mathbb{R}} \mid \alpha(x) > 0 \ \forall \alpha > 0 \text{ with } w(\alpha) < 0\}.$$

Here α ranges over the roots of T .

DL-Criterion. A DL-variety $X(w)$ is affine if there exists an element $x \in D(C, -w^{-1}C)$, such that $F^*x - wx \in C$.

Remark 5.1. It is not clear how close the Deligne–Lusztig criterion comes to being an equivalence. In [H^{*}], Haastert checks that for a split classical group, every conjugacy class of W contains elements which satisfy the DL-criterion. However, there are not enough elements of minimal length among his elements: there are elements w of minimal length in their conjugacy class such that there is no w' among Haastert's elements with $w \rightarrow_F w'$ (e.g. consider the root system D_ℓ and $w = t'$ below). Still, the method used below is modelled on Haastert's calculations. For G_0 of type G_2 , he shows that the DL-criterion is satisfied for all $w \in W$ and all q , except $q = 2$ and $w = s_1s_2s_1, s_2s_1s_2$, when it is not. We expect that these last two DL-varieties are not affine. It should be possible to check this with the help of the computer.

We now consider the root system of a split classical group. In [GP], Geck and Pfeiffer construct a subset of the Weyl group which contains enough elements of minimal length in their conjugacy class. To recall their result, we set up the notation as follows. Let $\mathbb{R}^\ell = (\mathbb{R}^\ell, (,))$ be the standard euclidian vector space with standard basis $\{e_1, \dots, e_\ell\}$. We recall the sets of simple roots and simple reflections. The extraneous elements below are introduced to give a reasonably uniform treatment of all types.

Type $A_{\ell-1}$ ($\ell \geq 2$): $\Delta = \{\alpha_1, \dots, \alpha_{\ell-1}\}$, where $\alpha_i = e_i - e_{i+1}$, $S = \{s_1, \dots, s_{\ell-1}\}$, where $s_i = s_{\alpha_i}$. Further we set $s'_i = 1$ for all $0 \leq i \leq \ell - 1$. The Weyl chamber is given by

$$C = \left\{ (x_1, \dots, x_\ell) \in \mathbb{R}^\ell \mid x_1 > x_2 > \dots > x_{\ell-1} > x_\ell, \sum_i x_i = 0 \right\}.$$

Type B_ℓ ($\ell \geq 2$): $\Delta = \{\alpha_0 = e_1, \alpha_1, \dots, \alpha_{\ell-1}\}$, $S = \{t, s_1, \dots, s_{\ell-1}\}$ where $t = s_{e_1}$, i.e., $t(e_i) = e_i, \forall i \neq 1$ and $t(e_1) = -e_1$. Further, we set $s'_0 = t$ and $s'_i = s_i s_{i-1} \dots s_1 t s_1 \dots s_{i-1} s_i$ for $1 \leq i \leq \ell - 1$. The Weyl chamber is given by

$$C = \{(x_1, \dots, x_\ell) \in \mathbb{R}^\ell \mid x_1 > 0, x_1 > x_2 > \dots > x_{\ell-1} > x_\ell\}.$$

Type D_ℓ ($\ell \geq 4$): $\Delta = \{\alpha_0 = e_1 + e_2, \alpha_1, \dots, \alpha_{\ell-1}\}$, $S = \{t', s_1, \dots, s_{\ell-1}\}$ where $t' = s_{e_1+e_2}$ is the reflection with $t'(e_1 + e_2) = -(e_1 + e_2)$. Further we set $s'_0 = t' s_1$ and $s'_i = s_{i+1} s_i \dots s_2 (t' \cdot s_1) s_2 \dots s_i s_{i+1}$ for $1 \leq i \leq \ell - 2$. The Weyl chamber is given by

$$C = \{(x_1, \dots, x_\ell) \in \mathbb{R}^\ell \mid x_1 + x_2 > 0, x_1 > x_2 > \dots > x_{\ell-1} > x_\ell\}.$$

For a decomposition $\lambda = (\lambda_1, \dots, \lambda_k)$ of the integer ℓ in the cases $A_{\ell-1}$ and B_ℓ , resp. of $\ell - 1$ in the case D_ℓ , and a vector of signs $\epsilon = (\epsilon_1, \dots, \epsilon_k) \in \{\pm 1\}^k$, let $w_{\lambda, \epsilon} = \prod_{i=1}^k w_{\lambda_i, \epsilon_i}$, where

$$w_{\lambda_i, \epsilon_i} := \begin{cases} s_{m_i} s_{m_i+1} \cdots s_{n_i-1} & \text{if } \epsilon_i = 1, \\ s'_{m_i-1} \cdot s_{m_i} s_{m_i+1} \cdots s_{n_i-1} & \text{if } \epsilon_i = -1. \end{cases}$$

Here we put $m_i = \sum_{j=1}^{i-1} \lambda_j + 1$, $n_i = \sum_{j=1}^i \lambda_j$, $i = 1, \dots, k$ in the cases $A_{\ell-1}$ and B_ℓ , resp. $m_i = \sum_{j=1}^{i-1} \lambda_j + 2$, $n_i = \sum_{j=1}^i \lambda_j + 1$ in the case D_ℓ . Since the elements $w_{\lambda_i, \epsilon_i}$ commute with each other, the above product makes sense.

Proposition 5.2. (See [GP], Prop. 2.3.) For each $w \in W$, there is a decomposition $\lambda = (\lambda_1, \dots, \lambda_k)$ of ℓ in the cases $A_{\ell-1}$ and B_ℓ , resp. of $\ell - 1$ in the case D_ℓ , and a vector of signs $\epsilon = (\epsilon_1, \dots, \epsilon_k) \in \{\pm 1\}^k$ such that $w \rightarrow_F \delta \cdot w_{\lambda, \epsilon}$, where $\delta = 1$ in the case of $A_{\ell-1}$ and B_ℓ and $\delta \in \{1, s_1, t'\}$ in the case of D_ℓ .

Remark 5.3. In the case $A_{\ell-1}$ the elements $\delta \cdot w_{\lambda, \epsilon}$ are all minimal in their conjugacy class; this is not true in the cases B_ℓ and D_ℓ . In general, not all elements minimal in their conjugacy class are of the form $\delta \cdot w_{\lambda, \epsilon}$.

We note that to prove the condition $\text{Aff}(G_0, q, w)$ for all q, w , it suffices to prove that for the elements of the form $\delta \cdot w_{\lambda, \epsilon}$ the corresponding DL-variety is affine. Indeed, if w is an element of minimal length in its conjugacy class, then by Proposition 5.2 we find $w' = \delta \cdot w_{\lambda, \epsilon}$ with $w \rightarrow_F w'$. Since then $\ell(w) = \ell(w')$, a repeated application of Lemma 2.6 shows that the DL-varieties $X(w)$ and $X(w')$ are universally homeomorphic. Hence the fact that $X(w')$ is affine implies that $X(w)$ is affine as well.

We will show that $X(w)$ is affine for elements $w = \delta \cdot w_{\lambda, \epsilon}$ by checking the DL-criterion for w . In the split case F -conjugacy is simply conjugacy and the action of the Frobenius F^* is simply the multiplication by q . We will in fact even show that we can find $x \in C$ such that $qx - wx \in C$.

Type $A_{\ell-1}$: We will use the following lemma.

Lemma 5.4. Let $w = s_m \cdot s_{m+1} \cdots s_{n-1}$. Let $x_1 > x_2 > \cdots > x_m > 0$ be positive real numbers. Then there exist $x_{m+1} > x_{m+2} > \cdots > x_{n+1} > 0$ with $x_m > x_{m+1}$, such that for any choice of $x_{n+2} > x_{n+3} > \cdots > x_{\ell-1} > x_\ell > 0$ with $x_{n+1} > x_{n+2}$, we have

$$(qx - wx, \alpha) > 0 \quad \forall \alpha \in \Delta \quad \text{and} \quad (qx - wx, \alpha_n) > x_{n+1}.$$

[If $n = \ell$, the last condition is interpreted as empty, and the other chains of inequalities are to be interpreted in the obvious way.]

Proof. We compute

$$wx = (x_1, \dots, x_{m-1}, x_n, x_m, x_{m+1}, \dots, x_{n-1}, x_{n+1}, x_{n+2}, \dots, x_\ell).$$

Thus we get for $i \leq m - 2$ and for $i \geq n + 1$,

$$(qx - wx, \alpha_i) = q(x_i - x_{i+1}) - (x_i - x_{i+1}) = (q - 1)(x_i - x_{i+1}).$$

Furthermore,

$$\begin{aligned}
 (qx - wx, \alpha_{m-1}) &= q(x_{m-1} - x_m) - (x_{m-1} - x_n), \\
 (qx - wx, \alpha_m) &= q(x_m - x_{m+1}) - (x_n - x_m), \\
 (qx - wx, \alpha_{m+1}) &= q(x_{m+1} - x_{m+2}) - (x_m - x_{m+1}), \\
 &\vdots \\
 (qx - wx, \alpha_{n-1}) &= q(x_{n-1} - x_n) - (x_{n-2} - x_{n-1}), \\
 (qx - wx, \alpha_n) &= q(x_n - x_{n+1}) - (x_{n-1} - x_{n+1}).
 \end{aligned} \tag{5.1}$$

We immediately see that $(qx - wx, \alpha_i) > 0 \forall i \leq m-2, \forall i \geq n+1$ for any $x = (x_1 > x_2 > \cdots > x_\ell > 0) \in \mathbb{R}^\ell$. For the remaining expressions, it suffices to treat the case $q = 2$. For $1 \leq i \leq n-m$ set $x_{m+i} := x_m - ia$ with $a > 0$. Then

$$\begin{aligned}
 (2x - wx, \alpha_{m-1}) &= 2(x_{m-1} - x_m) - (x_{m-1} - x_n) = (x_{m-1} - x_m) - (x_m - x_n) \\
 &= (x_{m-1} - x_m) - (n-m)a.
 \end{aligned}$$

This expression is positive if a is small enough. The inequality

$$(2x - wx, \alpha_m) = 2(x_m - x_{m+1}) - (x_n - x_m) > 0$$

is clearly satisfied since $x_m > x_n$. Since $(2x - wx, \alpha_i) = a > 0$ for $m+1 \leq i \leq n-1$, it remains to consider the term $(2x - wx, \alpha_n)$. But $(2x - wx, \alpha_n) = 2(x_n - x_{n+1}) - (x_{n-1} - x_{n+1}) > x_{n+1} > 0$ if

$$2(x_n - x_{n+1}) - x_{n-1} > 0.$$

Set $x_{n+1} = x_n - b$ with $0 < b < x_n$. Then $2(x_n - x_{n+1}) > x_{n-1}$, provided that $b > \frac{x_{n-1}}{2}$. If a is small enough, such that $2x_n > x_{n-1}$, such $b > 0$ exists. \square

Proposition 5.5. *Let $w = w_{\lambda, \epsilon} \in W$. Then there is an $x = (x_1 > x_2 > \cdots > x_\ell > 0) \in \mathbb{R}^\ell$ with $(qx - wx, \alpha) > 0$ for all $\alpha \in \Delta$.*

Proof. Let $w = w_{\lambda_1, \epsilon_1} \cdots w_{\lambda_k, \epsilon_k}$ and put $w_i = w_{\lambda_i, \epsilon_i}$. Note that the vector of signs ϵ does not affect this element. Set $x_1 = 1$ and apply successively Lemma 5.4 (starting with w_1). We have $(qx - wx, \alpha_k) = (qx - w_i x, \alpha_k) > 0$ for $k \in [m_i, n_i - 1]$. Further,

$$\begin{aligned}
 (qx - wx, \alpha_{n_i}) - (qx - w_i x, \alpha_{n_i}) &= -(x_{w^{-1}(n_i)} - x_{w^{-1}(n_i+1)}) + (x_{w_i^{-1}(n_i)} - x_{n_i+1}) \\
 &= -x_{n_i+1} + x_{w^{-1}(n_i+1)}.
 \end{aligned}$$

Thus $(qx - wx, \alpha_{n_i}) > 0$, since we arranged in Lemma 5.4 that $(qx - w_i x, \alpha_{n_i}) > x_{n_i+1}$. \square

Corollary 5.6. *There exists $x \in C$ with $(qx - wx, \alpha) > 0$ for all $\alpha \in \Delta$.*

Proof. We add to the x in Proposition 5.5 a multiple $r \cdot (1, 1, \dots, 1)$ such that $x + r \cdot (1, 1, \dots, 1) \in C$. \square

Type B_ℓ : In this case we use the following lemma.

Lemma 5.7. *Let $w = s_m \cdots s_{n-1}$ or $w = s'_{m-1} s_m \cdots s_{n-1}$. Let $x_1 > x_2 > \cdots > x_{m-1} > x_m > 0$ be positive real numbers with $x_{m-1} > 3x_m$ if $m \geq 2$. Then there exist $x_{m+1} > x_{m+2} > \cdots > x_{n+1} > 0$ with $x_m > x_{m+1}$ and $x_n > 3x_{n+1}$, such that for any choice of $x_{n+2} > x_{n+3} > \cdots > x_\ell > 0$ with $x_{n+1} > x_{n+2}$, we have*

$$(qx - wx, \alpha) > 0 \quad \forall \alpha \in \Delta \quad \text{and} \quad (qx - wx, \alpha_n) > 2x_{n+1}.$$

[If $n = \ell$, the last condition is interpreted as empty.]

Proof. The case of $w = s_m \cdots s_{n-1}$ is similar to the one treated in Lemma 5.4. We only have to check in addition that $(qx - wx, e_1) > 0$ which is clear.

So, let $w = s'_{m-1} s_m \cdots s_{n-1}$. Again, it suffices to consider the case $q = 2$. We compute

$$wx = (x_1, \dots, x_{m-1}, -x_n, x_m, x_{m+1}, \dots, x_{n-1}, x_{n+1}, x_{n+2}, \dots, x_\ell).$$

We get the same system of identities (5.1) as in the proof of Lemma 5.4 except for the first two, which now become

$$(2x - wx, \alpha_{m-1}) = 2(x_{m-1} - x_m) - (x_{m-1} + x_n) \quad (5.2)$$

and

$$(2x - wx, \alpha_m) = 2(x_m - x_{m+1}) + (x_n + x_m).$$

We also have to check that $(2x - wx, e_1) > 0$. This is easy since

$$(2x - wx, e_1) = \begin{cases} x_1 & \text{if } m \neq 1, \\ 2x_1 + x_n & \text{if } m = 1. \end{cases}$$

We only have to care of the first expression (5.2). We set $x_{m+i} := x_m - ia$ with $a > 0$ for $1 \leq i \leq n - m$ and write

$$\begin{aligned} (2x - wx, \alpha_{m-1}) &= 2(x_{m-1} - x_m) - (x_{m-1} + x_n) = (x_{m-1} - x_m) - (x_m + x_n) \\ &= (x_{m-1} - 3x_m) + (n - m)a. \end{aligned}$$

Since we have $x_{m-1} > 3x_m$, this term is positive. Finally, we have to show that

$$(2x - wx, \alpha_n) = 2(x_n - x_{n+1}) - (x_{n-1} - x_{n+1}) > 2x_{n+1}$$

and

$$x_n > 3x_{n+1}.$$

Write $x_{n+1} = x_n - b$ with $0 < b < x_n$. Then the first of the above inequalities becomes $2b > x_{n-1} + x_{n+1}$, i.e.,

$$3b > x_n + x_{n-1}$$

and the second becomes

$$3b > 2x_n.$$

Similarly as in Lemma 5.4, we can find b such that these inequalities are solvable. \square

Proposition 5.8. *Let $w = w_{\lambda, \epsilon} \in W$. Then there is an $x = (x_1 > x_2 > \cdots > x_\ell > 0) \in \mathbb{R}^\ell$ with $(qx - wx, \alpha) > 0$ for all $\alpha \in \Delta$.*

Proof. The proof is the same as in the case of $A_{\ell-1}$, except that we have

$$(qx - wx, \alpha_{n_i}) - (qx - w_i x, \alpha_{n_i}) = -x_{n_i+1} \pm x_{w^{-1}(n_i+1)}.$$

Thus $(qx - wx, \alpha_{n_i}) > 0$ since we made sure in Lemma 5.7 that $(qx - w_i x, \alpha_{n_i}) > 2x_{n_i+1}$. \square

Note that the x in Proposition 5.8 lies in C .

Type D_ℓ : In this case we prove the following proposition.

Proposition 5.9. *Let $w = \delta w_{\lambda, \epsilon} \in W$. Then there is an $x = (x_1 > x_2 > \cdots > x_\ell > 0) \in \mathbb{R}^\ell$ with $(qx - wx, \alpha) > 0$ for all $\alpha \in \Delta$.*

Proof. The element $s'_0 = s_1 t'$ is the reflection with $e_1 \mapsto -e_1$, $e_2 \mapsto -e_2$ and which fixes all other e_j . It follows that $s'_i(e_1) = -e_1$, $s'_i(e_{i+2}) = -e_{i+2}$ and $s'_i(e_j) = e_j$ for all $j \notin \{1, i+2\}$.

Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a decomposition of the closed interval $[2, \ell]$ and consider a vector of signs $\epsilon = (\epsilon_1, \dots, \epsilon_k)$. Set $|\epsilon| := \#\{i \mid \epsilon_i < 0\}$. Then one computes that the element $w = w_{\lambda, \epsilon}$ is given by

$$e_1 \mapsto (-1)^{|\epsilon|} e_1,$$

$$e_{m_1} \mapsto e_{m_1+1}, e_{m_1+1} \mapsto e_{m_1+2}, \dots, e_{n_1-1} \mapsto e_{n_1}, e_{n_1} \mapsto (-1)^{\epsilon_1} e_{m_1},$$

$$e_{m_2} \mapsto e_{m_2+1}, e_{m_2+1} \mapsto e_{m_2+2}, \dots, e_{n_2-1} \mapsto e_{n_2}, e_{n_2} \mapsto (-1)^{\epsilon_2} e_{m_2},$$

$$\vdots$$

$$e_{m_k} \mapsto e_{m_k+1}, e_{m_k+1} \mapsto e_{m_k+2}, \dots, e_{n_k-1} \mapsto e_{n_k}, e_{n_k} \mapsto (-1)^{\epsilon_k} e_{m_k}.$$

It follows that $w_{\lambda, \epsilon}$ corresponds to the element $w_{\tilde{\lambda}, \tilde{\epsilon}}$ of $W(B_\ell)$ with $\tilde{\lambda} = (1, \lambda)$ and $\tilde{\epsilon} = ((-1)^{|\epsilon|}, \epsilon)$.

If we multiply $w_{\lambda, \epsilon}$ by $\delta \in \{1, s_1, t'\}$ from the left, only the first factor $w_{\lambda_1, \epsilon_1}$ of $w_{\lambda, \epsilon}$ is affected. In particular, we may reduce by Lemma 5.7 to the case $\lambda = (\lambda_1, 1, \dots, 1)$.

Case. $\delta = 1$. Let $x \in \mathbb{R}^\ell$ be chosen as in Proposition 5.8. Then $(qx - wx, \alpha_i) > 0$ for all $i \geq 1$. So, we only have to ensure that $(qx - wx, \alpha_0) = q(x_1 + x_2) - ((-1)^{|\epsilon|} x_1 \pm x_{w^{-1}(2)}) > 0$ which is clearly satisfied.

Case. $\delta = s_1$.

Subcase. $m_1 = 2$. Then $w = \delta \cdot w_{\lambda_1, \epsilon_1}$ is given by

$$e_1 \mapsto (-1)^{|\epsilon|} e_2,$$

$$e_2 \mapsto e_3, e_3 \mapsto e_4, \dots, e_{n_1-1} \mapsto e_{n_1}, e_{n_1} \mapsto (-1)^{\epsilon_1} e_1.$$

If $|\epsilon|$ is even, this case is treated in Lemma 5.7. So, let $|\epsilon|$ be odd. We compute

$$wx = ((-1)^{\epsilon_1} x_{n_1}, -x_1, x_2, x_3, \dots, x_{n_1-1}, x_{n_1+1}, \dots, x_\ell).$$

Hence

$$(qx - wx, \alpha_0) = q(x_1 + x_2) + (x_1 + (-1)^{\epsilon_1+1} x_{n_1}) > 0,$$

$$(qx - wx, \alpha_1) = q(x_1 - x_2) - x_1 - (-1)^{\epsilon_1} x_{n_1},$$

$$(qx - wx, \alpha_2) = q(x_2 - x_3) + (x_1 + x_2) > 0,$$

$$(qx - wx, \alpha_3) = q(x_3 - x_4) - (x_2 - x_3),$$

$$\vdots$$

$$(qx - wx, \alpha_{n_1-1}) = q(x_{n_1-1} - x_{n_1}) - (x_{n_1-2} - x_{n_1-1}),$$

$$(qx - wx, \alpha_{n_1}) = q(x_{n_1} - x_{n_1+1}) - (x_{n_1-1} - x_{n_1+1}).$$

If ϵ_1 is even then we choose $x_1 > x_2 > \dots > x_n > 0$ in the following way. Let $x_2 > 0$ be arbitrary and set as in Lemma 5.7 $x_{2+i} = x_2 - ia$, $i = 0, \dots, n-3$, with $a > 0$ small enough. Further, let $x_n > 0$ be such that $x_{n-1} > 3x_n$. Finally choose $x_1 > x_2$ such that $x_1 - x_2 > x_2 - x_n$. One checks that the above expressions are positive. If ϵ_1 is odd then one chooses x similarly.

Subcase. $m_1 > 2$. In this case, one reduces by Lemma 5.7 to the situation of $w = s_1$ resp. $w = s_1 \cdot t$.

Case. $\delta = t'$.

Subcase. $m_1 = 2$. Then $w = \delta w_{\lambda_1, \epsilon_1}$ is given by

$$e_1 \mapsto (-1)^{|\epsilon|+1} e_2,$$

$$e_2 \mapsto e_3, e_3 \mapsto e_4, \dots, e_{n_1-1} \mapsto e_{n_1}, e_{n_1} \mapsto (-1)^{\epsilon_1+1} e_1.$$

These cases are already covered by the previous one.

Subcase. $m_1 > 2$. In this case, one reduces by Lemma 5.7 to the situation of $w = t'$ resp. $w = t' \cdot t$. \square

Note that the x in Proposition 5.9 lies in C .

Acknowledgments

We thank L. Illusie and Th. Zink for helpful discussions on ℓ -adic cohomology.

References

- [BR] C. Bonnafé, R. Rouquier, On the irreducibility of Deligne–Lusztig varieties, *C. R. Math. Acad. Sci. Paris* 343 (1) (2006) 37–39.
- [BR'] C. Bonnafé, R. Rouquier, Affineness of Deligne–Lusztig varieties for minimal length elements, *J. Algebra* 320 (3) (2008) 1200–1206.
- [Bou] N. Bourbaki, Lie Groups and Lie Algebras, Chapters 4–6. Elements of Mathematics, Springer-Verlag, Berlin, 2002, xii+300 pp.
- [DL] P. Deligne, G. Lusztig, Representations of reductive groups over finite fields, *Ann. of Math. (2)* 103 (1) (1976) 103–161.
- [DMR] F. Digne, J. Michel, R. Rouquier, Cohomologie des variétés de Deligne–Lusztig, *Adv. Math.* 209 (2) (2007) 749–822.
- [F] G. Faltings, Mumford-Stabilität in der algebraischen Geometrie, in: Proceedings of the International Congress of Mathematicians, vols. 1, 2, Zürich, 1994, Birkhäuser, Basel, 1995, pp. 648–655.
- [GP] M. Geck, G. Pfeiffer, On the irreducible characters of Hecke algebras, *Adv. Math.* 102 (1) (1993) 79–94.
- [GKP] M. Geck, S. Kim, G. Pfeiffer, Minimal length elements in twisted conjugacy classes of finite Coxeter groups, *J. Algebra* 229 (2) (2000) 570–600.
- [H] B. Haastert, Die Quasiaffinität der Deligne–Lusztig–Varietäten, *J. Algebra* 102 (1986) 186–193.
- [H'] B. Haastert, Über die Deligne–Lusztig Varietäten reductiver Gruppen, Diplomarbeit Universität, Bonn, 1984.
- [He] X. He, On the affineness of Deligne–Lusztig varieties, *J. Algebra* 320 (3) (2008) 1207–1219.
- [L] G. Lusztig, Representations of finite Chevalley groups, in: Expository Lectures from the CBMS Regional Conference Held at Madison, Wis., August 8–12, 1977, in: CBMS Reg. Conf. Ser. Math., vol. 39, American Mathematical Society, Providence, RI, 1978, v+48 pp.
- [L'] G. Lusztig, Coxeter orbits and eigenspaces of Frobenius, *Invent. Math.* 38 (2) (1976/77) 101–159.
- [M] J. Milne, Lectures on étale cohomology, <http://www.jmilne.org/math/>.
- [O] S. Orlik, The cohomology of period domains for reductive groups over finite fields, *Ann. Sci. École Norm. Sup. (4)* 34 (1) (2001) 63–77.
- [R] M. Rapoport, Period domains over finite and local fields, in: Algebraic Geometry, Part 1, Santa Cruz, 1995, in: Proc. Sympos. Pure Math., vol. 62, American Mathematical Society, Providence, RI, 1997, pp. 361–381.
- [S] J.-P. Serre, Galois Cohomology, Translated from the French by Patrick Ion and revised by the author. Corrected reprint of the 1997 English edition, Springer Monogr. Math., Springer-Verlag, Berlin, 2002, x+210 pp.
- [SGA1] Revêtements étales et groupe fondamental (SGA 1). Séminaire de géométrie algébrique du Bois Marie 1960–1961, Directed by A. Grothendieck. With two papers by M. Raynaud. Updated and annotated reprint of the 1971 original Lecture Notes in Math., vol. 224, Springer-Verlag, Berlin, 1971, Doc. Math. (Paris), vol. 3, Société Mathématique de France, Paris, 2003, xviii+327 pp.
- [SGA4'] P. Deligne, Cohomologie étale, Séminaire de Géométrie Algébrique du Bois-Marie SGA 4 $\frac{1}{2}$. Avec la collaboration de J.F. Boutot, A. Grothendieck, L. Illusie et J.L. Verdier, Lecture Notes in Math., vol. 569, Springer-Verlag, Berlin/New York, 1977, iv+312 pp.