



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



On unique factorization domains

Jim Coykendall^{a,*}, William W. Smith^b

^a Department of Mathematics, North Dakota State University, Fargo, ND 58105-5075, United States

^b Department of Mathematics, The University of North Carolina, Chapel Hill, NC 27599, United States

ARTICLE INFO

Article history:

Received 8 April 2009

Available online 26 February 2011

Communicated by Steven Dale Cutkosky

MSC:

13F15

Keywords:

Factorization

Unique factorization

ABSTRACT

In this paper we attempt to generalize the notion of “unique factorization domain” in the spirit of “half-factorial domain”. It is shown that this new generalization of UFD implies the now well-known notion of half-factorial domain. As a consequence, we discover that one of the standard axioms for unique factorization domains is slightly redundant.

© 2011 Published by Elsevier Inc.

1. Introduction and motivation

The notion of unique factorization is one that is central in the study of commutative algebra. A unique factorization domain (UFD) is an integral domain, R , where every nonzero nonunit can be factored uniquely. More formally we record the following standard definition.

Definition 1.1. We say that an integral domain, R , is a UFD if every nonzero nonunit in R can be factored into irreducible elements, and if we have

$$\alpha_1 \alpha_2 \cdots \alpha_n = \beta_1 \beta_2 \cdots \beta_m$$

with each α_i, β_j irreducible in R then

(a) $n = m$ and

(b) there is a $\sigma \in S_n$ such that $\alpha_i = u_i \beta_{\sigma(i)}$ for all $1 \leq i \leq n$ where each u_i is a unit of R .

* Corresponding author.

E-mail addresses: jim.coykendall@ndsu.edu (J. Coykendall), wwsmith@email.unc.edu (W.W. Smith).

We remark here that a domain with the property that every nonzero nonunit of R can be factored into irreducibles (or atoms) is said to be *atomic*. So a UFD is an atomic domain that satisfies conditions (a) and (b) above. We will use this terminology hereafter.

In 1960 the notion of the half-factorial property first appeared in a paper of Carlitz [3]. The terminology “half-factorial domain” first appeared when the results from [3] were generalized by Zaks [9]. Since that time many factorization properties have been investigated by a number of authors (see [1,2], among others). The half-factorial property has gotten a bit of attention as well (for example see [4,6–8]).

Historically, half-factorial domains have “half the axioms” of UFDs (namely axiom (a) from above). In a question of the second author, it was asked if there are nontrivial examples of “other half” factorial domains (that is, domains that have the “other half” of the axioms of UFDs: namely (b) above). To be more precise (and with apologies to the reader) we make the following definition.

Definition 1.2. We say that the atomic integral domain, R , is an OHFD if given

$$\alpha_1 \alpha_2 \cdots \alpha_n = \beta_1 \beta_2 \cdots \beta_n$$

with each α_i, β_j irreducible in R then there is a $\sigma \in S_n$ such that $\alpha_i = u_i \beta_{\sigma(i)}$ for all $1 \leq i \leq n$ where each u_i is a unit of R .

Intuitively speaking, an OHFD is an atomic domain where a given element may have many factorizations of different lengths, but for any positive integer n , the element has at most one factorization of length n .

Since its inception, the concept of half-factorial has borne much fruit. This lends much promise to the concept of OHFD introduced above. We will show, however, that this definition is only a temporary one. It is shown in the next section that any OHFD is an HFD. Since the two properties in tandem obviously imply UFD, it will follow that any OHFD is a UFD. We find it interesting that, as a consequence, the standard definition of UFD can be weakened considerably by essentially eliminating condition (a).

2. Properties of an OHFD

We note at the outset of this section that although the title is “properties of an OHFD”, the results that we present carry through in the context of monoids until Theorem 2.10. Indeed, the results numbered 2.1 through 2.9 only depend on the (multiplicative) monoid structure of our OHFD and hence could be considered results for an “OHFM”. In Theorem 2.10 and its corollaries, the additive structure of the OHFD comes into play (and so are results that apply strictly in the context of domains). In Example 2.13 we underscore the importance of the additive structure by presenting an example of an OHFM which does not have the HFM property.

To get to the heart of the factorization properties of an OHFD, we need to introduce the notion of “nondegenerate factorizations”. The motivation of this approach is simply the observation that any two factorizations of nonequal lengths

$$\pi_1 \pi_2 \cdots \pi_n = \xi_1 \xi_2 \cdots \xi_m$$

give rise to the nonequal length factorizations

$$\pi_1 \pi_2 \cdots \pi_n \alpha_1 \alpha_2 \cdots \alpha_t = \xi_1 \xi_2 \cdots \xi_m \alpha_1 \alpha_2 \cdots \alpha_t.$$

To streamline our study we make the following definition.

Definition 2.1. Let R be an integral domain. We say that a pair of irreducible factorizations

$$\pi_1 \pi_2 \cdots \pi_n = \xi_1 \xi_2 \cdots \xi_m$$

is mutually nondegenerate if the irreducibles $\pi_i \in R$ and $\xi_j \in R$ are pairwise non-associate.

It is easy to see that any two (nonunique) irreducible factorizations of an element of a ring can be reduced to a pair of mutually nondegenerate factorizations by canceling the associate pairs.

Definition 2.2. Let R be an atomic domain and $\pi_1 \in R$ an irreducible element. We say that π_1 is “long” (resp. “short”) if there is a pair of mutually nondegenerate irreducible factorizations

$$\pi_1 \pi_2 \cdots \pi_n = \xi_1 \xi_2 \cdots \xi_m$$

with $\pi_i, \xi_j \in R$ irreducible and $n > m$ (resp. $n < m$).

Although it is clear that prime elements are neither long nor short, it is not clear that assigning this label to a nonprime irreducible is well defined. We will show that for an OHFD this assignment makes sense.

Proposition 2.3. Suppose that R is an OHFD and let

$$\pi_1 \pi_2 \cdots \pi_k = \xi_1 \xi_2 \cdots \xi_m$$

and

$$\alpha_1 \alpha_2 \cdots \alpha_r = \beta_1 \beta_2 \cdots \beta_n$$

be two pairs of mutually nondegenerate irreducible factorizations with $k > m$ and $n > r$. Then each π_i is associate to some β_j and conversely (this holds as well as the α_i 's and ξ_j 's).

Proof. Let $a := n - r$ and $b := k - m$. Note that in the factorizations

$$\pi_1^a \pi_2^a \cdots \pi_k^a \alpha_1^b \alpha_2^b \cdots \alpha_r^b = \xi_1^a \xi_2^a \cdots \xi_m^a \beta_1^b \beta_2^b \cdots \beta_n^b$$

the lengths are the same (since $ak + br = am + bn$). Since R is an OHFD, each π_i must appear on the right side of the equality. Hence each π_i is an associate of some β_j (since all π_i 's and ξ_j 's are non-associate). By symmetry, it is also the case that each β_j is an associate of some π_i . The same argument shows that the α_i 's and ξ_j 's are associate. \square

Proposition 2.4. Let R be an OHFD that is not an HFD and $\pi \in R$ a nonprime irreducible. Then π cannot be both long and short.

Proof. Using the notation from above, assume that π_1 is both long and short. Replace α_1 by π_1 in the hypothesis of the above result. Hence we obtain that π_1 is associate to some ξ_i (and clearly π_1 is self-associate). This contradicts the mutual nondegeneracy of the first pair of factorizations. \square

Corollary 2.5. *Let R be an OHFD. Then R has only finitely many long (resp. short) irreducibles up to associates. What is more, given any pair of irreducible factorizations*

$$\pi_1 \pi_2 \cdots \pi_k = \xi_1 \xi_2 \cdots \xi_m$$

with $k > m$ and an arbitrary long (resp. short) irreducible β (resp. α), β is an associate of some π_i (resp. α is an associate of some ξ_i).

Proof. Again using the notation of the above we let $\{\pi_1, \pi_2, \dots, \pi_k\}$ be the (long) irreducibles involved in the factorization

$$\pi_1 \pi_2 \cdots \pi_k = \xi_1 \xi_2 \cdots \xi_m$$

with $k > m$.

Now suppose that β is an arbitrary long irreducible. By definition, β is involved in some pair of mutually nondegenerate, different length factorizations (with β on the “long” side). By the above, β is an associate of some π_i . The proof for short irreducibles is identical. \square

We now observe that any irreducible in an OHFD that is neither long nor short is necessarily prime.

Lemma 2.6. *Let R be an OHFD. If $x \in R$ is an irreducible element that is neither long nor short, then x is prime.*

Proof. Assume that $x \mid ab$ with $a, b \in R$. So there is a $c \in R$ such that

$$cx = ab.$$

We produce irreducible factorizations $c = c_1 c_2 \cdots c_r$, $b = b_1 b_2 \cdots b_s$, and $a = a_1 a_2 \cdots a_t$ and write

$$c_1 c_2 \cdots c_r x = a_1 a_2 \cdots a_t b_1 b_2 \cdots b_s.$$

Since x is neither long nor short there are two cases to consider. In the first case, the factorizations above are of equal length (in which case they are unique) and x is an associate of some b_i or a_j . In the second case the factorizations are not of equal length. In this case, since x is not associated to any long or short irreducible, the factorizations cannot be mutually nondegenerate. Hence (again) x is associated to some b_i or a_j . In either case, we get that $x \mid a$ or $x \mid b$. \square

Now that we have some preliminary results in hand, we will adopt a more convenient notation. We have not demanded so far that π_i is non-associate to π_j if $i \neq j$ (and indeed there is no reason to expect this to be true). We will now list our “canonical bad length factorizations” in a form more convenient for future computational purposes:

$$\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} = \xi_1^{b_1} \xi_2^{b_2} \cdots \xi_m^{b_m}$$

with $\sum_{i=1}^k a_i > \sum_{i=1}^m b_i$, and all irreducibles pairwise non-associate.

In the next result, we show that there is essentially one pair of “master factorizations” which itself produces all the nondegenerate factorizations of different length.

Definition 2.7. Let R be an OHFD, and let $\{\pi_1, \pi_2, \dots, \pi_k\}$ and $\{\xi_1, \xi_2, \dots, \xi_m\}$ respectively denote the sets of long and short irreducibles in R (up to associates). Among all pairs of factorizations

$$\pi_1^{a_1} \pi_2^{a_2} \dots \pi_k^{a_k} = \xi_1^{b_1} \xi_2^{b_2} \dots \xi_m^{b_m}$$

we select one with a_1 minimal and call this pair of factorizations the master factorizations (MF).

Remark 2.8. Note that if R is an OHFD then Corollary 2.5 shows that each $a_i, b_j > 0$.

Proposition 2.9. Let R be an OHFD and let

$$\pi_1^{a_1} \pi_2^{a_2} \dots \pi_k^{a_k} = \xi_1^{b_1} \xi_2^{b_2} \dots \xi_m^{b_m}$$

be the MF. Then any pair of mutually nondegenerate irreducible factorizations is a power of this MF. That is, any two mutually nondegenerate irreducible factorizations of different lengths are of the form (up to associates)

$$\pi_1^{a_1 t} \pi_2^{a_2 t} \dots \pi_k^{a_k t} = \xi_1^{b_1 t} \xi_2^{b_2 t} \dots \xi_m^{b_m t}$$

for some $t \geq 1$.

Proof. We know by Corollary 2.5 that any two mutually nondegenerate irreducible factorizations of different lengths must be of the form

$$\pi_1^{c_1} \pi_2^{c_2} \dots \pi_k^{c_k} = \xi_1^{d_1} \xi_2^{d_2} \dots \xi_m^{d_m}$$

with each $c_i, d_j > 0$.

We also have that $A := \sum_{i=1}^k a_i > B := \sum_{i=1}^m b_i$ and $C := \sum_{i=1}^k c_i > D := \sum_{i=1}^m d_i$. We apply the same technique as earlier by setting $s := C - D$ and $r := A - B$. We now look at the factorizations

$$\pi_1^{a_1 s} \pi_2^{a_2 s} \dots \pi_k^{a_k s} \xi_1^{d_1 r} \xi_2^{d_2 r} \dots \xi_m^{d_m r} = \pi_1^{c_1 r} \pi_2^{c_2 r} \dots \pi_k^{c_k r} \xi_1^{b_1 s} \xi_2^{b_2 s} \dots \xi_m^{b_m s}$$

which are of equal length as $sA + rD = AC - AD + AD - BD = AC - BC + BC - BD = rC + sB$. Since R is an OHFD we have that $sa_i = rc_i$ for all $1 \leq i \leq k$ and $sb_i = rd_i$ for all $1 \leq i \leq m$. In particular, $\frac{s}{r}a_1 = c_1$ and by the minimality of a_1 , we have that $\frac{s}{r} \geq 1$. From this it follows that $a_i \leq c_i$ for all $1 \leq i \leq k$ and $b_i \leq d_i$ for all $1 \leq i \leq m$.

We next claim that for all $1 \leq i \leq k$, a_i divides c_i and for all $1 \leq j \leq m$, b_j divides d_j (and what is more the quotients $\frac{c_i}{a_i}$ and $\frac{d_j}{b_j}$ are all the same). We simultaneously apply the Euclidean algorithm to all of the exponents above and obtain the systems of equations

$$c_i = q_i a_i + r_i \quad \text{and} \quad d_j = Q_j b_j + R_j$$

where each remainder r_i, R_j satisfies $0 \leq r_i < a_i$ and $0 \leq R_j < b_j$.

We again consider the factorizations

$$\pi_1^{c_1} \pi_2^{c_2} \dots \pi_k^{c_k} = \xi_1^{d_1} \xi_2^{d_2} \dots \xi_m^{d_m}.$$

Since each $a_i \leq c_i$ and $b_i \leq d_i$ we can divide the left side of the above equation by $\pi_1^{a_1} \pi_2^{a_2} \dots \pi_k^{a_k}$ and in tandem divide the right side by $\xi_1^{b_1} \xi_2^{b_2} \dots \xi_m^{b_m}$. After applying this simple algorithm a number of times, there is a first occurrence where at least one of the exponents of either π_i (for some i) or ξ_j (for some j) is equal to some r_i or R_j . This gives the factorizations

$$\pi_1^{v_1} \pi_2^{v_2} \cdots \pi_k^{v_k} = \xi_1^{w_1} \xi_2^{w_2} \cdots \xi_m^{w_m}$$

where one of the v_i 's is r_i or one of the w_i 's is R_i . If, at this step each v_i and w_i is 0, then we are done. If not, recall that we have shown that if the factorizations above are not of equal length, every v_i is of the form $\frac{s}{r}a_i$ and that every w_i is of the form $\frac{s}{r}b_i$ (with s and r chosen as above, $\frac{s}{r} \geq 1$), we obtain that

$$v_i \geq a_i > r_i \quad \text{and} \quad w_i \geq b_i > R_i$$

which is a contradiction. So at some point in the algorithm, we must have achieved equal length (and hence unique) factorizations. But this is clearly a contradiction since the irreducibles π_i are pairwise non-associate with the irreducibles ξ_j . Hence we must have $v_i = 0$ and $w_i = 0$ (for all i) at the same step of the algorithm. This means that $q_i = q_j = Q_i = Q_j$ for all i, j and $r_i = 0 = R_j$ for all i, j . This concludes the proof. \square

We now introduce the main theorem. At this juncture, our results depart from dependence only upon the multiplicative structure of our OHFD.

Theorem 2.10. *If R is an OHFD then R is an HFD.*

Proof. Assume that R is an OHFD that is not an HFD. In keeping with the notation above we will let our MF be

$$\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} = \xi_1^{b_1} \xi_2^{b_2} \cdots \xi_m^{b_m}$$

with $\sum_{i=1}^k a_i > \sum_{i=1}^m b_i$.

There are a number of cases to consider.

Case 1: $k \geq 2$ and $m \geq 2$.

In this case we consider the element

$$(\pi_1^{a_1} - \xi_1^{b_1})(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_2^{b_2} \xi_3^{b_3} \cdots \xi_m^{b_m}).$$

It is easy to see that π_1 divides the product displayed above. We also observe that π_1 divides neither $\xi_1^{b_1}$ nor $\xi_2^{b_2} \xi_3^{b_3} \cdots \xi_m^{b_m}$. Indeed, if π_1 divides $\xi_2^{b_2} \xi_3^{b_3} \cdots \xi_m^{b_m}$ then we have that there is a $c \in R$ such that

$$c\pi_1 = \xi_2^{b_2} \xi_3^{b_3} \cdots \xi_m^{b_m}$$

and since π_1 is not associated with any of the ξ_i 's, then these must be (after factoring c) factorizations of unequal lengths. Since the factorizations are mutually nondegenerate with respect to π_1 the left-hand side must be the “long” side. But now, Proposition 2.9 demands that ξ_1 must divide the right-hand side. So we now have that there is a $d \in R$ such that

$$d\xi_1 = \xi_2^{b_2} \xi_3^{b_3} \cdots \xi_m^{b_m}$$

but since ξ_1 cannot be long and short, this means that the factorizations above are of equal length or the factorizations are mutually degenerate. In either case, ξ_1 is associated with ξ_i for some $i > 1$ which is a contradiction. The proof that π_1 does not divide $\xi_1^{b_1}$ is similar.

With this in hand we obtain that

$$(\pi_1^{a_1} - \xi_1^{b_1})(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m}) = k\pi_1.$$

Factoring $(\pi_1^{a_1} - \xi_1^{b_1}) = \alpha_1 \alpha_2 \cdots \alpha_s$ and $(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m}) = \beta_1 \beta_2 \cdots \beta_t$, we obtain

$$\alpha_1 \alpha_2 \cdots \alpha_s \beta_1 \beta_2 \cdots \beta_t = k\pi_1$$

with α_i and β_j irreducible. The previous argument shows that π_1 is not associated to any of the α_i 's or β_j 's. So these must be factorizations of different lengths, and since π_1 is not associated to any irreducible on the left-hand side, this means that the right-hand side must be the “long” side.

We now apply Proposition 2.9 to obtain that ξ_1 must be associated with one of the irreducibles on the left side. In particular, this means that ξ_1 must divide either $(\pi_1^{a_1} - \xi_1^{b_1})$ or $(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m})$. Of course this means that ξ_1 must divide either $\pi_1^{a_1}$ or $\xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m}$. But if ξ_1 divides $\pi_1^{a_1}$ then $\pi_1^{a_1} = c\xi_1$ and these factorizations are of different length (after factoring c). Similar to the arguments before, we have a contradiction since π_i is not associated to π_1 if $i \neq 1$. Also if ξ_1 divides $\xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m}$ then $c\xi_1 = \xi_2^{b_2} \xi_3^{b_2} \cdots \xi_m^{b_m}$ and since ξ_1 is short, it must be associated to some ξ_i for $i > 1$ which is a contradiction. So we cannot possess MFs of these forms.

Case 2: $k \geq 2$ and $m = 1$ or $k = 1$ and $m \geq 2$.

These cases are symmetric and we will only show the case $k \geq 2$ and $m = 1$. In this case it is clear that $b_1 \geq 2$. First consider the MF

$$\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} = \xi_1^{b_1}.$$

And now consider the element

$$(\pi_1^{a_1} - \xi_1)(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_1^{b_1-1}).$$

As before, π_1 divides this product, but neither of the two elements in the product. If we (irreducibly) factor $(\pi_1^{a_1} - \xi_1) = \alpha_1 \alpha_2 \cdots \alpha_s$ and $(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_1^{b_1-1}) = \beta_1 \beta_2 \cdots \beta_t$ then we have

$$\alpha_1 \alpha_2 \cdots \alpha_s \beta_1 \beta_2 \cdots \beta_t = k\pi_1.$$

Since π_1 does not divide any of the irreducibles on the left side, the above must be factorizations of different lengths. Proposition 2.9 again shows that the right side of the equation is the long side and that ξ_1 must be associated to some (in fact all) nondegenerate irreducibles on the left. In particular, it must be the case that at least b_1 of the left irreducibles are associated to ξ_1 (as the short side of the pair of the master factorizations has b_1 factors of ξ_1). Note that since ξ_1 does not divide $(\pi_1^{a_1} - \xi_1)$ it cannot be associated to any α_i . Hence (at least) b_1 of the β_i 's must be associated to ξ_1 . This implies that $\xi_1^{b_1}$ divides $(\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k} - \xi_1^{b_1-1})$. Since $\xi_1^{b_1}$ divides $\pi_1^{a_1} \pi_2^{a_2} \cdots \pi_k^{a_k}$, it must divide $\xi_1^{b_1-1}$ which is a contradiction.

Case 3: $k = m = 1$.

In this final case, our MF is of the form

$$\pi^a = \xi^b$$

with $a > b \geq 2$. In this case we note that any R -linear combination of π and ξ is a nonunit (indeed, note that if \mathfrak{M} is a maximal ideal of R such that $\pi \in \mathfrak{M}$ then certainly $\xi \in \mathfrak{M}$).

We consider

$$(\pi - \xi)(\pi^{a-1} - \xi^{b-1}).$$

As before, π divides this product and we get

$$(\pi - \xi)(\pi^{a-1} - \xi^{b-1}) = k\pi,$$

and as before we factor $(\pi - \xi) = \alpha_1\alpha_2 \cdots \alpha_s$ and $(\pi^{a-1} - \xi^{b-1}) = \beta_1\beta_2 \cdots \beta_t$ to obtain

$$\alpha_1\alpha_2 \cdots \alpha_s\beta_1\beta_2 \cdots \beta_t = k\pi.$$

Note that, similarly to the previous cases, π cannot divide any of the irreducibles on the left (as, again, this forces π to be a divisor of either $(\pi - \xi)$ or $(\pi^{a-1} - \xi^{b-1})$ and the techniques of earlier cases show that this cannot happen). Hence these (after factoring k) are two factorizations of different lengths.

Proposition 2.9 shows that each α_i and β_j is divisible by ξ hence ξ divides $(\pi - \xi)$. Hence ξ divides π and this is a contradiction.

This exhausts the cases for the form of an MF, and hence there is no OHFD that is not an HFD. \square

Here is a rather striking corollary.

Corollary 2.11. *An integral domain is an OHFD if and only if it is a UFD.*

Proof. (\Leftarrow) is obvious. On the other hand, if R is an OHFD then R is an HFD and clearly these two properties in tandem imply UFD. \square

One interesting consequence of this is that it suffices to declare that an atomic integral domain is a UFD if any two equal length factorizations are the same (up to units and reordering). So in a certain sense, the condition (a) from the definition of UFD in the introduction is not needed.

Another useful result is that we can now retire the definition of OHFD (since the concept is equivalent to UFD). Before we dispose of it, however, we list an immediate corollary that is of some theoretical and aesthetic interest.

Corollary 2.12. *Any atomic domain that is not a UFD has an element with two distinct factorizations of the same length.*

Proof. If not then R is an OHFD and hence a UFD. \square

As a final note, it should be pointed out that although we have retired this “OHFD” notion for integral domains, this notion of “OHFM” for monoids can exist apart from the notion of “UFM” (see [5] for a far more complete discussion of factorization in the setting of monoids). It is of interest to once again point out that our results (until Theorem 2.10) can be couched in terms of monoid factorizations.

We conclude with an example highlighting contrast. This example, in conjunction with the proof of the main theorem, plainly illustrates some of the hazards that can appear in making the transition between the setting of monoids to the setting of integral domains.

Example 2.13. Consider the natural numbers \mathbb{N} and consider the additive submonoid of \mathbb{N} generated by $\{n, m\}$ where $2 \leq n < m$ and $\gcd(n, m) = 1$. To make comparisons with the earlier part of this paper easier, we now consider the multiplicative monoid generated by $a = e^n$ and $b = e^m$.

In the monoid $M = \langle a, b \rangle$, the only two irreducibles are the elements a and b and hence the only nondegenerate factorizations are of the form

$$a^x = b^y.$$

In fact, a is “long” and b is “short” since $a^m = b^n$ and $n < m$, and these particular factorizations ($a^m = b^n$) are easily seen to be the MF for this monoid and are also certainly examples of factorizations of the same element possessing different lengths.

To see that M is an OHFM we suppose that we have the factorizations

$$a^{k_1} b^{k_2} = a^{s_1} b^{s_2}$$

with $k_1 + k_2 = s_1 + s_2$.

This gives that $k_1 n + k_2 m = s_1 n + s_2 m$, which in turn yields

$$(k_1 - s_1)n = (s_2 - k_2)m.$$

Since n and m are relatively prime, we have that there is an $x \in \mathbb{Z}$ such that

$$nx = s_2 - k_2$$

and

$$mx = k_1 - s_1.$$

From this we obtain

$$(m - n)x = (k_1 + k_2) - (s_1 + s_2) = 0$$

since the factorizations are of the same length. Hence $x = 0$ (since $m \neq n$) and so $s_1 = k_1$ and $s_2 = k_2$. This gives the uniqueness of the factorization.

Acknowledgment

The authors thank the referee for very helpful comments.

References

- [1] D.D. Anderson, D.F. Anderson, M. Zafrullah, Factorization in integral domains, *J. Pure Appl. Algebra* 69 (1990) 1–19.
- [2] D.D. Anderson, D.F. Anderson, M. Zafrullah, Factorization in integral domains II, *J. Algebra* 152 (1992) 78–93.
- [3] L. Carlitz, A characterization of algebraic number fields with class number two, *Proc. Amer. Math. Soc.* 11 (1960) 391–392.
- [4] S. Chapman, J. Coykendall, Half-factorial domains, a survey, in: *Non-Noetherian Commutative Ring Theory*, Kluwer, 2000, pp. 97–115.
- [5] S.T. Chapman, A. Geroldinger, Krull domains and monoids, their sets of lengths, and associated combinatorial problems, in: *Lect. Notes Pure Appl. Math.*, vol. 189, Marcel Dekker, 1997, pp. 73–112.
- [6] S. Chapman, W.W. Smith, On the HFD, CHFD, and k -HFD properties in Dedekind domains, *Comm. Algebra* 20 (1992) 1955–1987.
- [7] J. Coykendall, A characterization of polynomial rings with the half-factorial property, in: *Lect. Notes Pure Appl. Math.*, vol. 189, Marcel Dekker, 1997, pp. 291–294.
- [8] J. Coykendall, On the integral closure of a half-factorial domain, *J. Pure Appl. Algebra* 180 (2003) 25–34.
- [9] A. Zaks, Half factorial domains, *Bull. Amer. Math. Soc.* 82 (1976) 721–723.