



ELSEVIER

Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



## Coverings of lura algebras: The standard case

Ibrahim Assem<sup>a</sup>, Juan Carlos Bustamante<sup>b</sup>, Patrick Le Meur<sup>c,\*</sup>

<sup>a</sup> Département de Mathématiques, Université de Sherbrooke, Sherbrooke, Québec, Canada J1K 2R1

<sup>b</sup> Departamento de Matemáticas, Universidad San Francisco de Quito, Cumbayá, Quito, Ecuador

<sup>c</sup> CMLA, ENS Cachan, CNRS, UniverSud, 61 Avenue du Président Wilson, F-94230 Cachan, France

### ARTICLE INFO

#### Article history:

Received 25 July 2008

Available online 23 October 2009

Communicated by Nicolás Andruskiewitsch

#### Keywords:

Representations of algebras

Covering theory

Galois coverings

Laura algebras

Hochschild cohomology

Simple connectedness

### ABSTRACT

In this paper, we study the covering theory of lura algebras. We prove that if a connected lura algebra is standard (that is, has a standard connecting component), then it has Galois coverings associated to the coverings of the connecting component. As a consequence, the first Hochschild cohomology group of a standard lura algebra vanishes if and only if it has no proper Galois coverings.

© 2009 Elsevier Inc. All rights reserved.

### Introduction

Introduced in 1945, the Hochschild cohomology groups are subtle and interesting invariants of associative algebras. The lower-dimensional groups have simple interpretations: for instance, the 0th group is the centre of the algebra, the 1st group can be thought of as the group of outer derivations of the algebra, while the 2nd and 3rd groups are related to the rigidity properties of the algebra. In [40, §3, Pb. 1], Skowroński has related the vanishing of the first Hochschild cohomology group  $\text{HH}^1(A)$  of an algebra  $A$  (with coefficients in the bimodule  ${}_A A_A$ ) to the simple connectedness of  $A$ . Recall that a basic and connected finite-dimensional algebra over an algebraically closed field  $k$  is *simply connected* if it has no proper Galois covering or, equivalently, if the fundamental group (in the sense of [33]) of any presentation is trivial. In particular, Skowroński posed the following problem: for which algebras  $A$  do we have  $\text{HH}^1(A) = 0$  if and only if  $A$  is simply connected? This problem has been the subject of several investigations: notably this equivalence holds true for al-

\* Corresponding author.

E-mail addresses: [ibrahim.assem@USherbrooke.ca](mailto:ibrahim.assem@USherbrooke.ca) (I. Assem), [jbustamante@ufsqu.edu.ec](mailto:jbustamante@ufsqu.edu.ec) (J.C. Bustamante), [patrick.lemeur@cmla.ens-cachan.fr](mailto:patrick.lemeur@cmla.ens-cachan.fr) (P. Le Meur).

gebras derived equivalent to hereditary algebras [31], weakly shod algebras [30] (see also [7]), large classes of selfinjective algebras [34] and schurian cluster-tilted algebras [10]. It was proved in [15] that, for a representation-finite algebra, the first Hochschild cohomology group vanishes if and only if its Auslander–Reiten quiver is simply connected. Note that if  $A$  is a representation-finite triangular algebra, then its Auslander–Reiten quiver is simply connected if and only if  $A$  has no proper Galois covering, that is,  $A$  is simply connected.

Here, we study this conjecture for lura algebras. These are defined as follows. Let  $\text{mod } A$  be the category of finitely generated right  $A$ -modules, and  $\text{ind } A$  be a full subcategory consisting of exactly one representative from each isomorphism class of indecomposable  $A$ -modules. The *left part*  $\mathcal{L}_A$  of  $\text{mod } A$  is the full subcategory of  $\text{ind } A$  consisting of those modules whose predecessors have projective dimension at most one, and the *right part*  $\mathcal{R}_A$  is defined dually. These classes were introduced in [24] in order to study the module categories of quasi-tilted algebras. Following [3,42], we say that  $A$  is *lura* provided  $\text{ind } A \setminus (\mathcal{L}_A \cup \mathcal{R}_A)$  has only finitely many objects. Part of the importance of lura algebras comes from the fact that this class contains (and generalises) the classes of representation-finite algebras, tilted, quasi-tilted and weakly shod algebras. Lura algebras have appeared naturally in the study of Auslander–Reiten components: an Auslander–Reiten component is called *quasi-directed* if it is generalised standard and almost all its modules are directed. It was shown in [3] that a lura algebra which is not quasi-tilted has a unique faithful convex quasi-directed Auslander–Reiten component (which is also the unique non-semiregular component). Conversely, any convex quasi-directed component occurs in this way [43]. The techniques used for the study of lura algebras were applied in [27] to obtain useful results on the infinite radical of the module category. Their representation dimension is at most three and this is a class of algebras with possibly infinite global dimension which satisfies the finitistic dimension conjecture [9]. Also, lura algebras have been characterised in terms of the Gabriel–Rojter measure as announced by Lanzilotta in the ICRA XI in Mexico, 2004 (see also [5]). For further properties of lura algebras we refer the reader to [3,4,6,8,42]. Here we concentrate on the conjecture that a lura algebra  $A$  is simply connected if and only if  $\text{HH}^1(A) = 0$ .

Our approach, already used in [30,31], uses coverings. Covering theory was introduced by Gabriel and his school (see, for instance, [14,21,36]) and consists in replacing an algebra by a locally bounded category, called its covering, which is sometimes easier to study. We recall that a tilted algebra is characterised by the existence of at least one, and at most two, *connecting components* (it has two if and only if it is concealed, in which case the connecting components are postprojective and preinjective) see [11]. If  $A$  is a lura not quasi-tilted algebra, then its unique faithful quasi-directed component is also called a connecting component (see [3]). Hence, by lura algebra *with connecting component*, we mean a connected lura algebra which is either tilted or not quasi-tilted. We call a lura algebra with connecting component *standard* provided its connecting components are all standard (it is known from [39] that the connecting components of concealed algebras are standard). This generalises the notion of standard representation-finite algebra (see [14]). Several classes of lura algebras are standard, notably tilted algebras or weakly shod algebras. Our first main theorem says that if  $\tilde{\Gamma} \rightarrow \Gamma$  is a Galois covering of the connecting component such that there exists a well-behaved covering functor  $k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  then it induces a covering of the algebra.

**Theorem A.** *Let  $A$  be a lura algebra with connecting component  $\Gamma$  and  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be a Galois covering with group  $G$  with respect to which there exists a well-behaved covering functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . Then there exists a covering functor  $F : A \rightarrow \tilde{A}$  whose fibres are in bijection with  $G$ . If moreover  $A$  is standard, then  $F$  is a Galois covering with group  $G$ .*

Note that if  $\Gamma$  is standard then there always exists a well-behaved covering functor  $p$ .

In order to prove Theorem A, we consider a more general situation. We first consider an Auslander–Reiten component, which contains a left section (in the sense of [1]) and show that to a Galois covering of this component such that there is a corresponding well-behaved functor corresponds a covering of its support algebra with nice properties, see Theorem 5.12 below. Applying this result to the connecting component of a lura algebra yields the required covering.

Because of the theorem, if  $A$  is standard, then we are able to work with Galois coverings which are notably easier to handle than covering functors. We prove that if  $A$  is standard lura, then any

Galois covering of the connecting component induces a Galois covering of  $A$ , with the same group. This allows us to prove our second main theorem, which settles the conjecture for standard lura algebras.

**Theorem B.** *Let  $A$  be a standard lura algebra, and  $\Gamma$  its connecting component(s). The following are equivalent:*

- (a)  $A$  has no proper Galois covering, that is,  $A$  is simply connected.
- (b)  $\text{HH}^1(A) = 0$ .
- (c)  $\Gamma$  is simply connected.
- (d) The orbit graph  $\mathcal{O}(\Gamma)$  is a tree.

Moreover, if these conditions are verified, then  $A$  is weakly shod.

If one drops the standard condition, then the above theorem may fail. Indeed, there are examples of non-standard representation-finite algebras which have no proper Galois covering and with non-zero first Hochschild cohomology group (see [14,15], or below). However, some implications are still true in Theorem B without assuming standardness. Indeed, we always have: (c) and (d) are equivalent and (c) implies (a) and (b).

Our paper is organised as follows. After a short preliminary section, we prove a few preparatory lemmata on covering functors in Section 2. In Section 3, we give examples of standard lura algebras. Section 4 is devoted to properties of tilting modules which are in the image of the push-down functor associated to a covering functor. In Section 5 we study the coverings of Auslander–Reiten components having left sections. The proof of Theorem A occupies Section 6. We concentrate on Galois coverings in Section 7, and prove Theorem B in Section 8.

## 1. Preliminaries

### Categories and modules

Throughout this paper,  $k$  denotes a fixed algebraically closed field. All our categories are locally bounded  $k$ -categories, in the sense of [14, 2.1]. We assume that all locally bounded  $k$ -categories are small and all functors are  $k$ -linear (the categories of finite-dimensional modules and their bounded derived categories are skeletally small).

Let  $F : \mathcal{E} \rightarrow \mathcal{B}$  be a  $k$ -linear functor and  $G$  be a group acting on  $\mathcal{E}$  and  $\mathcal{B}$  by automorphisms. Then  $F$  is called  $G$ -equivariant if  $F \circ g = g \circ F$  for every  $g \in G$ .

A basic finite-dimensional algebra  $A$  can be considered equivalently as a locally bounded  $k$ -category as follows: fix a complete set  $\{e_1, \dots, e_n\}$  of primitive orthogonal idempotents, then the object set of  $A$  is the set  $\{e_1, \dots, e_n\}$  and the morphisms space from  $e_i$  to  $e_j$  is  $e_j A e_i$ . The composition of morphisms is induced by the multiplication in  $A$ .

Let  $\mathcal{C}$  be a locally bounded  $k$ -category. We denote by  $\mathcal{C}_o$  its object class. A right  $\mathcal{C}$ -module  $M$  is a  $k$ -linear functor  $M : \mathcal{C}^{op} \rightarrow \text{MOD}k$ , where  $\text{MOD}k$  is the category of  $k$ -vector spaces. We write  $\text{MOD}\mathcal{C}$  for the category of  $\mathcal{C}$ -modules and  $\text{mod}\mathcal{C}$  for the full subcategory of the finite-dimensional  $\mathcal{C}$ -modules, that is, those modules  $M$  such that  $\sum_{x \in \mathcal{C}_o} \dim M(x) < \infty$ . If  $\mathcal{A}$  is a subcategory of  $\text{MOD}\mathcal{C}$ , we use the notation  $X \in \mathcal{A}$  to express that  $X$  is an object in  $\mathcal{A}$ . For every  $x \in \mathcal{C}_o$ , the indecomposable projective  $\mathcal{C}$ -module associated to  $x$  is  $\mathcal{C}(-, x)$ . The standard duality  $\text{Hom}_k(-, k)$  is denoted by  $D$ . Let  $M$  be a  $\mathcal{C}$ -module. If  $\mathcal{B}$  is a full subcategory of  $\mathcal{C}$ , then  $M|_{\mathcal{B}}$  is the induced  $\mathcal{B}$ -module. If  $\mathcal{X}$  is a subcategory of  $\text{mod}\mathcal{C}$ , then the  $\mathcal{X}$ -module  $\text{Hom}_{\mathcal{C}}(-, M)|_{\mathcal{X}}$  is denoted by  $\text{Hom}_{\mathcal{C}}(\mathcal{X}, M)$ . Also,  $\text{Hom}_{\mathcal{C}}(M, \mathcal{C})$  denotes the  $\mathcal{C}^{op}$ -module  $\text{Hom}_{\mathcal{C}}(M, \bigoplus_{x \in \mathcal{C}_o} \mathcal{C}(-, x))$  (if  $A = \mathcal{C}$  is a finite-dimensional algebra, this is just the left  $A$ -module  $\text{Hom}_A(M, A)$ ).

We let  $\text{ind}\mathcal{C}$  be a full subcategory of  $\text{mod}\mathcal{C}$  consisting of a complete set of representatives of the isomorphism classes of indecomposable  $\mathcal{C}$ -modules. We write  $\text{proj}\mathcal{C}$  and  $\text{inj}\mathcal{C}$  for the full subcategories of  $\text{ind}\mathcal{C}$  of projective and injective modules, respectively. Whenever we speak about an indecomposable  $\mathcal{C}$ -module, we always mean that it belongs to  $\text{ind}\mathcal{C}$ .

For a full subcategory  $\mathcal{A}$  of  $\text{mod } \mathcal{C}$ , we denote by  $\text{add } \mathcal{A}$  the full subcategory of  $\text{mod } \mathcal{C}$  with objects the direct sums of summands of modules in  $\mathcal{A}$ . If  $M$  is a module, then  $\text{add } M$  denotes  $\text{add}\{M\}$ .

The Auslander–Reiten translations in  $\text{mod } \mathcal{C}$  are denoted by  $\tau_{\mathcal{C}} = D\text{Tr}$  and  $\tau_{\mathcal{C}}^{-1} = \text{Tr}D$ . The Auslander–Reiten quiver of  $\mathcal{C}$  is denoted by  $\Gamma(\text{mod } \mathcal{C})$ . For a component  $\Gamma$  of  $\Gamma(\text{mod } \mathcal{C})$ , we denote by  $\mathcal{O}(\Gamma)$  its orbit graph (see [14, 4.2], or Section 8 below). The component  $\Gamma$  is *non-semiregular* if it contains both an injective and a projective module. It is *faithful* if its annihilator  $\text{Ann } \Gamma = \bigcap_{X \in \Gamma} \text{Ann } X$  is zero. Following [41], a component of  $\Gamma$  is *generalised standard* if  $\text{rad}^{\infty}(X, Y) = 0$  for every  $X, Y \in \Gamma$ . Denoting by  $k(\Gamma)$  the mesh category of  $\Gamma$  (see [14, 2.5]),  $\Gamma$  is *standard* if there exists an isomorphism of  $k$ -categories  $k(\Gamma) \xrightarrow{\sim} \text{ind } \Gamma$  which extends the identity on vertices, and which maps meshes to almost split sequences. Let  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be a morphism of translation quivers. Let  $\mathcal{X}$  be a full convex subquiver of  $\tilde{\Gamma}$ . We let  $k(\mathcal{X})$  be the full subcategory of  $k(\tilde{\Gamma})$  with objects the vertices in  $\mathcal{X}$ . Following [14, 3.1], a functor  $p : k(\mathcal{X}) \rightarrow \text{ind } \Gamma$  is *well-behaved* (with respect to  $\pi$ ) if it satisfies:

1.  $p(X) = \pi(X)$  for every  $X \in \mathcal{X}$ .
2. Let  $X \in \mathcal{X}$ . Let  $(u_i : Z_i \rightarrow X)_{i=1, \dots, t}$  be all the arrows in  $\mathcal{X}$  ending at  $X$  (or  $(v_j : X \rightarrow Y_j)_{j=1, \dots, s}$  be all the arrows in  $\mathcal{X}$  starting from  $X$ ), then the morphism  $[p(u_1) \ \dots \ p(u_t)] : \bigoplus_{i=1}^t p(Z_i) \rightarrow p(X)$  (or  $[p(v_1) \ \dots \ p(v_s)] : p(X) \rightarrow \bigoplus_{j=1}^s p(Y_j)$ , respectively) is irreducible.

Condition 2 above implies that if a mesh in  $\tilde{\Gamma}$  is contained in  $\mathcal{X}$ , then  $p$  maps this mesh to an almost split sequence.

For notions and results on modules, we refer the reader to [11]. For coverings and fundamental groups of translation quivers, we refer the reader to [14, §1]. Note that the translation quivers we use are not valued translation quivers and may have multiple arrows.

*Paths*

Let  $\mathcal{C}$  be a locally bounded  $k$ -category. Let  $X, Y$  be in  $\text{ind } \mathcal{C}$ . Following the convention used in [24], a *path*  $X \rightsquigarrow Y$  from  $X$  to  $Y$  in  $\text{ind } \mathcal{C}$  is a sequence of non-zero morphisms:

$$(\star) \quad X = X_0 \xrightarrow{f_1} X_1 \rightarrow \dots \rightarrow X_{t-1} \xrightarrow{f_t} X_t = Y \quad (t \geq 0),$$

where  $X_i \in \text{ind } \mathcal{C}$  for all  $i$ . We then say that  $X$  is a *predecessor* of  $Y$  and that  $Y$  is a *successor* of  $X$ . A path from  $X$  to  $X$  involving at least one non-isomorphism is a *cycle*. A module  $X \in \text{ind } \mathcal{C}$  which lies on no cycle is *directed*. If each  $f_i$  in  $(\star)$  is irreducible, we say that  $(\star)$  is a *path of irreducible morphisms* or a *path* in  $\Gamma(\text{mod } \mathcal{C})$ . A path  $(\star)$  of irreducible morphisms is *sectional* if  $\tau_{\mathcal{C}} X_{i+1} \neq X_{i-1}$  for all  $i$  with  $0 < i < t$ .

An indecomposable module  $M \in \mathcal{L}_A$  is *Ext-injective* in  $\text{add } \mathcal{L}_A$  if  $\text{Ext}_A^1(-, M)|_{\mathcal{L}_A} = 0$  (see [12]). This is the case if and only if  $\tau_A^{-1} M \notin \mathcal{L}_A$ .

The endomorphism algebra of the direct sum of the indecomposable projective modules lying in  $\mathcal{L}_A$  is called the *left support* of  $A$ . If  $A$  is lura with connecting component, then its left support is a product of tilted algebras (see [6, 4.4, 5.1]).

An algebra  $A$  is *weakly shod* if the length of any path in  $\text{ind } A$  from an injective to a projective is bounded [17]. Also,  $A$  is *quasi-tilted* if its global dimension  $\text{gl.dim } A$  is at most two and  $\text{ind } A = \mathcal{L}_A \cup \mathcal{R}_A$ , see [24].

**2. Covering functors**

A  $k$ -linear functor  $F : \mathcal{E} \rightarrow \mathcal{B}$  is a *covering functor* if (see [14, 3.1]):

1.  $F^{-1}(x) \neq \emptyset$  for every  $x \in \mathcal{B}_0$ .

2. For every  $x, y \in \mathcal{E}_0$ , the two following  $k$ -linear maps are bijective:

$$\bigoplus_{F(y')=F(y)} \mathcal{E}(x, y') \rightarrow \mathcal{B}(F(x), F(y)), \quad \text{and} \quad \bigoplus_{F(x')=F(x)} \mathcal{E}(x', y) \rightarrow \mathcal{B}(F(x), F(y)).$$

Following [21, §3],  $F$  is a *Galois covering with group*  $G$  if there exists a group morphism  $G \rightarrow \text{Aut}(\mathcal{E})$  such that  $G$  acts freely on  $\mathcal{E}_0$ ,  $F \circ g = F$  for every  $g \in G$  and the functor  $\mathcal{E}/G \rightarrow \mathcal{B}$  induced by  $F$  is an isomorphism. We refer the reader to [21, 3.1] for the definition of  $\mathcal{E}/G$ . Galois coverings are covering functors.

If  $F : \mathcal{E} \rightarrow \mathcal{B}$  is a covering functor, then  $F$  defines an adjoint pair  $(F_\lambda, F_*)$  of functors  $F_\lambda : \text{MOD } \mathcal{E} \rightarrow \text{MOD } \mathcal{B}$  and  $F_* : \text{MOD } \mathcal{B} \rightarrow \text{MOD } \mathcal{E}$  (see [14, 3.2]). The functor  $F_*$  is the *pull-up functor* and  $F_\lambda$  is the *push-down*. We recall their construction: If  $M \in \text{MOD } \mathcal{B}$ , then  $F_*M = M \circ F^{op}$ ; if  $M \in \text{MOD } \mathcal{E}$ , then  $F_\lambda M$  is the  $\mathcal{B}$ -module such that  $F_\lambda M(x) = \bigoplus_{F(\tilde{x})=x} M(\tilde{x})$ , for every  $x \in \mathcal{B}_0$ . Both  $F_\lambda$  and  $F_*$  are exact.

Let  $F : \mathcal{E} \rightarrow \mathcal{B}$  be a covering functor between locally bounded  $k$ -categories. We prove a few facts relative to  $F$ . Some are easy to prove in case  $F$  is a Galois covering. However, in general, the proofs are more complicated. This can be explained by the following fact:  $F^{op} : \mathcal{E}^{op} \rightarrow \mathcal{B}^{op}$  is also a covering functor, and  $DF_\lambda^{op} \simeq F_\lambda D$  if  $F$  is Galois. However, this isomorphism no longer exists in the general case of covering functors (see [14, 3.4], for instance).

As a motivation for the results in this section, we start with the following construction. We recall that the universal covering of a translation quiver  $\Gamma$  was introduced in [14, 1.2] using a homotopy relation denoted as  $H$ . We define  $\hat{H}$  to be the smallest equivalence relation containing  $H$  and satisfying the following additional relation: Let  $\alpha$  and  $\beta$  be two arrows in  $\Gamma$  having the same source and the same target, then  $\alpha$  and  $\beta$  are equivalent for  $\hat{H}$ . Using the construction of [14, 1.3] with respect to the relation  $\hat{H}$  we construct a covering of  $\Gamma$  which we call the *generic covering of*  $\Gamma$ . It is an immediate consequence of this definition and of [14, 1.3] that the generic covering is a Galois covering and is a quotient of the universal covering. They coincide if  $\Gamma$  has no multiple arrows (for example, if  $\Gamma$  is the Auslander–Reiten quiver of a representation-finite algebra).

The following proposition is mainly due to Riedtmann (see [36, 2.2]).

**Proposition 2.1.** *Let  $A$  be a basic finite-dimensional algebra. Let  $\Gamma$  be a component of  $\Gamma(\text{mod } A)$ . Let  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be the generic covering. Then there exists a well-behaved functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . If, moreover,  $\Gamma$  is generalised standard, then  $p$  is a covering functor.*

**Proof.** The functor  $p$  was constructed in [36, 2.2] for the stable part of the Auslander–Reiten quiver of a self-injective representation-finite algebra. The covering property was proved in [36, 2.3] under the same setting. The construction of  $p$  was generalised to any Auslander–Reiten component in [14, 3.1]. It is easily seen that the arguments given in [36, 2.3] to prove that  $p$  is a covering functor apply to the case of generalised standard components.  $\square$

Note that if  $\Gamma$  is a standard Auslander–Reiten component, then, by definition, there exists a well-behaved functor  $k(\Gamma) \rightarrow \text{ind } \Gamma$ . In particular, any covering of translation quivers  $p : \Gamma' \rightarrow \Gamma$  gives rise to a well-behaved covering functor  $k(\Gamma') \rightarrow \text{ind } \Gamma$  by composing the functors  $k(p) : k(\Gamma') \rightarrow k(\Gamma)$  and  $k(\Gamma) \rightarrow \text{ind } \Gamma$ .

The results of this section will be applied to covering functors as in 2.1. We now turn to the general situation where  $F : \mathcal{E} \rightarrow \mathcal{B}$  is a covering functor between locally bounded  $k$ -categories.

Since  $F_\lambda$  and  $F_*$  are exact, we still have an adjunction at the level of derived categories. Here and in the sequel,  $\mathcal{D}(\text{MOD } \mathcal{E})$  and  $\mathcal{D}^b(\text{mod } \mathcal{E})$  denote the derived category of  $\mathcal{E}$ -modules and the bounded derived category of finite-dimensional  $\mathcal{E}$ -modules, respectively. The following lemma is immediate. For a background on derived categories, we refer the reader to [22, Chap. III].

**Lemma 2.2.**  $F_\lambda$  and  $F$ , induce an adjoint pair  $(F_\lambda, F)$  of exact functors:

$$\mathcal{D}(\text{MOD } \mathcal{E}) \begin{array}{c} \xrightarrow{F_\lambda} \\ \xleftarrow{F} \end{array} \mathcal{D}(\text{MOD } \mathcal{B}).$$

Moreover  $F_\lambda(\mathcal{D}^b(\text{mod } \mathcal{E})) \subseteq \mathcal{D}^b(\text{mod } \mathcal{B})$ .

Let  $x \in \mathcal{E}_0$ . By condition 2 in the definition of a covering functor,  $F$  induces a canonical isomorphism  $F_\lambda(\mathcal{E}(-, x)) \xrightarrow{\sim} \mathcal{B}(-, F(x))$  of  $\mathcal{B}$ -modules (see [14, 3.2]). In the sequel, we always identify these two modules by means of this isomorphism. Using this identification we get the following result.

**Lemma 2.3.** Let  $M \in \mathcal{D}(\text{MOD } \mathcal{E})$ . Then  $F_\lambda$  induces two linear maps for every  $x \in \mathcal{B}_0$ :

$$\varphi_M : \bigoplus_{F(\tilde{x})=x} \mathcal{D}(\text{MOD } \mathcal{E})(M, \mathcal{E}(-, \tilde{x})) \rightarrow \mathcal{D}(\text{MOD } \mathcal{B})(F_\lambda M, \mathcal{B}(-, x)),$$

and

$$\psi_M : \bigoplus_{F(\tilde{x})=x} \mathcal{D}(\text{MOD } \mathcal{E})(\mathcal{E}(-, \tilde{x}), M) \rightarrow \mathcal{D}(\text{MOD } \mathcal{B})(\mathcal{B}(-, x), F_\lambda M).$$

These maps are functorial in  $M$ , and are bijective if  $M$  is quasi-isomorphic to a bounded complex of finite-dimensional projective modules (for example, if  $\text{gl.dim } \mathcal{E} < \infty$  and  $M \in \text{mod } \mathcal{E}$ ).

**Proof.** If  $M = P[l]$  where  $l \neq 0$  and  $P$  is a projective  $\mathcal{E}$ -module, then  $\varphi_M$  is bijective (because  $\text{Ext}_{\mathcal{E}}^{-l}(P, \mathcal{E}(-, \tilde{x})) = 0$ ). Also, if  $M$  is an indecomposable projective  $\mathcal{E}$ -module, then  $\varphi_M$  is bijective (because  $F$  is a covering functor). Finally, if  $M \rightarrow M' \rightarrow M'' \rightarrow M[1]$  is a triangle in  $\mathcal{D}(\text{MOD } \mathcal{E})$ , then  $\varphi_M, \varphi_{M'}$  and  $\varphi_{M''}$  are bijective as soon as two of them are so. Consequently,  $\varphi_M$  is bijective if  $M$  is quasi-isomorphic to a bounded complex of finite-dimensional projective  $\mathcal{E}$ -modules. The second map is handled similarly.  $\square$

In general,  $F_\lambda$  does not commute with the Auslander–Reiten translations. However, we have the following.

**Lemma 2.4.** Let  $X \in \text{ind } \mathcal{E}$  be such that  $F_\lambda X \in \text{ind } \mathcal{B}$  and  $\text{pd } X < \infty$ . Then  $\dim \tau_{\mathcal{E}} X = \dim \tau_{\mathcal{B}} F_\lambda X$ .

**Proof.** Let  $X \in \text{mod } \mathcal{E}$  be any module. Let  $P_1 \rightarrow P_0 \rightarrow X \rightarrow 0$  be a minimal projective presentation in  $\text{mod } \mathcal{E}$ . By [14, 3.2], we deduce that  $F_\lambda P_1 \rightarrow F_\lambda P_0 \rightarrow F_\lambda X \rightarrow 0$  is a minimal projective presentation in  $\text{mod } \mathcal{B}$ . So we have exact sequences in  $\text{mod } \mathcal{E}^{op}$  and  $\text{mod } \mathcal{B}^{op}$ , respectively:

$$0 \rightarrow \text{Hom}_{\mathcal{E}}(X, \mathcal{E}) \rightarrow \text{Hom}_{\mathcal{E}}(P_0, \mathcal{E}) \rightarrow \text{Hom}_{\mathcal{E}}(P_1, \mathcal{E}) \rightarrow \text{Tr}_{\mathcal{E}} X \rightarrow 0,$$

and

$$0 \rightarrow \text{Hom}_{\mathcal{B}}(F_\lambda X, \mathcal{B}) \rightarrow \text{Hom}_{\mathcal{B}}(F_\lambda P_0, \mathcal{B}) \rightarrow \text{Hom}_{\mathcal{B}}(F_\lambda P_1, \mathcal{B}) \rightarrow \text{Tr}_{\mathcal{B}} F_\lambda X \rightarrow 0.$$

Let  $X \in \text{mod } \mathcal{E}$  be of finite projective dimension, thus quasi-isomorphic to a bounded complex of finite-dimensional projective  $\mathcal{E}$ -modules. The bijections of 2.3 imply that  $\dim \text{Hom}_{\mathcal{E}}(X, \mathcal{E}) = \sum_{x \in \mathcal{E}_0} \dim \text{Hom}_{\mathcal{E}}(X, \mathcal{E}(-, x)) = \sum_{x \in \mathcal{B}_0} \dim \text{Hom}_{\mathcal{B}}(F_\lambda X, \mathcal{B}(-, x)) = \dim \text{Hom}_{\mathcal{B}}(F_\lambda X, \mathcal{B})$ . Using the above exact sequences, we deduce that  $\dim \text{Tr}_{\mathcal{E}} X = \dim \text{Tr}_{\mathcal{B}} F_\lambda X$ . Thus  $\dim \tau_{\mathcal{E}} X = \dim \tau_{\mathcal{B}} F_\lambda X$  if both  $X$  and  $F_\lambda X$  are indecomposable.  $\square$

### 3. Standard laura algebras

We now derive sufficient conditions for a laura algebra to be standard. Weakly shod algebras are particular cases of laura algebras. It is proved in [17, §4] that if  $A$  is weakly shod and not quasi-tilted, then  $A$  can be written as a one-point extension  $A = B[M]$  such that the connecting component of  $A$  can be recovered from  $M$  and from the connecting components of  $B$ . This motivates the following definition.

**Definition 3.1.** Let  $A$  be a laura algebra with connecting components. An indecomposable projective  $A$ -module  $P$  lying in a connecting component  $\Gamma$  is a *maximal projective* if it has an injective predecessor and no proper projective successor in  $\text{ind } A$ . Furthermore,  $A$  is a *maximal extension* of  $B$  if there exists a maximal projective  $P = eA$  such that  $B = (1 - e)A(1 - e)$  and  $A = B[M]$ , where  $M = \text{rad } P$ .

By definition, a maximal projective belongs to  $\mathcal{R}_A$ . In particular, by [7, 2.2], it is directed. The notions of *minimal injective* or *maximal coextension* are dual. If  $A$  is a tilted algebra which is the endomorphism algebra of a regular tilting module, then it has neither maximal projective, nor minimal injective (see [39]).

**Proposition 3.2.** Let  $A = B[M]$  be a maximal extension. Then  $B$  is a product of laura algebras with connecting components. Moreover, if every connected component of  $B$  is standard, then so is  $A$ .

**Proof.** By [4], every connected component of  $B$  is a laura algebra. Let  $P_m \in \text{ind } A$  be the maximal projective such that  $\text{rad } P_m = M$  and denote by  $\Gamma$  the component of  $\Gamma(\text{mod } A)$  in which  $P_m$  lies. So  $P_m \in \mathcal{R}_A \cap \Gamma$ . In particular,  $P_m$  is directed. Note that every proper predecessor of  $P_m$  is an indecomposable  $B$ -module.

Let us prove the first assertion. If it is false, then a connected component  $B'$  of  $B$  is quasi-tilted and not tilted (and, therefore, quasi-tilted of canonical type). Since  $A$  is connected, at least one indecomposable summand  $M'$  of  $M$  lies in  $\text{ind } B'$ . Assume first that  $M'$  is not directed. In particular,  $M' \in \Gamma$  implies that  $M' \notin \mathcal{L}_A \cup \mathcal{R}_A$ . Therefore there is a non-sectional path  $M' \rightsquigarrow P$  in  $\text{ind } A$  with  $P$  projective. If  $P = P_m$ , then there exists a non-sectional path  $M' \rightsquigarrow M''$  with  $M''$  an indecomposable summand of  $M = \text{rad } P_m$ . This is impossible because  $P_m$  is directed (see [25, Thm. 1 of §2]). So  $P \neq P_m$ . By maximality of  $P_m$ , the path  $M' \rightsquigarrow P$  is a non-sectional path in  $\text{ind } B'$  ending at a projective. So  $M' \notin \mathcal{R}_{B'}$ . On the other hand,  $M' \notin \mathcal{L}_A$  means that there exists a non-sectional path  $I \rightsquigarrow M'$  in  $\text{ind } A$ , where  $I$  is injective. By maximality of  $P_m$ , this is a non-sectional path in  $\text{ind } B'$ . For the same reason, we have  $\text{Hom}_A(P_m, I) = 0$ , so that  $I$  is injective as a  $B'$ -module. So  $M' \notin \mathcal{L}_{B'} \cup \mathcal{R}_{B'}$ . This is impossible because  $B'$  is quasi-tilted. Therefore  $M'$  is directed. Since  $B'$  is quasi-tilted of canonical type, the component  $\Gamma'$  of  $\Gamma(\text{mod } B')$  containing  $M'$  is either the unique postprojective or the unique preinjective component (see [32, Prop. 4.3]). Assume that  $\Gamma'$  is the unique postprojective component of  $\Gamma(\text{mod } B')$ . Then  $\Gamma' \subseteq \mathcal{L}_{B'} \setminus \mathcal{R}_{B'}$  (see [18, 5.2]). In particular, there exists a non-sectional path  $M' \rightsquigarrow P$  in  $\text{ind } B'$  with  $P$  projective. Since  $P_m$  is maximal, this is also a non-sectional path in  $\text{ind } A$ . Since  $P$  is projective and since  $M' \in \Gamma$ , we deduce that  $P \in \Gamma$  and that the path is refinable to a non-sectional path in  $\Gamma(\text{mod } A)$  and therefore in  $\Gamma(\text{mod } B')$  because  $P_m$  is maximal. Consequently,  $M'$  lies in the postprojective component  $\Gamma'$  of  $\Gamma(\text{mod } B')$  and is the starting point of a non-sectional path in  $\Gamma(\text{mod } B')$  ending at a projective. This is absurd. If  $\Gamma'$  is the unique preinjective component of  $\Gamma(\text{mod } B')$ , then, using dual arguments, we also get a contradiction. Thus,  $B'$  is either tilted or not quasi-tilted.

Now, we assume that every connected component of  $B$  is standard, and prove that  $A$  is standard. Later, in 5.7, we shall see that, if  $A$  is tilted, then its connecting components are standard. So assume that  $A$  is not tilted. Let  $\Gamma$  be the connecting component of  $\Gamma(\text{mod } A)$  and  $\Gamma'$  be the disjoint union of the connecting components of the Auslander-Reiten quivers of the connected components of  $B$ . We compare  $\Gamma$  and  $\Gamma'$ . More precisely, let  $\mathcal{X}$  be the full subquiver of  $\Gamma$  with vertices those modules which are not successors of  $P_m$ . So  $\mathcal{X}$  is a full subquiver of  $\Gamma(\text{mod } B)$  stable under predecessors in  $\Gamma(\text{mod } B)$ , and it contains  $\Gamma \setminus \mathcal{R}_A$ . We claim that  $\mathcal{X}$  is contained in  $\Gamma'$ . We prove a series of assertions.

**(a) The left supports of  $A$  and  $B$  coincide.** Indeed, we have  $\mathcal{L}_A \cap \text{ind} B \subseteq \mathcal{L}_B$  (see [4, 2.1]). On the other hand, if  $P \in \text{ind} B$  is a projective not lying in  $\mathcal{L}_A$ , then there is a non-sectional path  $I \rightsquigarrow P$  in  $\text{ind} A$  with  $I$  injective. Since  $P_m$  is maximal, this is a non-sectional path in  $\text{ind} B$ . For the same reason,  $\text{Hom}_A(P_m, I) = 0$ , so that  $I$  is injective as a  $B$ -module. So  $P \notin \mathcal{L}_B$ . Thus  $A$  and  $B$  have the same left support.

**(b) Let  $P \neq P_m$  be a projective lying in  $\Gamma$ . Then  $P \in \Gamma'$ .** Indeed, if there exists a path  $I \rightsquigarrow P$  in  $\Gamma$  with  $I$  injective, then the maximality of  $P_m$  implies that this path lies entirely in  $\text{ind} B$  and starts in an injective  $B$ -module. So  $P \in \Gamma'$ . If there is no such path, then  $P \in \mathcal{L}_A \cap \Gamma$ . So  $P$  lies in a connecting component of one of the components of the left support of  $A$ , which is also the left support of  $B$ . From [3, 5.4], we deduce that  $P$  lies in  $\Gamma'$ .

**(c) Let  $X \in \mathcal{X}$ . There exists  $m \geq 0$  such that  $\tau_A^m X \in \Gamma'$ .** By assumption on  $X$ , we have  $\tau_B X = \tau_A X$ . Assume first that  $\tau_A^m X = P$  for some  $m \geq 0$  and some projective  $P$ . So  $P \neq P_m$ . From (b), we get that  $P \in \Gamma'$ . Now assume that  $X$  is left stable and non-periodic. If  $X \in \mathcal{R}_A$ , there exists  $l \geq 0$  such that  $\tau_A^l X$  is Ext-projective in  $\mathcal{R}_A$ . Since  $X$  is left stable, we deduce that  $\tau_A^{l+1} X \in \Gamma \setminus \mathcal{R}_A$ . So assume that  $X \in \Gamma \setminus \mathcal{R}_A$ . Since  $A$  is laura, there exists  $m$  such that  $\tau_A^m X \in \Gamma \cap \mathcal{L}_A$ . So  $\tau_A^m X$  lies in one of the connecting components of the left support of  $A$ . So  $\tau_A^m X \in \Gamma'$  because the left supports of  $A$  and  $B$  are equal. Finally, assume that  $X$  is periodic. Then there exists a projective module  $P \in \Gamma$ , a periodic direct summand  $Y$  of  $\text{rad} P$ , and a path  $Y \rightsquigarrow X$  in  $\Gamma \setminus \mathcal{R}_A$ , and therefore in  $\Gamma(\text{mod} B)$ . Since  $Y$  is periodic, then  $P \neq P_m$  (otherwise  $P_m$  would be a proper successor of itself). Since  $P \in \Gamma'$ , we have  $Y \in \Gamma'$  and therefore  $X \in \Gamma'$ .

**(d)  $\mathcal{X}$  is contained in  $\Gamma'$ .** Indeed, we already know that  $\mathcal{X}$  is a full subquiver of  $\Gamma(\text{mod} B)$ . Also, we proved that for every  $X \in \mathcal{X}$ , there exists  $m \geq 0$  such that  $\tau_A^m X = \tau_B^m X \in \Gamma'$ . So  $\mathcal{X}$  is contained in  $\Gamma'$ .

We now show that  $\Gamma$  is standard. By hypothesis, there exists a well-behaved functor  $\varphi : k(\Gamma') \rightarrow \text{ind} \Gamma'$ . Since  $\mathcal{X}$  is a full subquiver of  $\Gamma'$  stable under predecessors in  $\Gamma(\text{mod} B)$ , there exists a well-behaved functor  $\psi : k(\mathcal{Y}) \rightarrow \text{ind} \Gamma$  where  $\mathcal{Y}$  is a full subquiver of  $\Gamma$  such that:

1.  $\mathcal{Y}$  contains  $\mathcal{X}$ .
2.  $\mathcal{Y}$  is stable under predecessors in  $\Gamma(\text{mod} A)$ .
3.  $\psi$  and  $\varphi$  coincide on  $\mathcal{X}$ .
4.  $\mathcal{Y}$  is maximal for these properties.

We show that  $\mathcal{Y} = \Gamma$ . Assume that  $\mathcal{Y} \neq \Gamma$ . Since  $\mathcal{Y}$  contains  $\mathcal{X}$ , it contains  $\Gamma \setminus \mathcal{R}_A$ , so there exists a source  $X$  in  $\Gamma \setminus \mathcal{Y}$ . If  $X$  is projective, then  $X = P_m$ . So  $\psi$  is defined on every indecomposable summand  $Y$  of  $\text{rad} P_m$ . Set  $\psi(X) = P_m$ . Let  $\alpha_1 : X_1 \rightarrow P_m, \dots, \alpha_t : X_t \rightarrow P_m$  be the arrows ending at  $X$ . Then  $X_1 \oplus \dots \oplus X_t = \text{rad} P_m$ , and let  $\psi(\alpha_i)$  be the inclusion  $X_i \hookrightarrow P_m$ . If  $X$  is not projective, then the mesh ending at  $X$  has the following shape:

$$\begin{array}{ccccc}
 & & & X_1 & & \\
 & & & \nearrow & & \\
 & & & u_1 & & \\
 & & & \rightarrow & & \\
 \tau_A X & & & & & X_1 & \xrightarrow{v_1} & X \\
 & & & & & \vdots & & \\
 & & & & & X_n & \xrightarrow{v_n} & X \\
 & & & & & \nwarrow & & \\
 & & & u_n & & & & \\
 & & & \rightarrow & & & & 
 \end{array} \tag{*}$$

Since  $X$  is a source of  $\Gamma \setminus \mathcal{Y}$ , then  $\psi$  is already defined on the full subquiver of the mesh consisting of all vertices except  $X$ . In particular, the following map is right minimal almost split:

$$[\psi(u_1) \quad \dots \quad \psi(u_n)]^t : \tau_A X \rightarrow X_1 \oplus \dots \oplus X_n.$$

Let  $\psi(X) = X$ , and  $[\psi(v_1) \quad \dots \quad \psi(v_n)] : X_1 \oplus \dots \oplus X_n \rightarrow X$  be the cokernel of the above map, following [14, 3.1, Ex. b]. Clearly, this construction contradicts the maximality of  $\mathcal{Y}$ . So  $\mathcal{Y} = \Gamma$  and

there exists a well-behaved functor  $\psi : k(\Gamma) \rightarrow \text{ind } \Gamma$  which is the identity on objects. The arguments in the proof of [14, 5.1] show that this is an isomorphism. So  $\Gamma$  is standard.  $\square$

Since weakly shod algebras are lura, it makes sense to speak of weakly shod algebras with connecting components. We have the following corollary.

**Corollary 3.3.** *Let  $A$  be a (connected) weakly shod algebra with connecting components, then  $A$  is standard.*

**Proof.** By [7, 3.3], there exists a sequence of full convex subcategories

$$C = A_0 \subsetneq A_1 \subsetneq \cdots \subsetneq A_m = A$$

with  $C$  tilted and, for each  $i \geq 0$ , the algebra  $A_{i+1}$  is a maximal extension of  $A_i$ . The result follows from 3.2 and induction because  $C$  is standard (see 5.7 below).  $\square$

The preceding result motivates the following definition, inspired from [7, 2.3].

**Definition 3.4.** Let  $A$  be a lura algebra. We say that  $A$  admits a *maximal filtration* if there exists a sequence

$$C = A_0 \subsetneq A_1 \subsetneq \cdots \subsetneq A_m = A \tag{f}$$

of full convex subcategories with  $C$  a product of representation-finite algebras and, for each  $i \geq 0$ , the algebra  $A_{i+1}$  is a maximal extension, or a maximal coextension, of  $A_i$ .

**Corollary 3.5.** *Let  $A$  be a lura algebra admitting a maximal filtration (f):*

- (a) *If  $C$  is a product of standard representation-finite algebras, then  $A$  is standard.*
- (b) *If the Auslander–Reiten quiver of every connected component of  $C$  is simply connected, then  $A$  is standard.*
- (c) *If  $\text{HH}^1(A) = 0$ , then  $A$  is standard.*

**Proof.** Statement (a) follows directly from 3.2.

(b) This follows from 3.2 and the fact that if a representation-finite connected algebra  $C$  has  $\text{HH}^1(C) = 0$ , or equivalently, if its Auslander–Reiten quiver is simply connected, then  $C$  is standard [15, 4.2].


(c) We use induction on the length  $m$  of a maximal filtration. If  $m = 0$ , then  $A$  is representation-finite and the result follows from [15, 4.2]. Assume that  $m \geq 1$  and that the statement holds for algebras admitting maximal filtrations of length less than  $m$ . Without loss of generality, we may assume that  $A = A_{m-1}[M]$  is a maximal extension. We claim that  $\text{Ext}_{A_{m-1}}^1(M, M) = 0$ . Indeed, if this is not the case, then there exists an indecomposable summand  $N$  of  $M$  such that  $\text{Ext}_{A_{m-1}}^1(M, N) \neq 0$ . Write  $M \simeq N \oplus N'$  and let  $P$  be the indecomposable projective such that  $M = \text{rad } P$ . Then  $N'$  is a submodule of  $P$  and  $L = P/N'$  is indecomposable. By [24, III.2.2(a)] we have  $\text{id } L \geq 2$ . But this contradicts the fact that  $L \in \mathcal{R}_A$  because it is a successor of the maximal projective  $P$ . So  $\text{Ext}_{A_{m-1}}^1(M, M) = 0$ . Applying [23, 5.3], the exact sequence

$$\text{HH}^1(A) \rightarrow \text{HH}^1(A_{m-1}) \rightarrow \text{Ext}_{A_{m-1}}^1(M, M)$$

yields  $\text{HH}^1(A_{m-1}) = 0$ . By the induction hypothesis,  $A_{m-1}$  is standard. By 3.2, so is  $A$ .  $\square$

**Examples 3.6.**


(a) Let  $A$  be the radical-square zero algebra given by the quiver



This is a lura algebra (see [3, 2.3]). Here and in the sequel, we denote by  $P_x, I_x$  and  $S_x$  the indecomposable projective, the indecomposable injective, and the simple module corresponding to the vertex  $x$ , respectively. Clearly  $P_1$  is maximal projective and  $I_5$  is minimal injective. Letting  $C$  be the full convex subcategory with objects  $\{2, 3, 4\}$  we see that

$$C \subseteq [S_4 \oplus S_4]C \subseteq A$$

is a maximal filtration. Since  $C$  is standard, so is  $A$ . Its connecting component is drawn below:



where the two copies of  $S_3$  are identified.

(b) Let  $B, C$  be products of standard lura algebras, and  $A$  an articulation of  $B, C$  (in the sense of [20]). Then  $A$  is lura with connecting components (see [20]). Using [20, 3.9] it is easy to check that  $A$  is standard.

The section motivates the following questions.

**Problem 1.** Which lura algebras admit maximal filtrations?

**Problem 2.** Assume that  $A$  is a lura algebra which does not admit a maximal filtration. If  $\text{HH}^1(A) = 0$ , do we have that  $A$  is standard?

**4. Tilting modules of the first kind with respect to covering functors**

For tilting theory, we refer to [11]. Let  $B$  be a product of tilted algebras and  $n$  be the rank of its Grothendieck group. In [31, Cor. 4.5], it is proved that tilting modules are of the first kind with respect to any Galois covering of  $B$ . More precisely, let  $F : \tilde{B} \rightarrow B$  be a Galois covering with group  $G$ , where  $\tilde{B}$  is locally bounded. Denote by  $\mathcal{T}$  the class of complexes  $T \in \mathcal{D}^b(\text{mod } B)$  such that:

1.  $T$  is multiplicity-free and has  $n$  indecomposable summands.
2.  $\mathcal{D}^b(\text{mod } B)(T, T[i]) = 0$  for every  $i \geq 1$  (so  $T$  is a *tilting complex* in the sense of [26]).
3.  $T$  generates the triangulated category  $\mathcal{D}^b(\text{mod } B)$ .

Any multiplicity-free tilting module lies in  $\mathcal{T}$ . It was proved in [31, §4] that for any  $T \in \mathcal{T}$  and for any indecomposable summand  $X$  of  $T$ , there exists  $\tilde{X} \in \mathcal{D}^b(\text{mod } \tilde{B})$  such that:

1.  $F_\lambda \tilde{X} \simeq X$ .
2.  ${}^g \tilde{X} \not\simeq {}^h \tilde{X}$  for  $g \neq h$ .
3. If  $Y \in \mathcal{D}^b(\text{mod } \tilde{B})$  is such that  $F_\lambda Y \simeq X$ , then  $Y \simeq {}^g \tilde{X}$  for some  $g \in G$ .

Given  $T \in \mathcal{T}$  and an indecomposable summand  $X$  of  $T$ , we fix  $\tilde{X} \in \mathcal{D}^b(\text{mod } \tilde{B})$  arbitrarily such that  $F_\lambda \tilde{X} \simeq X$ .

For later reference, we recall some facts. The following result was proved in [31, Cor. 4.5, Prop. 4.6, Lem. 4.8].

**Lemma 4.1.** *Let  $F : \tilde{B} \rightarrow B$  be a Galois covering with group  $G$ . Let  $T \in \text{mod } B$  be a multiplicity-free tilting module. Let  $T = T_1 \oplus \dots \oplus T_n$  be such that  $T_1, \dots, T_n$  are indecomposable. For every  $i$ , there exists  $\tilde{T}_i \in \text{ind } \tilde{B}$  such that  $F_\lambda \tilde{T}_i = T_i$ . Moreover:*

- (a)  ${}^g \tilde{T}_i \not\simeq {}^h \tilde{T}_j$  for  $(g, i) \neq (h, j)$ .
- (b)  $\text{pd } \tilde{T}_i \leq 1$  for every  $i$ .
- (c)  $\text{Ext}_B^1({}^g \tilde{T}_i, {}^h \tilde{T}_j) = 0$  for every  $g, h \in G, i, j \in \{1, \dots, n\}$ .
- (d) For every indecomposable projective  $\tilde{B}$ -module  $P$ , there exists an exact sequence  $0 \rightarrow P \rightarrow T^{(1)} \rightarrow T^{(2)} \rightarrow 0$  with  $T^{(1)}, T^{(2)}$  in  $\text{add}\{{}^g \tilde{T}_i \mid g \in G, i \in \{1, \dots, n\}\}$ .

We need similar facts about covering functors which need not be Galois. Thus we prove the following result.

**Proposition 4.2.** *Let  $F : \tilde{B} \rightarrow B$  be a Galois covering with group  $G$ , where  $\tilde{B}$  is locally bounded. With the above setting, let  $p : \tilde{B} \rightarrow B$  be a covering functor such that  $F(x) = p(x)$  for every  $x \in \tilde{B}_0$ . Let  $T \in \mathcal{T}$  and  $X$  be an indecomposable summand of  $T$ . Then:*

- (a) There exists an isomorphism  $p_\lambda({}^g \tilde{X}) \xrightarrow{\sim} X$ , for every  $g \in G$ .
- (b) If  $L \in \mathcal{D}^b(\text{mod } \tilde{B})$  is such that  $p_\lambda L \simeq X$ , then  $L \simeq {}^g \tilde{X}$  for some  $g \in G$ .
- (c) For every  $L \in \mathcal{D}^b(\text{mod } \tilde{B})$ , the following maps induced by  $p_\lambda$  and by the isomorphisms of (a) are linear bijections:

$$\varphi_{X,L} : \bigoplus_{g \in G} \mathcal{D}^b(\text{mod } \tilde{B})({}^g \tilde{X}, L) \xrightarrow{\sim} \mathcal{D}^b(\text{mod } B)(X, p_\lambda L),$$

and

$$\psi_{X,L} : \bigoplus_{g \in G} \mathcal{D}^b(\text{mod } \tilde{B})(L, {}^g \tilde{X}) \xrightarrow{\sim} \mathcal{D}^b(\text{mod } B)(p_\lambda L, X).$$

In order to prove the proposition, we need the following lemma. In case  $p$  is a Galois covering, the lemma was proved in [31, Lems. 4.2, 4.3] (see also [29, Lems. 3.2, 3.3]). For simplicity, we write  $\text{Hom}(X, Y)$  for the space of morphisms in the derived category.

**Lemma 4.3.** *Let  $T, T' \in \mathcal{T}$  be such that 4.2 holds true for  $T$  and for  $T'$ . Consider a triangle in  $\mathcal{D}^b(\text{mod } B)$ :*

$$X \rightarrow \bigoplus_{i=1}^t X'_i \rightarrow Y \rightarrow X[1], \tag{\Delta}$$

where  $X \in \text{add } T$  and  $X'_1, \dots, X'_t$  are indecomposable summands of  $T'$ . Assume that  $\text{Hom}(Y, X'_i[1]) = 0$  for all  $i$  (we do not assume that  $Y \in \text{add } T$  or  $Y \in \text{add } T'$ ). Then for every  $g \in G$ , there exist  $\tilde{Y} \in \mathcal{D}^b(\text{mod } \tilde{B})$  and  $g_1, \dots, g_t \in G$  such that the triangle  $\Delta$  is isomorphic to the image under  $p_\lambda$  of a triangle in  $\mathcal{D}^b(\text{mod } \tilde{B})$  as follows:

$${}^g\tilde{X} \rightarrow \bigoplus_{i=1}^t {}^{g_i}\tilde{X}'_i \rightarrow \tilde{Y} \rightarrow {}^g\tilde{X}[1].$$

Dually, consider a triangle in  $\mathcal{D}^b(\text{mod } B)$ :

$$Y \rightarrow \bigoplus_{i=1}^t X'_i \rightarrow X \rightarrow Y[1], \tag{\Delta'}$$

where  $X \in \text{add } T$  and  $X'_1, \dots, X'_t$  are indecomposable summands of  $T'$ . Assume that  $\text{Hom}(X'_i, Y[1]) = 0$  for all  $i$ . Then for every  $g \in G$ , there exist  $\tilde{Y} \in \mathcal{D}^b(\text{mod } \tilde{B})$  and  $g_1, \dots, g_t \in G$  such that the triangle  $\Delta'$  is isomorphic to the image under  $p_\lambda$  of a triangle in  $\mathcal{D}^b(\text{mod } \tilde{B})$  as follows:

$$\tilde{Y} \rightarrow \bigoplus_{i=1}^t {}^{g_i}\tilde{X}'_i \rightarrow {}^g\tilde{X} \rightarrow \tilde{Y}[1].$$

**Proof.** The proofs of [31, Lems. 4.2, 4.3] use the following key property of a Galois covering  $F : \tilde{B} \rightarrow B$  with group  $G$ . Given  $L, M \in \mathcal{D}^b(\text{mod } \tilde{B})$ , we have linear bijections induced by  $F_\lambda$ :

$$\bigoplus_{g \in G} \text{Hom}({}^g L, M) \xrightarrow{\sim} \text{Hom}(F_\lambda L, F_\lambda M) \quad \text{and} \quad \bigoplus_{g \in G} \text{Hom}(L, {}^g M) \xrightarrow{\sim} \text{Hom}(F_\lambda L, F_\lambda M).$$

Of course, these bijections no longer exist for a covering functor which is not Galois. However, using our hypothesis that 4.2 holds true for  $T$  and for  $T'$ , it is easy to check that the proofs of [31, Lems. 4.2, 4.3] still work in the present case. Whence the lemma.  $\square$

**Proof of 4.2.** We proceed in several steps.

**Step 1: If  $T = B$ , then 4.2 holds true.** The following facts follow from the definition of covering functors (see also [14, 3.2]):

1.  $Y \in \mathcal{D}^b(\text{mod } \tilde{B})$  is a projective module if and only if  $p_\lambda Y$  is a projective module.
2.  $p_\lambda(\tilde{B}(-, x)) \simeq F_\lambda(\tilde{B}(-, x)) \simeq B(-, F(x)) = B(-, p(x))$  for every  $x \in \tilde{B}_0$ .
3.  ${}^g\tilde{B}(-, x) = \tilde{B}(-, gx)$  for every  $x \in \tilde{B}_0$  and every  $g \in G$ .

Therefore 4.2 holds true for  $T = B$ .

Given an object  $X$  in a triangulated category, we write  $\langle X \rangle$  for the smallest additive full subcategory containing  $X$  which is stable under direct summands and shifts (in both directions).

**Step 2: If  $T, T' \in \mathcal{T}$  are such that  $T' \in \langle T \rangle$ , then 4.2 holds true for  $T$  if and only if it does for  $T'$ .** This follows directly from the compatibility of  $p_\lambda$  with the shift.

For the next step, consider the following situation. Assume that  $T, T' \in \mathcal{T}$  are such that:

1.  $T = M \oplus \bar{T}$ , where  $M$  is indecomposable.
2.  $T' = M' \oplus \bar{T}$ , where  $M'$  is indecomposable.
3. There exists a non-split triangle  $\Delta : M \xrightarrow{u} E \xrightarrow{v} M' \rightarrow M[1]$  where  $u$  is a left minimal  $\text{add } \bar{T}$ -approximation and  $v$  is a right minimal  $\text{add } \bar{T}$ -approximation.

**Step 3: If  $T, T' \in \mathcal{T}$  are as above, then 4.2 holds true for  $T$  if and only if it does for  $T'$ .** We prove that the condition is necessary. Clearly, it suffices to prove that the assertions (a), (b), and (c) of 4.2 are true for  $M'$ . For simplicity, we identify  $p_\lambda({}^g\tilde{X})$  and  $X$  via the isomorphism used to define  $\varphi_{X,-}$  and  $\psi_{X,-}$  for every indecomposable summand  $X$  of  $T$  and  $g \in G$ .

Let  $E = \bigoplus_{i=1}^t E_i$  with the  $E_i$  indecomposable. Recall from [31, Lem. 4.4] that  $\Delta$  is isomorphic to the image under  $F_\lambda$  of a triangle  $\tilde{\Delta}$  in  $\mathcal{D}^b(\text{mod } \tilde{B})$ :

$$\tilde{M} \xrightarrow{\tilde{u}} \bigoplus_{i=1}^t {}^g E_i \xrightarrow{\tilde{v}} {}^{g_0} \tilde{M}' \rightarrow \tilde{M}[1], \tag{\tilde{\Delta}}$$

for some  $g_0, g_1, \dots, g_t \in G$ . Moreover,  $\tilde{u}$  is a left minimal add  $\mathcal{X}$ -approximation and  $\tilde{v}$  is a right minimal add  $\mathcal{X}'$ -approximation, where  $\mathcal{X}$  and  $\mathcal{X}'$  are the following full subcategories of  $\mathcal{D}^b(\text{mod } \tilde{B})$ :

- $\mathcal{X} = \{{}^g \tilde{X} \mid g \in G, X \text{ an indecomposable summand of } T \text{ and } {}^g \tilde{X} \not\cong \tilde{M}\}$ .
- $\mathcal{X}' = \{{}^g \tilde{X}' \mid g \in G, X \text{ an indecomposable summand of } T' \text{ and } {}^g \tilde{X}' \not\cong \tilde{M}'\}$ .

Fix  $g \in G$ . Since 4.2 holds true for  $T$ , we apply 4.3 to construct a triangle  $\tilde{\Delta}' : {}^g \tilde{M} \xrightarrow{\tilde{u}'} \bigoplus_{i=1}^t {}^{g_i'} \tilde{E}_i \xrightarrow{\tilde{v}'}$   $Z_g \rightarrow {}^g \tilde{M}'[1]$  whose image under  $p_\lambda$  is isomorphic to  $\Delta$ . In particular,  $p_\lambda(Z_g) \simeq M'$ . For simplicity, assume that  $\Delta$  is equal to the image of  $\tilde{\Delta}$  under  $p_\lambda$ , and set  $\tilde{E}' = \bigoplus_{i=1}^t {}^{g_i'} \tilde{E}_i$ . Let us prove that  $Z_g \simeq {}^{g_0} \tilde{M}'$ . It suffices to prove that  $\tilde{\Delta}'$  and  ${}^g \tilde{\Delta}$  are isomorphic. For this purpose, we only need to prove that  $\tilde{u}'$  is a left minimal add  ${}^g \mathcal{X}$ -approximation. Let  $f : {}^g \tilde{M} \rightarrow {}^{g'} \tilde{Y}$  be non-zero, where  $Y$  is an indecomposable summand of  $T$  such that  ${}^{g'} \tilde{Y} \in {}^g \mathcal{X}$ . Since  $\varphi_{M, \tilde{M}}$  is bijective and since  $\text{End}(M) = k$ , we have  $Y \in \text{add } \tilde{T}$ . So we have a factorisation of  $p_\lambda(f)$  by  $u = p_\lambda(\tilde{u}')$ :

$$\begin{array}{ccc} M & \xrightarrow{u} & E \\ & \searrow p_\lambda(f) & \downarrow f' \\ & & Y. \end{array}$$


Since  $\psi_{Y, \tilde{E}_i}$  is bijective for every  $i$ , we have  $f' = \sum_{h \in G} p_\lambda(f'_h)$ , where  $(f'_h)_h \in \bigoplus_{h \in G} \text{Hom}(\tilde{E}, {}^h \tilde{Y})$ . So  $p_\lambda(f - f'_{g'} \tilde{u}') - \sum_{h \neq g'} p_\lambda(f'_h \tilde{u}') = 0$ . Using 4.2, we get  $f = f'_{g'} \tilde{u}'$ . Hence  $\tilde{u}'$  is a left add  ${}^g \mathcal{X}$ -approximation. On the other hand,  $\tilde{u}'$  is left minimal because  $u = p_\lambda(\tilde{u}')$  is left minimal and  $p_\lambda$  is exact. As explained above, these facts imply that  $Z_g \simeq {}^{g_0} \tilde{M}'$ . So  $p_\lambda({}^g \tilde{M}') \simeq M'$ , for every  $g \in G$ .

Let  $Y \in \mathcal{D}^b(\text{mod } \tilde{B})$ . Using the triangles  ${}^g \tilde{\Delta}$  ( $g \in G$ ) and using that 4.2 holds true for  $T$ , the maps  $\varphi_{M', Y}$  and  $\psi_{M', Y}$  are bijective (recall that Hom-functors are cohomological).

Finally, if  $Y \in \mathcal{D}^b(\text{mod } \tilde{B})$ , and if  $f : p_\lambda Y \rightarrow M'$  is an isomorphism, then  $f = \sum_{g \in G} p_\lambda(f_g)$  with  $(f_g)_g \in \bigoplus_{g \in G} \text{Hom}(Y, {}^g \tilde{M}')$ . Since  $p_\lambda Y$  and  $M'$  are indecomposable, there exists  $g_1 \in G$  such that  $p_\lambda(f_{g_1})$  is an isomorphism. Since  $p_\lambda$  is exact, we deduce that  $f_{g_1} : Y \rightarrow {}^{g_1} \tilde{M}'$  is an isomorphism. This finishes the proof of the assertion: 4.2 holds true for  $T'$  if it holds true for  $T$ . The converse implication is proved using similar arguments.

**Step 4: If  $T \in \mathcal{T}$ , then 4.2 holds true.** This follows directly from the three preceding steps, and from [31, Prop. 3.7].  $\square$

**Example 4.4.** Let  $B = kQ$  be the path algebra of the Kronecker quiver  $1 \begin{matrix} \xrightarrow{a} \\ \xrightarrow{b} \end{matrix} 2$ . There is a Galois covering  $F : \tilde{B} \rightarrow B$  with group  $\mathbb{Z}/2\mathbb{Z} = \{1, \sigma\}$ , where  $\tilde{B} = k\tilde{Q}$  is the path algebra of the following quiver:



and where  $F$  is the functor such that  $F(\sigma^i \alpha) = \alpha$  for every arrow  $\alpha$  and every  $i \in \{0, 1\}$ . On the other hand, there is a covering functor  $p : \tilde{B} \rightarrow B$  such that  $p(b) = p(\sigma b) = b$ ,  $p(a) = a$  and  $p(\sigma a) = a + b$ . The  $B$ -module  $T = e_2 B \oplus \tau_B^{-1}(e_1 B)$  is tilting. One checks easily that  $F_\lambda(e_2 \tilde{B}) = e_2 B$ ,  $F_\lambda(\tau_B^{-1}(e_1 \tilde{B})) = \tau_B^{-1}(e_1 B)$  and that  $p_\lambda(e_2 \tilde{B}) \simeq e_2 B$ ,  $p_\lambda(\tau_B^{-1}(e_1 \tilde{B})) \simeq \tau_B^{-1}(e_1 B)$ .

**5. Coverings of left sections**

Let  $A$  be a basic finite-dimensional  $k$ -algebra,  $\Gamma$  a component of  $\Gamma(\text{mod } A)$ ,  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  a Galois covering of translation quivers with group  $G$  such that there exists a well-behaved functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . A *left section* (see [1, 2.1]) in  $\Gamma$  is a full subquiver  $\Sigma$  such that:  $\Sigma$  is acyclic; it is convex in  $\Gamma$ ; and, for any  $x \in \Gamma$ , predecessor in  $\Gamma$  of some  $y \in \Sigma$ , there exists a unique  $n \geq 0$  such that  $\tau^{-n} x \in \Sigma$ . Assume that  $\Sigma$  is a left section in  $\Gamma$  and let  $B = A/\text{Ann } \Sigma$ . In this section, we construct a covering functor  $F : \tilde{B} \rightarrow B$  associated to  $p$  and a functor  $\varphi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{B}$ . Both  $F$  and  $\varphi$  are essential in the proofs of Theorems A and B.

By [1, Thm. A], the algebra  $B$  is a full convex subcategory of  $A$  and a product of tilted algebras and the components of  $\Sigma$  form complete slices in the connecting components of the connected components of  $B$ . Recall from [39] that a connected algebra  $B'$  is tilted if and only if its Auslander–Reiten quiver contains a so-called complete slice  $\Sigma'$ , that is, a class of indecomposable  $B'$ -modules such that: (1)  $U = \bigoplus_{X \in \Sigma'} X$  is sincere (that is,  $\text{Hom}_{B'}(P, U) \neq 0$  for any projective  $B'$ -module  $P$ ); (2)  $\Sigma'$  is convex in  $\text{ind } B'$ ; (3) If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is an almost split sequence, then at most one of  $L$  and  $N$  lies in  $\Sigma'$ . Moreover, if an indecomposable summand of  $M$  lies in  $\Sigma'$ , then either  $L$  or  $N$  lies in  $\Sigma'$ . Here we may assume that  $Q$  is a finite quiver with no oriented cycle and that  $T \in \text{mod } kQ$  is a tilting module such that  $B = \text{End}_{kQ}(T)$ . Any module  $X \in \text{mod } B$  defines the  $\Sigma$ -module  $\text{Hom}_B(\Sigma, X)$  which, as a functor, assigns the vector space  $\text{Hom}_B(E, X)$  to the object  $E$  of  $\Sigma$ . By the above properties of  $B$ , the map  $x \mapsto \text{Hom}_{kQ}(T, D(kQ e_x))$  defines an isomorphism of  $k$ -categories  $kQ \xrightarrow{\sim} \Sigma$ . We denote by  $\Gamma_{\leq \Sigma}$  the full subquiver of  $\Gamma$  generated by all the predecessors of  $\Sigma$  in  $\Gamma$ .

*The covering of the left section  $\Sigma$*

Let  $\tilde{\Sigma}$  be the full subcategory of  $k(\tilde{\Gamma})$  whose objects are the  $x \in k(\tilde{\Gamma})$  such that  $p(x) \in \Sigma$ . Therefore  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  induces a covering functor  $p : \tilde{\Sigma} \rightarrow \Sigma$ . Note that  $\tilde{\Sigma}$  and  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$  are stable under  $G$ , as subquivers of  $\tilde{\Gamma}$ . Since  $\Sigma$  is hereditary, so is  $\tilde{\Sigma}$ . Therefore we have  $\tilde{\Sigma} = k\tilde{Q}$  for some quiver  $\tilde{Q}$ . In particular, the isomorphism  $kQ \xrightarrow{\sim} \Sigma$  and the covering functor  $p : \tilde{\Sigma} \rightarrow \Sigma$  induce a covering functor  $q : k\tilde{Q} \rightarrow kQ$ .

*The covering functor of  $B$*

Since  $\pi$  and  $p$  coincide on vertices,  $\pi$  induces a Galois covering of quivers  $\pi : \tilde{Q} \rightarrow Q$  with group  $G$ . We write  $\pi : k\tilde{Q} \rightarrow kQ$  for the induced Galois covering with group  $G$ . Note that  $\tilde{Q}$  is a disjoint union of copies of the universal cover of  $Q$  because  $\tilde{\Gamma}$  is simply connected. Also, thanks to the Galois covering  $\pi : \tilde{Q} \rightarrow Q$  there is an action of  $G$  on  $\text{mod } k\tilde{Q}$ . Let  $T = T_1 \oplus \dots \oplus T_n$  be such that  $T_1, \dots, T_n$  are indecomposable and  $\tilde{B}$  be the full subcategory of  $\text{mod } k\tilde{Q}$  with objects the  ${}^g \tilde{T}_i$  (with  $i \in \{1, \dots, n\}$ ,  $g \in G$ , see 4.3).

**Lemma 5.1.** *The  $k$ -category  $\tilde{B}$  is locally bounded. The push-down functor  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$  induces a covering functor:*

$$F : \tilde{B} \rightarrow B,$$

$${}^g\tilde{T}_i \mapsto T_i = q_\lambda({}^g\tilde{T}_i).$$

Moreover, if  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  is a Galois covering with group  $\pi_1(\Gamma)$ , then so is  $F$ .

**Proof.** We apply the results of the preceding section to the covering functor  $q : k\tilde{Q} \rightarrow kQ$  and the Galois covering  $\pi : k\tilde{Q} \rightarrow kQ$ . The first assertion follows from 4.1 and 4.2, and the second from 4.2. The last assertion was proved in [29, Lem. 2.2].  $\square$

We also have a Galois covering  $\tilde{B} \rightarrow B$  induced by the push-down  $\pi_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$  (see [29, Lem. 2.2]). In particular, the covering functor  $F : \tilde{B} \rightarrow B$  and the Galois covering  $\tilde{B} \rightarrow B$  coincide on objects. Therefore we may apply the results of the preceding section to  $F$ .

In the sequel, we write  $\tilde{T}$  for the  $k\tilde{Q}$ -module  $\bigoplus\{{}^g\tilde{T}_i \mid i \in \{1, \dots, n\}, g \in G\}$ . Although  $\tilde{T}$  is not necessarily finite-dimensional, it follows from 4.2 that it induces a well-defined functor:

$$\text{Hom}_{k\tilde{Q}}(\tilde{T}, -) : \text{mod } k\tilde{Q} \rightarrow \text{mod } \tilde{B}.$$

More precisely, if  $X \in \text{mod } k\tilde{Q}$ , then  $\text{Hom}_{k\tilde{Q}}(\tilde{T}, X)$  is the  $\tilde{B}$ -module defined by  ${}^g\tilde{T}_i \mapsto \text{Hom}_{k\tilde{Q}}({}^g\tilde{T}_i, X)$ . In particular, an object  $x$  in  $\tilde{\Sigma} = k\tilde{Q}$  defines the injective  $k\tilde{Q}$ -module  $D(k\tilde{Q}(x, -))$  which gives rise to the  $\tilde{B}$ -module  $\text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(x, -)))$ . Therefore every  $\tilde{B}$ -module  $X$  defines a  $\tilde{\Sigma}$ -module:

$$\tilde{\Sigma}^{op} \rightarrow \text{mod } k,$$

$$x \mapsto \text{Hom}_{\tilde{B}}(\text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(x, -))), X).$$

For reasons that will become clear later, this module is denoted by  $\text{Hom}_{\tilde{B}}(\tilde{\Sigma}, X)$ . In this way, we get a functor  $\text{Hom}_{\tilde{B}}(\tilde{\Sigma}, -) : \text{mod } \tilde{B} \rightarrow \text{mod } \tilde{\Sigma}$ . We need the following result for later reference.

**Lemma 5.2.** *The following diagram commutes up to isomorphism of functors:*

$$\begin{array}{ccccc}
 \text{mod } k\tilde{Q} & \xrightarrow{\text{Hom}_{k\tilde{Q}}(\tilde{T}, -)} & \text{mod } \tilde{B} & \xrightarrow{\text{Hom}_{\tilde{B}}(\tilde{\Sigma}, -)} & \text{mod } \tilde{\Sigma} \\
 q_\lambda \downarrow & & \downarrow F_\lambda & & \downarrow p_\lambda \\
 \text{mod } kQ & \xrightarrow{\text{Hom}_{kQ}(T, -)} & \text{mod } B & \xrightarrow{\text{Hom}_B(\Sigma, -)} & \text{mod } \Sigma.
 \end{array}$$

Moreover:

- (a) *The two top horizontal arrows are  $G$ -equivariant.*
- (b) *If  $\theta : \text{mod } kQ \rightarrow \text{mod } \Sigma$  (or  $\tilde{\theta} : \text{mod } k\tilde{Q} \rightarrow \text{mod } \tilde{\Sigma}$ ) denotes the composition of the two bottom (or top) horizontal arrows, then it induces an equivalence from the full subcategory of injective  $kQ$ -modules (or injective  $k\tilde{Q}$ -modules) to the full subcategory of projective  $\Sigma$ -modules (or projective  $\tilde{\Sigma}$ -modules, respectively).*
- (c) *Let  $\tilde{\alpha} : \tilde{T} \rightarrow \tilde{J}$  be a surjective morphism between injective  $k\tilde{Q}$ -modules. Let  $\alpha : I \rightarrow J$  be equal to  $q_\lambda(\tilde{\alpha})$ . Then  $F_\lambda$  maps the connecting morphism  $\text{Hom}_{k\tilde{Q}}(\tilde{T}, \tilde{J}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{\alpha})$  to the connecting morphism  $\text{Hom}_{kQ}(T, J) \rightarrow \text{Ext}_{kQ}^1(T, \text{Ker } \alpha)$ .*

**Proof.** The commutativity of the diagram is an easy exercise on covering functors, and left to the reader.

(a) This follows from a direct computation.

(b) By tilting theory,  $\theta$  induces an equivalence (see [11, Chap. VIII, Thm. 3.5]):

$$\begin{aligned} \Phi : \text{inj } kQ &\rightarrow \text{proj } \Sigma, \\ I &\mapsto \text{Hom}_B(\Sigma, \text{Hom}_{kQ}(T, I)). \end{aligned}$$

Let  $I \in \text{inj } k\tilde{Q}$ . Then  $p_\lambda \tilde{\theta}(I) = \theta q_\lambda(I)$ . Moreover,  $q_\lambda$  maps indecomposable injective  $k\tilde{Q}$ -modules to indecomposable injective  $kQ$ -modules, because so does  $\pi_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$  (see 4.2). So  $p_\lambda \tilde{\theta}(I)$  is indecomposable projective, and therefore so is  $\tilde{\theta}(I)$  (see [14, 3.2]). Consequently,  $\tilde{\theta}$  induces the following functor:

$$\begin{aligned} \Psi : \text{inj } k\tilde{Q} &\rightarrow \text{proj } \tilde{\Sigma}, \\ I &\mapsto \text{Hom}_{\tilde{B}}(\tilde{\Sigma}, \text{Hom}_{k\tilde{Q}}(\tilde{T}, I)). \end{aligned}$$

So we have a commutative diagram:

$$\begin{array}{ccc} \text{inj } k\tilde{Q} & \xrightarrow{\Psi} & \text{proj } \tilde{\Sigma} \\ q_\lambda \downarrow & & \downarrow p_\lambda \\ \text{inj } kQ & \xrightarrow[\Phi]{\sim} & \text{proj } \Sigma. \end{array}$$

In this diagram,  $p_\lambda$ ,  $q_\lambda$  and  $\Phi$  are faithful. Hence, so is  $\Psi$ . Let  $I, J \in \text{inj } k\tilde{Q}$  and  $f : \Psi(I) \rightarrow \Psi(J)$ . Let  $h : q_\lambda I \rightarrow q_\lambda J$  be such that  $\Phi(h) = p_\lambda(f)$ . Using 4.2, we have  $h = \sum_{g \in G} q_\lambda(h_g)$ , where  $(h_g)_g \in \bigoplus_{g \in G} \text{Hom}_{k\tilde{Q}}(\tilde{I}, {}^g \tilde{J})$ . So  $p_\lambda(f) = \sum_{g \in G} p_\lambda \Psi(h_g)$ . Using 4.2 again, we deduce that  $f = \Psi(h_1)$ . So  $\Psi$  is full. Finally, we know from the preceding section that  $q_\lambda : \text{inj } k\tilde{Q} \rightarrow \text{inj } kQ$  is dense. Also, so is  $p_\lambda : \text{proj } \tilde{\Sigma} \rightarrow \text{proj } \Sigma$  (see [14, 3.2], for instance). Since  $\Phi$  is an equivalence, we deduce that  $\Psi$  is dense. Therefore  $\Psi$  is an equivalence.

(c) The push-down functors  $q_\lambda$  and  $F_\lambda$  are exact. So we have a commutative diagram up to isomorphism of functors:

$$\begin{array}{ccc} \mathcal{D}^b(\text{mod } k\tilde{Q}) & \xrightarrow{\mathbf{R}\text{Hom}_{k\tilde{Q}}(\tilde{T}, -)} & \mathcal{D}^b(\text{mod } \tilde{B}) \\ q_\lambda \downarrow & & \downarrow F_\lambda \\ \mathcal{D}^b(\text{mod } kQ) & \xrightarrow{\mathbf{R}\text{Hom}_{kQ}(T, -)} & \mathcal{D}^b(\text{mod } B). \end{array}$$

The statement follows from this diagram.  $\square$

We wish to construct a functor  $\varphi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{B}$ . We proceed in several steps:

1. Define a functor  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{mod } \tilde{B}$  where  $k(\tilde{\Gamma}_{\leq \tilde{\Sigma}})$  denotes the full subcategory of  $k(\tilde{\Gamma})$  with objects the vertices in  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$ .
2. Define  $\varphi$  on objects, so that it coincides with  $\varphi_0$  on predecessors of  $\tilde{\Sigma}$ .
3. Define  $\varphi$  on morphisms, so that it extends  $\varphi_0$ .

The functor  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{mod } \tilde{B}$

We first prove the following lemma. In the case of a Galois covering whose group acts freely on indecomposables, a corresponding result was proved in [21, 3.6]. We know that  $p : \tilde{\Sigma} \rightarrow \Sigma$  and  $F : \tilde{B} \rightarrow B$  are covering functors, and that the latter coincides on objects with a Galois covering  $\tilde{B} \rightarrow B$  with group  $G$ . Finally, if  $X \in \text{ind } B$  is a summand of a tilting  $B$ -module, then  $\tilde{X} \in \text{ind } \tilde{B}$  is such that  $p_\lambda(\tilde{X}) \simeq X$  (see 4.2).

**Lemma 5.3.** *Let  $X \in \Gamma_{\leq \Sigma}$  and  $g_0 \in G$ . If  $\tilde{u} : \tilde{E} \rightarrow {}^{g_0}\tilde{X}$  is right minimal almost split, then so is  $p_\lambda \tilde{u} : p_\lambda \tilde{E} \rightarrow X$ . Consequently,  $p_\lambda \tau_{\tilde{B}}({}^{g_0}\tilde{X}) \simeq \tau_B X$  if  $X$  is not projective.*

**Proof.** Notice that  $X$  is an indecomposable summand of some tilting  $B$ -module. So we may apply the results of 4.2. If  $X$  is projective, the assertion follows from [14, 3.2]. So we assume that  $X$ , and therefore  ${}^{g_0}\tilde{X}$ , are not projective. Let  $u : E \rightarrow X$  be right minimal almost split and  $E = E_1 \oplus \dots \oplus E_t$  be such that  $E_1, \dots, E_t$  are indecomposable. Since  $\Gamma_{\leq \Sigma}$  is acyclic (see [1, 2.2]), we have  $\text{Ext}_B^1(E, \tau_B X) = 0$ . Also, the linear map  $\bigoplus_{g \in G} \text{Hom}_{\tilde{B}}({}^g \tilde{E}_i, {}^{g_0}\tilde{X}) \rightarrow \text{Hom}_B(E_i, X)$  is bijective, for every  $i$  (see 4.2). Therefore we apply 4.3 to the exact sequence  $0 \rightarrow \tau_B X \rightarrow E \xrightarrow{u} X \rightarrow 0$ : There exist  $g_1, \dots, g_t \in G$  and morphisms  $\tilde{u}_i : {}^{g_i}\tilde{E}_i \rightarrow {}^{g_0}\tilde{X}$  ( $i \in \{1, \dots, t\}$ ) fitting into a commutative diagram whose vertical arrow on the left is an isomorphism:

$$\begin{array}{ccc}
 E & \xrightarrow{p_\lambda[\tilde{u}_1, \dots, \tilde{u}_t]} & X \\
 \sim \downarrow & & \parallel \\
 E & \xrightarrow{u} & X.
 \end{array}$$

We identify  $u$  and  $p_\lambda[\tilde{u}_1, \dots, \tilde{u}_t]$  via this diagram. Let  $i \in \{1, \dots, n\}$ . Then  $\tilde{u}_i : {}^{g_i}\tilde{E}_i \rightarrow {}^{g_0}\tilde{X}$  is not a retraction because  ${}^{g_i}\tilde{E}_i$  and  ${}^{g_0}\tilde{X}$  are non-isomorphic indecomposable modules. So  $\tilde{u}_i$  factors through  $\tilde{u}$ , for every  $i$ . Applying  $p_\lambda$  to each factorisation shows that  $u$  factors through  $p_\lambda(\tilde{u})$ . On the other hand,  $p_\lambda \tilde{u}$  is not a retraction because  $X$  is not a direct summand of  $E$ . So  $p_\lambda(\tilde{u})$  factors through  $u$ . The right minimality of  $u$  implies that the morphism  $u$  is a direct summand of  $p_\lambda(\tilde{u})$ . Finally, the following equality follows from 2.4:

$$\dim \text{Ker } u = \dim \tau_B X = \dim \tau_B p_\lambda {}^{g_0}\tilde{X} = \dim p_\lambda \tau_{\tilde{B}} {}^{g_0}\tilde{X} = \dim \text{Ker } p_\lambda(\tilde{u}).$$

So  $p_\lambda(\tilde{u})$  and  $u$  are isomorphic, and  $p_\lambda(\tilde{u})$  is right minimal almost split.  $\square$

Using the preceding lemma, we construct a functor  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{ind } \tilde{B}$ .

**Lemma 5.4.** *There exists a full and faithful functor,  $G$ -equivariant on vertices,  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{ind } \tilde{B}$ . This functor maps arrows in  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$  to irreducible maps, and meshes to almost split sequences. Moreover, it commutes with the translations and extends the canonical functor  $\tilde{\Sigma} \rightarrow \text{ind } \tilde{B}$  defined on the objects by  $x \mapsto \text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(x, -)))$ . Finally, the following diagram is commutative up to isomorphism of functors:*

$$\begin{array}{ccccc}
 k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) & \xrightarrow{\varphi_0} & \text{ind } \tilde{B} & \hookrightarrow & \text{mod } \tilde{B} \\
 p \downarrow & & & & \downarrow F_\lambda \\
 \text{ind}(\Gamma_{\leq \Sigma}) & \hookrightarrow & \text{ind } B & \hookrightarrow & \text{mod } B.
 \end{array}$$

**Proof. Step 1:** Clearly there is a functor  $\varphi_0 : \tilde{\Sigma} \rightarrow \text{mod } \tilde{B}$  given by  $\tilde{x} \mapsto \text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(\tilde{x}, -)))$ . Note that  $\tilde{\Sigma}$  (or  $\Sigma$ ) is naturally equivalent to the full subcategory of  $\text{mod } k\tilde{Q}$  (or  $\text{mod } kQ$ , respectively) consisting of the indecomposable injective modules. Therefore 5.2 shows that this functor is full and faithful, and that the following diagram commutes up to isomorphism:

$$\begin{array}{ccccc} \tilde{\Sigma} & \xrightarrow{\varphi_0} & \text{ind } \tilde{B} & \hookrightarrow & \text{mod } \tilde{B} \\ p \downarrow & & & & \downarrow F_\lambda \\ \Sigma & \hookrightarrow & \text{ind } B & \hookrightarrow & \text{mod } B. \end{array}$$

Note that  $\varphi_0(M)$  is indecomposable for every  $M$  because so is  $F_\lambda \varphi_0(M) = p(M)$ . The functor  $\varphi_0 : \tilde{\Sigma} \rightarrow \text{ind } \tilde{B}$  is  $G$ -equivariant on vertices: Indeed, for every  $g \in G$ , and every  $\tilde{x} \in \tilde{Q}_0$ , we have:

$$\begin{aligned} \varphi_0(g\tilde{x}) &= \text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(g\tilde{x}, -))) = \text{Hom}_{k\tilde{Q}}(\tilde{T}, {}^g D(k\tilde{Q}(\tilde{x}, -))) \\ &= {}^g \text{Hom}_{k\tilde{Q}}(\tilde{T}, D(k\tilde{Q}(\tilde{x}, -))) = {}^g \varphi_0(\tilde{x}). \end{aligned}$$

**Step 2:** If  $M \in k(\tilde{\Gamma}_{\leq \tilde{\Sigma}})$ , there exists a unique  $n \in \mathbb{N}$  such that  $\tau^{-n}M \in \tilde{\Sigma}$ . Let  $\varphi_0(M)$  be the  $\tilde{B}$ -module:

$$\varphi_0(M) = \tau_B^n \varphi_0(\tau^{-n}M).$$

It follows from 5.3 that  $F_\lambda \varphi_0(M) = p(M)$ . Also  $\varphi_0({}^g M) = {}^g \varphi_0(M)$  for every  $g \in G$  and for every vertex  $M$  because  $\tau$  commutes with the action of  $G$ .

**Step 3:** In order to define  $\varphi_0$  on morphisms, we construct inductively a sequence of  $G$ -invariant left sections  $\tilde{\Sigma}_i$  of  $\tilde{\Gamma}$  such that  $\tilde{\Sigma}_0 = \tilde{\Sigma}$ , such that  $\tilde{\Sigma}_{i+1} \setminus \tilde{\Sigma}_i$  consists of the  $G$ -orbit of a vertex, and such that, if  $\bigcup_{t=1}^i \tilde{\Sigma}_t$  denotes the full subcategory of the path category  $k\tilde{\Gamma}$  whose vertices are given by those of  $\tilde{\Sigma}_0, \dots, \tilde{\Sigma}_i$ , then  $k\tilde{\Gamma}_{\leq \tilde{\Sigma}} = \bigcup_{i \geq 0} \tilde{\Sigma}_i$ . Each inductive step defines a functor  $\varphi_0 : \bigcup_{t=1}^i \tilde{\Sigma}_t \rightarrow \text{ind } \tilde{B}$  which maps arrows to irreducible maps and extends the construction of the two preceding steps. This functor makes the following diagram commute:

$$\begin{array}{ccccc} \bigcup_{t=1}^i \tilde{\Sigma}_t & \xrightarrow{\varphi_0} & \text{ind } \tilde{B} & \hookrightarrow & \text{mod } \tilde{B} \\ \downarrow & & & & \downarrow F_\lambda \\ \text{ind}(\Gamma_{\leq \Sigma}) & \hookrightarrow & \text{ind } B & \hookrightarrow & \text{mod } B, \end{array}$$

where the vertical arrow on the left is induced by  $p$ . Assume that  $\varphi_0 : \bigcup_{t=1}^i \tilde{\Sigma}_t \rightarrow \text{ind } \tilde{B}$  has been defined for some  $i \geq 0$ . Since  $\tilde{\Sigma}_i$  is acyclic, it has a sink. Assume that all sinks are projective. First assume that  $P$  is a projective sink, and let  $\tilde{\Sigma}_{i+1}$  be equal to  $\tilde{\Sigma}_i \setminus \{gP \mid g \in G\}$ ; then  $\tilde{\Sigma}_{i+1}$  is a left section of  $\Gamma$ , and there is a unique  $\varphi_0 : \bigcup_{t=0}^{i+1} \tilde{\Sigma}_t \rightarrow \text{ind } \tilde{B}$  satisfying the required conditions. Now assume that there is a non-projective sink  $M$  in  $\tilde{\Sigma}_i$ . Then there exists a mesh in  $\tilde{\Gamma}$ :

$$\begin{array}{ccccc} & & N_1 & & \\ & v_1 \nearrow & & \searrow u_1 & \\ \tau M & & & & M, \\ & \searrow v_s & & \nearrow u_s & \\ & & N_s & & \end{array}$$

and  $M, N_1, \dots, N_s \in \tilde{\Sigma}_i$  because  $M \in \tilde{\Sigma}_i$  is a sink. In particular,  $\varphi_0(u_j)$  is defined, and  $F_\lambda \varphi_0(u_j) = p(u_j)|_B$  for every  $i$ . For simplicity, we write  $\varphi_0(u) = [\varphi_0(u_1) \ \dots \ \varphi_0(u_s)]$  and  $p(u) = [p(u_1) \ \dots \ p(u_s)]$ . Then  $\varphi_0(u)$  is right minimal almost split in  $\text{mod } \tilde{B}$ : Indeed, there exists a right minimal almost split morphism  $L \xrightarrow{w} \varphi_0(M)$ . Since  $\varphi_0(u) : \bigoplus_j \varphi_0(N_j) \rightarrow \varphi_0(M)$  is not a retraction (because each  $\varphi_0(u_j)$  is an irreducible morphism, by the induction hypothesis), there exists a morphism  $w' : \bigoplus_j \varphi_0(N_j) \rightarrow L$  such that  $\varphi_0(u) = ww'$ ; applying  $F_\lambda$ , we have  $F_\lambda \varphi_0(u) = F_\lambda(w)F_\lambda(w')$ ; but now  $F_\lambda \varphi_0(u) = p(u)|_B$  is right minimal almost split by construction, and so is  $F_\lambda(w)$  (see 5.3); hence,  $F_\lambda(w')$  is an isomorphism and therefore so is  $w'$  because  $F_\lambda$  is exact. We let  $[\varphi_0(v_1) \ \dots \ \varphi_0(v_s)]^t : \varphi_0(\tau M) \rightarrow \bigoplus_{j=1}^s \varphi_0(N_j)$  be the kernel of  $\varphi_0(u)$ . For simplicity, we set  $\varphi_0(v) = [\varphi_0(v_1) \ \dots \ \varphi_0(v_s)]^t$  and  $p(v) = [p(v_1) \ \dots \ p(v_s)]^t$ . We let  $\tilde{\Sigma}_{i+1} = (\tilde{\Sigma}_i \setminus \{gM \mid g \in G\}) \cup \{g\tau M \mid g \in G\}$ . Clearly,  $\tilde{\Sigma}_{i+1}$  is a left section. We now show that we may assume  $\varphi_0(v)$  to be taken such that  $F_\lambda \varphi_0(v) = p(u)|_B$ . Indeed, the commutative diagram with exact rows:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & F_\lambda \varphi_0(\tau M) & \xrightarrow{F_\lambda \varphi_0(v)} & F_\lambda \varphi_0(\bigoplus_{j=1}^s N_j) & \xrightarrow{F_\lambda \varphi_0(u)} & F_\lambda \varphi_0(M) \longrightarrow 0 \\
 & & & & \parallel & & \parallel \\
 0 & \longrightarrow & p(\tau M)|_B & \xrightarrow{p(v)|_B} & p(\bigoplus_{j=1}^s N_j)|_B & \xrightarrow{p(u)|_B} & p(M)|_B \longrightarrow 0
 \end{array}$$

gives an isomorphism  $F_\lambda \varphi_0(\tau M) \rightarrow p(\tau M)|_B$  making the left square commute. Since  $F_\lambda \varphi_0(\tau M) = p(\tau M)|_B$  is a brick (because it belongs to  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$ ), this isomorphism is the multiplication by a non-zero constant  $c$ . Hence,  $p(v)|_B = cF_\lambda \varphi_0(v)$ . Replacing  $\varphi_0(v)$  by  $c\varphi_0(v)$  does the trick. Thus, we have defined  $\varphi_0 : \bigcup_{t=1}^{i+1} \tilde{\Sigma}_t \rightarrow \text{ind } \tilde{B}$ . Clearly, the required conditions are satisfied. This induction gives a functor  $\varphi_0 : k\tilde{\Gamma}_{\leq \tilde{\Sigma}} \rightarrow \text{ind } \tilde{B}$  mapping arrows to irreducible maps and meshes to almost split sequences, and such that the following diagram commutes:

$$\begin{array}{ccccc}
 k\tilde{\Gamma}_{\leq \tilde{\Sigma}} & \xrightarrow{\varphi_0} & \text{ind } \tilde{B} & \hookrightarrow & \text{mod } \tilde{B} \\
 \downarrow & & & & \downarrow F_\lambda \\
 \text{ind } \Gamma_{\leq \Sigma} & \hookrightarrow & \text{ind } B & \hookrightarrow & \text{mod } B,
 \end{array}$$


where the vertical arrow on the left is induced by  $p$ . Since  $F_\lambda$  is faithful,  $\varphi_0$  induces a functor  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{ind } \tilde{B}$ . It is now clear that this functor satisfies the conditions of the lemma.  $\square$

It was shown in [1, 3.2] that the existence of a left section  $\Sigma$  in an Auslander–Reiten component  $\Gamma$  implies that  $\Gamma_{\leq \Sigma}$  is generalised standard. We now prove that it is standard.


**Corollary 5.5.** *Let  $A$  be a finite-dimensional  $k$ -algebra and  $\Gamma$  be a component of  $\Gamma(\text{mod } A)$  having a left section  $\Sigma$ . Then  $\Gamma_{\leq \Sigma}$  is standard.*

**Proof.** Let  $B = A/\text{Ann } \Sigma$ . Then  $B$  is a product of tilted algebras and the components of  $\Sigma$  form complete slices of the connecting components of the connected components of  $B$ . Let  $\Gamma'$  be the union of the components of  $\Gamma(\text{mod } B)$  intersecting  $\Sigma$ . The arguments of the proof of 5.4 show that there exists a full and faithful functor  $k(\Gamma'_{\leq \Sigma}) \rightarrow \text{ind } \Gamma'_{\leq \Sigma}$  extending the identity on vertices. So  $\Gamma_{\leq \Sigma} = \Gamma'_{\leq \Sigma}$  is standard.  $\square$


**Example 5.6.** Let  $A$  be the algebra given by the quiver




and the potential  $W = \delta\beta\alpha + \nu\mu\lambda$  (or, equivalently, by the relations  $\beta\alpha = 0, \delta\beta = 0, \alpha\delta = 0, \mu\lambda = 0, \nu\mu = 0$  and  $\lambda\nu = 0$ ). So  $A$  is a cluster-tilted algebra since it is the relation–extension (in the sense of [2]) of the tilted algebra of type  $\tilde{A}$  given by the quiver



bound by  $\beta\alpha = 0$  and  $\mu\lambda = 0$ . The transjective component  $\Gamma$  of  $\Gamma(\text{mod } A)$  is of the form



where vertices with the same label are identified. Then  $\Gamma$  admits a left section  $\Sigma = \{e, r, q, p, o, c\}$  and  $B = A/\text{Ann } \Sigma$  is the algebra given by the quiver:



with the inherited relations. As we have seen,  $\Gamma_{\leq \Sigma}$  is standard (and generalised standard) while  $\Gamma$  itself is not.

The following corollary seems to be well-known. However we have been unable to find a reference.

**Corollary 5.7.** *Let  $B$  be a tilted algebra and  $\Gamma$  a connecting component of  $B$ . Then  $\Gamma$  is standard.*

**Proof.** If  $B$  is concealed, this follows from [38, 2.4(11), p. 80]. Assume that  $B$  is not concealed. So  $\Gamma$  is the unique connecting component of  $B$ . Let  $\Sigma$  be a complete slice in  $\Gamma$ . As observed in 5.5, we have a full and faithful functor  $k(\Gamma_{\leq \Sigma}) \rightarrow \text{ind } \Gamma$  extending the identity on vertices. A dual construction extends this functor to a full and faithful functor  $k(\Gamma^{\geq}) \rightarrow \text{ind } \Gamma$  extending the identity on vertices. So  $\Gamma$  is standard.  $\square$

From now on, we identify  $\tilde{\Sigma}$  to a full subcategory of  $\text{mod } \tilde{B}$  by means of  $\varphi_0$ .

*Construction of  $\varphi$  on objects*

We prove that for any  $M \in \tilde{\Gamma}$ , there exists  $\varphi(M) \in \text{mod } \tilde{B}$  whose image under  $F_\lambda : \text{mod } \tilde{B} \rightarrow \text{mod } B$  coincides with  $p(M)|_B$ , in such a way that  $\varphi({}^g M) = {}^g \varphi(M)$ , for every  $g \in G$ . We define  $\mathcal{L}_\Sigma$  to be the full subcategory of  $\text{ind } B$  which consists of the predecessors of the complete slice  $\Sigma$ . Also a *minimal add  $\mathcal{L}_\Sigma$ -presentation* of a module  $R$  is a sequence of morphisms  $E_1 \rightarrow E_2 \rightarrow R$  where the morphism on the right is a minimal  $\text{add } \mathcal{L}_\Sigma$ -approximation and the one on the left is a minimal  $\text{add } \mathcal{L}_\Sigma$ -approximation of its kernel. Before constructing  $\varphi(M)$ , we prove some lemmata.

**Lemma 5.8.** *Let  $R \in \text{mod } B$  be a module with no direct summand in  $\mathcal{L}_\Sigma$ . There exists an exact sequence in  $\text{mod } B$ , which is a minimal  $\text{add } \mathcal{L}_\Sigma$ -presentation:*

$$0 \rightarrow E_1 \rightarrow E_2 \rightarrow R \rightarrow 0 \tag{*}$$

with  $E_1, E_2 \in \text{add } \Sigma$ . Moreover, the functor  $\text{Hom}_{kQ}(T, -)$  induces a bijection between the class of all such exact sequences, and the class of minimal injective copresentations:

$$0 \rightarrow \text{Tor}_1^B(R, T) \rightarrow I_1 \rightarrow I_2 \rightarrow 0.$$

Finally, there is an isomorphism in  $\text{mod } B$ :

$$R \simeq \text{Ext}_{kQ}^1(T, \text{Tor}_1^B(R, T)).$$

**Proof.** Let  $\mathcal{X}(T)$  be the torsion class induced by  $T$  in  $\text{mod } B$ . So  $R$  lies in  $\mathcal{X}(T)$  and has no direct summand in  $\Sigma$ . Therefore  $R$  is the epimorphic image of a module in  $\text{add } \Sigma$ . The first assertion then follows from [9, 2.2(d)].

Let  $f : I_1 \rightarrow I_2$  be the morphism between injective  $kQ$ -modules such that  $\text{Hom}_{kQ}(T, f)$  is equal to the morphism  $E_1 \rightarrow E_2$  in  $(*)$ . Because of the Brenner–Butler Theorem (see [11, Chap. VI, Thm. 3.8, p. 207]), the functor  $-\otimes_B T$  applied to  $(*)$  yields an injective copresentation in  $\text{mod } kQ$ :

$$0 \rightarrow \text{Tor}_1^B(R, T) \rightarrow I_1 \rightarrow I_2 \rightarrow 0.$$

The minimality of this copresentation follows from the minimality of  $E_2 \rightarrow R$ . With these arguments, it is straightforward to check that there is a well-defined bijection which carries the equivalence class of the exact sequence  $0 \rightarrow E_1 \rightarrow E_2 \rightarrow R \rightarrow 0$  to the equivalence class of the exact sequence  $0 \rightarrow \text{Tor}_1^B(R, T) \rightarrow I_1 \rightarrow I_2 \rightarrow 0$ .

The last assertion follows from the Brenner–Butler Theorem and the fact that  $R \in \mathcal{X}(T)$ .  $\square$

**Lemma 5.9.**  *$\text{add } \mathcal{L}_\Sigma$  is contravariantly finite in  $\text{mod } A$ . Therefore if  $X \in \Gamma \setminus \mathcal{L}_\Sigma$ , then  $X|_B$  lies in the torsion class induced by  $T$  in  $\text{mod } B$ .*

**Proof.** By [1, Thm. B], the algebra  $B$  is the endomorphism algebra of the indecomposable projective  $A$ -modules in  $\mathcal{L}_\Sigma$ . In particular, a projective  $B$ -module is projective as an  $A$ -module so the projective dimensions in  $\text{mod } A$  and in  $\text{mod } B$  coincide on  $\mathcal{L}_\Sigma$ . Also, by [1, Thm. B], all modules in  $\mathcal{L}_\Sigma$  have

projective dimension at most one as  $B$ -modules. Therefore  $\mathcal{L}_\Sigma \subseteq \mathcal{L}_A$ . Moreover,  $\bigoplus \Sigma$  is sincere as a  $B$ -module. Hence, [1, 8.2] implies that  $\text{add } \mathcal{L}_\Sigma$  is contravariantly finite in  $\text{mod } A$ . Let  $X \in \Gamma \setminus \mathcal{L}_\Sigma$ . Let  $P \twoheadrightarrow X|_B$  be a projective cover in  $\text{mod } B$ . As noticed above, we have  $P \in \text{add } \mathcal{L}_\Sigma$ . Therefore  $P \twoheadrightarrow X|_B$  factors through  $\text{add } \Sigma$ . Thus,  $X|_B$  lies in the torsion class.  $\square$

**Lemma 5.10.** *There exists a map  $\varphi : \tilde{\Gamma}_0 \rightarrow \text{mod } \tilde{B}$  extending  $\varphi_0$ , and such that  $F_\lambda(\varphi(M)) = p(M)|_B$ , for every  $M \in \tilde{\Gamma}$ . Moreover,  $\varphi({}^g M) = {}^g \varphi(M)$  for every  $g \in G$  and  $M \in \tilde{\Gamma}$ .*

**Proof.** Note that  $\varphi$  is already defined on  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$  because of 5.4. Let  $M \in \tilde{\Gamma} \setminus \tilde{\Gamma}_{\leq \tilde{\Sigma}}$ . Then  $p(M) \in \Gamma \setminus \Gamma_{\leq \Sigma} = \Gamma \setminus \mathcal{L}_\Sigma$ . By 5.9, the module  $p(M)|_B$  lies in the torsion class induced by  $T$  in  $\text{mod } B$ . So there is a decomposition in  $\text{mod } B$ :

$$p(M)|_B = R \oplus E,$$

where  $E \in \text{add } \Sigma$  and  $R$  has no indecomposable summand in  $\mathcal{L}_\Sigma$ . Also, fix a decomposition in  $\text{mod } \tilde{\Sigma}$ :

$$k(\tilde{\Gamma})(\tilde{\Sigma}, M) = \tilde{R} \oplus \tilde{P},$$

where  $\tilde{P}$  is projective and maximal for this property. Let  $\tilde{E} \in \text{add } \tilde{\Sigma}$  be such that  $\tilde{P} = k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E})$ .

We claim that  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$  maps  $\tilde{R}$  and  $\tilde{P}$  to  $\text{Hom}_B(\Sigma, R)$  and  $\text{Hom}_B(\Sigma, E)$  respectively. Indeed, since  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  is a covering functor inducing  $p : \tilde{\Sigma} \rightarrow \Sigma$ , the image of  $k(\tilde{\Gamma})(\tilde{\Sigma}, M)$  under  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$  is  $\text{Hom}_A(\Sigma, p(M)) \cong \text{Hom}_B(\Sigma, p(M)|_B)$  (functorially in  $M$ ). Moreover, the decomposition  $p(M)|_B = R \oplus E$  in  $\text{mod } B$  gives a decomposition  $\text{Hom}_B(\Sigma, p(M)) = \text{Hom}_B(\Sigma, R) \oplus \text{Hom}_B(\Sigma, E)$  in  $\text{mod } \Sigma$  where  $\text{Hom}_B(\Sigma, E)$  is projective and  $\text{Hom}_B(\Sigma, R)$  has no non-zero projective direct summand. The claim then follows from [14, 3.2].

In order to prove that  $R$  is the image of a  $\tilde{B}$ -module under  $F_\lambda$ , we consider a minimal projective presentation in  $\text{mod } \tilde{\Sigma}$ :

$$0 \rightarrow \tilde{P}_1 \rightarrow \tilde{P}_2 \rightarrow \tilde{R} \rightarrow 0.$$

Then there exists a morphism  $\tilde{f} : \tilde{I}_1 \rightarrow \tilde{I}_2$  between injective  $k\tilde{Q}$ -modules such that the morphism  $\tilde{P}_1 \rightarrow \tilde{P}_2$  equals  $\tilde{\theta}(\tilde{f})$  (here  $\tilde{\theta}$  is as in 5.2). Let  $f : I_1 \rightarrow I_2$  be the image of  $\tilde{f}$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$ . Hence, the image of  $\text{Ker } \tilde{f}$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$  is  $\text{Ker } f$ . Let  $P_1 \rightarrow P_2$  be the image of  $\tilde{\theta}(\tilde{f})$  under  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$ . Therefore the commutativity of the diagram in 5.2 and the fact that  $\text{Hom}_B(\Sigma, R)$  is the image of  $\tilde{R}$  under  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$  gives a minimal projective presentation in  $\text{mod } \Sigma$ :

$$0 \rightarrow P_1 \rightarrow P_2 \rightarrow \text{Hom}_B(\Sigma, R) \rightarrow 0.$$

On the other hand, 5.2 shows that  $P_1 \rightarrow P_2$  is equal to the following morphism in  $\text{mod } \Sigma$ :

$$\text{Hom}_B(\Sigma, \text{Hom}_{kQ}(T, I_1)) \xrightarrow{\text{Hom}_B(\Sigma, \text{Hom}_{kQ}(T, f))} \text{Hom}_B(\Sigma, \text{Hom}_{kQ}(T, I_2)).$$

Therefore we have a minimal  $\text{add } \mathcal{L}_\Sigma$ -presentation:

$$\text{Hom}_{kQ}(T, I_1) \xrightarrow{\text{Hom}_{kQ}(T, f)} \text{Hom}_{kQ}(T, I_2) \rightarrow R.$$

Because of 5.8, the sequence  $0 \rightarrow \text{Hom}_{kQ}(T, I_1) \rightarrow \text{Hom}_{kQ}(T, I_2) \rightarrow R \rightarrow 0$  is exact and  $\text{Ker } f = \text{Tor}_1^B(R, T)$ . In other words,  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$  maps  $\text{Ker } \tilde{f}$  to  $\text{Tor}_1^B(R, T)$ . Using 5.8 and the last diagram in the proof of 5.2, we get  $F_\lambda(\text{Ext}_{kQ}^1(\tilde{T}, \text{Ker } \tilde{f})) = R$ .

We give an explicit construction of  $\varphi$ . Let  $M \in k(\tilde{\Gamma})$ . We fix a minimal projective presentation in  $\text{mod } \tilde{\Sigma}$ :

$$0 \rightarrow \tilde{P}_1 \xrightarrow{\tilde{u}} \tilde{P}_2 \rightarrow \tilde{R} \rightarrow 0,$$

and injective  $k\tilde{Q}$ -modules  $\tilde{I}_1$  and  $\tilde{I}_2$ , together with a morphism  $\tilde{f} : \tilde{I}_1 \rightarrow \tilde{I}_2$  such that  $\tilde{u} = \tilde{\theta}(\tilde{f})$ . Then we let  $\varphi(M)$  be the following  $\tilde{B}$ -module:

$$\varphi(M) = \varphi_0(\tilde{E}) \oplus \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}),$$

where  $\varphi_0(\tilde{E}) = \varphi_0(\tilde{E}_1) \oplus \dots \oplus \varphi_0(\tilde{E}_s)$  if  $\tilde{E} = \tilde{E}_1 \oplus \dots \oplus \tilde{E}_s$  with  $\tilde{E}_1, \dots, \tilde{E}_s \in \tilde{\Sigma}$ . This finishes the construction of the map  $\varphi : \tilde{\Gamma}_0 \rightarrow \text{mod } \tilde{B}$ . We now prove the  $G$ -equivariance property. Let  $M \in k(\tilde{\Gamma})$  be a vertex and let  $g \in G$ . We keep the above notation  $\tilde{R}, \tilde{E}$ , etc. introduced for  $M$ , and we adopt the dashed notation  $\tilde{R}', \tilde{E}'$ , etc. for the corresponding objects associated to  ${}^gM$ . We have  $k(\tilde{\Gamma})(\tilde{\Sigma}, {}^gM) = {}^gk(\tilde{\Gamma})(\tilde{\Sigma}, M)$ . Indeed, the  $\tilde{\Sigma}$ -modules  $k(\tilde{\Gamma})(\tilde{\Sigma}, {}^gM)$  and  ${}^gk(\tilde{\Gamma})(\tilde{\Sigma}, M)$  are given by the functors  $X \mapsto k(\tilde{\Gamma})(X, {}^gM)$  and  $X \mapsto k(\tilde{\Gamma})({}^{g^{-1}}X, M)$  from  $\tilde{\Sigma}^{op}$  to  $\text{mod } k$ , respectively. These two functors coincide because  $G$  acts on  $k(\tilde{\Gamma})$ . Hence,  $\tilde{E}' = {}^g\tilde{E}$  and  $\tilde{R}' = {}^g\tilde{R}$ . Therefore any minimal projective presentation of  $\tilde{R}'$  in  $\text{mod } \tilde{\Sigma}$  is obtained from a minimal projective presentation of  $\tilde{R}$  by applying  $g$ . Since, moreover,  $\tilde{\theta}$  is  $G$ -equivariant (see 5.2), we deduce that  $\tilde{f}' = {}^g\tilde{f}$ . Finally, the  $G$ -action on  $\text{mod } k\tilde{Q}$  implies, as above, that  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } {}^g\tilde{f}) = \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, {}^g\text{Ker } \tilde{f}) = {}^g\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f})$ . From the construction of  $\varphi$ , we get  $\varphi({}^gM) = {}^g\varphi(M)$ .  $\square$

Construction of  $\varphi$  on morphisms

We complete the construction of  $\varphi$  by proving the following lemma.

**Lemma 5.11.** *Let  $u : M \rightarrow N$  be a morphism in  $k(\tilde{\Gamma})$ . Then there exists a unique morphism  $\varphi(u) : \varphi(M) \rightarrow \varphi(N)$  in  $\text{mod } \tilde{B}$ , such that  $F_\lambda(\varphi(u)) = p(u)|_B$ .*

**Proof.** Since  $F_\lambda$  is exact, it is faithful so the morphism  $\varphi(u)$  is unique. We prove its existence. By 5.4, we have constructed  $\varphi(u) = \varphi_0(u)$  in case  $N \in \tilde{\Gamma}_{\leq \tilde{\Sigma}}$ . So we may assume that  $N \in \tilde{\Gamma} \setminus \tilde{\Gamma}_{\leq \tilde{\Sigma}}$ . Since any path in  $\tilde{\Gamma}$  from a vertex in  $\tilde{\Gamma}_{\leq \tilde{\Sigma}}$  to  $N$  has a vertex in  $\tilde{\Sigma}$ , we may also assume that  $M \in (\tilde{\Gamma} \setminus \tilde{\Gamma}_{\leq \tilde{\Sigma}}) \cup \tilde{\Sigma}$ . The functor  $\varphi_0 : k(\tilde{\Gamma}_{\leq \tilde{\Sigma}}) \rightarrow \text{mod } \tilde{B}$  naturally extends to a unique functor  $\varphi_0 : \text{add}(k(\tilde{\Gamma}_{\leq \tilde{\Sigma}})) \rightarrow \text{mod } \tilde{B}$ , such that the following diagram commutes:

$$\begin{array}{ccc} \text{add}(k(\tilde{\Gamma}_{\leq \tilde{\Sigma}})) & \xrightarrow{\varphi_0} & \text{mod } \tilde{B} \\ \text{add } p \downarrow & & \downarrow F_\lambda \\ \text{add}(\text{ind } \Gamma_{\leq \Sigma}) & \hookrightarrow & \text{mod } B. \end{array}$$

We fix decompositions in  $\text{mod } \tilde{\Sigma}$  as in the proof of 5.10:

$$k(\tilde{\Gamma})(\tilde{\Sigma}, M) = \tilde{P} \oplus \tilde{R}, \quad \text{and} \quad k(\tilde{\Gamma})(\tilde{\Sigma}, N) = \tilde{P}' \oplus \tilde{R}'$$

where  $\tilde{P}, \tilde{P}'$  are projective and  $\tilde{R}, \tilde{R}'$  have no non-zero projective direct summand. We let  $\tilde{E}, \tilde{E}' \in \text{add } \tilde{\Sigma}$  be such that  $\tilde{P} = k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E})$  and  $\tilde{P}' = k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}')$ , respectively. Therefore the morphism  $k(\tilde{\Gamma})(\tilde{\Sigma}, u)$  can be written as:

$$k(\tilde{\Gamma})(\tilde{\Sigma}, u) = \begin{bmatrix} \tilde{u}_1 & 0 \\ \tilde{u}_2 & \tilde{u}_3 \end{bmatrix} : k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}) \oplus \tilde{R} \rightarrow k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}') \oplus \tilde{R}'.$$

Similarly, we fix decompositions in  $\text{mod } B$ :

$$p(M)|_B = E \oplus R, \quad p(N)|_B = E' \oplus R',$$

where  $E, E' \in \text{add } \Sigma$ , and  $R, R'$  have no direct summand in  $\Sigma$ . As above, the morphism  $p(u)|_B$  decomposes as:

$$p(u)|_B = \begin{bmatrix} u_1 & 0 \\ u_2 & u_3 \end{bmatrix} : E \oplus R \rightarrow E' \oplus R'.$$

Recall from the proof of 5.10 that  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$  maps  $k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}), \tilde{R}, k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}')$  and  $\tilde{R}'$  to  $\text{Hom}_B(\Sigma, E), \text{Hom}_B(\Sigma, R), \text{Hom}_B(\Sigma, E')$  and  $\text{Hom}_B(\Sigma, R')$ , respectively. As a consequence, it maps  $\tilde{u}_i$  to  $\text{Hom}_B(\Sigma, u_i)$ , for every  $i$ . As in the proof of 5.10, we have morphisms  $\tilde{f} : \tilde{I}_1 \rightarrow \tilde{I}_2$  and  $\tilde{f}' : \tilde{I}'_1 \rightarrow \tilde{I}'_2$  between injective  $k\tilde{Q}$ -modules and minimal projective presentations in  $\text{mod } \tilde{\Sigma}$ :

$$\tilde{\theta}(\tilde{I}_1) \xrightarrow{\tilde{\theta}(\tilde{f})} \tilde{\theta}(\tilde{I}_2) \rightarrow \tilde{R} \rightarrow 0 \quad \text{and} \quad \tilde{\theta}(\tilde{I}'_1) \xrightarrow{\tilde{\theta}(\tilde{f}')} \tilde{\theta}(\tilde{I}'_2) \xrightarrow{\tilde{v}} \tilde{R}' \rightarrow 0.$$

With these notations, we have:

$$\varphi(M) = \varphi_0(\tilde{E}) \oplus \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}) \quad \text{and} \quad \varphi(N) = \varphi_0(\tilde{E}') \oplus \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}').$$

Also, if  $M \in \tilde{\Sigma}$ , then  $\tilde{f} = 0$ , so  $\varphi(M) = \varphi_0(\tilde{E})$ .

It suffices to prove that each of  $u_1, u_2, u_3$  is the image under  $F_\lambda$  of some morphism  $\varphi_0(\tilde{E}) \rightarrow \varphi_0(\tilde{E}')$ ,  $\varphi_0(\tilde{E}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$  and  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$ , respectively. Clearly,  $u_1 : k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}) \rightarrow k(\tilde{\Gamma})(\tilde{\Sigma}, \tilde{E}')$  is induced by a morphism  $\tilde{E} \rightarrow \tilde{E}'$  in  $\text{add } \tilde{\Sigma}$ . This and 5.4 imply that  $u_1$  is the image under  $F_\lambda$  of a morphism  $\varphi_0(\tilde{E}) \rightarrow \varphi_0(\tilde{E}')$ . We now prove that  $u_2$  is the image under  $F_\lambda$  of a morphism  $\varphi_0(\tilde{E}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$ . Let  $f' : I'_1 \rightarrow I'_2$  be the image of  $\tilde{f}' : \tilde{I}'_1 \rightarrow \tilde{I}'_2$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$ . Therefore we have a minimal projective presentation in  $\text{mod } \Sigma$  (see 5.2 and the proof of 5.10):

$$0 \rightarrow \theta(I'_1) \xrightarrow{\theta(f')} \theta(I'_2) \rightarrow \text{Hom}_B(\Sigma, R') \rightarrow 0,$$

together with a minimal injective copresentation in  $\text{mod } kQ$ :

$$0 \rightarrow \text{Tor}_1^B(R', T) \rightarrow I'_1 \xrightarrow{f'} I'_2 \rightarrow 0.$$

Recall that  $\text{Tor}_1^B(R', T)$  is equal to the image of  $\text{Ker } \tilde{f}'$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$ . Therefore we have an exact sequence in  $\text{mod } B$ , which is also a minimal  $\text{add } \mathcal{L}_\Sigma$ -presentation:

$$0 \rightarrow \text{Hom}_{kQ}(T, I'_1) \rightarrow \text{Hom}_{kQ}(T, I'_2) \xrightarrow{v} R' \rightarrow 0,$$

where  $v$  is such that  $\text{Hom}_B(\Sigma, v)$  is the image of  $\tilde{v} : \tilde{\theta}(\tilde{I}'_2) \rightarrow \tilde{R}'$  under  $p_\lambda : \text{mod } \tilde{\Sigma} \rightarrow \text{mod } \Sigma$  (see the diagram in 5.2). The projective cover  $\tilde{v}$  of  $\tilde{R}'$  in  $\text{mod } \tilde{\Sigma}$  yields a morphism  $\tilde{\delta} : \tilde{I} \rightarrow \tilde{I}'_2$  in  $\text{mod } k\tilde{Q}$ , where  $\tilde{I}$  is the injective  $k\tilde{Q}$ -module such that  $\tilde{P} = \tilde{\theta}(\tilde{I})$ , and such that the following diagram of  $\text{mod } \tilde{\Sigma}$  commutes:

$$\begin{array}{ccc} & \tilde{P} & \\ & \swarrow \tilde{\theta}(\tilde{\delta}) & \downarrow \tilde{u}_2 \\ \tilde{\theta}(\tilde{I}'_2) & \xrightarrow{\tilde{v}} & \tilde{R}' \end{array}$$

Therefore if  $\delta : I \rightarrow I'_2$  denotes the image of  $\tilde{\delta} : \tilde{I} \rightarrow \tilde{I}'_2$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$ , then  $\text{Hom}_B(\Sigma, u_2)$  equals the composition  $\theta(I) \xrightarrow{\theta(\delta)} \theta(I'_2) \xrightarrow{\text{Hom}_B(\Sigma, \nu)} \text{Hom}_B(\Sigma, R')$ . This is an equality of morphisms in  $\text{mod } \Sigma$ , hence, of morphisms between contravariant functors from  $\text{add } \Sigma$  to  $\text{mod } k$ . Applying this equality to  $E$  yields that  $u_2$  equals the composition  $E \xrightarrow{\text{Hom}_{kQ}(T, \delta)} \text{Hom}_{kQ}(T, I'_2) \xrightarrow{\nu} R'$ . On the other hand, the morphism  $\text{Hom}_{kQ}(T, I'_2) \xrightarrow{\nu} R' = \text{Ext}_{kQ}^1(T, \text{Ker } f')$  is the connecting morphism of the sequence resulting from the application of  $\text{Hom}_{kQ}(T, -)$  to the exact sequence  $0 \rightarrow \text{Ker } f' \rightarrow I'_1 \xrightarrow{f'} I'_2 \rightarrow 0$ . Therefore 5.2 implies that  $\nu$  equals the image under  $F_\lambda$  of the connecting morphism of the sequence resulting from the application of  $\text{Hom}_{k\tilde{Q}}(\tilde{T}, -)$  to the exact sequence  $0 \rightarrow \text{Ker } \tilde{f}' \rightarrow \tilde{I}'_1 \rightarrow \tilde{I}'_2 \rightarrow 0$ . Consequently,  $u_2$  equals the image under  $F_\lambda$  of the composition  $\varphi_0(\tilde{E}) \xrightarrow{\text{Hom}_{k\tilde{Q}}(\tilde{T}, \delta)} \varphi_0(\text{Hom}_{kQ}(\tilde{T}, \tilde{I}'_2)) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$ .

It remains to prove that  $u_3 : R \rightarrow R'$  equals the image under  $F_\lambda$  of a morphism  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$  in  $\text{mod } \tilde{B}$ . Using the projective presentations of  $\tilde{R}$  and  $\tilde{R}'$ , we find morphisms  $\tilde{\alpha} : \tilde{I}_2 \rightarrow \tilde{I}'_2$  and  $\tilde{\beta} : \tilde{I}_1 \rightarrow \tilde{I}'_1$  such that the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \tilde{\theta}(\tilde{I}_1) & \xrightarrow{\tilde{\theta}(\tilde{f})} & \tilde{\theta}(\tilde{I}_2) & \longrightarrow & \tilde{R} \longrightarrow 0 \\ & & \tilde{\theta}(\tilde{\beta}) \downarrow & & \tilde{\theta}(\tilde{\alpha}) \downarrow & & \tilde{u}_3 \downarrow \\ 0 & \longrightarrow & \tilde{\theta}(\tilde{I}'_1) & \xrightarrow{\tilde{\theta}(\tilde{f}')} & \tilde{\theta}(\tilde{I}'_2) & \longrightarrow & \tilde{R}' \longrightarrow 0. \end{array}$$

Therefore there exists a morphism  $\tilde{\gamma} : \text{Ker } \tilde{f} \rightarrow \text{Ker } \tilde{f}'$  making the following diagram in  $\text{mod } k\tilde{Q}$  commute:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker } \tilde{f} & \longrightarrow & \tilde{I}_1 & \xrightarrow{\tilde{f}} & \tilde{I}_2 \longrightarrow 0 \\ & & \tilde{\gamma} \downarrow & & \tilde{\beta} \downarrow & & \tilde{\alpha} \downarrow \\ 0 & \longrightarrow & \text{Ker } \tilde{f}' & \longrightarrow & \tilde{I}'_1 & \xrightarrow{\tilde{f}'} & \tilde{I}'_2 \longrightarrow 0. \end{array}$$

We claim that the image of  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \tilde{\gamma}) : \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$  under  $F_\lambda$  equals  $u_3$ . Indeed, let  $\alpha, \beta, \gamma$  be the respective images of  $\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}$  under  $q_\lambda : \text{mod } k\tilde{Q} \rightarrow \text{mod } kQ$ . Then the image of  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \tilde{\gamma})$  under  $F_\lambda$  is equal to (see 5.2):

$$\text{Ext}_{kQ}^1(T, \gamma) : \text{Ext}_{kQ}^1(T, \text{Ker } f) \rightarrow \text{Ext}_{kQ}^1(T, \text{Ker } f').$$

On the other hand, we have two commutative diagrams in  $\text{mod } kQ$  and  $\text{mod } \tilde{B}$  respectively:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Ker } f = \text{Tor}_1^B(R, T) & \longrightarrow & I_1 & \longrightarrow & I_2 \longrightarrow 0 \\ & & \gamma \downarrow & & \beta \downarrow & & \alpha \downarrow \\ 0 & \longrightarrow & \text{Ker } f' = \text{Tor}_1^B(R', T) & \longrightarrow & I'_1 & \longrightarrow & I'_2 \longrightarrow 0, \end{array}$$

and

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \text{Hom}_{kQ}(T, I_1) & \longrightarrow & \text{Hom}_{kQ}(T, I_2) & \longrightarrow & R \longrightarrow 0 \\
 & & \text{Hom}_{kQ}(\tilde{T}, \beta) \downarrow & & \text{Hom}_{kQ}(\tilde{T}, \alpha) \downarrow & & u_3 \downarrow \\
 0 & \longrightarrow & \text{Hom}_{kQ}(T, I'_1) & \longrightarrow & \text{Hom}_{kQ}(T, I'_2) & \longrightarrow & R' \longrightarrow 0,
 \end{array}$$

from which it is straightforward to check that  $u_3 : R \rightarrow R'$  coincides with  $\text{Ext}_{kQ}^1(T, \gamma)$ . Thus,  $u_3$  is equal to the image under  $F_\lambda$  of the morphism  $\text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \tilde{\gamma}) : \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}) \rightarrow \text{Ext}_{k\tilde{Q}}^1(\tilde{T}, \text{Ker } \tilde{f}')$ . This completes the proof.  $\square$

We summarise our results in the following theorem.

**Theorem 5.12.** *Let  $A$  be a finite-dimensional  $k$ -algebra and  $\Gamma$  be a component of  $\Gamma(\text{mod } A)$  containing a left section  $\Sigma$ . Let  $B = A/\text{Ann } \Sigma$  and  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be a Galois covering with group  $G$  of translation quivers such that there exists a well-behaved functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . Then there exists a covering  $F : \tilde{B} \rightarrow B$  with  $\tilde{B}$  locally bounded and a functor  $\varphi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{B}$  which is  $G$ -equivariant on vertices and makes the following diagram commute:*

$$\begin{array}{ccc}
 k(\tilde{\Gamma}) & \xrightarrow{\varphi} & \text{mod } \tilde{B} \\
 p \downarrow & & \downarrow F_\lambda \\
 \text{ind } \Gamma & \xrightarrow{\text{Hom}_A(B, -)} & \text{mod } B.
 \end{array}$$

**Proof.** The functor  $\varphi$  is constructed as above. The  $G$ -equivariance on vertices follows from 5.10 and 5.11.  $\square$

**Corollary 5.13.** *If  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  is a Galois covering (with respect to the action of  $G$  on  $k(\tilde{\Gamma})$ ), then the functor  $\varphi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{B}$  of 5.12 is  $G$ -equivariant.*

**Proof.** We already know that  $\varphi$  is  $G$ -equivariant on objects. Also  $F : \tilde{B} \rightarrow B$  is a Galois covering with group  $G$  (see 5.1). Let  $f : M \rightarrow N$  be a morphism in  $k(\tilde{\Gamma})$ , and  $g \in G$ . Then  $\varphi({}^g f) : \varphi({}^g M) \rightarrow \varphi({}^g N)$  and  ${}^g \varphi(f) : {}^g \varphi(M) \rightarrow {}^g \varphi(N)$  are two morphisms in  $\text{mod } \tilde{B}$  such that  $F_\lambda(\varphi({}^g f)) = p({}^g f)|_B = p(f)|_B = F_\lambda({}^g \varphi(f))$  (recall that  $F_\lambda = F_\lambda \circ g$  for every  $g \in G$  because it is the push-down functor of a Galois covering with group  $G$ ). We deduce that  $\varphi({}^g f) = {}^g \varphi(f)$ .  $\square$

**6. The main theorem**

In this section we prove Theorem A. Assume that  $A$  is lura with connecting components. We use the following notation:

- $\Gamma$  is the connecting component of  $\Gamma(\text{mod } A)$  (if  $A$  is concealed we choose  $\Gamma$  to be the unique postprojective component), and  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  is a Galois covering with group  $G$  of translation quivers such that there exists a well-behaved covering functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . If  $\Gamma$  is standard, we assume that  $p$  equals the composition of  $k(\pi) : k(\tilde{\Gamma}) \rightarrow k(\Gamma)$  with some isomorphism  $k(\Gamma) \xrightarrow{\sim} \text{ind } \Gamma$ , so that  $p$  is a Galois covering with group  $G$ .
- $\Sigma$  is the full subcategory of  $\text{ind } \Gamma$  whose objects are the Ext-injective objects in  $\mathcal{L}_A$ .
- $B$  is the left support of  $A$ , that is,  $B$  is the endomorphism algebra of the direct sum of the indecomposable projective modules lying on  $\mathcal{L}_A$  (see Section 1).

Because of [6, 4.4, 5.1], the algebra  $B$  is a product of tilted algebras. Without loss of generality, we assume that:

- $B = \text{End}_{kQ}(T)$ , where  $T = T_1 \oplus \dots \oplus T_n$  is a multiplicity-free tilting  $kQ$ -module ( $T_i \in \text{ind}kQ$ ).
- $\Sigma$  is the full subcategory of  $\text{mod} B$  with objects the modules of the form  $\text{Hom}_{kQ}(T, D(kQe_x))$ ,  $x \in Q_0$ .

It follows from [1, 2.1 Ex. b] that  $\Sigma$  is a left section of  $\Gamma$ . So we may apply 5.12. The proof of Theorem A is done in the following steps: We first construct a locally bounded  $k$ -category  $\tilde{A}$  endowed with a free  $G$ -action in case  $A$  is standard; then we construct a covering functor  $F : \tilde{A} \rightarrow A$  extending the functor  $F : \tilde{B} \rightarrow B$  of 5.12 and satisfying the conditions of the theorem; we also construct a functor  $\Phi : k(\tilde{\Gamma}) \rightarrow \text{mod} \tilde{A}$  which extends the functor  $\varphi : k(\tilde{\Gamma}) \rightarrow \text{mod} \tilde{B}$  of 5.12; and finally we prove Theorem A.

The category  $\tilde{A}$

We need some notation. Let  $C$  be the full subcategory of  $\text{ind} A$  with objects the indecomposable projective  $A$ -modules not in  $\mathcal{L}_A$ . So  $C$  is a full subcategory of  $\text{ind} \Gamma$ . Let  $\tilde{C}$  be the full subcategory  $p^{-1}(C)$ , so that  $p$  induces a covering functor  $\tilde{C} \rightarrow C$ . If  $A$  is standard and  $p$  is Galois with group  $G$ , then  $p : \tilde{C} \rightarrow C$  is a Galois covering with group  $G$ . For every  $x \in \tilde{B}_0$ , let  $\tilde{P}_x$  be the corresponding indecomposable projective  $\tilde{B}$ -module. Also,  $P_x \in \text{mod} B$  denotes the indecomposable projective  $B$ -module associated to an object  $x \in B_0$ . We define the  $\tilde{C} - \tilde{B}$ -bimodule  $\tilde{M}$  to be the functor  $\tilde{C} \times \tilde{B}^{op} \rightarrow \text{mod} k$  such that for every  $\tilde{P} \in \tilde{C}_0$  and  $x \in \tilde{B}_0$

$$\tilde{P} \tilde{M}_x = \text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(\tilde{P})),$$

with obvious actions of  $\tilde{C}$  (using  $\varphi$ ) and  $\tilde{B}$ .

The following lemma defines  $\tilde{A}$  and its  $G$ -action in case  $A$  is standard.

**Lemma 6.1.** *Let  $\tilde{A} = \begin{bmatrix} \tilde{B} & 0 \\ \tilde{M} & \tilde{C} \end{bmatrix}$ . Then  $\tilde{A}$  is locally bounded and  $G$  acts freely on  $\tilde{A}$  if  $A$  is standard.*

**Proof.** We know that  $\tilde{B}$  and  $\tilde{C}$  are locally bounded. Let  $P \in \tilde{C}_0$ . We have the bijection of 4.2:

$$\bigoplus_{\tilde{x} \in \tilde{B}_0} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(\tilde{P})) \xrightarrow{\sim} \bigoplus_{x \in B_0} \text{Hom}_B(P_x, p(\tilde{P})|_B). \tag{i}$$

Since the right-hand side is finite-dimensional, then so is  $\bigoplus_{\tilde{x} \in \tilde{B}_0} \tilde{P} \tilde{M}_{\tilde{x}}$ , for every  $\tilde{P} \in \tilde{C}_0$ .

Now let  $P \in \tilde{C}_0$ , let  $\tilde{x} \in \tilde{B}_0$ , and let us prove that  $\bigoplus_{p(P')=P} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(P'))$  is finite-dimensional. By definition of  $p$ , we have  $p^{-1}(p(P)) = \{gP \mid g \in G\}$ . Also, we know from 5.12 that  $\varphi$  is  $G$ -equivariant on objects. Therefore:

$$\bigoplus_{p(P')=P} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(P')) = \bigoplus_{g \in G} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, {}^g \varphi(P)) = \bigoplus_{g \in G} \text{Hom}_{\tilde{B}}(g^{-1} \tilde{P}_{\tilde{x}}, \varphi(P)), \tag{ii}$$

where the last equality follows from the  $G$ -action on  $\text{mod} \tilde{B}$ . Applying 4.2 to the indecomposable projective  $\tilde{P}_{\tilde{x}}$  yields a bijection of vector spaces:

$$\bigoplus_{g \in G} \text{Hom}_{\tilde{B}}(g^{-1} \tilde{P}_{\tilde{x}}, {}^g \varphi(P)) \simeq \text{Hom}_B(P_{F(\tilde{x})}, F_\lambda \varphi(P)). \tag{iii}$$

From (ii) and (iii) we infer that  $\bigoplus_{p(P')=p(P)} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(P'))$  is finite-dimensional for every  $\tilde{x} \in \tilde{B}_0$  and  $P \in \tilde{C}_0$ . This shows that  $\tilde{A}$  is locally bounded.

Assume now that  $A$  is standard and that  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  is a Galois covering with group  $G$ . We define a free  $G$ -action on  $\tilde{A}$ . We already have a free  $G$ -action on  $\tilde{B}$  and on  $\tilde{C}$ . Also, for every  $\tilde{x} \in \tilde{B}_0$ ,  $\tilde{P} \in \tilde{C}_0$  and  $g \in G$ , we have an isomorphism of vector spaces:

$$\tilde{p}\tilde{M}_{\tilde{x}} = \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(P)) \xrightarrow{\sim} {}_g\tilde{p}\tilde{M}_{g\tilde{x}} = \text{Hom}_{\tilde{B}}({}^g\tilde{P}_{\tilde{x}}, \varphi({}^g\tilde{P})) \tag{*}$$

given by the  $G$ -action on  $\text{mod } \tilde{B}$  (recall that  $\varphi$  is  $G$ -equivariant on objects, and that  $\tilde{P}_{g\tilde{x}} = {}^g\tilde{P}_{\tilde{x}}$ ). We define the action of  $g$  on morphisms of  $\tilde{A}$  lying in  $\tilde{M}$  using this isomorphism. Since  $G$  acts on  $\text{mod } \tilde{B}$ , this defines a  $G$ -action on  $\tilde{A}$ , that is,  $g(vu) = g(v)g(u)$  whenever  $u$  and  $v$  are composable in  $\tilde{A}$ . Moreover,  $G$  acts freely on objects in  $\tilde{B}$  and in  $\tilde{C}$ . So we have a free  $G$ -action on  $\tilde{A}$ .  $\square$

The functor  $F : \tilde{A} \rightarrow A$

**Lemma 6.2.** *There exists a covering functor  $F : \tilde{A} \rightarrow A$  extending  $F : \tilde{B} \rightarrow B$ . If moreover  $A$  is standard, then  $F$  can be taken to be Galois with group  $G$ .*

**Proof.** Note that  $A = \begin{bmatrix} B & 0 \\ M & C \end{bmatrix}$  where  $M$  is the  $C$ – $B$ -bimodule such that  ${}_pM_x = \text{Hom}_B(P_x, P|_B)$  for every  $P \in C_0$  and  $x \in B_0$ . Let us define  $F : \tilde{A} \rightarrow A$  as follows:

- $F|_{\tilde{B}}$  coincides with the functor  $F : \tilde{B} \rightarrow B$ .
- $F|_{\tilde{C}}$  coincides with  $p : \tilde{C} \rightarrow C$ .
- Let  $x \in \tilde{B}_0$  and  $\tilde{P} \in \tilde{C}_0$ , then  $F : \tilde{p}\tilde{M}_x \rightarrow {}_{F(\tilde{P})}M_{F(x)}$  is the following map induced by  $F_\lambda$ :

$$\text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(\tilde{P})) \rightarrow \text{Hom}_B(P_{F(x)}, p(\tilde{P})|_B).$$

Since  $F_\lambda : \text{mod } \tilde{B} \rightarrow \text{mod } B$  is a functor and  $F_\lambda\varphi = p(-)|_B$  (see 5.12), we have defined a functor  $F : \tilde{A} \rightarrow A$ . We prove that  $F : \tilde{A} \rightarrow A$  is a covering functor. Since  $F : \tilde{B} \rightarrow B$  and  $p : \tilde{C} \rightarrow C$  are covering functors, the bijections (i), (ii) and (iii) in the proof of 6.1 show that for any  $\tilde{a} \in \tilde{B}_0$  and any  $\tilde{P} \in \tilde{C}_0$ , the two following maps induced by  $F_\lambda$  are isomorphisms:

$$\bigoplus_{F(\tilde{x})=F(\tilde{a})} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{x}}, \varphi(\tilde{P})) \rightarrow \text{Hom}_B(P_{F(\tilde{a})}, p(\tilde{P})|_B),$$

$$\bigoplus_{p(\tilde{Q})=p(\tilde{P})} \text{Hom}_{\tilde{B}}(\tilde{P}_{\tilde{a}}, \varphi(\tilde{Q})) \rightarrow \text{Hom}_B(P_{F(\tilde{a})}, p(\tilde{P})|_B).$$

So  $F$  is a covering functor. Assume now that  $A$  is standard. We may suppose that  $p$  is a Galois covering with group  $G$ . By 6.1, there is a free  $G$ -action on  $\tilde{A}$ . Moreover,  $F : \tilde{B} \rightarrow B$ , and therefore  $F_\lambda : \text{mod } \tilde{B} \rightarrow \text{mod } B$ , are  $G$ -equivariant, and so is  $p : \tilde{C} \rightarrow C$ , because it restricts the Galois covering  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . Therefore  $F : \tilde{A} \rightarrow A$  is  $G$ -equivariant. Finally, the fibres of  $F : \tilde{A} \rightarrow A$  on objects are the  $G$ -orbits in  $\tilde{A}_0$  because  $F : \tilde{B} \rightarrow B$  and  $p : \tilde{C} \rightarrow C$  are Galois coverings. Since  $F : \tilde{A} \rightarrow A$  is a covering functor, this implies that it is also a Galois covering with group  $G$  (see for instance the proof of [28, Prop. 6.1.37]).  $\square$

The functor  $\Phi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{A}$

We can write an  $\tilde{A}$ -module as a triple  $(K, L, f)$  where  $K \in \text{mod } \tilde{B}$ ,  $L \in \text{mod } \tilde{C}$  and  $f : L \otimes_{\tilde{C}} \tilde{M} \rightarrow K$  is a morphism of  $\tilde{B}$ -modules. Let  $\psi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{C}$  be the functor  $\psi : X \mapsto k(\tilde{\Gamma})(\tilde{C}, X)$ . Clearly, it is  $G$ -equivariant. Let  $L \in k(\tilde{\Gamma})$ . Then  $\psi(L) \otimes_{\tilde{C}} \tilde{M}$  is the  $\tilde{B}$ -module whose value at  $x \in \tilde{B}_0$  equals:

$$(\psi(L) \otimes_{\tilde{C}} \tilde{M})(x) = \left( \bigoplus_{\tilde{P} \in \tilde{C}_0} k(\tilde{\Gamma})(\tilde{P}, L) \otimes_k \text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(\tilde{P})) \right) / N,$$

where  $N$  is the following subspace:

$$N = \{ ff' \otimes u - f \otimes \varphi(f')u \mid f \in k(\tilde{\Gamma})(\tilde{P}, L), f' \in k(\tilde{\Gamma})(\tilde{P}', \tilde{P}), \\ u \in \text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(\tilde{P}')), \text{ for every } \tilde{P}, \tilde{P}' \in \tilde{C}_0 \}.$$

For every  $x \in \tilde{B}_0$  and  $\tilde{P} \in \tilde{C}_0$ , we have a  $k$ -linear map:

$$\eta_{L,x,P} : k(\tilde{\Gamma})(\tilde{P}, L) \otimes_k \text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(\tilde{P})) \rightarrow \text{Hom}_{\tilde{B}}(\tilde{P}_x, \varphi(L)) = \varphi(L)(x), \\ f \otimes u \mapsto \varphi(f)u.$$

It is not difficult to check that the family of maps  $(\eta_{L,x,\tilde{P}})_{L,x,\tilde{P}}$  defines a functorial morphism:

$$\eta : \psi(-) \otimes_{\tilde{C}} \tilde{M} \rightarrow \varphi.$$

Moreover, if  $\varphi$  is  $G$ -equivariant, then so is  $\eta$ . We let  $\Phi : k(\tilde{\Gamma}) \rightarrow \text{mod } \tilde{A}$  be the following functor:

$$\Phi : L \mapsto (\varphi(L), \psi(L), \eta_L).$$

*The main theorem*

**Theorem 6.3.** *Let  $A$  be a  $\lambda$ -algebra with connecting component  $\Gamma$ . Let  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be a Galois covering with group  $G$  such that there exists a well-behaved covering functor  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . Then there exist a covering functor  $F : \tilde{A} \rightarrow A$  where  $\tilde{A}$  is connected and locally bounded, and a commutative diagram:*

$$\begin{array}{ccc} k(\tilde{\Gamma}) & \xrightarrow{\Phi} & \text{mod } \tilde{A} \\ p \downarrow & & \downarrow F_\lambda \\ \text{ind } \Gamma & \hookrightarrow & \text{mod } A, \end{array}$$

where  $\Phi$  is faithful. If, moreover,  $A$  is standard, then  $F$  and  $p$  may be assumed to be Galois coverings with group  $G$ , and  $\Phi$  is then  $G$ -equivariant and full.

**Proof.** The commutativity of the above diagram follows from the one of 5.12 and from that of the diagram:

$$\begin{array}{ccc} k(\tilde{\Gamma}) & \xrightarrow{\psi} & \text{mod } \tilde{C} \\ p \downarrow & & \downarrow p_\lambda \\ \text{ind } \Gamma & \xrightarrow{X \mapsto \text{Hom}_A(C, X)} & \text{mod } C. \end{array}$$

Since  $F_\lambda \Phi = p$  and  $p$  is faithful, then  $\Phi$  is faithful. Therefore  $\Phi(k(\tilde{\Gamma}))$  is contained in a connected component  $\Omega$  of  $\text{mod } \tilde{A}$ .

We now prove that  $\tilde{A}$  is connected. Let  $x \in \tilde{A}_0$  and  $Q_x$  be the corresponding indecomposable projective  $\tilde{A}$ -module. If  $\tilde{F}_\lambda Q_x \in C_0$ , then, by construction,  $Q_x$  lies in the image of  $\Phi$ , so that  $Q_x \in \Omega$ . If  $\tilde{F}_\lambda Q_x \notin C_0$ , then  $F(x) \in B_0$  and  $x \in \tilde{B}_0$ . In this case, there is a non-zero morphism  $u : P_{F(x)} = \tilde{F}_\lambda Q_x \rightarrow E$  in  $\text{mod } B$ , where  $E \in \Sigma$ . Fix  $\tilde{E} \in p^{-1}(E)$  so that  $F_\lambda \Phi(\tilde{E}) = E$ . Since  $u$  is non-zero, 4.2 implies that there is a non-zero morphism  $Q_x \rightarrow {}^g \varphi(\tilde{E}) = \Phi({}^g \tilde{E})$  in  $\text{mod } \tilde{B}$  (recall that  $\varphi$  is  $G$ -equivariant on vertices). So  $Q_x \in \Omega$ , and  $\Omega$  contains all the indecomposable projective  $\tilde{A}$ -modules. This proves that  $\tilde{A}$  is connected.


It remains to prove that if  $A$  is standard, then  $\Phi$  is full,  $G$ -equivariant, and  $F$  is Galois with group  $G$ . In case  $A$  is standard, we suppose that  $p : k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$  is Galois with group  $G$ . Therefore  $\varphi$  is  $G$ -equivariant (see 5.13) and so is  $\eta$ . Hence,  $\Phi$  is  $G$ -equivariant. Also,  $F$  is Galois because of 6.2. We prove that  $\Phi$  is full. Given a morphism  $f : \Phi(L) \rightarrow \Phi(N)$ , there exists  $(f_g)_g \in \bigoplus_{g \in G} \text{Hom}_{k(\Gamma)}(L, {}^g N)$  such that  $F_\lambda(f) = \sum_g p(f_g)$  (because  $p$  is Galois). So  $F_\lambda(f - \Phi(f_1)) = \sum_{g \neq 1} F_\lambda(\Phi(f_g)) = 0$ . Since  $F$  is Galois with group  $G$  and since  $\Phi$  is  $G$ -equivariant, we get  $f = \Phi(f_1)$ . So  $\Phi$  is full and the theorem is proved.  $\square$

The following example of a non-standard representation-finite algebra due to Riedtmann shows that  $F$  needs not be a Galois covering.

**Example 6.4.** Assume that  $\text{char}(k) = 2$  and  $A$  is given by the bound quiver (see [14, §7, Ex. 14 bis] and [37]):

$$x \begin{matrix} \xleftarrow{\sigma} \\ \xrightarrow{\delta} \end{matrix} y \begin{matrix} \curvearrowright \rho \\ \curvearrowleft \rho \end{matrix}, \quad \rho^4 = 0, \quad \rho^2 = \delta\sigma, \quad \sigma\delta = \sigma\rho\delta.$$

Then  $A$  is representation-finite and not standard, with the following Auslander–Reiten quiver:



where the two copies of  $a, b, c, d, e$  and  $f$ , respectively, are identified. In this case, there exists a well-behaved covering functor associated to the universal cover  $\tilde{\Gamma}$  of  $\Gamma(\text{mod } A)$  (which is equal the generic covering). Here,  $G = \pi_1(\Gamma) \simeq \mathbb{Z}$  and  $\tilde{A}$  is the locally bounded  $k$ -category, given by the following bound quiver:

$$\begin{array}{ccccccc}
 \cdots & & y_{i-1} & \longrightarrow & y_i & \longrightarrow & y_{i+1} & \longrightarrow & y_{i+2} & \cdots \\
 & & \searrow & & \nearrow & & \searrow & & \nearrow & \\
 \cdots & & x_{i-1} & & x_i & & x_{i+1} & & x_{i+2} & \cdots \\
 & & \nearrow & & \searrow & & \nearrow & & \searrow & \\
 & & \delta_{i+1}\sigma_i = \rho_{i+1}\rho_i, & & \sigma_{i+1}\delta_i = 0, & & \text{for all } i, & & & 
 \end{array}$$

where  $\sigma_i$ ,  $\delta_i$  and  $\rho_i$  denote the arrows  $y_i \rightarrow x_{i+1}$ ,  $x_i \rightarrow y_{i+1}$ , and  $y_i \rightarrow y_{i+1}$ , respectively. Now the covering functor  $F : \tilde{A} \rightarrow A$  is as follows:

1.  $F(\rho_i) = \rho$  for every  $i$ ,
2.  $F(\sigma_i) = \sigma$  for every  $i \equiv 0, 1 \pmod{4}$ ,
3.  $F(\sigma_i) = \sigma + \sigma\rho$  for every  $i \equiv 2, 3 \pmod{4}$ ,
4.  $F(\delta_i) = \delta$  for every  $i \equiv 1, 3 \pmod{4}$ ,
5.  $F(\delta_i) = \delta + \rho\delta$ , for every  $i \equiv 0, 2 \pmod{4}$ .

Obviously,  $F$  is a covering functor which is not Galois. Actually, one can easily check that  $A$  is simply connected, that is, the fundamental group (in the sense of [33]) of any presentation of  $A$  is trivial. Hence,  $A$  has no proper Galois covering by a locally bounded and connected  $k$ -category.

The following corollary is a particular case of our main theorem. We state it for later purposes.

**Corollary 6.5.** *Let  $A$  be a standard lura algebra and let  $\Gamma$  be a connecting component. There exists a Galois covering  $F : \tilde{A} \rightarrow A$  with group  $\pi_1(\Gamma)$  where  $\tilde{A}$  is connected and locally bounded, together with a commutative diagram:*

$$\begin{array}{ccc}
 k(\tilde{\Gamma}) & \xrightarrow{\Phi} & \text{mod } \tilde{A} \\
 k(\pi) \downarrow & & \downarrow F_\lambda \\
 k(\Gamma) & \hookrightarrow & \text{mod } A,
 \end{array}$$

where  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  is the universal cover and where  $\Phi$  is full, faithful and  $\pi_1(\Gamma)$ -equivariant.

**Proof.** Since  $\Gamma$  is a standard component, there exists an isomorphism of categories  $k(\Gamma) \rightarrow \text{ind } \Gamma$  and the universal cover  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  induces a well-behaved functor  $k(\pi) : k(\tilde{\Gamma}) \rightarrow k(\Gamma)$  and therefore a well-behaved covering functor  $k(\tilde{\Gamma}) \rightarrow \text{ind } \Gamma$ . We then apply 6.3.  $\square$

We pose the following problems.

**Problem 3.** Does there exist a combinatorial characterisation of standardness for lura algebras (as happens for representation-finite algebras, see [13])?

**Problem 4.** Let  $A$  be a left supported algebra. Is it possible to construct coverings  $\tilde{A} \rightarrow A$  associated to the coverings of a component of  $\Gamma(\text{mod } A)$  containing the Ext-injective modules of  $\mathcal{L}_A$ ?

### 7. Galois coverings of the connecting component

**Theorem 7.1.** *Let  $A$  be a standard lura algebra, and  $p : \Gamma' \rightarrow \Gamma$  be a Galois covering with group  $G$  of a connecting component. Then there exist a Galois covering  $F' : A' \rightarrow A$  with group  $G$ , where  $A'$  is connected and locally bounded, and a commutative diagram:*

$$\begin{array}{ccc}
 k(\Gamma') & \xrightarrow{\Phi'} & \text{mod } A' \\
 k(p) \downarrow & & \downarrow F'_\lambda \\
 k(\Gamma) & \xrightarrow{j} & \text{mod } A,
 \end{array}$$

where  $\Phi'$  is full, faithful and  $G$ -equivariant.

**Proof.** Since  $A$  is standard, there exists a full and faithful functor  $j : k(\Gamma) \hookrightarrow \text{ind } A$  with image  $\text{ind } \Gamma$ , which maps meshes to almost split sequences. Let  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  be the universal cover. Then there exists a normal subgroup  $H \triangleleft \pi_1(\Gamma)$  such that  $\tilde{\Gamma}/H \simeq \Gamma'$  and  $G \simeq \pi_1(\Gamma)/H$ , and such that under these identifications, the following diagram commutes:

$$\begin{array}{ccc}
 \tilde{\Gamma} & & \\
 \pi \downarrow & \searrow q & \\
 & & \tilde{\Gamma}/H = \Gamma', \\
 & \swarrow p & \\
 \Gamma & & 
 \end{array}$$


where  $q$  is the projection. These identifications imply that  $p : \Gamma' \rightarrow \Gamma$  is induced by  $\pi : \tilde{\Gamma} \rightarrow \Gamma$  by factoring out by  $H$ . By 6.5, there exist a Galois covering  $F : \tilde{A} \rightarrow A$  with group  $\pi_1(\Gamma)$  and a commutative diagram:

$$\begin{array}{ccc}
 k(\tilde{\Gamma}) & \xrightarrow{\Phi} & \text{mod } \tilde{A} \\
 k(\pi) \downarrow & & \downarrow F_\lambda \\
 k(\Gamma) & \xrightarrow{j} & \text{mod } A,
 \end{array}$$

where  $\Phi$  is full, faithful and  $\pi_1(\Gamma)$ -equivariant. Setting  $A' = \tilde{A}/H$ , we deduce a Galois covering  $F' : A' \rightarrow A$  with group  $G$  and where  $A'$  is connected and locally bounded, making the following diagram commute:

$$\begin{array}{ccc}
 \tilde{A} & & \\
 F \downarrow & \searrow F'' & \\
 & & \tilde{A}/H = A', \\
 & \swarrow F' & \\
 A & & 
 \end{array}$$

where  $F''$  is the natural projection (and  $F'$  is deduced from  $F$  by factoring out by  $H$ ). Therefore we have a commutative diagram of solid arrows:



We prove the existence of the dotted arrow  $\Phi'$  such that  $\Phi'k(q) = F''_\lambda\Phi$ . For this purpose, recall that  $k(q)$  is a Galois covering with group  $H$ . Hence, it suffices to prove that  $F''_\lambda\Phi$  is  $H$ -invariant. Indeed, we have  $F''_\lambda\Phi'h = F''_\lambda h\Phi' = F''_\lambda\Phi'$ , for every  $h \in H$ , because  $\Phi$  is  $\pi_1(\Gamma)$ -equivariant and  $F''$  is a Galois covering with group  $H$ . Now, we prove that the whole diagram commutes. We have:

$$(F'_\lambda\Phi')k(q) = F'_\lambda F''_\lambda\Phi = F_\lambda\Phi = jk(\pi) = jk(p)k(q),$$

hence,  $F'_\lambda\Phi' = jk(p)$ . We prove next that  $\Phi'$  is full and faithful. Let  $f : X \rightarrow Y$  be a morphism in  $k(\Gamma')$  such that  $\Phi'(f) = 0$ . Fix  $\tilde{X}, \tilde{Y} \in k(\tilde{\Gamma})$  such that  $q(\tilde{X}) = X$  and  $q(\tilde{Y}) = Y$ . Since  $k(q)$  is Galois with group  $H$ , there exists  $(f_h)_{h \in H} \in \bigoplus_{h \in H} k(\tilde{\Gamma})(\tilde{X}, {}^h\tilde{Y})$  such that  $\sum_{h \in H} k(q)(f_h) = f$ . The commutativity of the diagram gives:

$$0 = \sum_{h \in H} F''_\lambda(\Phi(f_h)),$$

where  $(\Phi(f_h))_{h \in H} \in \bigoplus_{h \in H} \text{Hom}_{\tilde{A}}(\Phi(\tilde{X}), {}^h\Phi(\tilde{Y}))$  (recall that  $\Phi$  is  $\pi_1(\Gamma)$ -equivariant). Since  $F'' : \tilde{A} \rightarrow A'$  is Galois with group  $H$ , we deduce that  $\Phi(f_h) = 0$  for every  $h \in H$ , so that  $f_h = 0$  for every  $h \in H$ , because  $\Phi$  is faithful. Thus,  $f = \sum_{h \in H} k(q)(f_h) = 0$  and  $\Phi'$  is faithful. Let  $X, Y \in k(\Gamma')$  and  $u : \Phi'(X) \rightarrow \Phi'(Y)$  be a morphism in  $\text{mod } A'$ , and fix  $\tilde{X}, \tilde{Y} \in k(\tilde{\Gamma})$  as above. In particular,  $\Phi'(X) = F''_\lambda(\Phi(\tilde{X}))$  and  $\Phi'(Y) = F''_\lambda(\Phi(\tilde{Y}))$ . Therefore there exists  $(\tilde{u}_h)_{h \in H} \in \bigoplus_{h \in H} \text{Hom}_{\tilde{A}}(\Phi(\tilde{X}), {}^h\Phi(\tilde{Y}))$  such that  $u = \sum_{h \in H} F''_\lambda(\tilde{u}_h)$ . Since  $\Phi'$  is  $\pi_1(\Gamma)$ -equivariant, we have  $\text{Hom}_{\tilde{A}}(\Phi(\tilde{X}), {}^h\Phi(\tilde{Y})) = \text{Hom}_{\tilde{A}}(\Phi(\tilde{X}), \Phi({}^h\tilde{Y}))$ , for every  $h \in H$ . Since  $\Phi$  is full, there exists  $(\tilde{f}_h)_{h \in H} \in \bigoplus_{h \in H} k(\tilde{\Gamma})(\tilde{X}, {}^h\tilde{Y})$  such that  $\tilde{u}_h = \Phi(\tilde{f}_h)$  for every  $h \in H$ . Since  $k(q)$  is Galois with group  $H$ , we deduce that  $\sum_{h \in H} k(q)(\tilde{f}_h) \in k(\Gamma)(X, Y)$ . Moreover, we have:

$$\Phi' \left( \sum_{h \in H} k(q)(\tilde{f}_h) \right) = \sum_{h \in H} F''_\lambda\Phi(\tilde{f}_h) = \sum_{h \in H} F''_\lambda\tilde{u}_h = u,$$

whence the fullness of  $\Phi'$ . To finish, it remains to prove that  $\Phi'$  is  $G$ -equivariant. Let  $g \in G$  be the residual class of  $\sigma \in \pi_1(\Gamma)$  modulo  $H$ . We need to prove that  $\Phi' \circ g = g \circ \Phi'$ . We have  $\Phi' \circ g \circ k(q) = \Phi' \circ k(q) \circ \sigma$ , because  $q : \tilde{\Gamma} \rightarrow \Gamma' = \tilde{\Gamma}/H$  is the canonical projection. Hence,  $\Phi' \circ g \circ k(q) = F''_\lambda \circ \sigma \circ \Phi$ , because  $F''_\lambda \circ \Phi = \Phi' \circ k(q)$ , and  $\Phi$  is  $\pi_1(\Gamma)$ -equivariant. Since  $F''_\lambda \circ \sigma = g \circ F''_\lambda$  (because  $F''$  is deduced from  $F$  by factoring out by  $H$ ), we have  $\Phi' \circ g \circ k(q) = g \circ F''_\lambda \circ \Phi = g \circ \Phi' \circ k(q)$ , and so  $\Phi' \circ g = g \circ \Phi'$ . The proof is complete.  $\square$

**Corollary 7.2.** *In the situation of Theorem 7.1, the full subquiver  $\Omega$  of  $\Gamma(\text{mod } A')$  with vertex set equal to  $\{X \in \text{ind } A' \mid F'_\lambda X \in \Gamma\}$  is a faithful and generalised standard component of  $\Gamma(\text{mod } A')$ , isomorphic, as a translation quiver, to  $\Gamma'$ . Moreover, there exists a Galois covering of translation quivers  $\Gamma' \rightarrow \Gamma$  with group  $G$  extending the map  $X \mapsto F'_\lambda X$ .*

**Proof.** Since  $F'_\lambda \Phi' = jk(p)$ , the module  $\Phi'(X)$  is indecomposable and lies in  $\Omega$ , for every  $X \in \Gamma'$ . On the other hand if  $X \in \Omega$ , there exists  $X' \in \Gamma'$  such that  $F'_\lambda X = k(p)(X')$ . Therefore  $F'_\lambda X = F'_\lambda \Phi'(X')$ . Since  $X$  and  $\Phi'(X')$  are indecomposable, there exists  $g \in G$  such that  $X = {}^g \Phi'(X') = \Phi'({}^g X') \in \Phi'(\Gamma')$ . Thus, we have shown that:

(i)  $\Omega$  coincides with the full subquiver of  $\Gamma(\text{mod } A')$  with set of vertices  $\{\Phi'(X) \mid X \in \Gamma'\}$ .

Let  $X \xrightarrow{u} Y$  be an arrow in  $\Gamma'$ . Since  $F'_\lambda \Phi' = jk(p)$ , then  $F'_\lambda \Phi'(u)$  is an irreducible morphism between indecomposable  $A$ -modules. Using [30, Lem. 2.1], we deduce that  $\Phi'(u)$  is irreducible. This proves that:

(ii) The full subquiver of  $\Gamma(\text{mod } A')$  with set of vertices  $\{\Phi'(X) \mid X \in \Gamma'\}$  is contained in a connected component of  $\Gamma(\text{mod } A')$ .

Combining (i), (ii) and [30, Lem. 2.3], we deduce that  $\Omega$  is a component of  $\Gamma(\text{mod } A')$ . The same lemma shows that  $\Omega$  is faithful and generalised standard because so is  $\Gamma$ .

Let us prove that  $\Phi'$  induces an isomorphism between  $\Gamma'$  and  $\Omega$ . Since  $q : \tilde{\Gamma} \rightarrow \Gamma'$  is surjective on vertices and  $F''_\lambda \Phi = \Phi'k(q)$ , then  $X \in \Omega$  lies in the image of  $F''_\lambda$ . Also,  $k(q)$  and  $\Phi$  commute with the translation, and so does  $F''_\lambda$  (see [30, Lem. 2.1]). Hence  $\Phi'$  commutes with the translation. Finally  $k(q)$  maps meshes to meshes, and  $\Phi$  maps meshes to almost split sequences. So  $\Phi'$  maps meshes to almost split sequences (see [30, Lem. 2.2]). Therefore there exists a morphism of translation quivers  $\Gamma' \rightarrow \Omega$  extending the map  $X \mapsto \Phi'(X)$  on vertices. Since it is a bijection on vertices, it is an isomorphism  $\Gamma' \xrightarrow{\sim} \Omega$ .

Finally, the stabiliser  $G_X = \{g \in G \mid {}^g X \simeq X\}$  of  $X$  is trivial for every  $X \in \Omega$ , because  $G$  acts freely on  $\Gamma'$  and  $\Phi'$  is  $G$ -equivariant. Therefore there exists a Galois covering of translation quivers  $\Omega \rightarrow \Gamma'$  with group  $G$  and extending the map  $X \mapsto F'_\lambda(X)$  (see [21, 3.6]).  $\square$

**Corollary 7.3.** *In the situation of Theorem 7.1, if  $G$  is finite, then  $A'$  is a finite-dimensional standard lura algebra.*

**Proof.** Since  $G$  is finite,  $A'$  is finite-dimensional. By the preceding corollary,  $\Gamma'$  is generalised standard and faithful. Since  $\Gamma$  has only finitely many isomorphism classes of indecomposable modules lying on oriented cycles, the same is true for  $\Gamma'$ . Therefore  $\Gamma'$  is quasi-directed and faithful. Applying [35, 3.1] (or [43, Thm. 2]) shows that  $A'$  is a lura algebra with  $\Gamma'$  as a connecting component. Finally, the full and faithful functor  $\Phi' : k(\Gamma') \rightarrow \text{mod } A'$  with image equal to  $\text{ind } \Gamma'$  shows that  $\Gamma'$  is standard, that is,  $A'$  is standard.  $\square$

**Remark 7.4.** The above corollary may be compared with [8, Thm. 1.2] and [30, Thm. 3]. Indeed, if  $A'$  is a finite-dimensional algebra endowed with the free action of a (necessarily finite) group  $G$ , then the category  $A/G$  and the skew-group algebra  $A[G]$  are Morita equivalent.


We end this section with the following corollary:

**Corollary 7.5.** *In the situation of Theorem 7.1, if  $G$  is finite, then:*


- (a)  $A$  is tame if and only if  $A'$  is tame.
- (b)  $A$  is wild if and only if  $A'$  is wild.

**Proof.** This follows from Theorem 7.1 and from [3, 5.3(b)].  $\square$

**Example 7.6.** Consider the algebra  $A$  of 3.6(a). The connecting component  $\Gamma$  admits a Galois covering with group  $\mathbb{Z}/2\mathbb{Z}$  by the following translation quiver:



where the two copies of  $x$  are identified. With our construction, we get a Galois covering  $F : A' \rightarrow A$  with group  $\mathbb{Z}/2\mathbb{Z}$ , where  $A'$  is the radical square zero algebra with the following quiver:



Both  $A$  and  $A'$  are tame.

**8. Proof of Theorem B**

We recall the definition of the *orbit graph*  $\mathcal{O}(\Gamma)$  (see [14, 4.2]). Given a vertex  $x \in \Gamma$ , its  $\tau$ -orbit  $x^\tau$  is the set  $\{y \in \Gamma \mid y = \tau^l x, \text{ for some } l \in \mathbb{Z}\}$ . Also, we fix a polarisation  $\sigma$  in  $\Gamma$ . The periodic components of  $\Gamma$  are defined as follows. Consider the full translation subquiver of  $\Gamma$  with vertices the periodic vertices in  $\Gamma$ . To this subquiver, add a new arrow  $x \rightarrow \tau x$  for every vertex  $x$ . A *periodic component* of  $\Gamma$  is a connected component of the resulting quiver. Then:

1. The vertices of  $\mathcal{O}(\Gamma)$  are the periodic components of  $\Gamma$  and the  $\tau$ -orbits of the non-periodic vertices.
2. For each periodic component, there is a loop attached to the associated vertex in  $\mathcal{O}(\Gamma)$ .
3. Let  $u^\sigma$  be the  $\sigma$ -orbit of an arrow  $u : x \rightarrow y$ . If both  $x$  and  $y$  are non-periodic, then there is an edge between  $x^\tau$  and  $y^\tau$ . If  $x$  (or  $y$ ) is non-periodic and  $y$  (or  $x$ ) is periodic, then there is an edge between  $x^\tau$  (or  $y^\tau$ ) and the vertex associated to the periodic component containing  $y$  (or  $x$ , respectively). Otherwise, no arrow is associated to  $u^\sigma$ .

By [14, 4.2], the fundamental group of the orbit graph  $\mathcal{O}(\Gamma)$  is isomorphic to  $\pi_1(\Gamma)$ .

Throughout this section, we assume that  $A$  is standard lura, having  $\Gamma$  as a connecting component. We use the following lemmata:

**Lemma 8.1.** *If  $\mathcal{O}(\Gamma)$  is a tree, then  $A$  is weakly shod.*

**Proof.** If  $\mathcal{O}(\Gamma)$  is a tree, then  $\Gamma$  is simply connected (see [14, 4.1 and 4.2]). In particular,  $\Gamma$  has no oriented cycle. Hence,  $A$  is lura and its non-semiregular component (there is at most one) has no oriented cycles. So  $A$  is weakly shod ([17, 2.5]).  $\square$

**Lemma 8.2.** *Let  $A$  be a product of lura algebras with connecting components. If the orbit graph of any connecting component is a tree, then  $A$  is a product of simply connected algebras and  $\text{HH}^1(A) = 0$ .*

**Proof.** This follows from the preceding lemma and from [30, Cor. 2].  $\square$

We now prove Theorem B whose statement we recall for convenience.

**Theorem B.** *Let  $A$  be a standard lura algebra, and  $\Gamma$  its connecting component(s). The following are equivalent:*

- (a)  $A$  has no proper Galois covering, that is,  $A$  is simply connected.
- (b)  $\text{HH}^1(A) = 0$ .
- (c)  $\Gamma$  is simply connected.
- (d) The orbit graph  $\mathcal{O}(\Gamma)$  is a tree.


Moreover, if these conditions are verified, then  $A$  is weakly shod.

**Proof.** By [14, 4.1, 4.2] and the above lemma, (c) and (d) are equivalent and imply (a) and (b). If  $A$  is simply connected, then 6.5 implies  $\pi_1(\Gamma) = 1$ . So (a) implies (c). Finally, assume that  $\text{HH}^1(A) = 0$ . By 6.5, the algebra  $A$  admits a Galois covering with group  $\pi_1(\Gamma)$ . This group is free because of [14, 4.2]. On the other hand, the rank of  $\pi_1(\Gamma)$  is less than or equal to  $\dim \text{HH}^1(A)$  because of [19, Cor. 3]. Therefore  $\pi_1(\Gamma) = 1$ . So (b) implies (c). Thus the conditions are equivalent, and imply that  $A$  is weakly shod by 8.1.  $\square$

We illustrate Theorem B on the following examples. In particular, note that this theorem does not necessarily hold true if one drops standardness.

**Example 8.3.**

- (a) Let  $A$  be as in 3.6(a). Then  $A$  clearly admits a Galois covering with group a free group of rank 3 by a locally bounded  $k$ -category. It is given by the universal cover of the underlying graph of the ordinary quiver. So  $A$  is not simply connected. The orbit graph  $\mathcal{O}(\Gamma)$  of the connecting component  $\Gamma$  is as follows:



Then  $\pi_1(\Gamma)$  is free of rank 3. A straightforward computation gives  $\dim \text{HH}^1(A) = 7$  (see also [16, Thm. 1]).

- (b) Let  $A$  be as in 6.4. As already noticed,  $A$  is a simply connected representation-finite algebra. Also, it is not standard. The orbit graph of its Auslander–Reiten quiver is as follows:




Finally,  $A$  admits the following outer derivation, yielding a non-zero element in  $\text{HH}^1(A)$  (see [15, 4.2])


$$\begin{aligned}
 d : A &\rightarrow A, \\
 \sigma, \delta &\mapsto 0, \\
 \rho &\mapsto \rho^3.
 \end{aligned}$$

This example shows that Theorem B may fail if one drops standardness. Note that the definition of simple connectedness differs slightly from that used in [15, 4.3]: In [15], as in [14, §6], a representation-finite algebra is called simply connected if its Auslander–Reiten quiver is simply connected.

(c) Let  $A$  be given by the quiver:



bound by  $\beta\varepsilon = 0, \alpha\gamma = 0, \beta\delta = \alpha\delta, \delta\zeta = 0, \delta\eta = 0, \zeta\mu = \eta\lambda, \zeta\mu\nu = 0$  Then  $A$  is lura. Actually, it is right glued [5, 4.2]. The orbit graph  $\mathcal{O}(\Gamma)$  of its connecting component  $\Gamma$  is as follows:



It is a tree. Also,  $A$  is simply connected, and it is not hard to see that  $\text{HH}^1(A) = 0$  using, for instance, Happel’s long exact sequence (see [23, 5.3]).

We end with the following problem.

**Problem 5.** Let  $A$  be a non-standard lura algebra. How can the vanishing of  $\text{HH}^1(A)$  be expressed in terms of topological properties of  $A$ ?

**Acknowledgments**

The first author gratefully acknowledges financial support from the NSERC of Canada and the Université de Sherbrooke, the second from Universidad San Francisco de Quito and NSERC of Canada and the third from the NSERC of Canada and the École normale supérieure de Cachan. Part of the work presented in this paper was done during visits of the second and third author at the Université de Sherbrooke. They thank the algebra research group of Sherbrooke for their warm hospitality.

**References**

[1] I. Assem, Left sections and the left part of an Artin algebra, *Colloq. Math.* 116 (2) (2009) 273–300.  
 [2] I. Assem, T. Brüstle, R. Schiffler, Cluster-tilted algebras as trivial extensions, *Bull. London Math. Soc.* 40 (1) (2008) 151–162.  
 [3] I. Assem, F.U. Coelho, Two-sided gluings of tilted algebras, *J. Algebra* 269 (2) (2003) 456–479.  
 [4] I. Assem, F.U. Coelho, Endomorphism algebras of projective modules over lura algebras, *J. Algebra Appl.* 3 (1) (2004) 49–60.  
 [5] I. Assem, F.U. Coelho, M. Lanzilotta, D. Smith, S. Trepode, Algebras determined by their left and right parts, in: *Contemp. Math.*, vol. 376, 2005, pp. 13–47.  
 [6] I. Assem, F.U. Coelho, S. Trepode, The left and the right parts of a module category, *J. Algebra* 281 (2) (2004) 518–534.  
 [7] I. Assem, M. Lanzilotta, The simple connectedness of a tame weakly shod algebra, *Comm. Algebra* 32 (9) (2004) 3685–3701.  
 [8] I. Assem, M. Lanzilotta, M.J. Redondo, Laura skew group algebras, *Comm. Algebra* 35 (7) (2007) 2241–2257.  
 [9] I. Assem, M.I. Platzeck, S. Trepode, On the representation dimension of tilted and lura algebras, *J. Algebra* 296 (2) (2006) 426–439.  
 [10] I. Assem, M.J. Redondo, The first Hochschild cohomology group of a schurian cluster-tilted algebra, *Manuscripta Math.* 128 (3) (2009) 373–388.

- [11] I. Assem, D. Simson, A. Skowroński, Elements of the representation theory of associative algebras, in: London Math. Soc. Stud. Texts, vol. 65, Cambridge University Press, Cambridge, 2006, ix, 458 p.
- [12] M. Auslander, Sverre O. Smalø, Almost split sequences in subcategories, *J. Algebra* 69 (1981) 426–454.
- [13] R. Bautista, P. Gabriel, A.V. Roiter, L. Salmerón, Representation-finite algebras and multiplicative bases, *Invent. Math.* 81 (1985) 217–285.
- [14] K. Bongartz, P. Gabriel, Covering spaces in representation-theory, *Invent. Math.* 65 (1982) 331–378.
- [15] R.-O. Buchweitz, S. Liu, Hochschild cohomology and representation-finite algebras, *Proc. Lond. Math. Soc. Ser. III* 88 (2) (2004) 355–380.
- [16] C. Cibils, M. Saorín, The first Hochschild cohomology group of an algebra with coefficients in a bimodule, *J. Algebra* 237 (2001) 121–141.
- [17] F.U. Coelho, M.A. Lanzilotta, Weakly shod algebras, *J. Algebra* 265 (1) (2003) 379–403.
- [18] F.U. Coelho, A. Skowroński, On the Auslander–Reiten components of a quasitilted algebra, *Fund. Math.* 149 (1996) 67–82.
- [19] J.A. de la Peña, M. Saorín, On the first Hochschild cohomology group of an algebra, *Manuscripta Math.* 104 (4) (2001) 431–442.
- [20] J. Dionne, D. Smith, Articulations of algebras and their homological properties, *J. Algebra Appl.* 5 (3) (2006) 345–359.
- [21] P. Gabriel, The universal cover of a representation-finite algebra, in: *Lecture Notes in Math.*, vol. 903, Springer-Verlag, 1981, pp. 68–105.
- [22] S.I. Gelfand, Y.I. Manin, *Methods for Homological Algebra*, second ed., Springer Monogr. Math., Springer-Verlag, Berlin, 2003.
- [23] D. Happel, Hochschild cohomology of finite-dimensional algebras, in: *Séminaire d'algèbre P. Dubreil et M.-P. Malliavin, Proc., Paris/France 1987/1988*, in: *Lecture Notes in Math.*, vol. 1404, 1989, pp. 108–126.
- [24] D. Happel, I. Reiten, S.O. Smalø, Tilting in abelian categories and quasitilted algebras, *Mem. Amer. Math. Soc.* 575 (1996), 88 p.
- [25] D. Happel, C.M. Ringel, Directing projective modules, *Arch. Math.* 60 (3) (1993) 237–246.
- [26] B. Keller, D. Vossieck, Aisles in derived categories, in: *Deuxième Contact Franco-Belge en Algèbre, Faulx-les-Tombes, 1987*, in: *Bull. Soc. Math. Belg. Ser. A*, vol. 40, 1988, pp. 239–253.
- [27] M. Lanzilotta, D. Smith, Laura algebras and quasi-directed components, *Colloq. Math.* 105 (2) (2006) 179–196.
- [28] P. Le Meur, Revêtements galoisiens et groupe fondamental des algèbres de dimension finie, Thèse de doctorat de l'Université Montpellier 2, 2006, <http://tel.ccsd.cnrs.fr/tel-00011753>.
- [29] P. Le Meur, On Galois coverings and tilting modules, *J. Algebra* 319 (12) (2008) 4961–4999.
- [30] P. Le Meur, Galois coverings of weakly shod algebras, *Comm. Algebra* (2009), in press.
- [31] P. Le Meur, Topological invariants of piecewise hereditary algebras, *Trans. Amer. Math. Soc.* (2009), in press.
- [32] H. Lenzing, A. Skowroński, Quasi-tilted algebras of canonical type, *Colloq. Math.* 71 (2) (1996) 161–181.
- [33] R. Martínez-Villa, J.A. de la Peña, The universal cover of a quiver with relations, *J. Pure Appl. Algebra* 30 (1983) 277–292.
- [34] M.J. Redondo, Universal Galois coverings of self-injective algebras by repetitive algebras and Hochschild cohomology, *J. Algebra* 247 (2002) 332–364.
- [35] I. Reiten, A. Skowroński, Generalized double tilted algebras, *J. Math. Soc. Japan* 56 (1) (2004) 269–288.
- [36] C. Riedtmann, Algebren, Darstellungsköcher, Ueberlagerungen und zurück, *Comment. Math. Helv.* 55 (1980) 199–224.
- [37] C. Riedtmann, Many algebras with the same Auslander–Reiten quiver, *Bull. London Math. Soc.* 15 (1) (1983) 43–47.
- [38] C.M. Ringel, Tame Algebras and Integral Quadratic Forms, *Lecture Notes in Math.*, vol. 1099, Springer-Verlag Berlin, 1984.
- [39] C.M. Ringel, The regular components of the Auslander–Reiten quiver of a tilted algebra, *Chinese Ann. Math. Sci. Ser. B* 9 (1) (1988) 1–18.
- [40] A. Skowroński, Simply connected algebras and Hochschild cohomologies, in: *CMS Conf. Proc.*, vol. 14, Amer. Math. Soc., 1993, pp. 431–447.
- [41] A. Skowroński, Generalized standard Auslander–Reiten components, *J. Math. Soc. Japan* 46 (3) (1994) 517–543.
- [42] A. Skowroński, On Artin algebras with almost all indecomposable modules of projective or injective dimension at most one, *Cent. Eur. J. Math.* 1 (1) (2003) 108–122.
- [43] D. Smith, On generalized standard Auslander–Reiten components having only finitely many non-directing modules, *J. Algebra* 279 (2) (2004) 493–513.