

θ -Summation and Hardy Spaces¹

Ferenc Weisz

*Department of Numerical Analysis, Eötvös L. University, H-1117 Budapest,
 Pázmány P. sétány 1/D, Hungary
 E-mail: weisz@ludens.elte.hu*

Communicated by Rolf J. Nessel

Received July 21, 1999; accepted in revised form June 2, 2000;
 published online October 11, 2000

A general summability method of Fourier series and Fourier transforms is given with the help of an integrable function θ having integrable Fourier transform. Under some weak conditions on θ we show that the maximal operator of the θ -means of a distribution is bounded from $H_p(\mathbf{T})$ to $L_p(\mathbf{T})$ ($p_0 < p < \infty$) and is of weak type (1,1), where $H_p(\mathbf{T})$ is the classical Hardy space and $p_0 < 1$ is depending only on θ . As a consequence we obtain that the θ -means of a function $f \in L_1(\mathbf{T})$ converge a.e. to f . For the endpoint p_0 we get that the maximal operator is of weak type $(H_{p_0}(\mathbf{T}), L_{p_0}(\mathbf{T}))$. Moreover, we prove that the θ -means are uniformly bounded on the spaces $H_p(\mathbf{T})$ whenever $p_0 < p < \infty$ and are uniformly of weak type $(H_{p_0}(\mathbf{T}), H_{p_0}(\mathbf{T}))$. Thus, in the case $f \in H_p(\mathbf{T})$, the θ -means converge to f in $H_p(\mathbf{T})$ norm ($p_0 < p < \infty$). The same results are proved for the conjugate θ -means and for Fourier transforms, too. Some special cases of the θ -summation are considered, such as the Weierstrass, Picar, Bessel, Fejér, Riemann, de La Vallée-Poussin, Rogosinski and Riesz summations. © 2000 Academic Press

Key Words: Hardy spaces; p -atom; atomic decomposition; interpolation; Fourier transforms; θ -summation.

1. INTRODUCTION

The Hardy–Lorentz spaces $H_{p,q}(\mathbf{T})$ of distributions are introduced with the $L_{p,q}(\mathbf{T})$ Lorentz norm of the non-tangential maximal function. Of course, $H_p(\mathbf{T}) = H_{p,p}(\mathbf{T})$ are the usual Hardy spaces ($0 < p \leq \infty$).

Butzer and Nessel [3] and recently Bokor, Schipp, Szili and Vértesi [2, 11, 12, 16, 17] considered a general method of summation, the so-called θ -summability. The θ -means of Fourier transforms can be written in a

¹ This paper was written while the author was researching at Humboldt University in Berlin supported by Alexander von Humboldt Foundation. This research was also supported by the Hungarian Scientific Research Funds (OTKA) F019633, by the Foundation for Research and Improvement in Higher Education FKFP-0228/1999 and by the Széchenyi Professorship.

natural way as a singular integral of the Fourier transform of θ , $\hat{\theta}$ (see Butzer and Nessel [3]). They proved that if $\hat{\theta}$ can be estimated by a non-increasing integrable function, then the θ -means of a function $f \in L_1(\mathbf{R})$ converge a.e. to f . This convergence result is also proved there for the θ -means of Fourier series. As special cases they considered the Weierstrass, Picar, Bessel, Fejér, de La Vallée-Poussin and Riesz summations. For example, they verified that the Riesz means $\sigma_T^{\alpha, \gamma} f$ converge a.e. to f as $T \rightarrow \infty$ if $f \in L_1(\mathbf{R})$ and $\gamma = 1, 2$ (see also Stein and Weiss [14]).

The author [21] generalized this last result and proved that the maximal Riesz operator $\sigma_*^{\alpha, \gamma} := \sup_{T>0} |\sigma_T^{\alpha, \gamma}|$ is bounded from $H_p(\mathbf{R})$ to $L_p(\mathbf{R})$ provided that $0 < \alpha < \infty$, $1 \leq \gamma < \infty$, $1/(\min(\alpha, 1) + 1) < p < \infty$ and, moreover, it is of weak type $(1, 1)$, i.e.

$$\sup_{\rho>0} \rho \lambda(\sigma_*^{\alpha, \gamma} f > \rho) \leq C \|f\|_1 \quad (f \in L_1(\mathbf{R}))$$

(this last result for $\alpha = \gamma = 1$ can also be found in Zygmund [23] and Móricz [10]). This weak type inequality assures already the a.e. convergence of the Riesz means mentioned above.

In this paper we generalize these results. First we consider the θ -means of Fourier series and prove that the θ -means $U_n^\theta f$ of a function $f \in L_1(\mathbf{T})$ can be written also as a singular integral of f and $\hat{\theta}$ over \mathbf{R} . We introduce the maximal operator $U_*^\theta := \sup_{n \in \mathbf{N}} |U_n^\theta|$, the conjugate distribution \tilde{f} , the conjugate θ -means $\tilde{U}_n^\theta f$ and the conjugate maximal operator \tilde{U}_*^θ .

Under some weak conditions on θ and $\hat{\theta}$ we will show that the maximal operators U_*^θ and \tilde{U}_*^θ are bounded from $H_{p,q}(\mathbf{T})$ to $L_{p,q}(\mathbf{T})$ whenever $p_0 < p < \infty$, $0 < q \leq \infty$ and are of weak type $(1, 1)$. The parameter p_0 is less than 1 and depending on θ . For this endpoint we can verify that the preceding two maximal operators are of weak type $(H_{p_0}(\mathbf{T}), L_{p_0}(\mathbf{T}))$.

A usual density argument implies then that $U_n^\theta f \rightarrow f$ a.e. and $\tilde{U}_n^\theta f \rightarrow \tilde{f}$ a.e. as $n \rightarrow \infty$, provided that $f \in L_1(\mathbf{T})$. Note that \tilde{f} is not necessarily integrable whenever f is.

We will prove also that the operators U_n^θ and \tilde{U}_n^θ ($n \in \mathbf{N}$) are uniformly bounded in n from $H_{p,q}(\mathbf{T})$ to $H_{p,q}(\mathbf{T})$ ($p_0 < p < \infty$, $0 < q \leq \infty$) and are uniformly of weak type $(H_{p_0}(\mathbf{T}), H_{p_0}(\mathbf{T}))$. From this it follows that $U_n^\theta f \rightarrow f$ and $\tilde{U}_n^\theta f \rightarrow \tilde{f}$ in $H_{p,q}(\mathbf{T})$ norm (resp. in weak $H_{p_0}(\mathbf{T})$ norm) as $n \rightarrow \infty$, whenever $f \in H_{p,q}(\mathbf{T})$ ($p_0 < p < \infty$, $0 < q \leq \infty$) (resp. $f \in H_{p_0}(\mathbf{T})$).

As special case we investigate ten well known summability methods, amongst others the summations mentioned above.

We consider also the θ -means of Fourier transforms on the real line and prove all the results above in this context.

I thank the referees for reading the paper carefully and for their useful comments.

2. HARDY SPACES AND CONJUGATE FUNCTIONS

Let \mathbf{N} denote the none-negative integers, \mathbf{R} the real numbers; \mathbf{R}_+ the positive real numbers, $\mathbf{T} := [-\pi, \pi)$ and λ be the Lebesgue measure. We also use the notation $|I|$ for the Lebesgue measure of the set I . We briefly write $L_{p,q}(\mathbf{X})$ instead of the real Lorentz space $L_{p,q}(\mathbf{X}, \lambda)$ ($0 < p, q \leq \infty$) and its norm is denoted by $\|\cdot\|_{p,q}$ where $\mathbf{X} = \mathbf{T}$ or \mathbf{R} (for the exact definitions see e.g. Weisz [21] and the references there). We extend all functions on \mathbf{T} periodically to \mathbf{R} .

Let f be a distribution on $C^\infty(\mathbf{T})$. The n th Fourier coefficient is defined by $\hat{f}(n) := f(e^{-inx})$ where $i = \sqrt{-1}$. In special case, if f is an integrable function then

$$\hat{f}(n) = \frac{1}{2\pi} \int_{\mathbf{T}} f(x) e^{-inx} dx \quad (n \in \mathbf{N}).$$

The non-tangential maximal function of a distribution f is defined by

$$f^*(x) := \sup_{0 < r < 1} |f * P_r(x)|,$$

where $*$ denotes the convolution and

$$P_r(x) := \sum_{k=-\infty}^{\infty} r^{|k|} e^{ikx} = \frac{1-r^2}{1+r^2-2r \cos x} \quad (x \in \mathbf{T})$$

is the Poisson kernel.

For $0 < p, q \leq \infty$ the Hardy-Lorentz space $H_{p,q}(\mathbf{T})$ consists of all distributions f for which

$$\|f\|_{H_{p,q}(\mathbf{T})} := \|f^*\|_{p,q} < \infty.$$

Note that in case $p = q$ the usual definition of Hardy spaces $H_p(\mathbf{T}) = H_{p,p}(\mathbf{T})$ is obtained. For other equivalent definitions we call for Fefferman and Stein [5] and Stein [15]. Recall that $L_1(\mathbf{T}) \subset H_{1,\infty}(\mathbf{T})$, more exactly,

$$\|f\|_{H_{1,\infty}(\mathbf{T})} = \sup_{\rho > 0} \rho \lambda(f^* > \rho) \leq \|f\|_1 \quad (f \in L_1(\mathbf{T})). \quad (1)$$

Moreover,

$$H_{p,q}(\mathbf{T}) \sim L_{p,q}(\mathbf{T}) \quad (1 < p < \infty, 0 < q \leq \infty), \quad (2)$$

where \sim denotes the equivalence of the norms and spaces (see Fefferman and Stein [5], Stein [15], Fefferman, Riviere, Sagher [4]).

The following interpolation result concerning Hardy–Lorentz spaces will be used several times in this paper (see Fefferman, Riviere, Sagher [4] and also Weisz [19]).

THEOREM A. *If a sublinear (resp. linear) operator V is bounded from $H_{p_0}(\mathbf{T})$ to $L_{p_0}(\mathbf{T})$ (resp. to $H_{p_0}(\mathbf{T})$) and from $L_{p_1}(\mathbf{T})$ to $L_{p_1}(\mathbf{T})$ ($p_0 \leq 1 < p_1 \leq \infty$) then it is also bounded from $H_{p,q}(\mathbf{T})$ to $L_{p,q}(\mathbf{T})$ (resp. to $H_{p,q}(\mathbf{T})$) if $p_0 < p < p_1$ and $0 < q \leq \infty$.*

For a distribution

$$f \sim \sum_{k=-\infty}^{\infty} \hat{f}(k) e^{ikx}$$

the conjugate distribution is defined by

$$\tilde{f} \sim \sum_{k=-\infty}^{\infty} (-i \operatorname{sign} k) \hat{f}(k) e^{ikx}.$$

As is well known, if f is an integrable function then

$$\tilde{f}(x) = \text{p.v.} \frac{1}{\pi} \int_{\mathbf{T}} \frac{f(x-t)}{2 \tan(t/2)} dt := \lim_{\varepsilon \rightarrow 0} \frac{1}{\pi} \int_{\varepsilon < |t| < \pi} \frac{f(x-t)}{2 \tan(t/2)} dt.$$

Moreover, the conjugate function \tilde{f} does exist almost everywhere, but it is not integrable in general. It is easy to see that $(\tilde{f})^\sim = -f$.

Fefferman and Stein [5] verified that

$$\|f\|_{H_p(\mathbf{T})} \sim \|f\|_p + \|\tilde{f}\|_p \quad (0 < p < \infty). \quad (3)$$

3. θ -SUMMABILITY OF FOURIER SERIES

First we introduce the Fourier transform for an integrable function $f \in L_1(\mathbf{R})$ by

$$\hat{f}(u) = \frac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} f(x) e^{-iux} dx \quad (u \in \mathbf{R}).$$

The θ -summation was considered in Butzer and Nessel [3] and, more recently Bokor, Schipp, Szili and Vértesi [2, 11, 12, 16, 17] investigated the uniform convergence of the θ -means and some interpolation problems for continuous functions.

In what follows we suppose that $\theta \in L_1(\mathbf{R})$ is an even continuous function satisfying $\theta(0) = 1$, $\hat{\theta} \in L_1(\mathbf{R})$ and $\theta(\frac{\cdot}{n+1}) \in l_1$. Note that this last condition is satisfied if θ is non-increasing on \mathbf{R}_+ or if it has compact support.

Denote by $s_n f$ the n th partial sum of the Fourier series of a distribution f , namely,

$$s_n f(x) := \sum_{k=-n}^n \hat{f}(k) e^{ikx}.$$

The θ -means of a distribution f are defined by

$$U_n^\theta f(x) := \sum_{k=-\infty}^{\infty} \theta\left(\frac{k}{n+1}\right) \hat{f}(k) e^{ikx} = (f * K_n^\theta)(x) \quad (x \in \mathbf{T}), \quad (4)$$

where the K_n^θ kernels satisfy

$$K_n^\theta(t) := \sum_{k=-\infty}^{\infty} \theta\left(\frac{k}{n+1}\right) e^{ikt} = 1 + 2 \sum_{k=1}^{\infty} \theta\left(\frac{k}{n+1}\right) \cos(kt) \quad (t \in \mathbf{T}).$$

It is easy to see that if θ has bounded variation then the θ -summation is permanent, i.e. if $s_n f$ is convergent in some sense then $U_n^\theta f$ is also convergent and converges to the same limit.

Following Butzer and Nessel [3] and Schipp and Bokor [11] we verify a new characterization for the θ -means. We write $U_n^\theta f$ as a singular integral of f and the Fourier transform of θ over the whole real line.

LEMMA 1. *If $f \in L_1(\mathbf{T})$ then*

$$U_n^\theta f(x) = (n+1) \int_{-\infty}^{\infty} f(x-t) \hat{\theta}((n+1)t) dt \quad (n \in \mathbf{N}). \quad (5)$$

Proof. If $f(t) = e^{ikt}$ then

$$\begin{aligned} (n+1) \int_{-\infty}^{\infty} e^{ik(x-t)} \hat{\theta}((n+1)t) dt &= e^{ikx} \int_{-\infty}^{\infty} e^{-ikt/(n+1)} \hat{\theta}(t) dt \\ &= \theta\left(\frac{k}{n+1}\right) e^{ikx} = U_n^\theta f(x). \end{aligned}$$

Hence the lemma holds also for trigonometric polynomials. Let f be an arbitrary element from $L_1(\mathbf{T})$ and (f_k) be a sequence of trigonometric

polynomials such that $f_k \rightarrow f$ in $L_1(\mathbf{T})$ norm. The condition $\theta(\frac{\cdot}{n+1}) \in l_1$ implies that $K_n^\theta \in L_1(\mathbf{T})$. Since

$$U_n^\theta f(x) = \frac{1}{2\pi} \int_{\mathbf{T}} f(x-t) K_n^\theta(t) dt$$

for $f \in L_1(\mathbf{T})$, we can conclude that $U_n^\theta f_k \rightarrow U_n^\theta f$ in $L_1(\mathbf{T})$ norm as $k \rightarrow \infty$. On the other hand, $\hat{\theta} \in L_1(\mathbf{R})$, and so

$$(n+1) \int_{-\infty}^{\infty} f_k(x-t) \hat{\theta}((n+1)t) dt \rightarrow (n+1) \int_{-\infty}^{\infty} f(x-t) \hat{\theta}((n+1)t) dt$$

in $L_1(\mathbf{T})$ norm as $k \rightarrow \infty$. This finishes the proof of the lemma. ■

The *conjugate θ -means* of a distribution f are introduced by

$$\tilde{U}_n^\theta f(x) := \sum_{k=-\infty}^{\infty} \theta\left(\frac{k}{n+1}\right) (-i \operatorname{sign} k) \hat{f}(k) e^{ikx}.$$

The *maximal* and *conjugate maximal θ -operators* are defined by

$$U_*^\theta f := \sup_{n \in \mathbf{N}} |U_n^\theta f| \quad \text{and} \quad \tilde{U}_*^\theta f := \sup_{n \in \mathbf{N}} |\tilde{U}_n^\theta f|,$$

respectively. Our first boundedness result is the following

LEMMA 2. *The operator U_*^θ is bounded on $L_\infty(\mathbf{T})$.*

Proof. The characterization (5) implies that

$$\|U_n^\theta f\|_\infty \leq \|f\|_\infty \|\hat{\theta}\|_1$$

for all $n \in \mathbf{N}$, which shows the lemma. ■

In this paper the constants C are depending only on θ and the constants C_p (resp. $C_{p,q}$) are depending only on p and θ (resp. p, q and θ) and may denote different constants in different contexts.

4. THE BOUNDEDNESS OF THE MAXIMAL θ -OPERATOR

A *generalized interval* on \mathbf{T} is either an interval $I \subset \mathbf{T}$ or $I = [-\pi, x) \cup [y, \pi)$. A bounded measurable function a is a *p -atom* if there exists a generalized interval I such that

- (i) $\int_I a(x) x^j dx = 0$ where $j \in \mathbf{N}$ and $j \leq [1/p - 1]$, the integer part of $(1/p - 1)$,
- (ii) $\|a\|_\infty \leq |I|^{-1/p}$,
- (iii) $\{a \neq 0\} \subset I$.

We will use the following two theorems, the first one can be found in Weisz [21].

THEOREM B. *Suppose that the operator V is sublinear and, for some $0 < p \leq 1$,*

$$\int_{\mathbf{T} \setminus 8I} |Va|^p d\lambda \leq C_p \quad (6)$$

for every p -atom a where I is the support of the atom and $8I$ is the generalized interval with the same center as I and with length $8|I|$. If V is bounded from $L_{p_1}(\mathbf{T})$ to $L_{p_1}(\mathbf{T})$ for a fixed $1 < p_1 \leq \infty$ then

$$\|Vf\|_p \leq C_p \|f\|_{H_p(\mathbf{T})} \quad (f \in H_p(\mathbf{T})).$$

We formulate also a weak version of this theorem, which is an easy modification of a result in Long [8], so we sketch the proof, only.

THEOREM C. *Suppose that the operator V is sublinear and, for some $0 < p < 1$,*

$$\sup_{\rho > 0} \rho^p \lambda(\{|Va| > \rho\} \cap \{\mathbf{T} \setminus 8I\}) \leq C_p \quad (7)$$

for every p -atom a where I denotes again the support of the atom. If V is bounded from $L_{p_1}(\mathbf{T})$ to $L_{p_1}(\mathbf{T})$ for a fixed $1 < p_1 \leq \infty$ then

$$\|Vf\|_{p, \infty} \leq C_p \|f\|_{H_p(\mathbf{T})} \quad (f \in H_p(\mathbf{T})).$$

Proof. If (7) is satisfied without the intersection with $\{\mathbf{T} \setminus 8I\}$, then the result can be found in Long [8, p. 279]. Then

$$\begin{aligned} & \sup_{\rho > 0} \rho^p (\{|Va| > \rho\} \cap \{8I\}) \\ & \leq \int_{8I} |Va|^p d\lambda \leq C_p \left(\int_{\mathbf{T}} |Va|^{p_1} d\lambda \right)^{p/p_1} |I|^{1-p/p_1} \leq C_p, \end{aligned}$$

which proves the theorem. \blacksquare

Now we are ready to prove the boundedness of the maximal operator on the Hardy spaces. First we recall a known result, which was shown in another context. Taking into account (5) we can see that Torchinsky [18, p. 82–84] has proved essentially the next inequality.

PROPOSITION 1. *Assume that there is an even, on \mathbf{R}_+ non-increasing function η_0 such that $|\hat{\theta}| \leq \eta_0$. If $\eta_0 \in L_1(\mathbf{R})$ then*

$$\sup_{\rho > 0} \rho \lambda(U_*^\theta f > \rho) \leq C \|f\|_1 \quad (f \in L_1(\mathbf{T})). \quad (8)$$

It follows from Proposition 1 and Lemma 2 and by interpolation that

$$\|U_*^\theta f\|_p \leq C \|f\|_p \quad (f \in L_p(\mathbf{T}), 1 < p \leq \infty).$$

If we suppose a little bit more on η_0 then we can prove that U_*^θ is bounded from $H_1(\mathbf{T})$ to $L_1(\mathbf{T})$.

THEOREM 1. *Assume that there is an even, on \mathbf{R}_+ non-increasing function η_0 such that $|\hat{\theta}| \leq \eta_0$, $t\eta_0(t)$ is non-increasing on the interval $[1, \infty)$. If θ has compact support that is contained in $[-c, c]$ and if $\eta_0 \in L_1(\mathbf{R})$ then*

$$\|_*^\theta f\|_1 \leq cC \|f\|_{H_1(\mathbf{T})} \quad (f \in H_1(\mathbf{T})).$$

Proof. We will verify (6) for $p = 1$. Then Theorem 1 will follow from Lemma 2 and Theorem B.

To this end let a be an arbitrary 1-atom with support I and $2^{-K-1} < |I|/\pi \leq 2^{-K}$ ($K \in \mathbf{N}$). If t_0 is the center of I , then the center of $I' := I - t_0$ is zero. By changing the variables we can see that

$$\begin{aligned} \int_{\mathbf{T} \setminus 8I} |U_*^\theta a|^p d\lambda &= \int_{\mathbf{T} \setminus 8I} \sup_{n \in \mathbf{N}} \left| \int_I a(t) K_n^\theta(x-t) dt \right|^p dx \\ &= \int_{\mathbf{T} \setminus 8I'} \sup_{n \in \mathbf{N}} \left| \int_{I'} a'(t) K_n^\theta(x-t) dt \right|^p dx \\ &= \int_{\mathbf{T} \setminus 8I'} |U_*^\theta a'|^p d\lambda, \end{aligned}$$

where $a'(t) := a(t + t_0)$.

Hence we can suppose that the center of I is zero. In this case

$$[-\pi 2^{-K-2}, \pi 2^{-K-2}] \subset I \subset [-\pi 2^{-K-1}, \pi 2^{-K-1}].$$

First suppose that $x \geq 0$. Then

$$\begin{aligned} \int_{\{\mathbf{T} \setminus 8I\} \cap \{x \geq 0\}} |U_n^\theta a(x)| dx &\leq \sum_{i=2}^{2^K-1} \int_{\pi i 2^{-K}}^{\pi(i+1) 2^{-K}} \sup_{n+1 \geq r_i} |U_n^\theta a(x)| dx \\ &\quad + \sum_{i=2}^{2^K-1} \int_{\pi i 2^{-K}}^{\pi(i+1) 2^{-K}} \sup_{n+1 < r_i} |U_n^\theta a(x)| dx \\ &= (A) + (B), \end{aligned}$$

where $r_i := 2^K i^{-\alpha}$ ($i \in \mathbf{N}$) with $\alpha > 0$ chosen later.

By Lemma 1 and by the condition $|\hat{\theta}| \leq \eta_0$ we estimate $U_n^\theta a$ as follows:

$$|U_n^\theta a(x)| \leq (n+1) \int_{-\infty}^{\infty} |a(t)| \eta_0((n+1)(x-t)) dt.$$

Then, by the definition of the 1-atom,

$$\sup_{n+1 \geq r_i} |U_n^\theta a(x)| \leq 2^K r_i \sum_{k=-\infty}^{\infty} \int_{I+2k\pi}^{\infty} \eta_0(r_i(x-t)) dt := (A_1)(x) + (A_2)(x),$$

where (A_1) denotes the term $k=0$ and (A_2) the sum $\sum_{|k| \geq 1}^{\infty}$.

If $t \in I$ and $x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K}]$ for some $i=2, \dots, 2^K-1$, then

$$|x-t| \geq \pi i 2^{-K} - \pi 2^{-K-1} \geq \pi(i-1) 2^{-K}. \quad (9)$$

This implies

$$(A_1)(x) \leq C 2^K i^{-\alpha} \eta_0(i^{1-\alpha} \pi/2) \quad (x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K}]).$$

If $t \in I + 2k\pi$ for some $k \neq 0$ then $|x-t| \sim 2|k|\pi$. We have

$$(A_2)(x) \leq C 2^K i^{-\alpha} \sum_{k=1}^{\infty} \eta_0(2^{K+1} i^{-\alpha} \pi k) \leq C \int_0^{\infty} \eta_0 d\lambda \leq C.$$

Hence, in case $0 < \alpha < 1$,

$$(A)(x) \leq C + C 2^{-K} \sum_{i=2}^{2^K-1} 2^K i^{-\alpha} \eta_0(i^{1-\alpha} \pi/2) \leq C \int_0^{\infty} \eta_0 d\lambda \leq C.$$

Now let us consider (B) . Since $\text{supp } \theta \subset [-c, c]$ and θ is bounded, (4) implies

$$|U_n a(x)| \leq C \sum_{|k|=0}^{c(n+1)} |\hat{a}(k)|.$$

As

$$|\hat{a}(k)| = \left| \frac{1}{2\pi} \int_I a(x)(e^{-ikx} - 1) dx \right| \leq C \int_I |a(x)| |kx| dx \leq C |k| |I|$$

we obtain

$$\sup_{n+1 < r_i} |U_n a(x)| \leq c C r_i^2 2^{-K} \leq c C 2^K i^{-2\alpha}.$$

Therefore,

$$(B) \leq c C 2^{-K} \sum_{i=2}^{2^K-1} 2^{Ki-2\alpha}$$

which is bounded if $1/2 < \alpha < 1$. If $x < 0$ then

$$\begin{aligned} \int_{\{\mathbf{T} \setminus 8I\} \cap \{x < 0\}} |U_*^\theta a(x)| dx &\leq \sum_{i=-2}^{-(2^K-1)} \int_{\pi i 2^{-K}}^{\pi(i-1) 2^{-K}} \sup_{n+1 \geq r_i} |U_n^\theta a(x)| dx \\ &\quad + \sum_{i=-2}^{-(2^K-1)} \int_{\pi i 2^{-K}}^{\pi(i-1) 2^{-K}} \sup_{n+1 < r_i} |U_n^\theta a(x)| dx, \end{aligned}$$

where $r_i := 2^K |i|^{-\alpha}$. The inequality

$$\int_{\{\mathbf{T} \setminus 8I\} \cap \{x < 0\}} |U_*^\theta a(x)| dx \leq c C$$

can be proved exactly as above. The proof of the theorem is complete. \blacksquare

Remark. We can extend this result to $p < 1$ as follows. In addition to the conditions of Theorem 1 suppose that

$$\int_0^\infty t^{(1-p_0)(1+\varepsilon)/(2p_0-1)} \eta_0(t)^{p_0} dt < \infty \quad (10)$$

for some $1/2 < p_0 < 1$ and $\varepsilon > 0$. Then we can prove in the same way that

$$\|U_*^\theta f\|_{p_0} \leq C_{p_0} \|f\|_{H_{p_0}(\mathbf{T})} \quad (f \in H_{p_0}(\mathbf{T}))$$

and

$$\|U_*^\theta f\|_{p,q} \leq C_{p,q} \|f\|_{H_{p,q}(\mathbf{T})} \quad (f \in H_{p,q}(\mathbf{T}))$$

for every $p_0 < p < \infty$ and $0 < q \leq \infty$.

If $p_0 = 1$ then condition (10) reduces to the integrability of η_0 . Since $\hat{\theta}$ is bounded, we may suppose that η_0 is also bounded. It is easy to see that if (10) is satisfied for p_0 then it holds also for all $p_0 \leq p \leq 1$. The interval $[1, \infty)$ in Theorem 1 is a technical condition, only, we could change it to $[c, \infty)$ for any $c > 0$.

If we have some information about the derivatives of $\hat{\theta}$ we can prove an even sharper result. Let $\hat{\theta}^{(k)}$ be denote the k th derivative of $\hat{\theta}$.

THEOREM 2. *Assume that there are two even, on \mathbf{R}_+ non-increasing functions η_N and η_{N+1} such that $|\hat{\theta}^{(N)}| \leq \eta_N$, $0 \neq |\hat{\theta}^{(N+1)}| \leq \eta_{N+1}$, $t^{N+1}\eta_N(t)$ is non-increasing on $[1, \infty)$ and $t^{N+2}\eta_{N+1}(t)$ is non-decreasing on \mathbf{R}_+ ($N \in \mathbf{N}$). If $\eta_N, \eta_{N+1} \in L_{p_0}(\mathbf{R})$ for some $p_0 \leq 1/(N+1)$ then*

$$\|U_*^\theta f\|_{p_0} \leq C_{p_0} \|f\|_{H_{p_0}(\mathbf{T})} \quad (f \in H_{p_0}(\mathbf{T})) \quad (11)$$

and

$$\|U_*^\theta f\|_{p,q} \leq C_{p,q} \|f\|_{H_{p,q}(\mathbf{T})} \quad (f \in H_{p,q}(\mathbf{T})) \quad (12)$$

for every $p_0 < p < \infty$, $0 < q \leq \infty$. In particular, if $f \in L_1(\mathbf{T})$ then (8) holds. Moreover, if $\eta_N, \eta_{N+1} \in L_{p_0, \infty}(\mathbf{R})$ for some $p_0 \leq 1/(N+1)$ ($p_0 \neq 1$) then

$$\|U_*^\theta f\|_{p_0, \infty} \leq C_{p_0} \|f\|_{H_{p_0}(\mathbf{T})} \quad (f \in H_{p_0}(\mathbf{T})) \quad (13)$$

and (12) and (8) are valid.

Proof. First we show (6) for $p = p_0$. Let a be an arbitrary p_0 -atom with support I and $2^{-K-1} < |I|/\pi \leq 2^{-K}$ ($K \in \mathbf{N}$). As in the proof of Theorem 1 we can suppose that the center of I is zero. Let $A_0(x) := a(x)$ ($x \in \mathbf{R}$) and

$$A_j(x) := \int_{-\infty}^x A_{j-1}(t) dt \quad (x \in \mathbf{R}; j = 1, \dots, [1/p - 1] + 1).$$

By (i) of the definition of the atom we can show that $\text{supp } A_j \subset \bigcup_{k=-\infty}^{\infty} \{I + 2k\pi\}$ ($j = 1, \dots, [1/p - 1] + 1$). Moreover, by (ii),

$$\|A_j\|_{\infty} \leq |I|^{-1/p+j} \quad (j = 1, \dots, [1/p - 1] + 1). \quad (14)$$

Using Lemma 1 and integrating by parts we can see that

$$\begin{aligned} |U_n^\theta a(x)| &= (n+1)^{N+1} \left| \int_{-\infty}^{\infty} A_N(t) \hat{\theta}^{(N)}((n+1)(x-t)) dt \right| \\ &\leq (n+1)^{N+1} \int_{-\infty}^{\infty} |A_N(t)| \eta_N((n+1)(x-t)) dt. \end{aligned} \quad (15)$$

By the conditions of the theorem and (14),

$$\begin{aligned} \sup_{n+1 \geq 2^K} |U_n^\theta a(x)| &\leq 2^{K/p_0+K} \sum_{k=-\infty}^{\infty} \int_{I+2k\pi}^{\infty} \eta_N(2^K(x-t)) dt \\ &= (C)(x) + (D)(x), \end{aligned}$$

where (C) denotes the term $k=0$ and (B) the sum $\sum_{|k|=1}^{\infty}$.

We suppose again that $x \in [-\pi, \pi) \setminus 8I$ and $x \geq 0$. If $t \in I$ and $x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K})$ ($i=2, \dots, 2^K-1$), then (9) implies

$$(C)(x) \leq C 2^{K/p_0} \eta_{N+1}((i-1)\pi) \quad (x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K})),$$

thus

$$\begin{aligned} \int_{\{\mathbf{T} \setminus 8I\} \cap \{x \geq 0\}} (C)(x)^{p_0} dx &\leq C_{p_0} 2^{-K} \sum_{i=2}^{2^K-1} 2^K \eta_{N+1}^{p_0}((i-1)\pi) \\ &\leq C_{p_0} \int_0^\infty \eta_{N+1}^{p_0} d\lambda \leq C_{p_0}. \end{aligned}$$

If $t \in I + 2k\pi$ ($k \neq 0$) then

$$(D)^{p_0}(x) \leq C_{p_0} 2^K \sum_{k=1}^{\infty} \eta_{N+1}^{p_0}(2^{K+1}\pi k) \leq C_{p_0} \int_\varepsilon^\infty \eta_{N+1}^{p_0} d\lambda \leq C_{p_0}$$

and (6) is satisfied.

Similarly to (15) we can also obtain that

$$|U_n^\theta a(x)| \leq (n+1)^{N+2} \int_{-\infty}^{\infty} |A_{N+1}(t)| \eta_{N+1}((n+1)(x-t)) dt$$

and then $\sup_{n+1 < 2^K} |U_n^\theta a(x)|$ can be estimated in the same way as $\sup_{n+1 \geq 2^K} |U_n^\theta a(x)|$ above. The case $x < 0$ can be treated similarly. This proves inequality (11).

To prove (13) observe that

$$\begin{aligned} \rho^{p_0} \lambda(\{(C) > \rho\} \cap \{\mathbf{T} \setminus 8I\}) &= \rho^{p_0} \sum_{i \geq 1: \eta_{N+1}(i\pi) > \rho 2^{-K/p_0}} 2^{-K} \\ &\leq C_\varepsilon \rho^{p_0} 2^{-K} \lambda(\{\eta_{N+1} > \rho 2^{-K/p_0}\}) \\ &\leq C_\varepsilon \|\eta_{N+1}\|_{L_{p_0, \infty}(\mathbf{R})}^{p_0}. \end{aligned}$$

Obviously, (D) satisfies also (7). We can estimate $\sup_{n+1 < 2^K} |U_n^\theta a(x)|$ similarly, which shows (13). The inequality (12) follows from Theorem A. Applying (1) and (12) for $p=1$ and $q=\infty$, we conclude

$$\|U_*^\theta f\|_{1, \infty} \leq C \|f\|_{H_{1, \infty}} \leq C \|f\|_1$$

which shows (8). This finishes the proof of the theorem. ■

Remark. We can weaken the condition $t^{N+1}\eta_N(t) \searrow$ in Theorem 2 by

$$t^{N+1}\eta_N(t) \leq t_0^{N+1}\eta_N(t_0) \quad (t \geq t_0 \geq 1).$$

Of course we could replace $t^{N+2}\eta_{N+1}(t) \nearrow$ also by an analogous condition.

In the next theorem we show the boundedness of U_*^θ on Hardy spaces if $t^{N+2}\hat{\theta}^{(N+1)}(t)$ is bounded.

THEOREM 3. Assume that $0 \neq |t^{N+2}\hat{\theta}^{(N+1)}(t)| \leq C$ for some $N \in \mathbf{N}$. Then (12) holds for every $1/(N+2) < p < \infty$, $0 < q \leq \infty$ and

$$\|U_*^\theta f\|_{1/(N+2), \infty} \leq C_{1/(N+2)} \|f\|_{H_{1/(N+2)}(\mathbf{T})} \quad (f \in H_{1/(N+2)}(\mathbf{T})). \quad (16)$$

Especially, if $f \in L_1(\mathbf{T})$ then the weak type $(1, 1)$ inequality (8) holds.

Proof. First we show (12) for $1/(N+2) < p = q \leq 1/(N+1)$. The general case of (12) will follow from Theorem A. To this end, by Lemma 1 and Theorem B we have only to prove that condition (6) is satisfied for $1/(N+2) < p \leq 1/(N+1)$. Note that in this case $[1/p - 1] = N$.

Let a be an arbitrary p -atom with support I and $2^{-K-1} < |I|/\pi \leq 2^{-K}$ ($K \in \mathbf{N}$). We can suppose again that the center of I is zero. As in (15) we conclude

$$\begin{aligned} |U_n^\theta a(x)| &= (n+1)^{N+2} \left| \int_{-\infty}^{\infty} A_{N+1}(t) \hat{\theta}^{(N+1)}((n+1)(x-t)) dt \right| \\ &\leq |I|^{-1/p+N+1} \sum_{k=-\infty}^{\infty} \int_{I+2k\pi} |x-t|^{-(N+2)} dt \\ &:= (E)(x) + (F)(x), \end{aligned}$$

where (E) denotes again the term corresponding to $k=0$ and (F) the sum $\sum_{|k|=1}^{\infty}$.

If $t \in I$ and $x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K})$ for some $i=2, \dots, 2^K-1$, then (9) implies

$$(E)(x) \leq C 2^{K/p} i^{-(N+2)} \quad (x \in [\pi i 2^{-K}, \pi(i+1) 2^{-K})) \quad (17)$$

and so

$$\begin{aligned} \int_{\{\mathbf{T} \setminus 8I\} \cap \{x \geq 0\}} (E)(x)^p dx &\leq \sum_{i=2}^{2^K-1} \int_{\pi i 2^{-K}}^{\pi(i+1) 2^{-K}} (E)(x)^p dx \\ &\leq C_p 2^{-K} \sum_{i=2}^{2^K-1} 2^{Ki-(N+2)p} \leq C_p. \end{aligned}$$

If $t \in I + 2k\pi$ for some $k \neq 0$ then

$$(F)(x) \leq C 2^{-K(-1/p+N+2)} \sum_{k=1}^{\infty} k^{-(N+2)} \leq C$$

and (6) is satisfied automatically. If $x < 0$ then we can show (6) in the same way.

To prove (16) we have to check (7) for $p = 1/(N+2)$. Inequality (17) implies that

$$\rho^{1/(N+2)} \lambda(\{(E) > \rho\} \cap \{\mathbf{T} \setminus 8I\}) = \rho^{1/(N+2)} \sum_{i=1}^{(2^{K(N+2)} \rho^{-1})^{1/(N+2)}} 2^{-K} \leq 1.$$

Since (F) satisfies also (7), we have shown (16). Inequality (8) can be verified by interpolation as in Theorem 2. The proof of the theorem is complete. ■

Notice that by interpolation we get the inequality

$$\|f\|_{H_{p,q}(\mathbf{T})} \sim \|f\|_{p,q} + \|\tilde{f}\|_{p,q} \quad (0 < p < \infty, 0 < q \leq \infty)$$

from (3). Since $\|f\|_{H_{p,q}} \sim \|\tilde{f}\|_{H_{p,q}}$ and $\tilde{U}_n^\theta f = U_n^\theta \tilde{f}$, we can extend Theorems 2 and 3 easily to the conjugate maximal operators and to the θ -means as follows.

THEOREM 4. *Theorems 2 and 3 hold also for the operator \tilde{U}_*^θ instead of U_*^θ . Moreover, if we replace on the left hand side of the inequalities (11)–(13) and (16) the operator U_*^θ by U_n^θ or \tilde{U}_n^θ and the space $L_{p,q}$ by $H_{p,q}$, then these inequalities hold uniformly in n .*

Since the trigonometric polynomials are dense in $L_1(\mathbf{T})$ and in the Hardy spaces, the inequalities of Theorems 2-4 and the usual density argument (see Marcinkiewicz, Zygmund [9]) imply

COROLLARY 1. *Under the conditions of Theorem 2 or 3, $f \in L_1(\mathbf{T})$ implies*

$$U_n^\theta f \rightarrow f \text{ a.e.} \quad \text{and} \quad \tilde{U}_n^\theta f \rightarrow \tilde{f} \text{ a.e.} \quad \text{as } n \rightarrow \infty.$$

Moreover, if e.g. (12) is satisfied, then these two convergence hold in $H_{p,q}$ norm, whenever $f \in H_{p,q}$, if (13) is true, then in $H_{p_0, \infty}$ norm, whenever $f \in H_{p_0}$. From Theorems 3 and 4 we obtain similar convergence results, i.e. if (16) is satisfied.

Note that \tilde{f} is not necessarily integrable whenever f is.

5. APPLICATIONS TO VARIOUS SUMMABILITY METHODS

In this section we consider several summability methods introduced in the book of Butzer and Nessel [3] and some other popular ones as special cases of the θ -summation. Of course, there are a lot of other summability methods which could be considered as special cases. The elementary computations in the examples below are left to the reader.

Let C_0 consists of all continuous functions f , for which $\lim_{|x| \rightarrow \infty} f(x) = 0$. Butzer and Nessel [3] verified that if $\theta \in C_0$ and θ , θ' and $x\theta''(x)$ are integrable functions, then $\hat{\theta} \in L_1(\mathbf{R})$.

LEMMA 3. Suppose that $\theta \in L_1(\mathbf{R}) \cap C_0$ is even and each term of $(x^i \theta(x))^{(i+1)}$ is integrable for some $i \geq 0$. Then $\hat{\theta} \in L_1(\mathbf{R})$ and

$$|\hat{\theta}^{(i)}(x)| \leq \frac{C}{x^{i+1}} \quad (x \in (0, \infty)).$$

Proof. The integrability of $\hat{\theta}$ comes from the result above. By integrating by parts we have

$$\begin{aligned} |\hat{\theta}^{(i)}(x)| &= \left| \int_0^\infty t^i \theta(t) \cos tx \, dt \right| = \frac{1}{x} \left| \int_0^\infty (t^i \theta(t))' \sin tx \, dt \right| = \dots \\ &= \frac{1}{x^{i+1}} \left| [\theta(t) \cos tx]_0^\infty \right| + \frac{1}{x^{i+1}} \left| \int_0^\infty (t^i \theta(t))^{(i+1)} \cos tx \, dt \right|. \end{aligned}$$

Of course, in the last line probably cos have to be changed to sin. ■

Our first three examples satisfy the conditions of Lemma 3.

EXAMPLE 1. *Weierstrass summation.* Let $\theta_1(x) = e^{-|x|^\gamma}$ for some $0 < \gamma < \infty$. It is easy to see that $(x^i e^{-|x|^\gamma})^{(i+1)} \in L_1(\mathbf{R})$ for all $i \geq 0$. The θ -means are given by

$$U_n^{\theta_1} f(x) := \sum_{k=-\infty}^{\infty} e^{-(|k|/(n+1))^\gamma} \hat{f}(k) e^{ikx}.$$

Of course, we can take another index set than \mathbf{N} . For example we can change $(\frac{1}{n+1})^\gamma$ to t :

$$V_t^{\theta_1} f(x) := \sum_{k=-\infty}^{\infty} e^{-t|k|^\gamma} \hat{f}(k) e^{ikx} \quad (t \in (0, \infty)),$$

or e^{-t} by r :

$$W_r^{\theta_1} f(x) := \sum_{k=-\infty}^{\infty} r^{|k|^\gamma} \hat{f}(k) e^{ikx} \quad (r \in (0, 1)).$$

By Lemma 3, θ_1 satisfies the conditions of Theorem 3 for all $N \in \mathbf{N}$. This means e.g. that the operators $U_*^{\theta_1}$, $V_*^{\theta_1}$ and $W_*^{\theta_1}$ are bounded from $H_{p,q}(\mathbf{T})$ to $L_{p,q}(\mathbf{T})$ for every $0 < p < \infty$ and $0 < q \leq \infty$. Moreover, $U_n^{\theta_1} f \rightarrow f$ a.e. as $n \rightarrow \infty$, $V_t^{\theta_1} f \rightarrow f$ a.e. as $t \rightarrow 0$ and $W_r^{\theta_1} f \rightarrow f$ a.e. as $r \rightarrow 1$. If $\gamma = 1$ then this last result is the well known convergence of the Abel means.

EXAMPLE 2. *Picar summation.* Let $\theta_2(x) = (1 + |x|^\gamma)^{-1}$ for some $1 < \gamma < \infty$. One can check that $(x^i(1 + |x|^\gamma)^{-1})^{(i+1)} \in L_1(\mathbf{R})$ for all $i \geq 0$. The θ -means are given by

$$U_n^{\theta_2} f(x) := \sum_{k=-\infty}^{\infty} \frac{1}{1 + \left(\frac{|k|}{n+1}\right)^\theta} \hat{f}(k) e^{ikx}.$$

It follows from Lemma 3 that Theorems 3 and 4 and Corollary 1 hold for this summability method. For example, $U_*^{\theta_2}$ is bounded from $H_{p,q}(\mathbf{T})$ to $L_{p,q}(\mathbf{T})$ for every $0 < p < \infty$ and $0 < q \leq \infty$.

EXAMPLE 3. *Bessel assumption.* Let $\theta_3(x) = (1 + x^2)^{-\gamma/2}$ for some $1 < \gamma < \infty$. Again, $(x^i(1 + x^2)^{-\gamma/2})^{(i+1)} \in L_1(\mathbf{R})$ for all $i \geq 0$. The θ -means are given by

$$U_n^{\theta_3} f(x) := \sum_{k=-\infty}^{\infty} \frac{1}{\left(1 + \left(\frac{k}{n+1}\right)^2\right)^{\gamma/2}} \hat{f}(k) e^{ikx}.$$

Thus Theorems 3 and 4 and Corollary 1 hold again for every $0 < p < \infty$ and $0 < q \leq \infty$.

The next six θ -functions are special cases of Theorem 3.

EXAMPLE 4. Fejér summation. Let

$$\theta_4(x) = \begin{cases} 1 - |x| & \text{if } |x| \leq 1 \\ 0 & \text{if } |x| > 1. \end{cases}$$

$U_n^{\theta_4}$ is the n th Fejér operator:

$$U_n^{\theta_4} f(x) := \sum_{k=-n}^n \left(1 - \frac{|k|}{n+1}\right) \hat{f}(k) e^{ikx} = \frac{1}{n+1} \sum_{k=0}^n s_k f(x).$$

It is known that

$$\hat{\theta}_4(x) = \frac{1}{\sqrt{2\pi}} \left(\frac{\sin x/2}{x/2} \right)^2$$

and $|\hat{\theta}'_4(x)| \leq C/x^2$. Consequently, Theorems 3 and 4 and Corollary 1 hold for $N=0$.

EXAMPLE 5. Riemann summation. Let

$$\theta_5(x) = \left(\frac{\sin x/2}{x/2} \right)^2 = \sqrt{2\pi} \hat{\theta}_4(x).$$

Then $\hat{\theta}_5(x) = \sqrt{2\pi} \theta_4(x) = \sqrt{2\pi} \max(0, 1 - |x|)$ and so $|\hat{\theta}'_5(x)| = 1_{(-1,1)}(x) \leq C/x^2$. The Riemann means are given by

$$U_n^{\theta_5} f(x) := \sum_{k=-\infty}^{\infty} \left(\frac{\sin k/(2(n+1))}{k/(2(n+1))} \right)^2 \hat{f}(k) e^{ikx}.$$

If we change $1/(n+1)$ to μ then

$$V_{\mu}^{\theta_5} f(x) := \sum_{k=-\infty}^{\infty} \left(\frac{\sin k\mu/2}{k\mu/2} \right)^2 \hat{f}(k) e^{ikx} \quad (\mu \in (0, \infty)).$$

This yields that the operators $U_*^{\theta_5}$ and $V_*^{\theta_5}$ are bounded from $H_{p,q}(\mathbf{T})$ to $L_{p,q}(\mathbf{T})$ for every $1/2 < p < \infty$ and $0 < q \leq \infty$. Moreover, $U_n^{\theta_5} \rightarrow f$ a.e. as $n \rightarrow \infty$ and $V_{\mu}^{\theta_5} \rightarrow f$ a.e. as $\mu \rightarrow 0$. Note that the Riemann summation was considered in Bari [1], Zygmund [23], Gevorkyan [6, 7] and also in Weisz [20].

EXAMPLE 6. de La Vallée-Poussin summation. Let

$$\theta_6(x) = \begin{cases} 1 & \text{if } |x| \leq 1/2 \\ -2|x| + 2 & \text{if } 1/2 < |x| \leq 1 \\ 0 & \text{if } |x| > 1. \end{cases}$$

One can show that $U_{2n+1}^{\theta_6} f = 2U_{2n+1}^{\theta_4} f - U_n^{\theta_4} f$ and since $\theta_6(x) = 2\theta_4(x) - \theta_4(2x)$, we have $|\hat{\theta}'_6(x)| \leq C/x^2$ (cf. Schipp and Bokor [11]). Hence we get again Theorems 3 and 4 and Corollary 1 for $N=0$. Note that we could generalize this summation if we take in the definition of θ_6 another number than $1/2$.

EXAMPLE 7. *Rogosinski summation.* Let

$$\theta_7(x) = \begin{cases} \cos \pi x/2 & \text{if } |x| \leq 1 \\ 0 & \text{if } |x| > 1 \end{cases}$$

and

$$U_n^{\theta_7} f(x) := \sum_{k=-n}^n \cos\left(\frac{\pi |k|}{2(n+1)}\right) \hat{f}(k) e^{ikx}.$$

Since

$$\hat{\theta}_7(x) = \frac{\sin(x - \pi/2)}{2(x^2 - (\pi/2)^2)}$$

(see e.g. Schipp and Bokor [11]), we can verify that $|\hat{\theta}'_7(x)| \leq C/x^2$ and so we obtain Theorems 3 and 4 and Corollary 1 for $N=0$.

EXAMPLE 8. *Jackson-de La Vallée-Poussin summation.* Let

$$\theta_8(x) = \begin{cases} 1 - 3x^2/2 + 3|x|^3/4 & \text{if } |x| \leq 1 \\ (2 - |x|)^3/4 & \text{if } 1 < |x| \leq 2 \\ 0 & \text{if } |x| > 2. \end{cases}$$

One can find in Butzer and Nessel [3] that

$$\hat{\theta}_8(x) = \frac{3}{\sqrt{8\pi}} \left(\frac{\sin x/2}{x/2} \right)^4.$$

Therefore we can show by elementary computations that $|\hat{\theta}_8^{(i)}(x)| \leq C/x^4$ for $i=0, 1, 2, 3$. Consequently, we have the corresponding results with $N=2$.

EXAMPLE 9. *The Summation method of cardinal B-splines.* For $m \geq 2$ let

$$M_m(x) := \frac{1}{(m-1)!} \sum_{k=0}^l (-1)^k \binom{m}{k} (x-k)^{m-1} \\ (x \in [l, l+1), l=0, 1, \dots, m-1)$$

and

$$\theta_9(x) = \frac{M_m(m/2 + mx/2)}{M_m(m/2)}.$$

It is shown in Schipp and Bokor [11] that θ_9 is even and

$$\hat{\theta}_9(x) = \frac{1}{\pi m M_m(m/2)} \left(\frac{\sin x/m}{x/m} \right)^m.$$

It is easy to see that $|\hat{\theta}_9^{(i)}(x)| \leq C/x^m$ for $i=0, 1, \dots, m-1$. Thus Theorems 3 and 4 and Corollary 1 hold for $N=m-2$.

The next example satisfies the conditions of Theorem 2.

EXAMPLE 10. *Riesz summation.* Let

$$\theta_{10}(x) = \begin{cases} (1 - |x|^\gamma)^\alpha & \text{if } |x| \leq 1 \\ 0 & \text{if } |x| > 1 \end{cases}$$

for some $0 < \alpha \leq 1 \leq \gamma < \infty$. The Riesz operators are given by

$$U_n^{\theta_{10}} f(x) := \sum_{k=-n}^n \left(1 - \left| \frac{k}{n+1} \right|^\gamma \right)^\alpha \hat{f}(k) e^{ikx}.$$

We proved in [21] that

$$|\hat{\theta}_{10}(x)|, |\hat{\theta}'_{10}(x)| \leq C/x^{\alpha+1}$$

As $0 < \alpha \leq 1$, in Theorem 2 we have $N=0$. It is easy to see that $C/x^{\alpha+1} \in L_{p_0}[\varepsilon, \infty)$ if and only if $p_0 > 1/(\alpha+1)$ and $C/x^{\alpha+1} \in L_{p_0, \infty}[\varepsilon, \infty)$ if and only if $p_0 \geq 1/(\alpha+1)$. Consequently, (12) holds for $1/(\alpha+1) < p < \infty$ and (13) for $1/(\alpha+1) \leq p_0 < \infty$ and the endpoints are the same in Theorem 4 and Corollary 1.

6. θ -SUMMATION OF FOURIER TRANSFORMS

In this section we summarize briefly the above results for Fourier transforms. First we introduce the Hardy spaces on the real line.

The *Fourier transform* of a tempered distribution f is denoted by \hat{f} . The non-tangential maximal function of a tempered distribution is defined by

$$f^*(x) := \sup_{t>0} |(f * P_t)(x)|,$$

where

$$P_t(x) := \frac{ct}{t^2 + x^2} \quad (t > 0, x \in \mathbf{R})$$

is the non-periodic Poisson kernel.

The *Hardy–Lorentz space* $H_{p,q}(\mathbf{R})$ ($0 < p, q \leq \infty$) consists of all tempered distributions f for which

$$\|f\|_{H_{p,q}(\mathbf{R})} := \|f^*\|_{p,q} < \infty.$$

For a tempered distribution $f \in H_p(\mathbf{R})$ ($0 < p < \infty$) the *Hilbert transform* or the *conjugate distribution* \tilde{f} is defined by

$$\tilde{f} := f * \Phi,$$

where

$$\hat{\Phi}(u) = -i \operatorname{sign} u, \quad \Phi(x) = \frac{1}{\pi x}.$$

We remark that the analogues of (1), (2) and (3) and the analogues of Theorem A, B and C are true in this case (cf. Weisz [21] and the references there).

For $f \in L_p(\mathbf{R})$ ($1 \leq p \leq 2$) the θ -means are defined by

$$U_T^\theta f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \theta\left(\frac{t}{T}\right) \hat{f}(t) e^{ixt} dt = (f * K_T^\theta)(x),$$

where

$$K_T^\theta(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} \theta\left(\frac{t}{T}\right) e^{ixt} dt = T\hat{\theta}(Tx).$$

Thus the θ -means can be rewritten as

$$U_T^\theta f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(t) T\hat{\theta}(T(x-t)) dt.$$

We extend the definition of the θ -means to tempered distributions as follows:

$$U_T^\theta f := f * K_T^\theta \quad (T > 0).$$

One can show that $U_T^\theta f$ is well defined for all tempered distributions $f \in H_p(\mathbf{R})$ ($0 < p \leq \infty$) and for all functions $f \in L_p(\mathbf{R})$ ($1 \leq p \leq \infty$) (cf. Stein [15]). The definition of the *conjugate θ -means* is

$$\tilde{U}_T^\theta f := \tilde{f} * K_T^\theta \quad (T > 0).$$

The *maximal* and *conjugate maximal θ -operators* are introduced by

$$U_*^\theta f := \sup_{T>0} |U_T^\theta f| \quad \text{and} \quad \tilde{U}_*^\theta f := \sup_{T>0} |\tilde{U}_T^\theta f|,$$

respectively.

We can prove all the results of Section 4 also for tempered distributions and Fourier transforms and for the Hardy spaces $H_{p,q}(\mathbf{R})$. We do not formulate exactly the theorems and proofs, because they are almost the same as in Section 4.

THEOREM 5. *Theorems 1–4, Proposition 1 and Corollary 1 hold also for the operators U^θ acting on tempered distributions and defined in this section and for the Hardy spaces $H_{p,q}(\mathbf{R})$.*

Note that the applications of Section 5 are also special cases of the θ -summation of Fourier transforms, the details are left to the reader.

REFERENCES

1. N. K. Bari, "Trigonometric Series," Fizmatgiz, Moskva, 1961. [In Russian]
2. J. Bokor and F. Schipp, Approximate identification in Laguerre and Kautz bases, *Automatica* **34** (1998), 463–468.
3. P. L. Butzer and R. J. Nessel, "Fourier Analysis and Approximation," Birkhäuser, Basel, 1971.
4. C. Fefferman, N. M. Rivière, and Y. Sagher, Interpolation between H^p spaces the real method, *Trans. Amer. Math. Soc.* **191** (1974), 75–81.
5. C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.* **129** (1972), 137–194.
6. G. G. Gevorkyan, On the uniqueness of trigonometric series, *Mat. Sb.* **180** (1989), 1462–1474. [In Russian].
7. G. G. Gevorkyan, On the trigonometric series that are summable by Riemann's method, *Mat. Zametki* **52** (1992), 17–34. [In Russian]
8. R. Long, "Martingale Spaces and Inequalities," Peking University Press and Vieweg, 1993.
9. J. Marcinkiewicz and Z. Zygmund, On the summability of double Fourier series, *Fund. Math.* **32** (1939), 122–132.
10. F. Móricz, The maximal Fejér operator for Fourier transforms of functions in Hardy spaces, *Acta Sci. Math. (Szeged)* **62** (1996), 537–555.
11. F. Schipp and J. Bokor, L^∞ system approximation algorithms generated by φ summations, *Automatica* **33** (1997), 2019–2024.

12. F. Schipp and L. Szili, Approximation in \mathcal{H}_∞ -norm, in "Approximation Theory and Function Series" Vol. 5, pp. 307–320, Bolyai Soc. Math. Studies, Budapest, 1996.
13. F. Schipp, W. R. Wade, P. Simon, and J. Pál, "Walsh Series: An Introduction to Dyadic Harmonic Analysis," Hilger, Bristol/New York, 1990.
14. E. M. Stein and G. Weiss, "Introduction to Fourier Analysis on Euclidean Spaces," Princeton Univ. Press, Princeton, 1971.
15. E. M. Stein, "Harmonic Analysis: Real-Variable Methods, Orthogonality and Oscillatory Integrals," Princeton Univ. Press, Princeton, 1993.
16. L. Szili and P. Vértesi, On uniform convergence of sequences of certain linear operators, *Acta Math. Hungar.*, to appear.
17. L. Szili, On the summability of trigonometric interpolation processes, *Acta Math. Hungar.*, to appear.
18. A. Torchinsky, "Real-Variable Methods in Harmonic Analysis," Academic Press, New York, 1986.
19. F. Weisz, "Martingale Hardy Spaces and Their Applications in Fourier-Analysis," Lecture Notes in Mathematics, Vol. 1568, Springer-Verlag, Berlin, 1994.
20. F. Weisz, Riemann summability of two-parameter trigonometric-Fourier series, *East J. Approx.* **3** (1997), 403–418.
21. F. Weisz, Riesz means of Fourier transforms and Fourier series on Hardy spaces, *Studia Math.* **131** (1998), 253–270.
22. N. Wiener, "The Fourier Integral and Certain of Its Applications," Dover, New York, 1959.
23. A. Zygmund, "Trigonometric Series," Cambridge Press, London, 1959.